Zoo 506 current paper solve
sir Arslan

Zoo506 ka paper meri old file sy boht km arha he .lekin achi baat ye h k new question jo arhy hen wo repeated hen so un ki file bn gei he is file ko achy sy yaad kry our quiz k liy old file sy all mcq yaad kry mcq usi file sy arhy hen ,is sy apka paper 75% to conferm hojay ga sath me aman khan ki quiz ki file jo 2 huwy hen wo krly to 85% tk conferm hojay ga is sy zeada b..

1. List the insect born diseases?

- Malaria
- Bubonic plague
- Typhoid
- Typhus
- Yellow fever

2. Clean culture?

The destruction of vegetable trash and weeds is known as cultural clean

3. Heat treatment?

Most insects cease reproduction and soon die at temperature from 100 to 110F

- ➢ The temperature of their surrounding should be raised to 120 or higher to kill them with heat artificially within a few hour
- ➢ The length of period for effective treatment is more dependent upon the
- Time to heat the grain
- Clothing
- Other material in which insects are located
- ➢ Then upon the heat resistance of the pest themselves because no insect can survive more than an hour if actually exposed to a temperature of 120F
- ➢ Bales and bags are difficult to heat through.

4. Hypo pharynx?

Near the base of labium on its inner or mouth side there is frequently a fleshy swelling, more or less covered by bristles or hairs which is called hypo pharynx, lingua or tongue. it varies greatly in size and form

5. Symptoms of red cotton bug?

Red cotton bug: Dysdercus cingulatus

- Family : Pyrrhocoridae
- Order : Hemiptera

Symptom of damage

- Red stained lint and rotting bolls.
- Inner boll wall with warty growth or water soaked spots
- Young bolls abort and turn dark brown
- The bacterium Nematospora gossypii enters the site of injury and stains the fiber.

Sir Arslan

6. Hopper burn cause?

Brown plant hopper is the most destructive pest of rice .. both nymph and adult feed on paddy, sugercan and grasses by sucking cell sap which turn the leaves yellow.
A heavy infestation produce symptoms of "hopper burn" i.e. leaves become dry and brown after insect feeding and patched of burned plants are often lodged.

Damage cause by termits?

- They enter through cut ends of setts & feed on the soft tissue.
- The tunnel is filled with the soil.
- The termites attack setts, shoots, canes and also stubbles
- Entire shoot dries up and can be pulled out easily

7. Physical control methods?

Physical factor controlling insects are following

- Heat
- Cold
- Light
- Moisture
- Electricity

8. Enlist the plants require pollination?

Smyrna fig, clovers, melons, tomatoes, beans, peas,and most of common fruits require insects for their pollination

9. Agronomic practice in cultural control?

1. Optimize growing conditions for the crop
2. Anything that increases a crop's competitive edge will result in increased tolerance to pests often resulting in reduced pesticide use
3. Create unfavorable conditions for the pest

10. What is cotton?

Cotton is a soft, fluffy staple fiber that grows in boll, or protective

Cotton (Gossypium spp), the king of fibers, usually commercial referred as white gold
and one of the important commercial crops, plays a pivotal role in human civilization, economic, political and social affairs of world.

11. Rice ear head bug?

Rice ear- head bug

The adult of Leptocorisa oratorius is green, light brown or mixed yellow in color The adult are slender and about 20mm long.

Sir Arslan

Symptoms of Damage

- The pest is essentially diurnal with highest activities in the morning and in the evening, seeking shelter during hotter parts of day
- Many generation are completed in a year
- Sucking of the grain sap causes empty or partially filled and chaffy grains and enhances subsequent fungal and bacterial infection

Adult and nymph both suck the sap of developing rice grains at the milking stage and cause considerable yield loss

Management

- he population can be suppressed by killing the bug by using light traps
- Collection of adult bug using net
- Destroy weed to remove alternate host
- Keep on hanging the cattle urine soaked gunny bags or cow dung wrapped cloth in the field to attract the bug.
- Synchronize rice planting to maintain simultaneous crop maturity in the field in an area for equal distribution of bugs in all fields
- Conserve predator of rice bug like tiger beetle, Cicindela sexpunctata by using chemical pesticide judiciously
- Spray after at least 10 bug per 100 panicles with pesticide like carbaryl (Carbaryl 5% dust@25kg/ha)
- Spray Fenvalerate 20 EC 0.5ml/liter of water

12. Scientific name of early shoot borer?

Chilo infuscatelus

Sir Arslan

13. Pest management?

The stud of principles and experimental approaches to the science that underlins the development of working integrated pest management systems

Types of insect control

- A-NATURAL CONTROL
- B-ARTIFICIAL CONTROL: Measures for the artificial control of insects may be divided into six groups:
- (1) mechanical control
- (2) cultural control
- (3) biological control,
- (4) legislative control
- (5) physical control
- (6) chemical control.

14. Strategies of pest management?

techniques and methods to keep pest populations below economically injurious levels

- Each pest management technique must be environmentally sound and compatible with producer objectives

Utilizes all suitable pest management Ways

- Pesticide Control
- Cultural Control
- Mechanical Control
- Sanitary Control
- Natural Control
- Biological Control
- Host Plant Resistance

15. Hairy spider belongs to ?

Class: Arachnida

Order: Araneae

Family: Theraphosidae

16. Cat and dog flea can live for 100 days without eating

17. Scientific name of top borer?

Scirpophaga nivella

18. Sryphid larva act as biological agent for?

19. Define transmission?

Transmission is the passing of a pathogen causing communicable disease from an infected host

individual or group to a particular individual or group .Regardless of whether the other individual was previously infected

Sir Arslan

20. Economic threshold?

"Pest Population at which a grower must take action to prevent a pest populations from reaching the economic injury level"

21. Define pulvilus, pulvilli,empodium?

At the end of the last a pair of claws is generally found, and between them a sort of pad or cushion, the pulvillus

Sometimes there are three of these, in which case the outer ones are called the pulvilli and the middle one the empodium

22. Intelligent cultivation?

Intelligent cultivation and use of fertilizers is well as pruning to remove injured or diseased parts will aid in insect control

23. **Arthropoda classes**?

5 Classes of Arthropods

1. Class #1- Hexapoda: Insects, six, jointed appendages
2. Class #2- Chilopoda- Centipedes, one pair of legs per body segment, multiple body segments
3. Class #3- Diplapoda- Millipedes, two pairs of legs per body segment, multiple body segments
4. Class #4- Crustacea- Crayfish, five pairs of jointed appendages
5. Class #5- Arachnida- Spiders, four pairs of jointed appendages

24. Insect pest management?

The stud of principles and experimental approaches to the science that underpins the development of working integrated pest management systems

25. Yellow rice stem borer?

Yellow Rice Stem Borer

- The yellow stem borer is a specific pest of rice
- The caterpillar alone are destructive, when full grown, they measure about 20mm and are dirty white or greenish yellow having brown head and pronotum

Symptoms of Damage

- The larval feeding damage may cause death of the central leaf whorl at the vegetative stage, which is known as dead heart.
- Damage at the reproductive stage causes ear devoid of grain, which is known as white head.

Management

- The removal and destruction of stubble at the time of the first ploughing after harvesting the crop
- Ploughing and flooding the field is also effective in killing the larvae
- Clipping of tips of seedling before transplanting can reduce the carryover of eggs to the field.
- Use trichocard with 50,000-1,00,000 Trichogramma per ha after 3-4 weeks

Sir Arslan

- The field showing more than 5% dead hearts should be sprayed 25kg of Cartap hydrochloride 4G or 15 kg of Fipronil 0.3 G per ha in standing water in the field.

26. Write the name of thorax segments?

The thorax has its three segments usually quite clearly marked
1-Prothorax
2-Mesothorax
3-Metathorax

27. Write the damage and symptoms of fruit borer?

American bollworm/Fruit borer: Helicoverpa armiger
Family: Noctuidae

- **Order: Lepidoptera**
- **It is one of the important major insect of cotton crop**

- **Symptoms of damage**
- Bolls showing regular, circular bore holes
- Larvae seen feeding on the boll by thrusting their heads alone inside and leaving the rest of the body outside

Presence of granular faecal pellets outside the bore hole.

- A single larva can damage 30-40 bolls**.**

MANAGEMENT

Avoid continuous cropping of cotton both during winter and summer seasonsin the same area as well as rationing

- Avoid monocropping. Growing of less preferred crops like greengram, blackgram, soyabean, castor, sorghum etc., along with the cotton as intercrop or border crop or alternate crop to reduce the pest infestation
- Removal and destruction of crop residues to avoid carry over of the pest to the next season, and avoiding extended period of crop growth by continuous irrigation.
- Optimizing the use of nitrogenous fertilizers which will not favor the multiplication of the pest.
- Judicious water management for the crop to prevent excessive vegetative growth and larval harbourage
- Releasing predator Chrysoperla carnea @ 1, 00, 000/ha at 6th, 13th and 14th week after sowing.
- During bolling and maturation stage, apply any one of the following insecticides (1000

l of spray fluid/ha):

28. Write the use of pesticides?

Pesticide Control

- Pesticides can to be used in an IPM program, however only as a last resort
- Pesticides are to be used when there is no risk of environmental damage or when benefits outweigh the risks

Sir Arslan

- Use pesticides only when other control practices aren't available, economical or practical
- Important aspect to use pesticides:
- Must monitor pest populations in the field.
- Identify the pest
- Compare pest population and the economic threshold
- Life stage susceptible to pesticide?
- Crop stage and preventable loss

29. What is cerci?

A pair of antenna like structure (sometimes short sometimes long) present at the end of abdomen are known as cerci.

30. Write commercial products produce by insects?

- 1-Silk
- 2-Honey
- 3-Beeswax
- 4-Shellac

31. What are repellents?explain some repllents.

. *Repellents are those materials used to keep insects away from crops, animals and man*

2. Various coal-tar and pine products are rather general insect repellents.

3. Naphthalene is a clothes moth repellent and oil of citronella is a common ingredient of preparations used to keep mosquitoes away from people

32. Differentiate between intelligent cultivation and destruction of vegetables? (5)

- Intelligent cultivation and use of fertilizers is well as pruning to remove injured or diseased parts will aid in insect control
- The destruction of vegetable trash and weeds, frequently called clean culture, is also

an important factor, especially in the control of hibernating insects

33. Difference beween the exoskeleton of mollusks and arthropods?

In the possession of this external skeleton these animals have a seeming resemblance to the shells

(Mollusca) But the materials of which the skeleton is composed are quite different Being largely

calcium carbonate in the Mollusca And chitin which somewhat resembles horn in its nature,

sometimes with calcareous salts·deposited in it, in the Arthropoda .The simple members of arthropod

have bilaterally symmetrical body, though this condition is concealed somewhat by secondary changes

in many of the group .The possession of a bilaterally symmetrical body consisting of a series of

Sir Arslan

segments, an exoskeleton of chitin and the presence of jointed legs are, then, distinctive features of
the arthropods .To separate the various groups of arthropods, other characters must be Used Aside
from several small sections not often seen, there are five large and important divisions which call for
recognition

34. Why strains of lants are used in cultural control?

- Varieties or strains of some plants have been found more resistant to insect attack than

others

- To develop resistant strains and to prove their value, however, frequently require long,

expensive research in plant genetics

- Instances in which some success has been obtained will be mentioned under the insect

pests concerned

35. Management strategies of rice grasshopper?

Fish cum Rice farming Larva of the banded blister beetle *(Mylabris phalerata*) attack the eggs of the rice grasshopper .Trimming the bund and field sanitation reduces the grasshopper population .Spray chlorpyriphos 2.5 lit Chlorpyriphos 20EC in 250 liter of water per ha.

36. Allergy

(hay fever, asthma, etc.) are frequently caused by exposure of susceptible persons to dried insect scales and hairs

Sir Arslan