

MTH 100
Quiz
2

Question # 1 of 15 (**Start time: 12:28:18 PM, 12 January 2020**)

Total Marks: 1

If I added -5 to each coordinate point $(-2, 3)$, what would be my new coordinates?

Select the correct option

☐	$(7, 2)$
☒	$(-7, -2)$
☐	$(3, 2)$
☐	$(10, -15)$

Question # 2 of 15 (Start time: 12:29:35 PM, 12 January 2020)

The range of the cosine function is between _____.

Select the correct option

- | | |
|----------------------------------|------------|
| ☐ | 0 and -1 |
| ☐ | 0 and 1 |
| ☐ | -1 and 0.5 |
| ☒ | -1 and 1 |

4



The sum formula for $\cos(A + B)$ is _____.

Select the correct option

- | | |
|----------------------------------|---------------------------------|
| ☐ | $\cos A \cos B + \sin A \sin B$ |
| ☐ | $\sin A \cos B - \cos A \sin B$ |
| ☐ | $\sin A \cos B + \cos A \sin B$ |
| ☒ | $\cos A \cos B - \sin A \sin B$ |

Click to Sa

4



MTH100:Quiz No. 2

Question # 4 of 15 (Start time: 12:31:18 PM, 12 January 2020)

Determine the slope of the line that passes through (2, 2) and (5, 8).

Select the correct option

☐	10/7
☒	1/2
☐	-2
☐	2

Question # 5 of 15 (Start time: 12:32:33 PM, 12 January 2020)

In the cosine function, the minimum value of _____ is taken when $\theta = \pm 180^\circ$, $\pm 540^\circ$ and so on.

Select the correct option

☒	-1
☐	-0.5
☐	0
☐	1

Click to Save Answer

The value of $\sin 135^\circ$ is same as the value of

Select the correct option

[Reload Math Equations](#)

- | | |
|----------------------------------|-------------------|
| ☐ | $-\sin 135^\circ$ |
| ☐ | $-\sin 45^\circ$ |
| ☒ | $\sin 45^\circ$ |
| ☐ | $-\sin 90^\circ$ |

[Click to Save Answer & Move to Next Question](#)

If I subtracted 3 from each coordinate point $(-2, 3)$, what would be my new coordinates?

Select the correct option

- | | |
|----------------------------------|-----------|
| ☐ | $(5, 0)$ |
| ☐ | $(-5, 0)$ |
| ☒ | $(1, 0)$ |
| ☐ | $(-1, 0)$ |

[Click to Save Answer & Move to Next Question](#)

$\tan \theta =$

Select the correct option

[Reload Math Equations](#)

☐	$\frac{\cos \theta}{\sin \theta}$
☒	$\frac{\sin \theta}{\cos \theta}$
☐	$\frac{\cot \theta}{\sin \theta}$
☐	$\frac{\cot \theta}{\cos \theta}$

[Click to Save Answer & Move to Next Question](#)

Question 7 of 10 (Start time: 12:00:10 PM, 12 January 2020) Total marks: 1

In the cosine function, the maximum value of _____ is taken when $\theta = 0^\circ$, $\pm 360^\circ$, $\pm 720^\circ$ and so on.

Select the correct option

[Reload Math Equations](#)

☐	-1
☐	0
☐	0.5
☒	1

[Click to Save Answer & Move to Next Question](#)

12:36 PM 1/12/2020

The domain of inverse sine function and inverse cosine function is same.

Select the correct option

☒	True
☐	False

Click to Save Answer & Move to Next Question



The tangent function is periodic, but its period is _____ degrees.

Select the correct option

☐	60
☐	90
☒	180
☐	360

$\sin \theta$ is negative in the _____ and the fourth quadrant.

Select the correct option

 Reload Math Equations

☐	first
☐	second
☒	third
☐	fourth



$\cos(-\theta) = \dots\dots$

Select the correct option

 Reload Math Equations

☐	$\sec(\theta)$
☒	$\cos \theta$
☐	$-\cos \theta$
☐	$-\cos(-\theta)$



The amplitude of the function is the height from the rest value to the extreme value

Select the correct option

☒	True
☐	False

Click to Save Answer & Move to Next Question