

STA-301 Statistics & Probability Update MCQS For Mid Term Solve By Vu Topper RM

85% To 100% Marks

وَتُعَزُّ مَنْ تَشَاءُ وَتُذِلُّ مَنْ تَشَاءُ

VU
TOPPER RM

Our Services

- Assignments
- Quiz & Gdb's
- LMS Handling
- Online Classes
- Lectures Views
- Project Handling
- Important Notes

پہلے ہمارے سٹوڈنٹس کے رزلٹ دیکھیں
پھر ہم سے سروسز لیں



www.youtube.com/@vutopperm



0322 4021365
0316 4980094

Contact us

Professional Online Academy

Result

**85% To
100%**

بري صحبت سے تٹھائی بہتر ہے اور تٹھائی سے نیک صحبت بہتر ہے

For More Help Contact What's app 03224021365

Which of the following method of skewness is a pure number/unitless?

- A. Fisher
- B. Bowleys
- C. Sheppard
- D. Pearson**

In a linear regression, the degree of scatter of the observed values about the regression line is measured by:

- A. Standard error of estimate**
- B. Median
- C. Mean
- D. Mean deviation

The first moment about mean is:

- A. 0**
- B. 1
- C. 2
- D. All of the above

If a distribution has positive skewness, in what order (highest to lowest) will the averages be?

- A. median, mode, mean
- B. mean, median, mode**
- C. mean, mode, median
- D. mode, median, mean

If we calculate the probability of an event and it turns out to be -0.7 , we know that:

- A. we made a mistake**
- B. the event is probably going to happen
- C. . the probability of it not happening is $.3$
- D. the event is probably not going to happen

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

For More Help Contact What's app 03224021365

The number of ways in which four books can be arranged on a shelf are:

- 4
- 6
- 12
- 24**

Relationships among the averages

- A. $HM \leq GM \leq AM$**
- B. $GM \leq HM \leq AM$
- C. None of the above
- D. $HM \geq GM \geq AM$

Statistics branches include

- A. Industry Statistics
- B. Mathematical Statistics
- C. Applied Statistics
- D. Both Applied and Mathematical**

In a frequency distribution the categories must

- A. Be of the same size
- B. Mathematical Statistics
- C. Have at least 5 observations
- D. Be mutually exclusive**

If the relative frequencies are used, the cumulative frequency polygon rises _____

- A. From the value 0 at left to the value 1 at right**
- B. From the value 1 at left to the value 0 at right
- C. From the value 0 at right to the value 1 at left
- D. None of these

According to the property of parameter, it is

- A. Continuous

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

For More Help Contact What's app 03224021365

B. Constant

C. Discrete

D. Variable

Data arrangement in order of magnitude is called:

A. Grouped data

B. Arrayed data

C. Frequency distribution

D. Raw data

From the following table, what is the Upper limit of the 3 class?

Class	Frequency
30.0 - 32.9	2
33.0 - 35.9	4
36.0 - 38.9	14
39.0 - 41.9	8
42.0 - 44.9	2

A. 42.0

B. 33.0

C. 38.9

D. 35.9

Which one of the following is equal to mean in a symmetrical distribution:

A. 50th Percentile

B. 5th Decile

C. All of these

D. 2nd Quartile

Temperature, in Fahrenheit, is an example of a variable measured on the following scale:

A. Ordinal

B. Nominal

C. Interval

D. Ratio

If in a set of discrete values, 50 percent values are greater than 25, then

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Q2 is

A. 20

B. 75

C. 25

D. 50

In a histogram the area of each rectangle is proportional to:

A. Frequency of the corresponding class interval

B. Cumulative frequency of the corresponding class interval

C. The class mark of the corresponding class interval

D. The class size of the corresponding class interval

Half of the observations are always larger than:

A. Mode

B. Median

C. Mean

D. Total

If the GM of a set of two observations is 10 and its HM is 8. then the AM of the set of observations is

A. 64

B. 100

C. 7.5

D. 12.5

The total of frequency up to an upper class limit or boundary is known as:

A. cumulative frequency

B. frequency polygon

C. average frequency

D. frequency distribution

A scale in which objects are rank ordered according to how much of the

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

attribute they possess, with equal space between objects.

A. Nominal

B. Interval

C. Ratio

D. Ordinal

For the given set of observations 1,4,4,4 and 7, it can be said that the:

A. Mean=Median=Mode

B. None of the above

C. Mean>Median>Mode

D. mean is larger than either median or mode

Quota sampling is an example of:

A. None of these

B. Random sampling

C. Simple random sampling

D. Non random sampling

How many percent corresponds to Q3 if you will interpret it?

A. 75%

B. 25%

C. 50%

D. 90%

If the sum of the reciprocal of the values of a data is 256.7 and $n=23$, then Harmonic mean will be:

A. 0.0789

B. 0.0624

C. 0.0895

D. 0.0569

The mean of a distribution is 23, the median is 24 and the mode is 25.5. It is most likely that this distribution is:

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

- A. Symmetrical
- B. Asymptotic
- C. Positively Skewed**
- D. Negatively Skewed

In symmetrical distribution:

- A. mean=median=mode**
- B. mean=median=mode
- C. Mean>median>mode
- D. mean<median<mode

100% correct answer required.

A chart consists of horizontal or vertical bars of equal width and lengths proportional to values they represent

- A. multiple bar chart
- B. pie chart
- C. simple bar chart**
- D. component bar chart

For the digit 40, leaf will be:

- A. 1
- B. 2
- C. 4**
- D. 0

The mean is obtained by

- A. Most frequent value in the data set
- B. Dividing the sum of all observations by their number**
- C. Divides a set of data into two equal parts
- D. None of these

The beginnings of a cumulative frequency distribution are presented below. What is the next number in the Cumulative Frequency column?

Classes	Frequency	Cumulative Frequency
6.1 to 8	18	18
8.1 to 10	22	40
10.1 to 12	30	70
12.1 to 14	25	95

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

For More Help Contact What's app 03224021365

A. 2

B. 0

C. 3

D. 1

From the table given below, how many students obtained marks 80 or above: Marks

Marks	50-59	60-69	70-79	80-89	90-100
Frequency	5	9	5	6	3

A. 8

B. 5

C. 3

D. 15

What is/are the mode for the following data:

1,1,2,2,5,5,7,7

A. 1,2

B. No mode in the data

C. 1,2,5

D. 1,2,5,7

A list of pulse rates is: 70, 64, 70, 80, 74, 92. What is the mode for this list?

A. 80

B. 72

C. 76

D. 70

If we want to calculate annual increase in a firm then which of the following average will be used?

A. Geometric mean

B. Mode

C. Mean

D. Harmonic mean

If you have 26.95 – 29.95 class boundaries then the mid point will be:

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

A. 31.45

B. 30.45

C. 28.45

D. 29.45

Find the mean of the values 2,4,6,8.

A. 4

B. 20

C. 10

D. 5

A number of fishes in a pond is an example of:

A. Finite population

B. Infinite population

C. Hypothetical population

D. None of these

In a frequency distribution, which of the following is sometimes referred to as the class mark?

A. frequency

B. cumulative frequency

C. relative frequency

D. class midpoint

What is the mode in the word TOMATO.

A. M

B. O

C. T and O

D. T

100% correct answer required.

In a shoes store, the average size is used as:

A. GM

B. Mode

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

C. Mean

D. Median

What is mode for the following set of data: 1, m, d, n, 2, d, 2, d, s, 5, 5, 7

A. 2

B. 2.d.5

C. 5

D. d

Village patwari collecting the data about cotton crops is the example of:

A. Secondary data

B. Primary data

C. Quantitative data

D. Numerical data

Which scale of measurement is more suitable in the following example? The number of minutes of overtime work that employees perform.

A. Nominal

B. Ordinal

C. Ratio

D. Interval

In how many different ways can the letters of the word 'LEADING' be arranged in such a way that the vowels always come together?

A. 5040

B. 720

C. 650

D. 860

If $A = \{1, 2, 3, 4\}$ and $B = \{3, 4, 5, 6\}$ then $A \cup B$ will be:

A. $\{3, 4\}$

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

For More Help Contact What's app 03224021365

B. {1,2,3,4,5,6} ok

C. {1,2}

D. {2,1}

A set which is the sub-set of every set is

A. Super Set

B. Empty Set ok

C. Universal Set

D. Power Set

If $A = \{1,2,3,4\}$ and $B = \{3,4,5,6\}$ then $A \cap B$ will be:

A. {2}

B. {1,2}

C. {3,4}

D. {1}

In a coin tossing example, we have only two possible outcomes, a head or a tail; this is an example of :

A. Equally likely events

B. None of these

C. Both of these

D. Mutually exclusive events

The Probability of a sure event is:

A. 0

B. 1

C. 8

D. 0.5

What are the chances that no two boys are sitting together for a photograph if there are 5 girls and 2 boys?

A. $1/21$

B. $4/7$

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

C. $2/7$

D. $5/7$ ok

In a class, there are 15 boys and 10 girls. Three students are selected at random. The probability that 1 girl and 2 boys are selected, is:

A. $27/112$

B. $21/46$ ok

C. $3/25$

D. $1/50$

For two mutually exclusive events A and B, $P(A) = 0.1$ and $P(B) = 0.4$, then $P(A \cup B)$ is:

A. 0.1

B. 0.5

C. 0.3

D. 0.4

Two dice are thrown simultaneously. What is the probability of getting two numbers whose product is even?

A. $1/2$

B. $3/4$

C. $3/8$

D. $5/16$

If $P(B) = 0.75$, $P(A) = 0.35$ and $P(A \text{ and } B) = 0.20$, then $P(B | A)$ is:

A. 0.57

B. 0.27

C. 0.32

D. 0.41

The probability of drawing a red Jack from well-shuffled pack of 52 playing cards is:

A. $4/52$

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

B. 26/50

C. 2/52

D. 13/52

Order of the power set of a set of order n is

A. n^2

B. 2^n ok

C. n

D. n^2

Two cards are drawn together from a pack of 52 cards. The probability that one is a spade and one is a heart, is:

A. 2/20

B. 29/30

C. 47/100

D. 13/102

17 students are present in a class. In how many ways, can they be made to stand in 2 circles of 8 and 9 students?

A. $17C8 \times 8! \times 9!$ ok

B. $8! \times 7!$

C. $17C9 \times 8! \times 7!$

D. $17C9 \times 9! \times 8!$

By using identity law $A \cup S$ is equal to:

A. None of them

B. $A \cap S$

C. S

D. A

What is the Cartesian product of $A = \{1, 2\}$ and $B = \{a, b\}$?

A. $\{(1, a), (2, a), (1, b), (2, b)\}$

B. $\{(1, 1), (a, a), (2, a), (1, b)\}$

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

- C. $\{(1, a), (1, b), (2, a), (b, b)\}$
- D. $\{(1, 1), (2, 2), (a, a), (b, b)\}$

We use the General Rule of Multiplication to combine:

- A. Events that total more than 1.00
- B. Mutually exclusive events
- C. Events those are not independent**
- D. Events based on subjective probabilities

What does an each outcome in the sample space regarded as

- A. Element
- B. Sample point
- C. Both a & b** ok
- D. None of the above

A card is drawn from a pack of 52 cards. The probability of getting a queen of club or a king of heart is:

- A. $1/13$
- B. $1/26$** ok
- C. $2/13$
- D. $1/50$

A standard deck of 52 cards is shuffled. What is the probability of choosing a card which is not a diamond:

- A. $1/13$
- B. $5/52$
- C. $39/53$**
- D. $13/50$

Complement of the event B will be

- A. $P(B)$
- B. $P(B) - 1$
- C. $1 - P(B)$**

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

D. $1 + P(B)$

If A and B are mutually exclusive events and $P(A) = 4/36$ and $P(B) = 6/36$, then $P(A \cup B)$ will be

A. $5/36$

B. $11/36$

C. $9/36$

D. $10/36$

An urn contains 10 articles, 2 having minor Defects and three have major defects. Determine the probability that an article selected at random has minor defect.

A. $1/2$

B. $1/4$

C. $1/3$

D. $1/5$

If a die is rolled, what is the probability of getting an even number greater than 2?

A. $5/6$

B. $2/3$

C. $1/3$

D. $1/2$

Two mutually exclusive (disjoint) events:

A. are always independent

B. have the same probability

C. have no effect on the occurrence of each other can be independent

D. cannot occur together

The probability of drawing 'zero white' ball from a bag containing 4 red, 8 black and 3 white balls is:

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

A. 12/15

B. 1/2

C. 8/15

D. 0

When a fair coin is tossed, what is the probability of getting head.

A. All

B. 1

C. 1/2

D. 0

If $P(A) = 0.45$, $P(B) = 0.35$ and $P(B \text{ and } A) = 0.25$, then $P(B | A)$ is:

A. 0.801

B. 0.750

C. 0.751

D. 0.555

If a box contains two red and three yellow balloons then probability of yellow balloons will be equal to:

A. 2/5

B. 1/5

C. 3/5

D. 4/5

When three coins are tossed simultaneously, $P(3 \text{ Heads})$ is:

A. 3/8

B. 2/8

C. 1/8

D. 4/8

When two coins are tossed simultaneously, what are the chances of getting at least one tail?

A. 1/5

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

B. $1/4$

C. $4/5$

D. $3/4$

Which of the following is NOT a possible probability?

A. 1

B. 0

C. 1.25

D. $25/100$

A group consists of three persons, A, B, & C. How many ways are there of selecting a group of two persons out of these three?

A. 2

B. 6

C. 3 **ok**

D. 1

Which of the following best expresses the General Addition Rule?

A. None of the above.

B. $P(A \text{ or } B) = P(A) + P(B)$

C. $P(A \text{ or } B) = P(A) + P(B) + P(A \text{ and } B)$

D. $P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$

A bag contains 4 white, 5 red and 6 blue balls. Three balls are drawn at random from the bag. The probability that all of them are red, is:

A. $2/77$

B. $2/91$

C. $1/22$

D. $3/22$

When the union of mutually exclusive events is equal to entire sample space, it is called:

A. Mutually exclusive events

B. Random events

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

For More Help Contact What's app 03224021365

C. Exhaustive events

D. Equally likely events

The occurrence of sum of 11 when a pair of die is thrown is an example of:

A. Compound event

B. Simple event

C. Mutually exclusive event

D. Random event

If a coin is tossed 3 times then probability that three tails appear simultaneously is equal to:

A. $1/8$

B. $1/4$

C. $1/6$

D. $1/2$

$0!$ is equal to:

A. 0

B. 1

C. 0 and 1

D. None

A graphical device used to list all possibilities of a sequence of outcomes in systematic way is called:

A. Pie diagram

B. Venn diagram

C. Probability histogram

D. Tree diagram

If a player well shuffles the pack of 52 playing cards, then the probability of a red card from 52 playing cards is:

A. $2/52$

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

For More Help Contact What's app 03224021365

B. 26/52

C. 4/25

D. 1/25

Consider a set $S = \{a, b, c, d, e\}$. Which set will make the PARTITION of S ?

A. $\{ \}$ and $\{a, b, c, d, e\}$

B. $\{a, b\}$ and $\{c, d, e\}$

C. $\{a, b, c\}$ and $\{c, d, e\}$

D. $\{a, b\}$, $\{c, d\}$ and $\{a, e\}$

If $\text{median} < \text{mid-quartile range} < \text{midrange}$ then distribution will be:

A. Symmetrical

B. Negatively skewed

C. Positively skewed

D. Zero skewed

Find the mean of the values 2,4,7,8,9

A. 6

B. 5

C. 3

D. 30

For a positively skewed distribution median is To/than the mid quartile range

A. None of the above

B. Equal

C. Greater

D. Less

Which of the following measures based on all observations?

A. Standard deviation

B. Mean deviation

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

C. None of these

D. Mean deviation and Standard deviation

The mean of a distribution is 25, the mode is 25 and the standard deviation is 5, then the coefficient of skewness will be:

A. Less than zero

B. None of the above

C. Greater than zero

D. Equal to zero

Dividing mean deviation by mean we get a pure number known as

A. Co-efficient of quartile deviation

B. Co-efficient of mean deviation

C. None of the above

D. Co-efficient

If coefficient of skewness is equal to "0" then distribution will be:

A. Positively skewed

B. Symmetrical

C. Negatively skewed

D. Asymmetrical

Bowley's coefficient of skewness is expressed in terms of:

A. Square of units

B. Different units

C. Same units

D. Independent of units

When the peak of the curve becomes relatively high, it is called:

A. Hetrokurtic

B. Platykurtic

C. Mesokurtic

D. Leptokurtic

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

For a symmetrical data set mean value is 150 and standard deviation 25. 68% values will lie between

A. (125,175)

B. (110,190)

C. (100,200)

D. (125,180)

First moment about origin is always equal to:

A. Mean

B. Standard Deviation

C. Variance

D. Zero

For a particular data set the Pearson's coefficient of skewness is greater than zero. What will be the shape of distribution?

A. Symmetrical

B. Positively Skewed

C. Negatively Skewed

D. None of the above

In a left skewed distribution:

A. $X_m - Q_3$ greater than $Q_1 - X_0$

B. $X_m - Q_3$ less and equal to $Q_1 - X_0$

C. $X_m - Q_3$ less than $Q_1 - X_0$

D. $X_m - Q_3$ greater and equal to $Q_1 - X_0$

Which of the rule gives no information about the probability of observing a value within one standard deviation of the mean?

A. Permutation rule

B. Combination rule

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

- C. Empirical rule
- D. Chebychev's rule**

If in a simple linear regression model it is assumed that the intercept parameter is equal to 0:

- A. The slope of the line will also be equal to 0
- B. The regression line will pass through the point (0,10)
- C. The regression line will pass through the origin** ok
- D. The regression line will pass through the point (0,-10)

Pearson's coefficient of skewness cannot be used when standard deviation is:

- A. 0**
- B. 3
- C. 1.5
- D. 1

Which of the following method of skewness based on mean, median, mode?

- A. Pearson
- B. Bowleys**
- C. Fisher
- D. Sheppred

Kurtosis is used to represent the degree of peakedness/flatness about a:

- A. Trimodal distribution
- B. Multinomial distribution
- C. Unimodal distribution** ok
- D. Bimodal distribution

Pearson coefficient of skewness is defined as:

- A. $SD / (\text{Mean} - \text{Mode})$
- B. $(\text{Median} - \text{Mode}) / SD$

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

C. SD/(Median-Mode)

D. (Mean-Mode)/SD

If we have mean=6, median =5 and standard deviation = 4 then the Pearson's Coefficient of Skewness will be:

A. 0.5

B. 0.25

C. 0.7

D. 0.75

Putting the value of "k" which is greater than one in Chebyshev's inequality we get the proportion of values that lies in that range.

A. Minimum

B. Optimum

C. Maximum

D. None of the above

Standard Deviation is measure of dispersion.

A. A pure number

B. A relative

C. An absolute

D. None

Given the least squares regression line $\hat{Y} = 2 + 0.3x$:

A. the relationship between x and y is negative

B. None of the above

C. the relationship between x and y is positive.

D. as x decreases, so does y

Which of the following method of skewness based on quartiles?

A. Bowleys

B. Pearson

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

- C. Fisher
- D. Sheppred

When the mean deviation is calculated by taking deviations around median; then co-efficient of mean deviation is obtained dividing mean deviation by:

- A. Mode
- B. Mean
- C. Median**
- D. All above

The distribution will be symmetrical. IF

- A. None of the above
- B. $\text{median} > \text{midquartilerange} > \text{midrange}$
- C. $\text{median} = \text{midquartilerange} = \text{midrange}$**
- D. $\text{median} < \text{midquartilerange} < \text{midrange}$

In a five number summary, for a symmetrical distribution:

- A. Distance between Q1 and median equals to distance from median to Q3**
- B. Distance from X_0 to Q1 equals to the distance from Q3 to X_m
- C. Median, the mid-quartile range and the midrange would be equal.
- D. All of above

For empirical rule approximately all observations fall within

- A. $\bar{X} \pm 0.5\sigma$
- B. $\bar{X} \pm 3\sigma$
- C. $\bar{X} \pm 2\sigma$**
- D. $\bar{X} \pm 1\sigma$

According to Chebychev's rule, at least 8/9 of the data values will fall within:

- A. (Mean-3S.D, Mean+3S.D)**

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

For More Help Contact What's app 03224021365

- B. None of these
- C. (Mean-2S.D, Mean+2S.D)
- D. (Mean-S.D, Mean+S.D)

Moment ratio b_2 is used to measure:

- A. Symmetry
- B. Kurtosis** ok
- C. Mean
- D. Skewness

Standard Deviation of 64 is:

- A. 8**
- B. 3
- C. 4
- D. 10

If median > mid-quartile range > midrange then distribution will be:

- A. Zero skewed
- B. Symmetrical
- C. Positively skewed**
- D. Negatively skewed

When a researcher want to compare intensity of symptoms when different doses are administered. In this case, "intensity of symptoms" will be treated as:

- A. Discrete variable
- B. Independent variable
- C. Dependent variable**
- D. Quantitative variable

When our data set contains a few very high or a few very low values we use as a representative average.

- A. All above

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

For More Help Contact What's app 03224021365

B. Median

C. Mean

D. Mode

In a five number summary, which of the following is not used for data summarization?

A. the mean

B. the 25th percentile

C. the largest value

D. the smallest value

According to Empirical rule, approximately 95% of the measurements will fall within:

A. (Mean-2S.D, Mean+2S.D)

B. (Mean-S.D, Mean+S.D)

C. None of these

D. (Mean-3S.D, Mean+3S.D)

For a Normal distribution, b_2 (moment ratio) will be:

A. Equal to zero

B. Equal to 3

C. Greater than 3

D. Less than 3

The distribution will be negatively skewed if:

A. Q_2 closer to Q_3

B. $Q_1 = Q_3$

C. Q_1 closer to Q_2

D. $Q_1 = Q_2$

For a non-symmetric (Skewed) distributions median is than/to mean

A. Equal

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

B. Greater

C. Greater or less (Not equal)

D. Less

In a Box and Whisker plot, if the median line is closer to the left of the box then distribution will be:

A. Positively skewed

B. Symmetrical

C. Negatively skewed

D. Zero skewed

Question No:1

(Marks:1)

Vu-Topper RM

Positive square root of variance is known as:

A. Rang

B. Quartile deviation

C. Standard deviation

Page 91

ok

D. only (a) &(c)

Question No:2

(Marks:1)

Vu-Topper RM

Given $P(A) = 0.4$, $P(B) = 0.5$ and $P(A \cup B) = 0.9$, then:?

A. A and B are independent events

B. A and B are equally likely events

C. A and B are mutually exclusive events

ok

D. A and B are not mutually exclusive events

Question No:3

(Marks:1)

Vu-Topper RM

Which is appropriate average for finding the average speed of a car:

A. Mean

B. Mode

C. Both

D. None of these

Question No:4

(Marks:1)

Vu-Topper RM

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

The-----is often the preferred measure of central tendency if the data are severely skewed.

A. The median

B. The Mode

Question No:5

(Marks:1)

Vu-Topper RM

In measures of relative dispersion, unit of measurement remains:

A. Different

B. Free of units

ok

C. Square of units

D. same

Question No:6

(Marks:1)

Vu-Topper RM

Frequency curve is:

A. Asymmetric to x axis

B. Asymmetric to y axis

C. Non-Asymptotic to y-axis

D. Non-Asymptotic to y-axis

Question No:7

(Marks:1)

Vu-Topper RM

Parameter is aquantity.

A. Constant

B. Variable

Question No:8

(Marks:1)

Vu-Topper RM

What does the set comprising all possible outcomes of an experiment known as?

A. Sure event

ok

B. Null event

C. Elementary event

D. None of the these

Question No:9

(Marks:1)

Vu-Topper RM

بري صحبت سے تنہائي بہتر ہے اور تنہائي سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

A histogram is consisting of a set of a set of adjacent rectangles whose bases are marked off by:

A. Class boundaries Page 32

B. Class limits

C. Class frequency

D. Class marks

Question No:10

(Marks:1)

Vu-Topper RM

The middle value of an ordered array of numbers is the

A. Mean

B. Median Page 59

C. Mode

D. Midpoint

Question No:11

(Marks:1)

Vu-Topper RM

If Mean =25 & S.D is 5 then C.V is:

A. 100%

B. 20% Page 88

C. 10%

D. 25%

Question No:12

(Marks:1)

Vu-Topper RM

You connect the mid-points of rectangles in a histogram by a series of lines that also touches the x-axis from both ends, you will get:

A. Ogive

B. Frequency polygon

C. Frequency curve Page 38

D. Histogram

Question No:13

(Marks:1)

Vu-Topper RM

The conditional probability $P(A/B)$ is:

A. $P(A \cap B)/P(B)$ Page 154

B. $P(A \cap B)/P(A)$

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

- C. $P(A \cup B)/P(B)$
D. $P(A \cup B)/P(A)$

Question No:14

(Marks:1)

Vu-Topper RM

Chebyshev's inequality does not hold for $k = ?$

- A. 3
B. 2
C. 1

D. 0

Page 94

ok

Question No:15

(Marks:1)

Vu-Topper RM

According to Empirical rule, approximately 68% of the measurements will fall within:

- A. (Mean – S.D, Mean + S.D)
B. (Mean – 2S.D, Mean + 2S.D)
C. (Mean – 3S.D, Mean + 3S.D)
D. None of these

Page 90

ok

Question No:16

(Marks:1)

Vu-Topper RM

For Platykurtic distribution, b_2 (moment ratio) will be:

- A. Greater than 3
B. Less than 3
C. Equal to 3
D. Equal to zero

Page 114

ok

Question No:17

(Marks:1)

Vu-Topper RM

There are two broad categories of data, which are:

- A. Weighted and Un-weighted
B. Grouped and Un-grouped

C. Qualitative and Quantitative

Page 16

D. Primary and Secondary

Question No:18

(Marks:1)

Vu-Topper RM

بري صحبت سے تنہائي بہتر ہے اور تنہائي سے نيک صحبت بہتر ہے

For More Help Contact What's app 03224021365

In a linear regression, $Y=a+bX$, the variable “Y” will always:

A. A random variable

Page 117

ok

B. A non-random variable

C. Qualitative variable

D. Quantitative variable

Question No:19

(Marks:1)

Vu-Topper RM

In regression line $Y= a+bX$, X is called:

A. Dependent variable

B. Independent variable

Page 116

C. Explained variable

D. Regress and

Question No:20

(Marks:1)

Vu-Topper RM

The extremely positively skewed curve is also known as:

A. Frequency curve

B. U-shaped curve

C. J-shaped curve

D. Reverse J-shaped curve

Page 35

Question No:21

(Marks:1)

Vu-Topper RM

An event that contains more than one sample points is called:

A. Mutually exclusive event

B. Not mutually exclusive event

C. Hyper event

D. Compound event

Page 140

ok

Question No:22

(Marks:1)

Vu-Topper RM

Direct personal investigation is-----when the area to be covered is vast.

A. Costly

B. Time-consuming

C. Both

Page 6

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:23

(Marks:1)

Vu-Topper RM

According to this empirical rule, approximately how much values will fall within (Mean – 3S.D, Mean + 3S.D)?

A. 100%

Page 90

B. 95%

C. 75%

D. 68%

Question No:24

(Marks:1)

Vu-Topper RM

For a perfect symmetric curve distance from X_o to Q_1 is to the distance from Q_3 to X_m .

A. Less

B. Equal

Page 92

ok

C. Greater

D. None of the above

Question No:25

(Marks:1)

Vu-Topper RM

In a linear regression, best fitted line is obtained through:

A. Method of moment

B. Method of likelihood

C. Method of least square

Page 122

ok

D. Method of semi average

Question No:26

(Marks:1)

Vu-Topper RM

When all the values falling in a class are equal to the mid point of the class interval is called

A. Random error

B. Unbiased Error

C. Biased Error

D. Grouping Error

Page 56

Question No:27

(Marks:1)

Vu-Topper RM

بري صحبت سے تنہائي بہتر ہے اور تنہائي سے نيک صحبت بہتر ہے

For More Help Contact What's app 03224021365

For a Leptokurtic distribution, b_2 (moment ratio) will be:

A. Greater than 3 **Page 114**

B. Less than 3

C. Equal to 3

D. Equal to zero

Question No:28

(Marks:1)

Vu-Topper RM

Which scale will you use to measure the temperature?

A. Nominal scale

B. Interval scale **Page 4**

C. Ratio scale

D. Ordinal scale

Question No:29

(Marks:1)

Vu-Topper RM

The number of classes in a frequency distribution depends upon:

A. Sample

B. Population **Page 29**

C. Range

D. Average

Question No:30

(Marks:1)

Vu-Topper RM

Variance is expressed in units as the units of data set.

A. Squared **Page 90** **ok**

B. Cube

C. Single

D. Same

Question No:31

(Marks:1)

Vu-Topper RM

Which of the following will be used to draw an OGIVE?

A. A cumulative frequency distribution **Page 43**

B. A joint frequency distribution

C. A frequency distribution

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

D. A relative frequency distribution

Question No:32

(Marks:1)

Vu-Topper RM

If the Coefficient of variance = 50% and standard deviation= 2, what will be the value of mean(u)?

A. 4 **Page 93**

B. 5

C. 8

D. 10

Question No:33

(Marks:1)

Vu-Topper RM

In uni-modal distribution, if mode is less than mean

A. Positively Skewed **Page 98**

B. Negatively Skewed

C. Symmetrical

D. Symmetrical

Question No:34

(Marks:1)

Vu-Topper RM

Dispersion means the that exists in a data set.

A. Similiarty

B. Variability **Page 51**

C. Strength

D. Weakness

Question No:35

(Marks:1)

Vu-Topper RM

The _____ is the value you calculate when you want the arithmetic average:

A. Mode

B. Median

C. Mean **Page 58**

D. All above

Question No:36

(Marks:1)

Vu-Topper RM

بري صحبت سے تنہائي بہتر ہے اور تنہائي سے نيک صحبت بہتر ہے

For More Help Contact What's app 03224021365

When coin is tossed, the sample space consist of:

A. 2 outcomes **Page 145**

B. 4 outcomes

C. 6 outcomes

D. 8 outcomes

Question No:37

(Marks:1)

Vu-Topper RM

Sometimes mean deviation can also be caluculated around:

A. Quartiles

B. Deciles

C. Median

Page 89

ok

D. None of the above

Question No:38

(Marks:1)

Vu-Topper RM

A teacher asked 10 of her students how many books they had read in the last 12 months. Their answers were as follow: 12,13,19,6,7,15,25,21,12

The stem part is:

A. 12,6,7,10

B. 0,1,2

Page 47

C. 1,2,3,7,6

D. 25,21,12

Question No:39

(Marks:1)

Vu-Topper RM

In case of frequency distribution, the second quartile is given by the formula:

A. $l + 2h/f(n/2-c)$

B. $l + f/h(n/4-c)$

C. $l + f/h(2n/4-c)$

D. $l + h/f(2n/4-c)$

Page 69

Question No:40

(Marks:1)

Vu-Topper RM

Quartile Deviation is also defined as:

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

A. Interquartile Range

B. Semi range

C. Semi interquartile range Page 85

D. range

Question No:41

(Marks:1)

Vu-Topper RM

The types of frequency distribution are:

A. 3

B. 4 Page 38 ok

C. 5

D. 2

Question No:42

(Marks:1)

Vu-Topper RM

There are 30 people in a group. If all shake hands with one another , how many handshakes are possible?

A. 435 ok

B. 370

C. 291

D. 870

Question No:43

(Marks:1)

Vu-Topper RM

Which of the following is a subset of population?

A. Distribution

B. Sample

C. Data

D. Set

Question No:44

(Marks:1)

Vu-Topper RM

The data which has undergone any statistical treatment is called:

A. Primary data

B. Secondary Data Page 11

C. Qualitative data

D. Quantitative Data

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:45

(Marks:1)

Vu-Topper RM

A coin is tossed 4 times in succession. What is the probability that at least one head occurs?

A. 16/15

B. 15/16

C. 2/16

D. 1/16

Question No:46

(Marks:1)

Vu-Topper RM

From the following table;

The value of $F(x=3)$ will be:

A. 1/36

B. 36/36

C. 35/36

Question No:47

(Marks:1)

Vu-Topper RM

When a fair die is rolled, then sample space consists of:

A. 2 outcomes

ok

B. 16 outcomes

C. 6 outcomes

D. 36 outcomes

Question No:48

(Marks:1)

Vu-Topper RM

Positive correlation COEFFICIENT “r” will fall within the range:

A. $-1 < r < 0$

B. $-1 < r < 1$

C. All

Question No:49

(Marks:1)

Vu-Topper RM

In a lottery, there are 10 prizes and 25 blanks. A lottery is drawn at random. What is the probability of getting a prize?

A. 1/10

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

B. $2/5$

C. $5/7$

D. $2/7$ ok

Question No:50

(Marks:1)

Vu-Topper RM

If one event is not affected by the outcome of the other event, the two events are said to be:

A. Independent

B. Dependent

C. Mutually Exclusive

D. Not Mutually Exclusive

Question No:51

(Marks:1)

Vu-Topper RM

P (A union B) is equal to:

A. $P(A) - P(B)$.

B. $P(A) + P(B)$.

C. $P(A) + P(B) - P(A \cap B)$

D. $P(A) + P(B) + P(A \cap B)$

Question No:52

(Marks:1)

Vu-Topper RM

In a box, there are 8 red, 7 blue and 6 green balls. One ball is picked up randomly. What is the probability that it is neither red nor green?

A. $1/3$

B. $3/4$

C. $7/19$

D. $8/21$

Question No:53

(Marks:1)

Vu-Topper RM

A fair coin is tossed three times. What is the probability that at least one head appears?

A. $1/8$

B. $7/8$ ok

C. $4/8$

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

D. 6/8

Question No:54

(Marks:1)

Vu-Topper RM

If we roll a die then probability of an even number will be

A. 1

B. 3/6 ok

C. 4/6

D. 2/6

Question No:55

(Marks:1)

Vu-Topper RM

If $f(x)$ is a continuous probability function, then $P(X = 2)$ is:

A. 1

B. 0

C. $\frac{1}{2}$

D. 2

Question No:56

(Marks:1)

Vu-Topper RM

In regression line $Y=a+bX$, Y is called:

A. Dependent variable ok

B. Independent variable

C. Explanatory variable

D. Regressor

Question No:57

(Marks:1)

Vu-Topper RM

If A and B are mutually exclusive events with $P(A) = 0.25$ and $P(B) = 0.50$, Then $P(A \text{ or } B) =$

A. 0.25

B. 0.75 ok

C. 0.50

D. 1

Question No:58

(Marks:1)

Vu-Topper RM

In a 52 well shuffled pack of 52 playing cards, the probability of

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

drawing any one diamond card is

- A. $1/52$
- B. $4/52$
- C. $13/52$**
- D. $52/52$

Question No:59

(Marks:1)

Vu-Topper RM

Probability of a sure event is

- A. 8
- B. 1**
- C. 0
- D. 0.5

Question No:60

(Marks:1)

Vu-Topper RM

If $Y=3X+5$, then S.D of Y is equal to

- A. 9 s.d(x)
- B. 3 s.d(x)**
- C. s.d(x)+5
- D. 3s.d(x)+5

ok

Question No:61

(Marks:1)

Vu-Topper RM

The probability of drawing a red queen card from well-shuffled pack of 52 playing cards is

- A. $4/52$
- B. $2/52$**
- C. $13/52$
- D. $26/52$

ok

Question No:62

(Marks:1)

Vu-Topper RM

If $P(B|A) = 0.25$ and $P(A \text{ and } B) = 0.20$, then $P(A)$ is

- A. 0.05
- B. 0.80**

ok

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

- C. 0.95
- D. 0.75

Question No:63

(Marks:1)

Vu-Topper RM

When a coin is tossed 3 times, the probability of getting 3 tails is

- A. 1/8**
- B. 3/8
- C. 3/6
- D. 2/8

Question No:64

(Marks:1)

Vu-Topper RM

In how many ways, a team of 11 players can be chosen from a total of 16 players?

- A. 4368** **ok**
- B. 2426
- C. 5400
- D. 2680

Question No:65

(Marks:1)

Vu-Topper RM

The standard deviation of c (constant) is equal to:

- A. C
- B. c square
- C. 0** **ok**
- D. 2c

Question No:66

(Marks:1)

Vu-Topper RM

If P (E) is the probability that an event will occur, which of the following must be false:

- A. P(E)= - 1**
- B. P(E)=1
- C. P(E)=1/2
- D. P(E)=1/3

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:67

(Marks:1)

Vu-Topper RM

Let E and F be events associated with the same experiment. Suppose the E and F are independent and that $P(E) = 1/4$ and $P(F) = 1/2$ Then $P(E \cup F)$ is:

- A. $1/8$
- B. $3/4$
- C. $7/8$
- D. $5/8$

Question No:68

(Marks:1)

Vu-Topper RM

A student solved 25 questions from first 50 questions of a book to be solved. The probability that he will solve the remaining all questions is:

- A. 0.25
- B. 0.5
- C. 1
- D. 0

Question No:69

(Marks:1)

Vu-Topper RM

If $Y = bX$, then variance of Y is

- A. $b^2 \text{ var}(x)$
- B. $\text{var}(x)$
- C. $b \text{ var}(x)$
- D. $b \text{ square root var}(x)$

Question No:70

(Marks:1)

Vu-Topper RM

The classical definition of probability is not applicable when the assumption of does not hold:

- A. Exhaustive events
- B. Mutually exclusive events
- C. Equally likely events
- D. Independent events

Question No:71

(Marks:1)

Vu-Topper RM

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

In scatter diagram, the variable plotted along Y-axis is:

- A. Independent variable
- B. Dependent variable** ok
- C. Continuous variable
- D. Discrete variable

Question No:72

(Marks:1)

Vu-Topper RM

Which of the following measures of dispersion are based on deviations from the mean?

- A. Variance
- B. Standard deviation
- C. Mean deviation
- D. All of these** ok

Question No:73

(Marks:1)

Vu-Topper RM

What does it mean when a data set has a standard deviation equal to zero?

- A. All values of the data appear with the same frequency.
- B. The mean of the data is also zero.**
- C. All of the data have the same value.
- D. There are no data to begin with.

Question No:74

(Marks:1)

Vu-Topper RM

A set of possible values that a random variable can assume and their associated probabilities of occurrence are referred to as _____.

- A. Probability distribution**
- B. The expected return
- C. The standard deviation
- D. Coefficient of variation

Question No:75

(Marks:1)

Vu-Topper RM

Which of the following can never be probability of an event?

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

- A. 0
- B. 1
- C. 0.5
- D. -0.5**

Question No:76

(Marks:1)

Vu-Topper RM

The standard deviation of -1, -1, -1, -1 will be

- A. 1
- B. -1
- C. 0** **ok**
- D. Does not exist

Question No:77

(Marks:1)

Vu-Topper RM

Which formula represents the probability of the complement of event A:

- A. $1 + P(A)$
- B. $1 - P(A)$**
- C. $P(A)$
- D. $P(A) - 1$

Question No:78

(Marks:1)

Vu-Topper RM

The Special Rule of Addition is used to combine:

- A. Independent Events
- B. Mutually Exclusive Events**
- C. Events that total more than 1.00
- D. Events based on subjective probabilities

Question No:79

(Marks:1)

Vu-Topper RM

Set which is the sub-set of every set is

- A. Empty Set**
- B. Power Set
- C. Universal Set
- D. Super Set

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:80

(Marks:1)

Vu-Topper RM

$E(4X + 5) =$ _____

A. 12 E (X)

B. 4 E (X) + 5

C. 16 E (X) + 5

D. 16 E (X)

Question No:81

(Marks:1)

Vu-Topper RM

When two dice are rolled the number of possible sample points is :

A. 6

B. 12

C. 24

D. 36

Question No:82

(Marks:1)

Vu-Topper RM

If two events A and B are not mutually exclusive then

A. $P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$

ok

B. $P(A \text{ or } B) = P(A) + P(B)$

C. $P(A \text{ or } B) = P(A) \times P(B)$

D. $P(A \text{ or } B) = P(A) + P(B)$

Question No:83

(Marks:1)

Vu-Topper RM

Evaluate $(10-4)!$

A. 1000

B. 720

C. 480

D. 32

Question No:84

(Marks:1)

Vu-Topper RM

When E is an impossible event, then $P(E)$ is:

A. 0

B. 1

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

- C. 2
- D. 0.5

Question No:85

(Marks:1)

Vu-Topper RM

When we toss a coin, we get only:

- A. 1 outcome**
- B. 2 outcome
- C. 3 outcome
- D. 4 outcome

Question No:86

(Marks:1)

Vu-Topper RM

For exhaustive events, the $P(A \cup B \cup C)$ is equal to:

- A. $P(A)$
- B. $P(S)$**
- C. $P(A) * P(B) * P(C)$
- D. $P(B)$

Question No:87

(Marks:1)

Vu-Topper RM

A student solved 25 questions from first 50 questions of a book to be solved. The probability that he will solve the remaining all questions is:

- A. 0.25
- B. 0.5**
- C. 1
- D. 0

Question No:88

(Marks:1)

Vu-Topper RM

A set of possible values that a random variable can assume and their associated probabilities of occurrence are referred to as _____.

- A. Probability distribution**
- B. The expected return
- C. The standard deviation
- D. Coefficient of variation

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:89

(Marks:1)

Vu-Topper RM

If we roll a die then probability of getting a '6' will be

A. 2/6

B. 1/6

C. 4/6

D. 1

Question No:90

(Marks:1)

Vu-Topper RM

If $P(A) = 0.45$, $P(B) = 0.35$, and $P(A \text{ and } B) = 0.25$, then $P(A | B)$ is:

A. 1.4

B. 1.8

C. 0.714 ok

D. 0.556

Question No:91

(Marks:1)

Vu-Topper RM

Which of the following is not a measure of central tendency?

A. Percentile

B. Quartile

C. Standard deviation

D. Mode

Question No:92

(Marks:1)

Vu-Topper RM

Random experiment can be repeated any no. of times under the..... conditions.

A. Different

B. Similar

Question No:93

(Marks:1)

Vu-Topper RM

The simultaneous occurrence of two events is called:

A. Joint probability

B. Subjective probability

C. Prior probability

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

D. Conditional probability

Question No:94

(Marks:1)

Vu-Topper RM

The sum of squared deviation from mean is:

- A. Minimum
- B. Maximum
- C. Zero**
- D. Undefined

Question No:95

(Marks:1)

Vu-Topper RM

A frequency curve touches x-axis:

- A. No
- B. Yes**
- C. Some times
- D. None of these

Question No:96

(Marks:1)

Vu-Topper RM

Harmonic means is better than other means if the data are for:

- A. Ratios or proportion**
- B. Heights or Lengths
- C. Binary values like 0 &1
- D. Speed of rates

Question No:97

(Marks:1)

Vu-Topper RM

Quantiles are:

- A. A range of scores which might contain the population value
- B. Points on a distribution which split it into equal sized portions**
- C. Summary values for the entire population
- D. The difference the top and bottom 5% of scores

Question No:98

(Marks:1)

Vu-Topper RM

The free hand frequency curve is actually a:

- A. Scientific concept

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

For More Help Contact What's app 03224021365

B. Theoretically concept

- C. Mathematical concept
- D. Probabilistic concept

Question No:99

(Marks:1)

Vu-Topper RM

If any value in the data is 0 then it is not possible to have:

A. Harmonic mean

- B. Arithmetic mean
- C. Mode
- D. Median

Question No:100

(Marks:1)

Vu-Topper RM

Value of harmonic mean depends on:

A. All the observations

- B. Both a and b
- C. Few observations
- D. Extreme values

Question No:101

(Marks:1)

Vu-Topper RM

The middle value of an ordered array of number is the:

A. Mid-point

B. Median

- C. Mode
- D. Mean

Question No:102

(Marks:1)

Vu-Topper RM

In case of an open-ended class:

- A. A median can-not be computed
- B. The arithmetic mean and median will always be exactly equal
- C. A mean can-not be computed

D. The distribution is always positively skewed

Question No:103

(Marks:1)

Vu-Topper RM

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Which one is the formula of mid quartile range:

A. $(Q1+Q3)/2$

B. $Q3-Q1$

C. $(Q1-Q3)/2$

D. $(Q3-Q1)/2$

Question No:104

(Marks:1)

Vu-Topper RM

In a cumulative frequency polygon, the cumulative frequency of each class is plotted against:

A. Mid-point

B. Upper class limit

C. Lower class boundary

D. Upper class boundary

Question No:105

(Marks:1)

Vu-Topper RM

The number of classes in a frequency distribution generally should be:

A. Between five and twenty

ok

B. Between ten and twenty

C. More than five

D. Less than five

Question No:106

(Marks:1)

Vu-Topper RM

Find the medians of the set of numbers 1,2,3,4,5,6,7,8,9 and 10:

A. 55

B. 1

C. 5.5

D. 10

Question No:107

(Marks:1)

Vu-Topper RM

Relationship among the averages:

A. $GM \leq HM \leq AM$

B. $HM \geq GM \geq AM$

C. $AM \leq HM \leq GM$

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

D. $AM \geq GM \geq HM$

Question No:108

(Marks:1)

Vu-Topper RM

Which of the following comes first to make frequency distribution:

A. Range

B. Tally mark

C. Class interval

D. No. of groups

Question No:109

(Marks:1)

Vu-Topper RM

It is recommended that the number of classes in a frequency distribution be between:

A. 10 and 20

B. 5 and 20

C. 5 and 15

D. 6 and 20

Question No:110

(Marks:1)

Vu-Topper RM

The number of times each values appears is called the value's:

A. Frequency

B. Mode

C. Range

D. Standard deviation

Question No:111

(Marks:1)

Vu-Topper RM

A tabular management for classifying data into different group is called:

A. Class mark

B. Arithmetic mean

C. Frequency distribution

D. Standard deviation

Question No:112

(Marks:1)

Vu-Topper RM

Serious disadvantage of using range as a measure of dispersion is that it is based on only:

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

- A. Minimum Values
- B. Maximum Values
- C. Both Minimum and Maximum values
- D. None of the above**

Question No:113 (Marks:1)

Vu-Topper RM

Frequency of a variable is always in:

- A. Fraction form
- B. Percentage form
- C. Less than form
- D. Integer form**

Question No:114

(Marks:1)

Vu-Topper RM

5C_5 is equal to

- A. 5
- B. 1**
- C. 10
- D. 24

Question No:115

(Marks:1)

Vu-Topper RM

If $A = \{1,2,3,4\}$ and $B = \{3,4,5,6\}$ then $A - B$ will be:

- A. **$\{1,2\}$**
- B. $\{3,4\}$
- C. $\{3,2,1\}$
- D. $\{1,2,3,4,5,6\}$

Question No:116

(Marks:1)

Vu-Topper RM

Difference between the largest and the smallest data values is called

- A. variance
- B. interquartile range
- C. range**
- D. coefficient of variation

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:117

(Marks:1)

Vu-Topper RM

A list of 7 pulse rates is: 70, 64, 80, 74, 92, 96, 98. What is the median for this list?

A. 70

B. 80

C. 92

D. 98

Question No:118

(Marks:1)

Vu-Topper RM

The value of $10C_9$:

A. 45

B. 35

C. 35

D. 10

ok

Question No:119

(Marks:1)

Vu-Topper RM

The most frequent value in the data is called

A. Mean

B. Median

C. Mode

D. Harmonic mean

Question No:120

(Marks:1)

Vu-Topper RM

Calculate range for the following data: 22, 22, 30, 32, 37, 48, 60, 88, 90.

A. 22

B. 90

C. 37

D. 68

Question No:121

(Marks:1)

Vu-Topper RM

What is the median of this set of numbers: 4, 6, 7, 9, 2000000?

A. 9

B. 6

بري صحبت سے تنہائي بہتر ہے اور تنہائي سے نيک صحبت بہتر ہے

For More Help Contact What's app 03224021365

C. 7.5

D. 7

Question No:122

(Marks:1)

Vu-Topper RM

The value of the middle term in a ranked (ordered) data set is called the:

A. Mode

B. Mean

C. Median

D. Harmonic mean

Question No:123

(Marks:1)

Vu-Topper RM

The median is _____.

A. The highest number

B. The middle point

C. The average

D. Affected by extreme scores

Question No:124

(Marks:1)

Vu-Topper RM

The Mode of 8, 5, 7, 10, 15, 21, 5, 7, 2, 5 is

A. 8

B. 5

C. 7

D. 21

Question No:125

(Marks:1)

Vu-Topper RM

Let $A = \{1, 2, 3, 4\}$ and $B = \{3, 4, 5, 6\}$ Then $A \cap B$:

A. {3,4} **ok**

B. {1,4}

C. {3,5}

D. {3,6}

Question No:126

(Marks:1)

Vu-Topper RM

The range of the scores 29, 3, 143, 27, 99 is:

بري صحبت سے تنہائي بہتر ہے اور تنہائي سے نيک صحبت بہتر ہے

For More Help Contact What's app 03224021365

A. 140

B. 143

C. 146

D. 70

Question No:127

(Marks:1)

Vu-Topper RM

In how many ways, a team of 11 players can be chosen from a total of 16 players?

A. 4368

B. 2426

C. 5400

D. 2680

Question No:128

(Marks:1)

Vu-Topper RM

What is the mean of this set of numbers: 4, 6, 7, 9, 200000?

A. 7.5

B. 7

C. 400,005.2

D. 4

Question No:129

(Marks:1)

Vu-Topper RM

In case of frequency distribution, the median is given by the formula:

A. $I + \frac{h}{f} (n/2 - 2c)$

B. $I + \frac{h}{f} (n/2 - c)$

C. $I + \frac{f}{h} (n/2 - c)$

D. $I + \frac{f}{h} (n/4 - c)$

Question No:130

(Marks:1)

Vu-Topper RM

The sum of squared deviations from mean is:

A. Maximum

B. Minimum

C. Zero

D. Undefined

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:131

(Marks:1)

Vu-Topper RM

In a week the prices of a bag of rice were 350,280,340,290,320,310,300.

A. 320

B. 315

C. 300

D. 420

Question No:132

(Marks:1)

Vu-Topper RM

Calculate range for the following data: 10, 32, 33, 34, 37, 42, 55, 58, 70

A. 50

B. 60

C. 40

D. 20

Question No:133

(Marks:1)

Vu-Topper RM

When the frequency distribution or curve departs from symmetry, is called

A. Skewed

B. Positively skewed

C. Negatively skewed

D. None of these

Question No:134

(Marks:1)

Vu-Topper RM

Measure of central tendency is used to measure:

A. Average

B. Variability

C. Location

D. Both Average and Location

Question No:135

(Marks:1)

Vu-Topper RM

Component bar charts are used when data is divided into:

A. Parts

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

B. Groups

C. Circles

D. None of these

Question No:136

(Marks:1)

Vu-Topper RM

In a Box and Whisker plot, right end of the box is referred as:

A. First quartile

B. Second quartile

C. Third quartile

D. Mode

Question No:137

(Marks:1)

Vu-Topper RM

Fourth moment about mean provides information about the _____ of the distribution.

A. Centre

B. Dispersion

C. Symmetry

D. Kurtosis

Page 114

Question No:138

(Marks:1)

Vu-Topper RM

Let A and B are two dependent events such that

$P(A)=1/4$, $P(A/B)=1/2$ and $P(B/A)=2/3$.

Find $P(A \cap B)$.

A. $1/8$

B. $1/6$

C. $2/3$

D. $1/4$

Question No:139

(Marks:1)

Vu-Topper RM

Consider a set $A = \{1, 2, 3\}$. What is the number of subsets of A?

A. 3

B. 6

C. 8 **Google** **ok**

بري صحبت سے تنہائي بہتر ہے اور تنہائي سے نيک صحبت بہتر ہے

For More Help Contact What's app 03224021365

D.9

Question No:140

(Marks:1)

Vu-Topper RM

The stem for the following data is:

22, 45, 36, 15, 14, 12, 14, 14, 17, 21, 24, 24, 25, 25, 26, 26, 27, 29, 31, 34, 35

A. 1,2,3,4

B. 1,2,3,4,5

C. 11, 12, 13, 14, 15

D. 10, 20, 30, 40, 50

Question No:141

(Marks:1)

Vu-Topper RM

An event that contains only one sample points is called:

A. Simple event

ok

B. Normal event

C. Compound event

D. Mutually exclusive event

Question No:142

(Marks:1)

Vu-Topper RM

Frequency distribution is considered as negatively skewed if all values of distribution moves to

A. lower tail

ok

B. median tail

C. variance tail

D. upper tail

Question No:143

(Marks:1)

Vu-Topper RM

Which of the following is NOT a common measure of central tendency?

A. Mode

B. Range

ok

C. Median

D. Mean

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:144

(Marks:1)

Vu-Topper RM

From the table given below, how many students obtained marks between 60 and 69?

Marks	f	Mid-Points
50-59	5	54.5
60-69	7	64.5
70-79	8	74.5
80-89	5	84.5

- A. 64.5
- B. 12
- C. 60
- D. 7**

Question No:145

(Marks:1)

Vu-Topper RM

If the first and third quartiles are 22, 16 and 56,36 respectively, then the quartile deviation is:

- A. 17.1**
- B. 30.5
- C. 50.5
- D. 51.3

Question No:146

(Marks:1)

Vu-Topper RM

Adding all the squared deviations taken from mean and dividing by the number of observations, we get:

- A. Standard Deviation
- B. Variance** Google ok
- C. Mean Deviation
- D. None of the above

Question No:147

(Marks:1)

Vu-Topper RM

Standard deviation divided by mean is known as:

- A. Co-efficient of standard deviation
- B. Co-efficient of variation** ok

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

C. None

D. Both Co-efficient of standard deviation and Co-efficient of variation

Question No:148

(Marks:1)

Vu-Topper RM

5C5 equals to:

A. 5

B. 1

ok

C. 10

D. 15

Question No:149

(Marks:1)

Vu-Topper RM

Which of the measure of dispersion is used to compare variation between two series?

A. C.V.

ok

B. Q.D.

C. M.D.

D. S.D.

Question No:150

(Marks:1)

Vu-Topper RM

If $Y=3X+5$, then S.D of Y is equal to

A. 9 s. d(x)

B. 3 s. d(x)

C. s. d(x)+5

D. 3 s. d(x) + 5

Question No:151

(Marks:1)

Vu-Topper RM

Which of the following technique is not used to represent the bivariate qualitative data?

A. Component Bar Chart

B. Multiple Bar Chart

C. Line Chart

D. Pie Chart

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:152

(Marks:1)

Vu-Topper RM

When a frequency distribution involves “open-end” classes, then which average is appropriate?

- A. Mean
- B. Mode
- C. Median**
- D. None of these

Page 62

ok

Question No:153

(Marks:1)

Vu-Topper RM

Harmonic mean is extremely useful in averaging _ types of data.

- A. Ratios
- B. Rates
- C. Both ratios and rates**
- D. None of the above

Question No:154

(Marks:1)

Vu-Topper RM

According to Empirical rule, approximately how much values will fall within (Mean-3S.D, Mean+3S.D)?

- A. 100%**
- B. 95%
- C. 75%
- D. 68%

Page 90

ok

Question No:155

(Marks:1)

Vu-Topper RM

What is probability of drawing two clubs from a well shuffled pack of 52 cards?

- A. 13/51
- B. 1/17**
- C. 1/26
- D. 13/17

ok

Question No:156

(Marks:1)

Vu-Topper RM

بري صحبت سے تنہائي بہتر ہے اور تنہائي سے نيک صحبت بہتر ہے

For More Help Contact What's app 03224021365

Which pair of measure cannot be calculated when one of numbers in the series is zero?

- A. G.M. and A.M.
- B. H.M. and A.M.
- C. G.M. and H.M.**
- D. None of these

Question No:157

(Marks:1)

Vu-Topper RM

Which of the following techniques is used to predict the value of one variable on the basis of other variables?

- A. Correlation analysis
- B. Coefficient of correlation
- C. Covariance
- D. Regression analysis**

ok

Question No:158

(Marks:1)

Vu-Topper RM

A bag contains 12 red balls and 12 blue balls. A ball is drawn at random. The probability that ball drawn is red is

- A. 1/2**
- B. 5/11
- C. 6/10
- D. 1

Question No:159

(Marks:1)

Vu-Topper RM

If the GM of a set of two observations is 10 and its HM is 8, then the AM of the set of observations is

- A. 100
- B. 12.5**
- C. 64
- D. 7.5

Question No:160

(Marks:1)

Vu-Topper RM

If $A = \{H, T\}$ then which of the following is power set of A?

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

- A. $\{\{\}, \{H, T\}\}$
- B. $\{\{H\}, \{T\}, \{H, T\}\}$
- C. $\{\{H\}, \{T\}, \{H, T\}, \{T, H\}\}$
- D. $\{\{\}, \{H\}, \{T\}, \{H, T\}\}$**

ok

Question No:161

(Marks:1)

Vu-Topper RM

The total number of observations, which are below a certain value are known as

- A. class boundaries
- B. class marks
- C. cumulative frequency**
- D. variances

Question No:162

(Marks:1)

Vu-Topper RM

Histogram can be drawn only for:

- A. Discrete frequency distribution
- B. Continuous frequency distribution**
- C. Continuous frequency distribution
- D. Relative frequency distribution

Question No:163

(Marks:1)

Vu-Topper RM

Classification is the process of arranging data according to:

- A. one characteristic
- B. Two or more characteristic
- C. Similar characteristic**
- D. None of these

Question No:164

(Marks:1)

Vu-Topper RM

Which of the following, measures the dispersion around mean?

- A. Mean deviation
- B. Standard deviation**
- C. Mean deviation and Standard deviation
- D. None of these

ok

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:165

(Marks:1)

Vu-Topper RM

What is mode for the following set of data: 1,1,2,2,5,5,7

- A. 1
- B. 1,2
- C. 1,2,5**
- D. no mode in the data

Question No:166

(Marks:1)

Vu-Topper RM

Which of the following averages give information about central value in the distribution?

- A. Mean
- B. Median** **ok**
- C. Mode
- D. Harmonic mean

Question No:167

(Marks:1)

Vu-Topper RM

In a Pie diagram, the sector of a circle is obtained by:

- A. $(\text{component part}/\text{total}) \times 100$
- B. $(\text{component part}/\text{total}) \times 360$**
- C. $(\text{component part}/\text{total}) \times 180$
- D. $(\text{component part}/\text{total}) \times 300$

Question No:168

(Marks:1)

Vu-Topper RM

Relationship among the averages

- A. $HM > GM > AM$
- B. $AM > GM > HM$**
- C. $GM < HM < AM$
- D. $AM > HM < GM$

Question No:169

(Marks:1)

Vu-Topper RM

Which of the scale is best to use for measuring the salary of an employee?

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

A. nominal

B. ordinal

C. interval

ok

D. ratio

Question No:170

(Marks:1)

Vu-Topper RM

The mean of a distribution is 30, the mode is 24 and the standard deviation is 4, then the coefficient of skewness will be:

A. Less than zero

B. Equal to zero

C. Greater than zero

D. None of the above

Question No:171

(Marks:1)

Vu-Topper RM

Smaller standard error of estimate shows:

A. Data points are very far to the line

B. Data points are close to the line

ok

C. There is no difference between line and points

D. Difference is additive

Question No:172

(Marks:1)

Vu-Topper RM

Data arranged in ascending or descending order of magnitude is called:

A. Ungrouped data

B. Grouped data

C. Discrete frequency distribution

D. Arrayed data

Question No:173

(Marks:1)

Vu-Topper RM

A circle in which sectors represents various quantities is called:

A. Histogram

B. Frequency Polygon

C. Pie Chart

ok

D. Component Bar Chart

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:174

(Marks:1)

Vu-Topper RM

Sum of the absolute deviations of the values is least when deviations are taken from:

- A. Mean
- B. Median**
- C. Mode
- D. G.M

Question No:175

(Marks:1)

Vu-Topper RM

Chebyshev's inequality is valid for the data set

- A. Sample**
- B. Entire population
- C. Both sample and entire population
- D. None of the above

Question No:176

(Marks:1)

Vu-Topper RM

Which of the following terms best describes data that were originally collected at an earlier time by a different person for a different purpose?

- A. Primary data
- B. Secondary data**
- C. Experimental data
- D. Field notes

Question No:177

(Marks:1)

Vu-Topper RM

Statistics deals with

- A. Individuals
- B. Isolated items
- C. Observations
- D. Aggregates of facts**

Question No:178

(Marks:1)

Vu-Topper RM

If a box contains six red, three blue and five pink ties then probability of

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

For More Help Contact What's app 03224021365

blue ties will be equal to:

A. 1/14

B. 3/14 ok

C. 5/14

D. 6/14

Question No:179

(Marks:1)

Vu-Topper RM

Which of the rule is applied to any data set, regardless shape of the frequency distribution?

A. Chebychev's rule

B. Empirical rule

C. Combination rule

D. Permutation rule

Question No:180

(Marks:1)

Vu-Topper RM

Which average is used in the situation where the number of floors in the buildings at the center of a city?

A. Mean

B. Median

C. Mode

D. Variance

Question No:181

(Marks:1)

Vu-Topper RM

Rankings of the finishes of competitors in a foot race is an example of a(n)_____.

A. ratio scale

B. ordinal scale

C. nominal scale

D. interval scale

Question No:182

(Marks:1)

Vu-Topper RM

Among 18 articles, six having minor Defects and three have major defects. Determine the probability that an article selected at random has

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

major defect.

A. 1/6

B. 1/5

C. 0.25

D. 0.11

Question No:183

(Marks:1)

Vu-Topper RM

A series of data with exclusive classes along with the corresponding frequencies is called:

A. Discrete frequency distribution

B. Continuous frequency distribution

C. Percentage frequency distribution

D. Cumulative frequency distribution

Question No:184

(Marks:1)

Vu-Topper RM

Using the following table, calculate $P(X < 2)$

X	0	1	2	3
f(x)	1/8	3/8	3/8	1/8

A. 1/8

B. 3/8

C. 4/8

D. 7/8

Question No:185

(Marks:1)

Vu-Topper RM

If a distribution has two modes then the distribution is called:

A. Uni-Modal

B. Bi-Modal

C. Tri-Modal

D. Multi-Modal

Question No:186

(Marks:1)

Vu-Topper RM

When two dice are rolled. What is the probability that total is at least 12:

A. $\frac{1}{36}$

بري صحبت سے تنہائي بہتر ہے اور تنہائي سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

B. $\frac{2}{36}$

ok

C. $\frac{12}{36}$

D. $\frac{36}{36}$

Question No:187

(Marks:1)

Vu-Topper RM

Correlation COEFFICIENT measures:

A. Dispersion

B. Skewness

C. Degree of linear relationship between two random variables

Page 128

ok

D. Dependence of one variable to another variable

Question No:188

(Marks:1)

Vu-Topper RM

When two dice are rolled the number of possible sample points are:

A. 6

B. 12

C. 24

D. 36 Confirm $6*6 = 36$

ok

Question No:189

(Marks:1)

Vu-Topper RM

The number of elements in the Power set $P(S)$ of the set $S = [[\Phi] , 1, [2, 3]]$ is:

A. 3

B. 4

C. 8 ($2^n = 2^3 = 8$)

ok

D. 6

Question No:190

(Marks:1)

Vu-Topper RM

In how many ways can a group of 5 men and 2 women be made out of a total of 7 men and 3 women?

A. 63

ok

B. 54

C. 86

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

D. 156

Question No:191

(Marks:1)

Vu-Topper RM

If the median of two observations is 8, then mean of these two observations will be:

A. 7

B. 9

C. 8 ok

D. 6

Question No:192

(Marks:1)

Vu-Topper RM

A fair die is rolled. Probability of getting even face given that face is less than 5 is given by:

A. $\frac{1}{2}$ ok

B. 5

C. 2

D. 6

Question No:193

(Marks:1)

Vu-Topper RM

The first and third quartiles are 22.16 and 56.36 respectively, then the quartile deviation is:

A. 17.1 ($(56.36 - 22.16)/2 = 17.1$)

B. 30.5

C. 50.5

D. 51.3

Question No:194

(Marks:1)

Vu-Topper RM

Consider a set $A = \{4, 6, 8, 10\}$. What is the number of subsets of A?

A. 2

B. 8

C. 10

D. 16 ($2^4 = 16$) ok

بري صحبت سے تنہائي بہتر ہے اور تنہائي سے نيک صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:195

(Marks:1)

Vu-Topper RM

Suitable average for averaging the shoe sizes for children is:

- A. Mean
- B. Mode**
- C. Median
- D. Geometric Mean

Question No:196

(Marks:1)

Vu-Topper RM

If we flip a coin five times, then possible outcomes of the sample space are:

- A. 2
- B. 4
- C. 16
- D. 32 ($2^5 = 32$)**

Question No:197

(Marks:1)

Vu-Topper RM

For a symmetrical distribution having 10 values the mean is 20. Which one of the following is the mode of the distribution?

- A. 20 (mean = mode = median)**
- B. 10
- C. 5
- D. 15

Question No:198

(Marks:1)

Vu-Topper RM

Let A and B are two dependent events such that $P(B)=1/3$, $P(A/B)=1/2$ and $P(B/A)=1/3$. Find $P(A \cap B)$.

- A. $1/6$ ($1/2 * 1/3 = 1/6$)**
- B. $1/9$
- C. $1/2$
- D. $1/3$

Question No:199

(Marks:1)

Vu-Topper RM

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

A bag contains 12 red balls and 12 blue balls. A ball is drawn at random. The probability that ball drawn is red is:

A. $1/2$ ($12/24 = 12$)

B. $5/11$

C. $6/10$

D. 1

Question No:200

(Marks:1)

Vu-Topper RM

Alpha is the probability of

A. Rejecting H_0

B. Accepting H_0

C. Rejecting H_1

D. Accepting H_1

Question No:201

(Marks:1)

Vu-Topper RM

What type of data is collected in population census?

A. Two Types

B. Four

C. Six

Question No:202

(Marks:1)

Vu-Topper RM

The collection of all outcomes for an experiment is called

A. A sample spaces

B. the intersection of events

C. joint probability

D. population

Question No:203

(Marks:1)

Vu-Topper RM

Which of the graph is used for a time series data:

A. Frequency curve

B. Frequency polygon

C. Histogram

D. Histogram

بري صحبت سے تنہائي بہتر ہے اور تنہائي سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:204

(Marks:1)

Vu-Topper RM

The value that has half of the observations above it and half the observations below it is known as:

- A. Mean
- B. Median**
- C. Mode
- D. Standard deviation

Question No:205

(Marks:1)

Vu-Topper RM

The height of a student is 60 inches. This is an example of

- A. Continuous data**
- B. Qualitative data
- C. Categorical data
- D. Discrete data

Page 9

Question No:206

(Marks:1)

Vu-Topper RM

If the both tails of the distribution are equal, then distribution is called:

- A. J-shaped
- B. Symmetrical**
- C. Positively Skewed
- D. Negatively Skewed

Question No:207

(Marks:1)

Vu-Topper RM

Ranking scale also include the properties of which scale?

- A. Nominal scale**
- B. Interval scale
- C. Ratio scale
- D. All of these

Question No:208

(Marks:1)

Vu-Topper RM

Range of the values -2.50, -3.70, -4.80, -3.10, -9.70, -2.20, -8.90, -1.60, 0.60 is

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

- A. 10.03
- B. **10.30**
- C. 9.10
- D. 9.00

Question No:209

(Marks:1)

Vu-Topper RM

If the standard deviation of a population is 5.5, the population variance is:

- A. 5.5
- B. 31
- C. 25
- D. **30.25**

Question No:210

(Marks:1)

Vu-Topper RM

Range of the values -10, -19, -9, -15, -28, -26, -25 is:

- A. +18
- B. -18
- C. -19
- D. **+19**

Question No:211

(Marks:1)

Vu-Topper RM

Which one of the following is less than median for a symmetrical distribution?

- A. 50percentile
- B. 51 percentiles
- C. 2quartile
- D. **4decile**

Question No:212

(Marks:1)

Vu-Topper RM

Sum of absolute deviations of the values is least when deviations are taken from

- A. **Mean**
- B. Median

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

- C. Mode
- D. gm.

Question No:213

(Marks:1)

Vu-Topper RM

Statistic is a numerical quantity, which is calculated from

- A. Data**
- B. Observation
- C. Sample
- D. population

Question No:214

(Marks:1)

Vu-Topper RM

The branch of Statistics that is concerned with the procedures and methodology for obtaining valid conclusions is called:

- A. descriptive
- B. advance
- C. inferential**
- D. sample

Question No:215

(Marks:1)

Vu-Topper RM

How to find the class midpoint?

- A. Half the sum of upper-class limit and lower-class limit
- B. Find the difference between consecutive lower limits**
- C. Count the number of observations in the class
- D. Divide the class frequency by the number of observe

Question No:216

(Marks:1)

Vu-Topper RM

For given data, discuss the shape of the distribution: X f 0.2 8 1.2 15 2.2 23 3.2 40

- A. Positively skewed
- B. Negatively skewed**
- C. Symmetric curve
- D. U- Shaped curve

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:217

(Marks:1)

Vu-Topper RM

if '2' is a leading digit in 24335, then what are the trailing digits in the observation to display a 'Stem-and –Leaf display'.

A. 4 335

B. 4335

C. 43

Question No:218

(Marks:1)

Vu-Topper RM

A frequency polygon is obtained by plotting the class frequencies against what?

A. class boundary

B. cumulative frequency

C. relative frequency

D. Mid-point

Question No:219

(Marks:1)

Vu-Topper RM

When more values are lying at the start of the distribution, it is:

A. u shape

B. positive

C. negative

D. symmetrical

Question No:220

(Marks:1)

Vu-Topper RM

The data for an ogive is found in which distribution:

A. A cumulative frequency distribution

B. A joint frequency distribution

C. A frequency distribution

D. A relative frequency distribution

Question No:221

(Marks:1)

Vu-Topper RM

Which one of the following is greater than median for a symmetrical distribution:

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

A. 1st Decile

B. 7th Decile

ok

C. 44th Percentile

D. 14th Percentile

Question No:222

(Marks:1)

Vu-Topper RM

Data classified by attributes are called:

A. Grouped data

B. Qualitative data

C. Quantitative data

D. **Arrayed data**

Question No:223

(Marks:1)

Vu-Topper RM

As a general rule, statisticians tend to use which of the following number of classes when arranging the data

A. Fewer than 5

B. **Between 5 & 20**

C. Between 8 & 15

D. More than 20

Question No:224

(Marks:1)

Vu-Topper RM

A quantity obtained by applying certain rule or formula is known as

A. **Estimate**

B. Estimator

Question No:225

(Marks:1)

Vu-Topper RM

The F-distribution always ranges from:

A. 0 to 1

B. 0 to -8

C. **-8 to +8**

D. 0 to +8

Question No:226

(Marks:1)

Vu-Topper RM

بري صحبت سے تنہائي بہتر ہے اور تنہائي سے نيک صحبت بہتر ہے

For More Help Contact What's app 03224021365

To find the estimate of a parameter..... methods are used.

- A. Two
- B. Three
- C. Four
- D. **Many**

Question No:227

(Marks:1)

Vu-Topper RM

A failing student is passed by an examiner. It is an example of:

- A. Type I error
- B. **Type II error**
- C. Correct decision
- D. No information regarding student exams

Question No:228

(Marks:1)

Vu-Topper RM

For two mutually exclusive events A and B, $P(A) = 0.2$ and $P(B) = 0.4$, then $P(A \cup B)$ is:

- A. 0.8
- B. 0.2
- C. **0.6**
- D. 0.5

$$P(A \cup B) = P(A) + P(B) = 0.2 + 0.4 = 0.6$$

ok

Question No:229

(Marks:1)

Vu-Topper RM

An urn contains 4 red balls and 6 green balls. A sample of 4 balls is selected from the urn without replacement. It is the example of:

- A. Binomial distribution
- B. **Hypergeometric distribution**
- C. Poisson distribution
- D. Exponential distribution

Page 219

Question No:230

(Marks:1)

Vu-Topper RM

If $P(A \cap B) = 0.12$ $P(A) = 0.3$, find $P(B)$ where 'A' and 'B' are independent:

- A. 0.1

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

- B. 0.2
- C. 0.3
- D. 0.4**

Question No:231

(Marks:1)

Vu-Topper RM

The mean deviation of the normal distribution is approximately:

- A. $7/8$ of the S.D
- B. $4/5$ of the S.D
- C. $3/4$ of the S.D**
- D. $1/2$ of the S.D

Question No:232

(Marks:1)

Vu-Topper RM

The conditional probability of the event A when event B has occurred is denoted by:

- A. $P(A/B)$** **ok**
- B. $P(A + B)$
- C. $P()$
- D. $P(A - B)$

Question No:233

(Marks:1)

Vu-Topper RM

The probability of an event is always:

- A. less than 1
- B. between 0 and 1**
- C. greater than 1

Question No:234

(Marks:1)

Vu-Topper RM

Symbolically, a conditional probability is:

- A. $P(AB)$
- B. $P(A/B)$** **ok**
- C. $P(A)$
- D. $P(A \cup B)$

Question No:235

(Marks:1)

Vu-Topper RM

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

If $P(A) = 0.3$ and $P(B) = 0.5$, find $P(A/B)$ where 'A' and 'B' are independent:

- A. 0.3
- B. 0.5
- C. 0.8
- D. **0.15**

Question No:236

(Marks:1)

Vu-Topper RM

The probability of an event cannot be

- A. 1
- B. 0.5
- C. 0.3
- D. **-0.5**

Question No:237

(Marks:1)

Vu-Topper RM

A set of possible values that a random variable can assume and their associated probabilities of occurrence are referred to as _____.

- A. **Probability distribution**
- B. The expected return
- C. The standard deviation
- D. Coefficient of variation

Question No:238

(Marks:1)

Vu-Topper RM

The probability of drawing any one spade card is:

- A. $1/52$
- B. $4/52$
- C. **$13/52$**
- D. $52/52$

Question No:239

(Marks:1)

Vu-Topper RM

The function abbreviated to d.f. is also called the.....

- A. Probability density function
- B. **Probability distribution function**

Page 172

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

- C. Commutative distribution function
- D. Discrete function

Question No:240

(Marks:1)

Vu-Topper RM

A discrete probability function $f(x)$ is always:

- A. Zero
- B. **One** Page 172
- C. Negative
- D. non-negative

Question No:241

(Marks:1)

Vu-Topper RM

In the FA examination, 24 candidates offered Statistics. If the probability of passing the subject be $1/3$, what will be the mean of the distribution?

- A. 7
- B. **8**
- C. 6
- D. 5

Question No:242

(Marks:1)

Vu-Topper RM

If the values of variables are increasing or decreasing in the same direction then such kind of correlation is referred as

- A. Zero Correlation
- B. Perfect Correlation
- C. **Positive Correlation**
- D. Negative Correlation

Question No:243

(Marks:1)

Vu-Topper RM

The best measure of variation is

- A. Range
- B. Quartile deviation
- C. **Variance**
- D. Coefficient of variance

Question No:244

(Marks:1)

Vu-Topper RM

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Ms. Christian calculated a correlation coefficient of .75. Which of the following reflects the best interpretation of this?

- A. Weak negative.
- B. Strong negative.
- C. Weak positive.
- D. Strong positive.**

Question No:245

(Marks:1)

Vu-Topper RM

.....use the division of a circle into different sectors.

- A. Line graph
- B. Sector graphs**
- C. Frequency Polygon
- D. Conversion Graphs

Question No:246

(Marks:1)

Vu-Topper RM

The measurement of measure of degree of to which any two variables vary together is called

- A. Regression Coefficient
- B. Correlation
- C. Both (a) and (b)
- D. None of these**

Question No:247

(Marks:1)

Vu-Topper RM

Analysis of Variance (ANOVA) is a test for equality of:

- A. Variances**
- B. Means
- C. Proportions
- D. only two parameters

Question No:248

(Marks:1)

Vu-Topper RM

If strength of the association between X and Y is very weak, then $r = ?$

- A. $r = -1$
- B. $r = 0$**

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

- C. $r = 1$
- D. $r = 2$

Question No:249

(Marks:1)

Vu-Topper RM

In the central tendency Mean, Median and Mode

- A. **Mean is better than Median**
- B. Median is better than Mode
- C. Mean is better than Mode
- D. All of these are true

Question No:250

(Marks:1)

Vu-Topper RM

The degree to which numerical data tend to spread about an average is called

- A. **The dispersion**
- B. Standard deviation
- C. Correlation
- D. None of these

Question No:251

(Marks:1)

Vu-Topper RM

..... graphs are similar to bar graphs.

- A. **Column**
- B. Line
- C. Conversion
- D. sector

Question No:252

(Marks:1)

Vu-Topper RM

A pattern of variation of a time series that repeats every year is called:

- A. Cyclical
- B. **Seasonal**
- C. Trend
- D. Secular

Question No:253

(Marks:1)

Vu-Topper RM

Assume that a population consists of 7 similar containers having the

بري صحبت سے تڻهائي بهتر هے اور تڻهائي سے نيگ صحبت بهتر هے

For More Help Contact What's app 03224021365

following weights (kg): 9.8, 10.2, 10.4, 9.8, 10.0, 10.2, 9.6 What is the second moment about mean?

- A. 0.262 kg
- B. **0.069kg**
- C. 0.521 kg
- D. 0.313kg

Question No:254

(Marks:1)

Vu-Topper RM

If the graph is very much scattered, then what can be the suitable value of r ?

- A. $r = -0.9$
- B. $r = -0.5$
- C. $r = 0.1$
- D. **$r=0.8$**

Question No:255

(Marks:1)

Vu-Topper RM

A list of pulse rates is 70. 64. 70. 80. 74, 92. What is the mode for this list?

- A. **70**
- B. 80
- C. 90
- D. 100

Question No:256

(Marks:1)

Vu-Topper RM

If the mean of two observations is 11.5. then the median of these two observations will be

- A. 10.5
- B. **11.5**
- C. 12.5
- D. 13.5

Question No:257

(Marks:1)

Vu-Topper RM

A coin is tossed and a single 6-sided 11.4 is rolled. Find the probability

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

of landing on the head side of the coin and rolling a 3 on the die_

A. 1/12

B. 2/12

C. 3/12

D. 4/12

Question No:258

(Marks:1)

Vu-Topper RM

When two dice are rolled What is the probability that total is at least 12

A. 1/36

B. 2/36

C. 3/36

D. 4/36

Question No:259

(Marks:1)

Vu-Topper RM

In a lottery there are 10 prizes and 25 blanks. A lottery is drawn at random What is the probability of getting a prize?

A. 1/7

B. 2/7

C. 3/7

D. 4/7

Question No:260

(Marks:1)

Vu-Topper RM

A graph of a cumulative frequency distribution is called

A. Ogive Curve

B. Frequency Polygon

C. Pie Chart

D. Bar Chart

Question No:261

(Marks:1)

Vu-Topper RM

A Histogram contains a set of

A. Adjacent Rectangles

B. Non-adjacent Rectangles

C. Adjacent Squares

بري صحبت سے تنہائي بہتر ہے اور تنہائي سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

D. Adjacent Triangles

Question No:262

(Marks:1)

Vu-Topper RM

In a Pie chart one can calculate the angles for each sector by the following formula

A. $(\text{Component part} / \text{Total}) \times 100$

B. $(\text{Component part} / \text{Total}) \times \pi$

C. $(\text{Total} / \text{Component part}) \times 360$

D. $(\text{Component part} / \text{Total}) \times 360$

Question No:263

(Marks:1)

Vu-Topper RM

A frequency polygon is constructed by plotting frequency of the class interval and the

A. The upper limit of the class

B. The lower limit of the class

C. Mid value of the class

D. None of the above

Question No:264

(Marks:1)

Vu-Topper RM

A frequency polygon is a closed figure of

A. Two sides

B. Three sides

C. Many sides

D. None of these

Question No:265

(Marks:1)

Vu-Topper RM

De-cumulative frequency is presented by

A. More than Ogive

B. Less than Ogive

C. Equal to Ogive

D. None of these

Question No:266

(Marks:1)

Vu-Topper RM

In a histogram, the area of each rectangle is proportional to

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

- A. The class mark of the corresponding class interval
- B. The class size of the corresponding class interval
- C. Frequency of the corresponding class interval**
- D. Cumulative Frequency of the corresponding class interval

Question No:267

(Marks:1)

Vu-Topper RM

A frequency polygon curve touches the x-axis

- A. Yes**
- B. No
- C. Some times
- D. None of the above

Question No:268

(Marks:1)

Vu-Topper RM

Which of the following considerations for setting up classes in a frequency distribution is correct?

- A. Class widths can be different
- B. Classes should not overlap**
- C. Open ended classes only at extremes
- D. The lower limit of the first class should not be an even multiple of the class width

Question No:269

(Marks:1)

Vu-Topper RM

What are the members in the right column of a frequency distribution table called?

- A. Class frequency**
- B. Interval frequency
- C. Ordinal frequency
- D. Number frequency

Question No:270

(Marks:1)

Vu-Topper RM

A variable is any characteristic which can assume_____ values.

- A. Different**
- B. Similar

بري صحبت سے تنہائي بہتر ہے اور تنہائي سے نيک صحبت بہتر ہے

For More Help Contact What's app 03224021365

- C. Fixed
- D. Assumed

Question No:271

(Marks:1)

Vu-Topper RM

A ---- variable is a variable whose values can theoretically take on an infinite number of values within a given range of values. a. Continuous

A. Discrete

- B. Random
- C. Both a and b

Question No:272

(Marks:1)

Vu-Topper RM

The magnitude of the class is the

- A. The product of lower limit and upper
- B. The sum of lower limit and upper
- C. The difference of upper limit and lower limit**
- D. None

Question No:273

(Marks:1)

Vu-Topper RM

The classes in which the lower limit or the upper limit is not specified are known as:

- A. Open end classes**
- B. Close end classes
- C. Inclusive classes
- D. Exclusive classes

Question No:274

(Marks:1)

Vu-Topper RM

Classes in which upper limits are excluded from the respective classes and are included in the immediate next class are:

- A. Open end classes
- B. Close end classes
- C. Inclusive classes
- D. Exclusive classes**

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:275

(Marks:1)

Vu-Topper RM

The number of observations in a particular class is called:

- A. Width of the class
- B. Class mark
- C. Frequency**
- D. None of the above

Question No:276

(Marks:1)

Vu-Topper RM

If the class mid points in a frequency distribution of age of a group of persons are 25, 32, 39, 46, 53 and 60. The size of class interval is:

- A. 5
- B. 7**
- C. 8
- D. 6

Question No:277

(Marks:1)

Vu-Topper RM

If the mid points of the classes are 16, 24, 32, 40, and so on, then the magnitude of the class interval is:

- A. 8**
- B. 9
- C. 7
- D. 6

Question No:278

(Marks:1)

Vu-Topper RM

A pie diagram is also called:

- A. Pictogram
- B. Angular diagram**
- C. Line diagram
- D. Bar diagram

Question No:279

(Marks:1)

Vu-Topper RM

The most commonly used device of presenting business and economic data is:

بري صحبت سے تنہائي بہتر ہے اور تنہائي سے نيک صحبت بہتر ہے

For More Help Contact What's app 03224021365

- A. Pie diagrams
- B. Pictograms
- C. Bar diagrams**
- D. Line diagrams

Question No:280

(Marks:1)

Vu-Topper RM

Type of bar diagram is:

- A. Pictogram
- B. Sub divided diagram**
- C. Line diagrams
- D. Pie diagram

Question No:281

(Marks:1)

Vu-Topper RM

The algebraic sum of deviations from mean is:

- A. Zero**
- B. One
- C. Two
- D. Five

Question No:282

(Marks:1)

Vu-Topper RM

If an observation in the data set is zero, then its geometric mean will be:

- A. Zero**
- B. One
- C. Two
- D. Five

Question No:283

(Marks:1)

Vu-Topper RM

Find the mode from these test results: 90, 80, 77, 86, 90, 91, 77, 66, 69, 65, 43, 65, 75, 43, 90.?

- A. 90** **ok**
- B. 80
- C. 70
- D. 60

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:284

(Marks:1)

Vu-Topper RM

The number of classes in a frequency distribution is obtained by dividing the range of variable by the:

- A. Total frequency
- B. Class interval**
- C. Mid-point
- D. Relative frequency

ok

Question No:285

(Marks:1)

Vu-Topper RM

If a curve has longer tail to the right, it is called:

- A. positive skew**
- B. negative skew

Question No:286

(Marks:1)

Vu-Topper RM

Histogram and histogram are:

- A. Always same
- B. Not same**
- C. Off and on same
- D. Randomly same

Question No:287

(Marks:1)

Vu-Topper RM

If $Q_1=62$ and $Q_3=87$, then the mid-quartile range will be:

- A. 74.4**
- B. 70.5
- C. 35.5
- D. 68.4

Question No:288

(Marks:1)

Vu-Topper RM

The measure of Dispersion can never be:

- A. Positive
- B. Negative**
- C. 0

بري صحبت سے تنہائي بہتر ہے اور تنہائي سے نيک صحبت بہتر ہے

For More Help Contact What's app 03224021365

D. 1

Question No:289

(Marks:1)

Vu-Topper RM

Data must be arranged either in ascending or descending order if some want to compute

A. Mode

B. Median

C. Geometric Mean

D. Harmonic Mean

Question No:290

(Marks:1)

Vu-Topper RM

Mean deviation is a measure of dispersion in which deviations are taken around the:

A. Mean

ok

B. First Quartile

C. Third Quartile

D. None of the above

Question No:291

(Marks:1)

Vu-Topper RM

The concept of a five-number summary directly linked with the concept of.

A. Polygon curve

B. Frequency curve

C. Box and whisker plot

D. Scatter plot

Question No:292

(Marks:1)

Vu-Topper RM

Value of the harmonic mean is lower than-----

A. Arithmetic Mean

B. Geometric Mean

C. Both arithmetic mean & geometric mean

D. None of the above

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:293

(Marks:1)

Vu-Topper RM

A graph of plotted points which shows the relationship between two sets of data is known as:

- A. Venn Diagram
- B. Polygon Curve
- C. Histogram diagram

D. Scatter diagram

Page 121

ok

Question No:294

(Marks:1)

Vu-Topper RM

For certain distribution, A.M=136.75, Median= 148.37 and Mode= 152.80, then the distribution will be:

- A. Positively skewed
- B. Negatively skewed (mean < median < mode)**
- C. Symmetrical
- D. Extremely negative J shaped

Question No:295

(Marks:1)

Vu-Topper RM

When two coins are tossed the probability of at least one head is:

- A. $1/4$
- B. $2/4$
- C. $3/4$ ((H,H),(H,T),(T,H),(T,T) = $3/4$)**
- D. 1

Question No:296

(Marks:1)

Vu-Topper RM

The deviation of a distribution from symmetry is called:

- A. Kurtosis
- B. Skewness**
- C. Dispersion
- D. Flatness

Page 101

ok

Question No:297

(Marks:1)

Vu-Topper RM

if a box contains two red and three green balls then probability of red balls will be equal to:

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

A. 2/3

B. 3/4

C. 2/5 (red balls/ total balls = 2/5) ok

D. 1/5

Question No:298 (Marks:1)

Vu-Topper RM

Co-efficient of standard deviation is:

A. An absolute measure of Dispersion

B. A relative measure of dispersion Page 93 ok

C. Both

D. None

Question No:299 (Marks:1)

Vu-Topper RM

If any value in the data is zero, then it is not possible to have?

A. Harmonic Mean Page 77

B. Arithmetic Mean

C. Median

D. Mode

Question No:300 (Marks:1)

Vu-Topper RM

If the grading of diabetes is classified as mild, moderate and severe the scale of measurement used is:

A. Interval

B. Nominal

C. Ordinal ok

D. Ratio

Question No:301 (Marks:1)

Vu-Topper RM

_____ determines the shape of the frequency distribution without drawing graph of frequency distribution.

A. Probability theory

B. Random number theory

C. Scatter diagram

D. Five number theory Page 99 ok

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:302

(Marks:1)

Vu-Topper RM

A sector diagram is also called?

- A. Bar diagram
- B. Histogram
- C. Historigram

D. Pie diagarm

Page 23

Question No:303

(Marks:1)

Vu-Topper RM

_____ is the measure of average which can have more than one value.

- A. Harmonic Mean
- B. Geometric Mean
- C. Median

D. Mode

Question No:304

(Marks:1)

Vu-Topper RM

If $\mu=3.82$ and $S.D(X)=1.2$, then $C.V(X)$ will be:

- A. 20.482
- B. 24.896
- C. 31.412**
- D. 26.451

Question No:305

(Marks:1)

Vu-Topper RM

When the peak value of the curve becomes relatively high, it is called:

- A. Mesokurtic
- B. Leptokurtic**
- C. Platykurtic
- D. hetrokurtic

Page 114

Question No:306

(Marks:1)

Vu-Topper RM

A relative measure of dispersion is one that is expressed in the form of:

- A. Ratio
- B. Co-efficient

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

C. Percentage

D. All above

Question No:307

(Marks:1)

Vu-Topper RM

In case of positively skewed distribution:

A. $Q1 + Q2 = 2 \text{ median} < 0$

B. $Q1 + Q2 = 2 \text{ median} > 0$

Page 111

ok

C. $Q1 + Q2 = 2 \text{ median} = 0$

D. None of these

Question No:308

(Marks:1)

Vu-Topper RM

By using method of least square in a linear regression, sum of square of the vertical distances between the points and fitted line is always:

A. Zero

B. Minimum

Page 124

ok

C. Maximum

D. All of these

Question No:309

(Marks:1)

Vu-Topper RM

If the outcome of one event affects the outcome of another, then the events are said to be:

A. Dependent Events

Page 162

B. Mutually Exclusive

C. Independent Events

D. Not Mutually Exclusive Events

Question No:310

(Marks:1)

Vu-Topper RM

Quartile Deviation is based on:

A. All values

B. Not all values

Page 84

C. Extreme Values

D. Smallest values

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:311

(Marks:1)

Vu-Topper RM

In a Box and Whisker plot, a line which divides the box into two equal parts is referred to as:

A. Mean

B. Median

Page 107

C. Mode

D. Range

Question No:312

(Marks:1)

Vu-Topper RM

Data in the Population census report is:

A. Grouped Data

B. Secondary Data

C. Primary Data

D. Array Data

Question No:313

(Marks:1)

Vu-Topper RM

For a symmetrical distribution, b_1 is always:

A. Less than 1

B. Greater than 1

C. Equal to 0

Page 119

D. Less or equal to 1

Question No:314

(Marks:1)

Vu-Topper RM

When we smooth a frequency polygon, it becomes:

A. OGIVE

B. Pie chart

C. Bar chart

D. Frequency Curve

Page 37

Question No:315

(Marks:1)

Vu-Topper RM

The suitable digram to represent the data relating to the monthly expenditure on different items by a family is:

A. Historigram

بري صحبت سے تنہائي بہتر ہے اور تنہائي سے نيک صحبت بہتر ہے

For More Help Contact What's app 03224021365

- B. Histogram
- C. Multiple bar diagram
- D. Pie diagram**

Question No:316

(Marks:1)

Vu-Topper RM

When data are classified according to a single characteristic, it is called:

- A. (Quantitative classification
- B. Qualitative classification
- C. Area classification
- D. Simple classification**

Question No:317

(Marks:1)

Vu-Topper RM

Classification of data by attributes is called:

- A. Quantitative classification
- B. Chronological classification
- C. Qualitative classification**
- D. Geographical classification

Question No:318

(Marks:1)

Vu-Topper RM

Classification of data according to location or areas is called:

- A. Qualitative classification
- B. Quantitative classification
- C. Geographical classification**
- D. Chronological classification

Question No:319

(Marks:1)

Vu-Topper RM

Classification is applicable in case of:

- A. Normal characters
- B. Quantitative characters
- C. Qualitative characters
- D. Both (b) and (c)**

Question No:320

(Marks:1)

Vu-Topper RM

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

In classification, the data are arranged according to:

- A. Similarities**
- B. Differences
- C. Percentages
- D. Ratios

Question No:321

(Marks:1)

Vu-Topper RM

When data are arranged at regular interval of time, the classification is called:

- A. Qualitative
- B. Quantitative
- C. Chronological**
- D. Geographical

Question No:322

(Marks:1)

Vu-Topper RM

When an attribute has more than three levels it is called:

- A. Manifold-division**
- B. Dichotomy
- C. One-way
- D. Bivariate

Question No:323

(Marks:1)

Vu-Topper RM

The number of tally sheet count for each value or a group is called:

- A. Class limit
- B. Class width
- C. Class boundary
- D. Frequency**

Question No:324

(Marks:1)

Vu-Topper RM

The frequency distribution according to individual variate values is called:

- A. Discrete frequency distribution**
- B. Cumulative frequency distribution

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

For More Help Contact What's app 03224021365

- C. Percentage frequency distribution
- D. Continuous frequency distribution

Question No:325

(Marks:1)

Vu-Topper RM

A series arranged according to each and every item is known as:

- A. Discrete series
- B. Continuous series
- C. Individual series**
- D. Time series

Question No:326

(Marks:1)

Vu-Topper RM

The largest and the smallest values of any given class of a frequency distribution are called:

- A. Class Intervals
- B. Class marks
- C. Class boundaries
- D. Class limits**

Question No:327

(Marks:1)

Vu-Topper RM

If there are no gaps between consecutive classes, the limits are called:

- A. Class limits
- B. Class boundaries**
- C. Class intervals
- D. Class marks

Question No:328

(Marks:1)

Vu-Topper RM

Class boundaries are also called:

- A. Mathematical limits**
- B. Arithmetic limits
- C. Geometric limits
- D. Qualitative limits

Question No:329

(Marks:1)

Vu-Topper RM

بري صحبت سے تنہائي بہتر ہے اور تنہائي سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

The average of lower- and upper-class limits is called:

- A. Class boundary
- B. Class frequency
- C. Class mark**
- D. Class limit

Question No:330

(Marks:1)

Vu-Topper RM

The lower- and upper-class limits are 20 and 30, the midpoints of the class is:

- A. 20
- B. 25**
- C. 30
- D. 50

Question No:331

(Marks:1)

Vu-Topper RM

A frequency distribution that contains a class with limits of "10 and under 20" would have a midpoint:

- A. 10
- B. 14.9
- C. 15**
- D. 20

Question No:332

(Marks:1)

Vu-Topper RM

If the number of workers in a factory is 128 and maximum and minimum hourly wages are 100 and 20 respectively. For the frequency distribution of hourly wages, the class interval is:

- A. 8
- B. 9
- C. 10**
- D. 80

Question No:333

(Marks:1)

Vu-Topper RM

Total angle of the pie-chart is:

بري صحبت سے تنہائي بہتر ہے اور تنہائي سے نيک صحبت بہتر ہے

For More Help Contact What's app 03224021365

- A. 45
- B. 90
- C. 180
- D. 360**

Question No:334

(Marks:1)

Vu-Topper RM

Cumulative frequency polygon can be used for the calculation of:

- A. Mean
- B. Median**
- C. Mode
- D. Geometric mean

Question No:335

(Marks:1)

Vu-Topper RM

The amount of hump of a distribution is called:

- A. Kurtosis**
- B. Symmetry
- C. Dispersion
- D. Skewness

Page 109 ok

Question No:336

(Marks:1)

Vu-Topper RM

When there is no correlation then:

- A. $R=2$
- B. $R=0$**
- C. $R=1$
- D. $R=-1$

Google ok

Question No:337

(Marks:1)

Vu-Topper RM

Skewness is based on quartiles. It indicates that in a symmetrical distribution first and third quartiles are equi-distant from the median

Bowley's Coefficient Google

Question No:338

(Marks:1)

Vu-Topper RM

Formula for Co-efficient of Quartile Deviation is:

$Q_3 - Q_1 / Q_3 + Q_1$ Google

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:339

(Marks:1)

Vu-Topper RM

In simple linear regression, which of the following statements indicates there is no linear relationship between the variables x and y ?

- A. none of the above
- B. sum of square of errors is 0
- C. Coefficient of determination is -1.0

D. Coefficient of correlation is 0

Google

ok

Question No:340

(Marks:1)

Vu-Topper RM

Quartile deviation is used as a measure of dispersion when we use _____ as a measure of central tendency

Mean

Question No:341

(Marks:1)

Vu-Topper RM

The measure of dispersion which uses only two observations is called:

Range

Google

Question No:342

(Marks:1)

Vu-Topper RM

For a symmetrical data set mean value is 150 and standard deviation 25. 95% values will lie between

125,175

Question No:343

(Marks:1)

Vu-Topper RM

The suitable average for the qualitative data is:

Median

Google

Question No:344

(Marks:1)

Vu-Topper RM

Geometric Mean gives the equal weightage to _____ values

- A. Larger Values
- B. Smaller Values

C. All the

observations

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

For More Help Contact What's app 03224021365

Google

D. Both smaller and larger values

Question No:345

(Marks:1)

Vu-Topper RM

Which average gives the more weightage to the smaller values?

Harmonic mean

Google

Question No:346

(Marks:1)

Vu-Topper RM

In a right skewed distribution,

A. $X_m - Q_3$ less than $Q_1 - X_0$

B. $X_m - Q_3$ grater than $Q_1 - X_0$

ok

C. $X_m - Q_3$ less and equal to $Q_1 - X_0$

D. $X_m - Q_3$ greater and equal to $Q_1 - X_0$

Question No:347

(Marks:1)

Vu-Topper RM

Moment ratio b_1 is used to measure:

Mean Google

Question No:348

(Marks:1)

Vu-Topper RM

In a Box and Whisker plot, Box is divided into:

A. One port

B. Two ports

ok

C. Four ports

D. Three ports

Question No:349

(Marks:1)

Vu-Topper RM

Which of the rule is applied to the frequency distribution that is mound-shaped and symmetric?

A. Empirical rule

Google

ok

B. Combination rule

C. Chebychev's rule

D. Combination rule

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:350

(Marks:1)

Vu-Topper RM

The variance of a sample of 81 observations is equal to 64; The standard deviation of these observations will be:

A. 8 Google ok

B. 256

C. 4096

D. 6561

Question No:351

(Marks:1)

Vu-Topper RM

The measures used to calculate the variation present among the observations in the unit of the variable is called:

Relative measures of dispersion

Google

Question No:352

(Marks:1)

Vu-Topper RM

Standard deviation is calculated from the Harmonic Mean.

A. Never Google

B. Always

C. Sometimes

D. None of these

Question No:353

(Marks:1)

Vu-Topper RM

Larger the quartile deviation:

Greater is the scatter of values

Question No:354

(Marks:1)

Vu-Topper RM

Given the N values in a series, the geometric mean is:

A. The third root of the product of N values.

B. The fourth root of the product of N values.

C. The square root of the product of N values.

D. The Nth root of the product of N positive values.

ok

Question No:355

(Marks:1)

Vu-Topper RM

Harmonic mean is particularly useful in computing

بري صحبت سے تنہائي بہتر ہے اور تنہائي سے نيک صحبت بہتر ہے

For More Help Contact What's app 03224021365

Average rate

Question No:356

(Marks:1)

Vu-Topper RM

Co-efficient of Quartile Deviation is a:

Pure Number

Question No:357

(Marks:1)

Vu-Topper RM

The measures used to calculate the variation present among the observations relative to their average is called:

Coefficient of kurtosis

Google

Question No:358

(Marks:1)

Vu-Topper RM

Which of the following measures based on all observations?

A. Standard deviation

B. Mean deviation

C. None of these

D. Mean deviation and Standard deviation

Question No:359

(Marks:1)

Vu-Topper RM

Quartile Deviation measures the spread of data around:

A. Mode

B. Median

Google

C. Arithmetic Mean

D. Geometric Mean

Question No:360

(Marks:1)

Vu-Topper RM

In a linear regression, $Y=a+bX$, the variable "X" will always:

A. A random variable

B. Qualitative variable

C. Quantitative variable

ok

D. A non-random variable

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:361

(Marks:1)

Vu-Topper RM

The variable plotted on the horizontal or X-axis in a scatter diagram, is called

- A. Scatter variable
- B. Dependent variable
- C. Correlation variable

D. independent variable.

Google

ok

Question No:362

(Marks:1)

Vu-Topper RM

When the curve is flat-topped, it is called:

- A. Hetrokurtic
- B. Leptokurtic
- C. Mesokurtic

D. Platykurt

Google

ok

Question No:363

(Marks:1)

Vu-Topper RM

Given the least squares regression line $\hat{Y} = 5 - 2x$:

The relationship between x and y is positive

Question No:364

(Marks:1)

Vu-Topper RM

The mean of a distribution is 24, the mode is 25 and the standard deviation is 5, then the coefficient of skewness will be:

A. Equal to zero

B. Less than zero

ok

C. Greater than zero

D. None of the above

Question No:365

(Marks:1)

Vu-Topper RM

Third moment about mean provides information about the _____ of the distribution.

- A. Centre
- B. Kurtosis
- C. Dispersion

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

D. Symmetry

ok

Question No:366

(Marks:1)

Vu-Topper RM

Relative measure of dispersion corresponding to mean deviation is:

Co-efficient of Mean Deviation

Question No:367

(Marks:1)

Vu-Topper RM

If any of the value in the data set is negative then it is impossible to compute:

A. Arithmetic Mean

B. Harmonic Mean

C. Geometric Mean

D. None of the these

Question No:368

(Marks:1)

Vu-Topper RM

Formula for quartile deviation is:

Q.D= $q_3 - q_1 / 2$

Question No:369

(Marks:1)

Vu-Topper RM

Relative measures of dispersion can be used for:

Comparison of two data sets

Question No:370

(Marks:1)

Vu-Topper RM

Empirical rule is considered when the data is

A. Skewed

B. Symmetrical

C. Positively skewed

D. Negatively skewed

Page 89

ok

Question No:371

(Marks:1)

Vu-Topper RM

The standard deviation is always _____ than mean deviation.

Greater

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:372

(Marks:1)

Vu-Topper RM

A five-number summary consists of:

- A. X_m , Q1, Mode, Q3, and X_0
- B. X_m , Q1, Mean, Q3, and X_0
- C. X_0 , Q1, Median, Q2, and X_m
- D. X_0 , Q1, Median, Q3, and X_m**

Page 92

ok

Question No:373

(Marks:1)

Vu-Topper RM

If a distribution has negative skewness, in what order (lowest to highest) will the averages be?

Mean, median, mode

Question No:374

(Marks:1)

Vu-Topper RM

If any of the value in data set is zero then it is not possible to compute

- A. Mode
- B. Median
- C. Mean

D. Harmonic Mean

Question No:375

(Marks:1)

Vu-Topper RM

Measure of dispersion is used to measure:

- A. Interdependence between variables**
- B. How much values are clustered around mean
- C. How much values are disperse from the mean
- D. Dependence of one variable upon other variable

Question No:376

(Marks:1)

Vu-Topper RM

Range i.e. (maximum value - minimum value) for a symmetrical distribution is approximately equal to

- A. σ
- B. 3σ
- C. 4σ

D. 6σ

ok

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:377

(Marks:1)

Vu-Topper RM

If all the points in a scatter diagram lie on the least squares regression line, then the coefficient of correlation:

- A. 1
- B. 0
- C. -1

D. -1to+1

ok

Question No:378

(Marks:1)

Vu-Topper RM

b² is used to measure the:

- A. None
- B. Both
- C. Skewness of the distribution

D. Kurtosis of the distribution

Page 114

ok

Question No:379

(Marks:1)

Vu-Topper RM

Mean Deviation, Variance and Standard Deviation of the values 4, 4, 4, 4, 4 is

4

Question No:380

(Marks:1)

Vu-Topper RM

For a particular data set the Pearson's coefficient of skewness is less than zero. What will be the shape of distribution?

- A. Symmetrical
- B. Positively Skewed

C. Negatively Skewed

ok

D. None of the above

Question No:381

(Marks:1)

Vu-Topper RM

Which one of the following is a meso-kurtic curve?

- A. Negatively skewed
- B. Positively skewed
- C. J-

shaped

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:382

(Marks:1)

Vu-Topper RM

Which method is used for obtaining the relative frequencies?

Dividing the frequency by total number of frequencies

Question No:383

(Marks:1)

Vu-Topper RM

Relative dispersion is expressed in terms of:

Ratio

Question No:384

(Marks:1)

Vu-Topper RM

To find the average speed of a journey which is the appropriate measure of central tendency.

A. Mode

B. Weighted Mean

ok

C. Geometric Mean

D. Harmonic mean

Question No:385

(Marks:1)

Vu-Topper RM

When a researcher want to compare intensity of symptoms when different doses are administered. In this case," different doses" will be treated as:

A. Qualitative variable

B. Continuous variable

C. Dependent variable

D. Independent variable

ok

Question No:386

(Marks:1)

Vu-Topper RM

For the given data 20, 13, 27, 0, -8 G. M will be:

A. Negative

B. Positive

C. Zero

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

D. Undefined

Question No:387

(Marks:1)

Vu-Topper RM

For the given data 2, 3, 7, 0, -8 G. M will be:

A. Negative

B. Positive

C. Zero

Page 75

D. Undefined

Question No:388

(Marks:1)

Vu-Topper RM

Historiogram and histogram are:

A. Always same

B. Not same

ok

C. Off and on same

D. Randomly same

Question No:389

(Marks:1)

Vu-Topper RM

Relationship among the averages

A. $HM \geq GM \geq AM$

B. $AM \geq GM \geq HM$

C. $GM \geq HM \geq AM$

D. $AM \geq HM \geq GM$

Question No:390

(Marks:1)

Vu-Topper RM

In a group frequency distribution, corrected moments are calculated by the method of:

A. Sheppard's correction

B. Continuity correction

C. Pearson's correction

ok

D. Bowley's correction

Question No:391

(Marks:1)

Vu-Topper RM

بري صحبت سے تنہائي بہتر ہے اور تنہائي سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

A tabular arrangement for classifying data into different groups is called:

- A. Standard deviation
- B. Frequency distribution**
- C. Class
- D. Arithmetic Mean

Question No:392

(Marks:1)

Vu-Topper RM

Pearson's coefficient of skewness is equal to:

mean – mode / standard deviation

Page 104

Question No:393

(Marks:1)

Vu-Topper RM

r is a pure number that lies between:

- A. None
- B. 0 to ∞
- C. 0 to 1
- D. -1 and +1**

Google

ok

Question No:394

(Marks:1)

Vu-Topper RM

When A and B are two non-empty and mutually exclusive events, then:

- A. $P(A \cap B) = P(A) + P(B)$
- B. $P(A \cup B) = P(A) + P(B)$**
- C. $P(A \cup B) = P(A) \cdot P(B)$
- D. $P(A \cap B) = P(A) \cdot P(B)$

Question No:395

(Marks:1)

Vu-Topper RM

If a fair coin is tossed 2 times then probability that two heads appear is equal to:

- A. 1/3
- B. 1/2

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

C. 1/5

D. 1/4

ok

Question No:396

(Marks:1)

Vu-Topper RM

If one card is selected at random from a deck of 52 playing cards, what is the probability that the card is a club or a face card or both?

A. 22/52

ok

B. 52/22

C. 21/52

D. 20/52

Question No:397

(Marks:1)

Vu-Topper RM

In how many different ways can the letters of the word 'OPTICAL' be arranged so that the vowels always come together?

A. 720

B. 2450

C. 1560

D. 950

Question No:398

(Marks:1)

Vu-Topper RM

In how many different ways can the letters of the word 'CORPORATION' be arranged so that the vowels always come together?

A. 720

B. 640

C. 50400

D. 970

Question No:399

(Marks:1)

Vu-Topper RM

The probability of a jack card from a well shuffled pack of 52 playing cards will be:

A. 13/52

B. 4/52

ok

بري صحبت سے تنہائي بہتر ہے اور تنہائي سے نيک صحبت بہتر ہے

For More Help Contact What's app 03224021365

- C. $1/52$
D. $1/26$

Question No:400

(Marks:1)

Vu-Topper RM

If $A = \{10, 11\}$ then which of the following is power set of A?

- A. $\{\{\}, \{10, 11\}\}$
B. $\{\{10\}, \{11\}, \{10, 11\}\}$
C. $\{\{\}, \{10\}, \{11\}, \{10, 11\}\}$ ok
D. $\{\{10\}, \{11\}, \{10, 11\}, \{11, 10\}\}$

Question No:401

(Marks:1)

Vu-Topper RM

10! equals to:

- A. 362800
B. 3628800 ok
C. 362280
D. 3622880

Question No:402

(Marks:1)

Vu-Topper RM

In a multiplication theorem P (A and B) equals (when events are not independent):

- A. $P(A) P(B)$
B. $P(A) + P(B)$
C. $P(A) * P(A|B)$ ok
D. $P(A) + P(B) - P(A \cup B)$

Question No:403

(Marks:1)

Vu-Topper RM

In a drawer there are 5 black socks and 3 green socks. Two socks are picked randomly one after the other without replacement. What is the possibility that both the socks are black?

- A. $5/14$** ok
B. $5/8$
C. $8/5$
D. $3/16$

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:404

(Marks:1)

Vu-Topper RM

The probability of an event always lies between:

A. 0 and ∞

B. 0 and 1 ok

C. 1 and + 1

D. $-\infty$ and $+\infty$

Question No:405

(Marks:1)

Vu-Topper RM

Let A and B are two independent events. If $P(A \cap B) = 0.12$ and $P(A) = 0.3$ then find $P(B)$.

A. 0.4

B. 0.3

C. 0.2

D. 0.1

Question No:406

(Marks:1)

Vu-Topper RM

A set that contains all possible outcomes of a system is known as:

A. Finite set

B. Infinite set

C. Universal set ok

D. None of these

Question No:407

(Marks:1)

Vu-Topper RM

In a probability distribution, the sum of the probabilities is equal to:

A. 0

B. 1 ok

C. 0.1

D. 1.5

Question No:408

(Marks:1)

Vu-Topper RM

If $P(A|B) = 0.3$ and $P(B) = 0.8$, then:

A. $P(A) = 0.24$

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

- B. $P(B|A)=0.7$
- C. $P(A \cup B)=0.5$
- D. $P(A \cap B)=0.24$

Question No:409

(Marks:1)

Vu-Topper RM

A bag contains 10 white, 2 orange and 3 black balls. What is the probability of black balls?

- A. $2/15$
- B. $3/15$**
- C. $5/15$
- D. $10/15$

Question No:410

(Marks:1)

Vu-Topper RM

When an event is as likely to occur as other, it is called:

- A. Normal event
- B. Equally likely event** **ok**
- C. Mutually exclusive event
- D. Not mutually exclusive event

Question No:411

(Marks:1)

Vu-Topper RM

Two dice are tossed. The probability that the total score is a prime number is:

- A. $7/9$
- B. $1/6$**
- C. $1/2$
- D. $5/12$

Question No:412

(Marks:1)

Vu-Topper RM

If a player well shuffles the pack of 52 playing cards, then the probability of a black card from 52 playing cards is:

- A. $1/2$** **ok**
- B. $1/13$

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

- C. $1/52$
- D. $13/52$

Question No:413

(Marks:1)

Vu-Topper RM

Find the number of subsets of the following set. $\{x \mid x \text{ is a day of the week}\}$

- A. 7
- B. 14
- C. 49
- D. 128**

ok

Question No:414

(Marks:1)

Vu-Topper RM

If we roll a die four times, then possible outcomes of the sample space are:

- A. 6
- B. 36
- C. 216
- D. 1296**

Question No:415

(Marks:1)

Vu-Topper RM

Probability of an impossible event is always:

- A. Zero**
- B. Less than one
- C. Greater than one
- D. Between one and zero

ok

Question No:416

(Marks:1)

Vu-Topper RM

If we roll three fair dice then the total number of outcomes are:

- A. 18
- B. 36
- C. 216**
- D. 1296

بري صحبت سے تنہائي بہتر ہے اور تنہائي سے نيک صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:417

(Marks:1)

Vu-Topper RM

If we roll a die then probability of getting a '2' will be

A. 7/9

B. 1/6 ok

C. 1/2

D. 5/12

Question No:418

(Marks:1)

Vu-Topper RM

In which approach to probability the outcomes are equally likely to occur?

A. Relative Frequency

B. Classical Probability ok

C. Objective Probability

D. Subjective Probability

Question No:419

(Marks:1)

Vu-Topper RM

Which of the following pairs of A and B events are mutually exclusive?

A. The numbers above 100;

B. The numbers less than -200

C. Both

D. None of these

Question No:420

(Marks:1)

Vu-Topper RM

Total number of words formed by 2 vowels and 3 consonants taken from 4 vowels and 5 consonants is equal to

A. 60

B. 120

C. 520

D. 720 ok

Question No:421

(Marks:1)

Vu-Topper RM

If we have three events A, B and C, then for exhaustive events $P(A \cup B \cup C)$ will be equal to:

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

- A. $P(S)$** **ok**
B. $P(B)$
C. $P(A)$
D. $P(A) * P(B) * P(C)$

Question No:422

(Marks:1)

Vu-Topper RM

A club consists of four members. How many ways are there of selecting three officers: president, secretary and treasurer?

- A. 3
B. 6
C. 20
D. 24 **ok**

Question No:423

(Marks:1)

Vu-Topper RM

From a pack of 52 cards, two cards are drawn together at random. What is the probability of both the cards being kings?

- A. $1/122$
B. $1/15$
C. $25/57$
D. $1/221$ **ok**

Question No:424

(Marks:1)

Vu-Topper RM

A letter is chosen at random from the word STATISTICS. The probability of getting a vowel is:

- A. $3/10$** **ok**
B. $4/10$
C. $5/10$
D. $6/10$

Question No:425

(Marks:1)

Vu-Topper RM

$P(\text{event}) = \text{No. of favorable outcomes} / \text{total no. of outcomes}$ is a definition of:

- A. Normal event** **ok**

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

- B. Binomial event
- C. Objective Approach
- D. Subjective Approach

Question No:426

(Marks:1)

Vu-Topper RM

When a coin is tossed, then sample space consists of:

- A. 2 outcomes** ok
- B. 4 outcomes
- C. 6 outcomes
- D. 8 outcomes

Question No:427

(Marks:1)

Vu-Topper RM

If a player selected a card from a well shuffles the pack of 52 playing cards then the probability of number 10 card is equals to:

- A. $1/52$
- B. $10/52$
- C. $4/52$** ok
- D. $26/52$

Question No:428

(Marks:1)

Vu-Topper RM

If $P(A) = 0.7$ and $P(B) = 0.2$ then find $P(A \cup B)$ where A and B are mutually exclusive events.

- A. 0.1
- B. 0.5
- C. 0.9**
- D. 0.14

Question No:429

(Marks:1)

Vu-Topper RM

For the independent events C and D, if $P(C) = 0.25$ and $P(D) = 0.40$ then $P(C \text{ and } D) = ?$

- A. 0.1**
- B. 0.15
- C. 0.50

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

D. 0.65

Question No:430

(Marks:1)

Vu-Topper RM

If A, B and C are three events and only one event of them must occur then $P(A) + P(B) + P(C)$:

A. 0

B. 1 ok

C. $P(S)=1$

D. sample space

Question No:431

(Marks:1)

Vu-Topper RM

If two fair dice are thrown, the probability of getting a double six is:

A. $1/6$

B. $1/12$

C. $1/36$ ok

D. $2/36$

Question No:432

(Marks:1)

Vu-Topper RM

In a Venn diagram, the overlap between two circles represents:

A. The union of two sets

B. The intersection of two sets ok

C. The elements that are in either of two sets

D. The difference between the number of elements in two sets

Question No:433

(Marks:1)

Vu-Topper RM

A bag contains 3 red balls, 4 green balls, and 5 blue balls. One ball is taken from the bag and then replaced. Another ball is taken from the bag. What is the probability that the first ball is red and the second ball is blue?

A. $6/72$

B. $5/48$

C. $10/90$

D. $12/98$

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

Question No:434

(Marks:1)

Vu-Topper RM

From a group of 7 men and 6 women, five persons are to be selected to form a committee so that at least 3 men are there on the committee. In how many ways can it be done?

A. 645

B. 564

C. 756 ok

D. 865

Question No:435

(Marks:1)

Vu-Topper RM

If we want to calculate average speed of a car then which of the following average will be used?

A. Mode

B. Mean

C. Harmonic mean

D. Geometric mean

Question No:436

(Marks:1)

Vu-Topper RM

5C5 is equal to

A. 1

B. 5

C. 10

D. 24

Question No:437

(Marks:1)

Vu-Topper RM

Harmonic mean is better than other means if the data are for

A. Speed or rates

B. Heights or lengths

C. Ratios or proportions

D. Binary values like 0 & 1

Question No:438

(Marks:1)

Vu-Topper RM

The reciprocal of the arithmetic mean of the reciprocals of the values is

بري صحبت سے تنہائي بہتر ہے اور تنہائي سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

defined as:

- A. weighted mean
- B. weighted mean
- C. harmonic mean**
- D. geometric mean

Google

Question No:439

(Marks:1)

Vu-Topper RM

Second moment about mean will be equal to:

- A. 0
- B. Mean
- C. Median
- D. Variance**

Google

ok

Question No:440

(Marks:1)

Vu-Topper RM

For a positively skewed distribution mid range is To/than the mid quartile range

- A. Less
- B. Equal
- C. Greater**
- D. None of the above

Google

ok

Question No:441

(Marks:1)

Vu-Topper RM

Sum of the deviations taken from the mean is equal to:

- A. One
- B. Zero**
- C. Postive number
- D. Negative Number

Google

ok

Visit My YouTube Channel

For More Important Notes
Channel Name = #VuTopperRM

VuTopperRM

برى صحت سے تٲهائى بهتر هے اور تٲهائى سے نيك صحت بهتر هے

For More Help Contact What's app 03224021365