Download Solved Mid/Final Term
Past Papers by Waqar Siddhu
in PDF form
VUNET.BLOGSPOT.COM



Facebook.com/vunets



Question No : 41 of 52 Marks: 2 (Budgeted Time 4 Min)

Is there any difference between distribution function and cumulative distribution function of discrete r.v. X?

Answer (Please click here to Add Answer)

VuAnswers.com

The distribution function of a random variable X, denoted by F(x), is defined by F(x) = P(X < x).
The function F(x) gives the probability of the event that X takes a value LESS THAN OR EQUAL TO a specified value x. The distribution function is abbreviated to *d.f.* and is also called the *cumulative distribution function (cdf)* as it is the cumulative probability function of the random variable X from the smallest value upto a *specific* value x.
Let us illustrate this concept with the help of the same example that we have been considering --- that of the probability distribution of the sum of the dots when two fair dice are thrown.

Made by: Waqar Siddhu

Question No : 42 of 52 Marks: 2 (Budgeted Time 4 Min)

What is meant by sampling distribution of sample proportion?

Answer (Please click here to Add Answer)

VuAnswers.com

The probability distribution of any statistic (such as the mean, the standard deviation, the proportion of successes in a sample, etc.) is known as its sampling distribution.

The sample proportion has different values in different samples.


Made by: Waqar Siddhu

Question No : 43 of 52 Marks: 2 (Budgeted Time 4 Min)

If an automobile is driven on the average no more than 16000 Km per year then formulate the null and alternative hypothesis.

Answer (Please click here to Add Answer)

VuAnswers.com




Made by: Waqar Siddhu

Question No : 44 of 52 Marks: 2 (Budgeted Time 4 Min)

While applying the chi-square goodness of fit test, if an expected frequency in any category is less than 5, then what we do to solve this problem.

Answer (Please click here to Add Answer)

VuAnswers.com




Made by: Waqar Siddhu

Question No : 45 of 52 Marks: 3 (Budgeted Time 6 Min)

Write a short note on Quota Sampling.

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

Question No : 46 of 52 Marks: 3 (Budgeted Time 6 Min)

Write down the properties of hypergeometric distribution.

Answer (Please click here to Add Answer)

VuAnswers.com

The probability of success changes on each trial.

· The successive trials are not independent.

· The experiment is repeated a fixed number of times.

Made by: Waqar Siddhu

Question No : 47 of 52 Marks: 3 (Budgeted Time 6 Min)


Write down critical region for the following hypothesis at 5% level of significance.

$H_0 : \mu = 75$

$H_1 : \mu < 75$

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu


Question No : 48 of 52 Marks: 3 (Budgeted Time 6 Min)

For the following table we want to test the independence of smoking pattern and marital status. What will be the degrees of freedom for the chi-square test of independence?

Marital Status	Smoking Pattern			Total
	Total self-restraint	Only at times	Regular Smoker	
Single	67	213	74	354

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

Married	411	633	129	1173
Widowed	85	51	7	143
Divorced	27	60	15	102
Total	590	957	225	1772

Answer (Please click here to Add Answer)

VuAnswers.com

Made by: Waqar Siddhu


Question No : 49 of 52 Marks: 5 (Budgeted Time 10 Min)

Answer the following question using the given stem and leaf plot of the ages of the people who attended a musical event.

Stem	Leaf
1	2 2 3 3 4 4 5 5 5 5 5 5 6 6 6 6 6 6 7 7 7 7 7 8 8 9 9
2	3 3 4 8
3	6

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu


Question No : 49 of 52 Marks: 5 (Budgeted Time 10 Min)

2	3 3 4 8
3	6
4	5 8

a) How many people attended the event?
b) What is the age of youngest attendee?

Answer (Please click here to Add Answer)

VuAnswers.com




Made by: Waqar Siddhu

Question No : 49 of 52 Marks: 5 (Budgeted Time 10 Min)

a) How many people attended the event?
b) What is the age of youngest attendee?
c) What was the age of oldest person attending event?
d) Which age group was more widely represented?
e) How many people were above 30?

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

Question No : 50 of 52 Marks: 5 (Budgeted Time 10 Min)

If $\bar{X}$ is normally distributed with $\mu_{\bar{X}} = \mu = 24000$ and $\sigma_{\bar{X}} = 412.20$

then find out the $P(\bar{X} > 24500)$.

Answer (Please click here to Add Answer)

VuAnswers.com

Made by: Waqar Siddhu

Question No : 51 of 52 Marks: 5 (Budgeted Time 10 Min)

If we have $\bar{X} = 15$ and $\sum_{i=1}^{10}(X_i - \bar{X})^2 = 56$

Then, find the 90% confidence interval for population variance (σ^2).

Answer (Please click here to Add Answer)

VuAnswers.com


Made by: Waqar Siddhu

Question No : 52 of 52 Marks: 5 (Budgeted Time 10 Min)

A research worker wishes to estimate the mean of a population using a sample sufficiently large that the probability will be 0.95 that the sample mean will not differ from the true mean by more than 25 percent of the standard deviation. How large a sample should be taken?

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

What does quartile deviation measure in a distribution?

Answer (Please click here to Add Answer)

VuAnswers.com

Normal | Arial | 12 | 100%

Made by: Waqar Siddhu

Start Time: 9:48 AM

In a random sample of 1000 women in a locality, 224 women said that they use Aerial washing powder. What is the point estimator and point estimate of the proportion of the women who use Aerial washing powder?

Answer (Please click here to Add Answer)

VuAnswers.com

Normal Arial 12 100%

Made by: Waqar Siddhu

Start Time: 9:48 AM

What is the impact of level of confidence (1-α) on the value of $Z_{\alpha/2}$?

Answer (Please click here to Add Answer)

VuAnswers.com

Normal Arial 12 100%

Made by: Waqar Siddhu

Formulate the hypothesis in case of goodness of fit test.

Answer (Please click here to Add Answer)

VuAnswers.com

Normal | Arial | 12 | 100%

Made by: Waqar Siddhu

Start Time: 9:48 AM

44

Why the median is suitable average for the below data set? Explain.

Monthly income (in rupees)	No. of workers
Less than 2000/-	100
2000/-to 2999/-	300

Answer (Please click here to Add Answer)

VuAnswers.com

Made by: Waqar Siddhu

(in rupees)	
Less than 2000/-	100
2000/-to 2999/-	300
3000/- to 3999/-	500
4000/- to 4999/-	250
5000/- and above	50

Answer (Please click here to Add Answer)

VuAnswers.com

Made by: Waqar Siddhu

If Z is a standard normal random variable with mean 0 and variance 1, then find the Lower quartile.

Answer (Please click here to Add Answer)

VuAnswers.com

Normal | Arial | 12 | 100%

Made by: Waqar Siddhu

Write down critical region for the following hypothesis at 5% level of significance.

$H_0 : \mu = 75$

$H_1 : \mu < 75$

Answer (Please click here to Add Answer)

VuAnswers.com

Normal Arial 12 100%

Made by: Waqar Siddhu

Start Time: 9:48 AM

47

The manager of a bottling plant is anxious to reduce the variability in net weight of fruit bottled. Over a long period, the standard deviation has been 15.2 gm. A new machine is introduced and the net weights (in grams) in 10 randomly selected bottles (all of the same nominal weight) are 987, 966, 955, 977, 981, 967, 975, 980, 953, and 972.
State null and alternative hypothesis that machine has a better performance? Also write down test-statistics about the hypothesis.

Answer (Please click here to Add Answer)

VuAnswers.com

Normal Arial 12 100%

Made by: Waqar Siddhu

Start Time: 9:48 AM

Find the coefficient of standard deviation from the following data:
Life in Hours (X): 130, 150, 180, 250, 345

Answer (Please click here to Add Answer)

VuAnswers.com

Normal Arial 12 100%

Made by: Waqar Siddhu

Start Time: 9:48 AM

What is the probability that a poker hand of 5 cards contain (i) exactly 2 aces (ii) exactly 1 ace?
(Use hypergeometric distribution)

Answer (Please click here to Add Answer)

VuAnswers.com

Normal Arial 12 100%

Made by: Waqar Siddhu

Start Time: 9:48 AM

If n=22, $\sum\left(d-\bar{d}\right)^2 = 270, \bar{d} = 4$

Then, using a 0.05 level of significance level, test the hypothesis $H_0 : \mu_d = 0$ against $H_1 : \mu_d \neq 0$

Answer (Please click here to Add Answer)

VuAnswers.com

Made by: Waqar Siddhu

A company launched new layout of its website. After a survey, 62 of 115 visitors liked the new layout while 59 of 135 visitors liked the old layout of the website. Company claims that new layout did not improve the visitor's liking about the website.

By using the critical value $Z < -Z_{0.05} = -1.645$, verify the company's claim:

$H_o: P_1 - P_2 \geq 0$

$H_1: P_1 - P_2 < 0$

Answer (Please click here to Add Answer)

VuAnswers.com

Made by: Waqar Siddhu

Question No : 41 of 52 Marks: 2 (Budgeted Time 4 Min)

Name the measures of dispersion that are not based on all the values.

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

Question No : 42 of 52 Marks: 2 (Budgeted Time 4 Min)

When we use two-tailed test?

Answer (Please click here to Add Answer)

VuAnswers.com

100%

Normal Arial 12

test :
A hypothesis test in which rejection of the null hypothesis occurs for values of the test statistic in either tail of the sampling distribution.

Made by: Waqar Siddhu

Question No : 43 of 52 Marks: 2 (Budgeted Time 4 Min)

For a sample data n = 15, calculate $t_{\frac{\alpha(v)}{2}}$ for $\alpha = 0.10$.

Answer (Please click here to Add Answer)

VuAnswers.com

Made by: Waqar Siddhu

Question No : 44 of 52 Marks: 2 (Budgeted Time 4 Min)

Suppose we want to determine the proportions of smokers and non smoker in a city? In this situation what type of distribution we can use?

Answer (Please click here to Add Answer)

VuAnswers.com

following proportions, we are dealing with a BINOMIAL situation:

- Proportion of smokers in a city smoker → success, non-smokers → failure.
- Proportion of literates in a community → literacy rate, literate → success, illiterate → failure.
- Proportion of males in a city → *sex ratio*).

Made by: Waqar Siddhu

Question No : 45 of 52 Marks: 3 (Budgeted Time 6 Min)

If mean of a distribution is 20 and standard deviation is 2. Find out $\mu \pm 2\sigma$ limits by applying empirical rule. What percent of data will lies between these two limits?

Answer (Please click here to Add Answer)

VuAnswers.com

calculate app keroo hun

According to this empirical rule:

• Approximately 68% of the measurements will fall within 1 standard deviation of the mean, i.e. within the interval (–X – S, –X + S)

• Approximately 95% of the measurements will fall within 2 standard deviations of the mean, i.e. within the interval (–X – 2S, –X + 2S).

• Approximately 100% (practically all) of the measurements will fall within 3 standard deviations of the mean, i.e. within the interval (–X – 3S, –X + 3S).

Made by: Waqar Siddhu

Question No : 46 of 52 Marks: 3 (Budgeted Time 6 Min)

Suppose X is a random variable having Poisson distribution with its parameter value 3, find value of P(X=1).

Answer (Please click here to Add Answer)

VuAnswers.com

The Poisson distribution has only one parameter μ > 0.

μ=3 (X=1)

$$\lim_{\substack{n\to\infty \\ p\to 0}} b(x;n,p) = \frac{e^{-\mu}\mu^{x}}{x!}, \quad x = 0,1,2,\ldots,\infty$$

where e = 2.71828.

& p is 0.05 or less, n is 20 or more,

calculate easly now..



Made by: Waqar Siddhu

Question No : 47 of 52 Marks: 3 (Budgeted Time 6 Min)

If X = 341, n = 634, p_0 = 0.50 then find the z-test statistic for proportion.

Answer (Please click here to Add Answer)

VuAnswers.com

Step 3:
Test statistic:

$$Z = \frac{X \pm \frac{1}{2} - n\,p_0}{\sqrt{n\,p_0(1-p_0)}}$$

Step 4:
Computation:

Here np0 = 634 (0.50) = 317
and X = 341
Hence X > np0 so use X - ½

$$\text{So} \quad Z = \frac{341 - \frac{1}{2} - 317}{\sqrt{634(0.50)(0.50)}} = \frac{23.5}{12.59}$$

$$= 1.87$$

Made by: Waqar Siddhu

Question No : 48 of 52 Marks: 3 (Budgeted Time 6 Min)

A random sample of size n is drawn from normal population with mean 5 and variance σ^2
If s=2.5, $\bar{x}$ =7and t=3, then what is the values of n?

Answer (Please click here to Add Answer)

VuAnswers.com

The test-statistic to be used is

$$t = \frac{\bar{X} - \mu_0}{s/\sqrt{n}}$$

hare mue = 5...

ider uder lai aj ker laiana now. n ko

ans shid 1.92 ai ga aprox 2 i htink not sure :) becoz stat ae :p

Made by: Waqar Siddhu

Question No : 49 of 52 Marks: 5 (Budgeted Time 10 Min)

A random variable X is normally distributed with $\mu = 50$ *and* $\sigma^2 = 25$. Find the probability of X larger than 54.

Answer (Please click here to Add Answer)

VuAnswers.com

The test-statistic to be used is

$$Z = \frac{\bar{X} - \mu_0}{\text{sigma}}$$

hare give sigma sqr, take under root

sigma= 5 mue alsi give and X= 54 we find X>54

(54-50)/5= 4/5

its half ques see these type of ques 70 % slove now.. 30 % see table etc

Made by: Waqar Siddhu

Question No : 50 of 52 Marks: 5 (Budgeted Time 10 Min)

Find the coefficient of variation (C.V) for the following price of a commodity.
Price (X): 8, 13, 18, 23, 30

Answer (Please click here to Add Answer)

VuAnswers.com

COEFFICIENT OF VARIATION

$$C.V. = \frac{S}{\overline{X}} \times 100$$

Shortcut Formula for Ungroup data

$$S = \sqrt{\left\{\frac{\sum x^2}{n} - \left(\frac{\sum x}{n}\right)^2\right\}}$$

mean ka farmula laga laian X bar ki jaga.... =X/n
small n is 5

Made by: Waqar Siddhu

Question No : 51 of 52 Marks: 5 (Budgeted Time 10 Min)

The given data is $n = 1150, x = 450, p = 0.39, H_0 : p_0 = 0.3, \alpha = 0.01$

Test the stated hypothesis. (*Use table value of* $z = \pm 2.58$)

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

Question No : 52 of 52 Marks: 5 (Budgeted Time 10 Min)

Given, $n_1 = n_2 = 16, s_1^2 = 50, s_2^2 = 16$, Construct a 90% confidence interval for the variance ratio σ_1^2 / σ_2^2.

Answer (Please click here to Add Answer)

VuAnswers.com

Normal Arial 12 100%

regards i have no idea abou F test and this ques in the F test...
remmber me in ur pryaer after 3 hr slove 1 paper.. :)

Made by: Waqar Siddhu

Question No : 41 of 52 Marks: 2 (Budgeted Time 4 Min)

Describe the Poisson distribution as the limiting form of the binomial distribution.

Answer (Please click here to Add Answer)

VuAnswers.com




Made by: Waqar Siddhu

Question No : 42 of 52 Marks: 2 (Budgeted Time 4 Min)

Define Disjoint Sets.

Answer (Please click here to Add Answer)

VuAnswers.com



Question No : 43 of 52 Marks: 2 (Budgeted Time 4 Min)

What is acceptance region?

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

Question No : 44 of 52 Marks: 2 (Budgeted Time 4 Min)

If there are K treatments and R rows in a Randomized Complete Block Ddesign then calculate the total number of experimental units used.

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

Question No : 45 of 52 Marks: 3 (Budgeted Time 6 Min)

In how many ways a three-person committee can be formed from a group of ten persons? (Use the formula)

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

Question No : 46 of 52 Marks: 3 (Budgeted Time 6 Min)

From the given data, calculate mean and standard deviation of sampling distribution of mean if the sampling is done with replacement.

$N = 120, n = 64, \mu = 50, \sigma = 2$

Answer (Please click here to Add Answer)

VuAnswers.com

Made by: Waqar Siddhu

Question No : 47 of 52 Marks: 3 (Budgeted Time 6 Min)

Construct 90% confidence interval for the difference in means $\mu_1 - \mu_2$ in case of paired observations, where

$\bar{d} = 1.8$, $s_d = 1.32$, $t_{0.05(9)} = 1.833$

Answer (Please click here to Add Answer)

VuAnswers.com

Made by: Waqar Siddhu

Question No : 48 of 52 Marks: 3 (Budgeted Time 6 Min)

For a data X : $N(\mu,\sigma^2)$. Two unbiased estimators T_1 and T_2 have following variances

$$V(T_1)=\frac{11\sigma^2}{25}$$

$$V(T_2)=\frac{9\sigma^2}{25}$$

Answer (Please click here to Add Answer)

VuAnswers.com

Made by: Waqar Siddhu

Question No : 49 of 52 Marks: 5 (Budgeted Time 10 Min)

A man travels on car from Lahore to karachi on Motor way in 8 stages of equal intervals. The speed of the car in the various stages was observed to be 110, 116, 120, 114, 115,112, 120, 117 kilometers per hour. Find the average speed at which the car travels.

Answer (Please click here to Add Answer)

VuAnswers.com

Made by: Waqar Siddhu

Question No : 50 of 52 Marks: 5 (Budgeted Time 10 Min)

A random sample of size three is drawn without replacement from the population consisting of four numbers 4, 5, 5, 7. Sampling distribution of sample means is calculated as below;

Sample Means $(\bar{X})$	$f(\bar{X})$
14/3	1/4
16/3	2/4
17/3	1/4

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

Question No : 51 of 52 Marks: 5 (Budgeted Time 10 Min)

If $n = 1150, x = 450, p = 0.39, H_0 : p_0 = 0.5, and\ \alpha = 0.05$

Test the stated hypothesis.

(*Use table value of* $z = \pm 1.96$)

Answer (Please click here to Add Answer)

VuAnswers.com

Made by: Waqar Siddhu

Question No : 52 of 52 Marks: 5 (Budgeted Time 10 Min)

A personal manager is interested in trying to determine whether absenteeism is greater on some specific day of the week or not? His records for the past year show the following sample distributions.

Day of the week:	Monday	Tuesday	Wednesday	Thursday	Friday
No. of absentees:	66	57	54	48	75

Test goodness of fit.

Where test statistic $chi-square = \sum \frac{(o_i - e_i)^2}{e_i} = 7.50$

Answer (Please click here to Add Answer)

VuAnswers.com

Made by: Waqar Siddhu

In how many branches, estimation can be divided? Also write down the names of the branches.

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

A Sample has following values.
101, 99, 111, 199, 150, 140
Find out the maximum likelihood estimate of population mean.

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

Write down critical region for the following hypothesis.


$H_0 : \sigma^2 = 20$

$H_1 : \sigma^2 < 20$

Where, $\alpha = 0.01$ and n=10

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

What is the concept of extrapolation in regression analysis?

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

If Z is a standard normal random variable with mean 0 and variance 1, then find the Lower quartile.

Answer (Please click here to Add Answer)

VuAnswers.com

Normal | Arial | 12 | 100%

Made by: Waqar Siddhu

For the following table we want to test the independence of smoking pattern and marital status. What will be the degrees of freedom for the chi-square test of independence?

Marital Status	Smoking Pattern			Total
	Total self-restraint	Only at times	Regular Smoker	
Single	67	213	74	354

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

Find the proportion for the X_1 and X_2, Where X_1 =300 with sample size= n_1 = 400 and X_2 = 200 with sample size = n_2 =300.

Answer (Please click here to Add Answer)

VuAnswers.com


Made by: Waqar Siddhu

Calculate class boundaries from the following data.

Age Group	f
20-29	2
30-39	4
40-49	5

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

A certain type of storage battery lasts on the average 3.0 years, with a standard deviation of 0.5 year. Assuming that the battery lives are normally distributed, find the probability that a given battery will last less than 2.3 years.

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

For the Given information, what is your conclusion in testing the indicated null hypothesis?

$n_1 = 13, n_2 = 41, s_1^2 = 6.3, s_2^2 = 15.6$, $H_0 : \sigma_1^2 = \sigma_2^2$ and $H_1 : \sigma_2^2 > \sigma_1^2$

Answer (Please click here to Add Answer)

VuAnswers.com


Normal Arial 12 100%

Made by: Waqar Siddhu

In a random sample of 200 persons having their lunch at the University cafeteria on meatless day it was observed that 30 percent preferred vegetable dishes.Find 95% confidence interval for p.

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

Answer the following questions:

i. Which method has been used for colleting the data in the following case?
"An investigator collects the information personally from the individuals concerned".

i. Find the sampling error when the population mean is 15.785 and sample mean is 15.6.

ii. Write down any two types of nonrandom sampling

iii. What is the singular of strata?

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

Question No : 41 of 52 Marks: 2 (Budgeted Time 4 Min)

Write down the name of methods/techniques that are used to represent the quantitative discrete data.

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

Question No : 42 of 52 Marks: 2 (Budgeted Time 4 Min)

A coin is tossed 900 times and heads appear 490 times. State the null and alternative hypotheses to show that the coin is unbiased.

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

Question No : 43 of 52 Marks: 2 (Budgeted Time 4 Min)

Write down the names of two types of experimental designs.

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

Question No : 44 of 52 Marks: 2 (Budgeted Time 4 Min)

What are the mean and variance of binomial distribution?

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

Question No : 45 of 52 Marks: 3 (Budgeted Time 6 Min)

Find the probability of drawing a white ball from a bag containing 4 red, 8 black and 3 white balls.

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

Question No : 46 of 52 Marks: 3 (Budgeted Time 6 Min)

Write down the properties of sampling distribution of proportion $\hat{p}$, when sampling is performed *without* replacement.

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

Question No : 47 of 52 Marks: 3 (Budgeted Time 6 Min)

If $X = 255, n = 500, p_0 = 0.60$ then find the z-test statistic for proportion.

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

Question No : 48 of 52 Marks: 3 (Budgeted Time 6 Min)

What is the impact of degrees of freedom on chi- square distribution?

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

Question No : 49 of 52 Marks: 5 (Budgeted Time 10 Min)

Find quartile deviation for the data given below:
18, 53, 45, 28, 39, 29, 23, 40 and 21

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

Question No : 50 of 52 Marks: 5 (Budgeted Time 10 Min)

A population consists of N=5 values 1, 2, 3, 5, 6.A sample size of n=3 is selected from the population without replacement, calculate sampling distribution of sample proportions for even numbers.

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

Question No : 51 of 52 Marks: 5 (Budgeted Time 10 Min)

Write down the testing procedure in case of goodness of fit test.

Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

Question No : 52 of 52 Marks: 5 (Budgeted Time 10 Min)

If we have n=634 and $\hat{p}$ =0.459, where $Z_{\frac{0.01}{2}}$=2.58, then find the 99% confidence interval for population proportion.


Answer (Please click here to Add Answer)

VuAnswers.com



Made by: Waqar Siddhu

For More Solutions or Short Lectures (Playlist) Visit our Youtube Channel



VUNET.BLOGSPOT.Com