MTH644
Measure Theory

FINAL TERM MODULE: 47 TO 93

Topic # 47

Theorem: Interval (a,∞) is measurable $\forall\ a \in R$.

Proof:

Taking test set say $A \subseteq R$ and we only need to prove:

$\because\ m^*(A) = m^*(A \cap U) + m^*(A \cap U')$

$\because\ m^*(a,b) = l(a,b) = b-a$

$$m^*(A) \geq m^*[A \cap (a,\infty)] + m^*[A \cap (-\infty, a]] \longrightarrow (1)$$

Let $A_\alpha = A \cap (a,\infty)$ and $A_\beta = A \cap (-\infty, a]$

$\therefore (1) \Longrightarrow m^*(A) \geq m^*(A_\alpha) + m^*(A_\beta) \longrightarrow (2)$

Case I: $m^*(A) = +\infty$, then (2) is obviously.

Case II: if $m^*(A) < \infty$ and for $A \subseteq R$ $\exists$ a seq. $\{I_n'\}_{n=1}^{\infty}$ such that $A \subseteq \bigcup_{n=1}^{\infty} I_n' \longrightarrow (A)$

$\forall\ \varepsilon > 0$, Then there exists another such sequence $\{I_n\}$ such that

$(A) \Rightarrow m^*(A) \leq m^*\left(\bigcup_{n=1}^{\infty} I_n^*\right)$ "

$A \subseteq \bigcup_{n=1}^{\infty} I_n$ and $m^*(A) + \varepsilon \geq m^*\left(\bigcup_{n=1}^{\infty} I_n'\right) \longrightarrow (a)$

$\because$ on R; $m^*(I_n) = l(I_n)$

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

$\therefore m^*\left(\bigcup_{n=1}^{\infty} I_n\right) = \sum_{n=1}^{\infty} l(I_n)$ → not disjoint

(a) $\Rightarrow \sum_{n=1}^{\infty} l(I_n) < m^*(A) + \epsilon$ → (3)

Let $M_n = I_n \cap (a, \infty)$ and $N_n = I_n \cap (-\infty, a]$

$M_n \cap N_n = \{I_n \cap (a,\infty)\} \cap \{I_n \cap (-\infty, a]\}$

$= I_n \cap \{(a,\infty) \cap (-\infty, a]\}$

$M_n \cap N_n = I_n \cap \phi = \phi$

Now $M_n \cup N_n = \{I_n \cap (a,\infty)\} \cup \{I_n \cap (-\infty, a]\}$

$= I_n \cap \{(a,\infty) \cup (-\infty, a]\}$

$= I_n \cap R$

$M_n \cup N_n = I_n$

$\Rightarrow l(I_n) = l(M_n) + l(N_n)$ → (4)

$\because M_n \cap N_n = \phi \ \forall\, n \in \mathbb{N}$

$\because A_\alpha = A \cap (a,\infty)$ and $A \subseteq \bigcup_{n=1}^{\infty} I_n$

$\Rightarrow A_\alpha \subseteq \left(\bigcup_{n=1}^{\infty} I_n\right) \cap (a,\infty)$

$\Rightarrow A_\alpha \subseteq (I_1 \cup I_2 \cup \dots) \cap (a,\infty)$

$\Rightarrow A_\alpha \subseteq \{I_1 \cap (a,\infty)\} \cup \{I_2 \cap (a,\infty)\} \cup \dots$

$\Rightarrow A_\alpha \subseteq \bigcup_{n=1}^{\infty} \{I_n \cap (a,\infty)\}$

$\Rightarrow A_\alpha \subseteq \bigcup_{n=1}^{\infty} M_n$

$\Rightarrow m(A_\alpha) \le m^*\left(\bigcup_{n=1}^{\infty} M_n\right)$ → (c)

Similarly $A_\beta \subseteq \bigcup_{n=1}^{\infty} N_n$

$\Rightarrow m(A_\beta) \le m^*\left(\bigcup_{n=1}^{\infty} N_n\right)$ → (d)

Adding (c) and (d)

$m(A_\alpha) + m(A_\beta) \le m^*\left(\bigcup_{n=1}^{\infty} M_n\right) + m^*\left(\bigcup_{n=1}^{\infty} N_n\right)$

$\le \sum_{n=1}^{\infty} m^*(M_n) + \sum_{n=1}^{\infty} m^*(N_n)$

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

$$m^*(A_\alpha) + m^*(A_\beta) \leq \sum_{n=1}^{\infty} l(M_n) + \sum_{n=1}^{\infty} l(N_n)$$

$$\leq \sum_{n=1}^{\infty} (l(M_n) + l(N_n))$$

$$m^*(A_\alpha) + m^*(A_\beta) \leq \sum_{n=1}^{\infty} l(I_n) \quad \text{From (4)}$$

$$\Rightarrow m^*(A_\alpha) + m^*(A_\beta) < m^*(A) + \varepsilon \quad \text{From (3)}$$

$\because \varepsilon \to 0$ arbitrarily;

$$\therefore m^*(A_\alpha) + m^*(A_\beta) \leq m^*(A)$$

or $m^*(A) \geq m^*(A_\alpha) + m^*(A_\beta)$

Hence proved; (a, ∞) is measurable.

Topic #48

Theorem: Open interval $(a,b) \subset \mathbb{R}$ is measurable.

proof: $\because (a, \infty)$ is measurable

$\Rightarrow (a,\infty)^c = (-\infty, a]$ is also measurable.

And $(-\infty, b - \frac{1}{n}]$ is also measurable $\forall\ n \in \mathbb{N}$ as $n \to \infty$

and $(-\infty, b) = \bigcup_{n=1}^{\infty} (-\infty, b - \frac{1}{n}]$ is also measurable

because being countable union of measurable set

$\Rightarrow (a,\infty) \cap (-\infty, b) = (a,b)$ i.e also measurable.

being intersection of two measurable set

Open set in $\mathbb{R}$:

Any open set in $\mathbb{R}$ say U can be written as countable union of disjoint open intervals

i.e $U = \bigcup_{n=1}^{\infty} \{I_n\}$ and each open intervals is

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

measurable and countable union of measurable set is measurable.

Topic #49

Theorem: A subset $\phi \neq E \subset \mathbb{R}$ is measurable $\Leftrightarrow$ for given $\varepsilon > 0$, $\exists$ an open set $U \supseteq E$ with $m^*(U-E) < \varepsilon$.

proof: Suppose E is measurable

Case I: If $m^*(E) < \infty$, $\forall\ \phi \neq E \subset \mathbb{R}$, there exists an open set U such that $E \subset U$ and for any $\varepsilon > 0$, $m^*(U) < m^*(E) + \varepsilon \longrightarrow (a)$

$E \subset U \Rightarrow U = E \cup (U-E)$

$\Rightarrow m^*(U) = m^*(E) + m^*(U-E)$

$\Rightarrow m^*(U-E) = m^*(U) - m^*(E) < \varepsilon$. from (a)



Case II: If $m^*(E) = \infty$

$\mathbb{R} = \bigcup_{n=1}^{\infty} I_n$, where $\ell(I_n) < \infty\ \forall\ n \in \mathbb{N}$

and $I_j \cap I_k = \phi$, $j \neq k$

put $E_n = E \cap I_n \Rightarrow E_n \subseteq I_n\ \forall\ n \in \mathbb{N}$

$\Rightarrow m^*(E_n) \leq m^*(I_n) = \ell(I_n) < \infty$



By case I $\forall\ E_n$, $\exists$ an open set say "V_n" such that $E_n \subseteq V_n$, for $\varepsilon > 0$ and $m^*(V_n - E_n) < \frac{\varepsilon}{2^n} \longrightarrow (b)$

Also $V = \bigcup_{n=1}^{\infty} V_n$, an open set.

$\Rightarrow \bigcup_{n=1}^{\infty} E_n = \bigcup_{n=1}^{\infty} (E \cap I_n) = E \cap (\bigcup_{n=1}^{\infty} I_n) = E \cap \mathbb{R} = E$

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

$E = \bigcup_{n=1}^{\infty} E_n \subseteq \bigcup_{n=1}^{\infty} V_n = V$ $\quad \because E_n \subseteq V_n \ \forall\, n \in \mathbb{N}$

$\Rightarrow V - E = \bigcup_{n=1}^{\infty} V_n - \bigcup_{n=1}^{\infty} E_n \subseteq \bigcup_{n=1}^{\infty} (V_n - E_n)$

$\Rightarrow m^*(V-E) \leq \sum_{n=1}^{\infty} m^*(V_n - E_n)$

From (b) $\Rightarrow m^*(V-E) \leq \sum_{n=1}^{\infty} \frac{\varepsilon}{2^n}$

$m^*(V-E) < \varepsilon\left(\sum_{n=1}^{\infty} \frac{1}{2^n}\right) = \varepsilon(1)$

$m^*(V-E) \leq \varepsilon$

Conversely, let $\varepsilon = \frac{1}{n}$ (say) and

$\exists$ an open set $O_n \supseteq E$ such that $E \subseteq O_n \ \forall\, n \in \mathbb{N}$

and $m^*(O_n - E) < \varepsilon = \frac{1}{n} \longrightarrow (A)$

We will prove "E" is measurable.

put $G_1 = \bigcap_{n=1}^{\infty} O_n \Rightarrow$ "G_1" is measurable being countable intersection of measurable open set.

$E \subseteq O_n \ \forall\, n$ (given)

and $E \subseteq \bigcap_{n=1}^{\infty} O_n = G_1$

$G_1 - E \subseteq O_n - E$

$\Rightarrow m^*(G_1 - E) \leq m^*(O_n - E) < \frac{1}{n} = \varepsilon$

$\because \varepsilon \to 0$ arbitrary $m^*(G_1 - E) = 0$.

$\Rightarrow G_1 - E$ is measurable $\Rightarrow (G_1 - E)'$ is also measurable.



$(V_1 \cup V_2) = (E_1 \cup E_2)$



$(V_1 - E_1) \cup (V_2 - E_2)$

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

$\therefore E = G_1 - (G_1 - E) = G_1 \cap (G_1 - E)'$ is measurable being intersection of two measurable set.

Topic #50

Theorem: For a subset "E" of R and given $\varepsilon > 0$ $\exists$ an open set $O \supseteq E$ and $m^*(O-E) < \varepsilon \Leftrightarrow \exists$ a closed set $F \subseteq E$ and $m^*(E-F) < \varepsilon$.

proof: For $\phi \neq E \subseteq R \Rightarrow E' \subseteq R$ $\exists$ for given $\varepsilon > 0$

$\exists$ an open set $O \supseteq E$ and

$m^*(O-E) < \varepsilon \longrightarrow (1)$



$O \supseteq E' \Rightarrow E'' \supseteq O' = F$ (say)

i.e. closed set

$\Rightarrow E \supseteq F \Rightarrow m^*(E-F) = m^*(E \cap F')$

$m^*(E-F) = m^*(F' \cap E)$

$m^*(E-F) = m^*(O-E') < \varepsilon$ from (1)

$\Rightarrow m^*(E-F) < \varepsilon$

Conversely for the set $E' \subseteq R$, $\exists$ a closed set "F" and $\varepsilon > 0$ such that $F \subseteq E'$ and $m^*(E'-F) < \varepsilon \longrightarrow (2)$ (given)

$F \subseteq E' \Rightarrow E = E'' \subseteq F' = O$ (open set) say.

$\Rightarrow m^*(O-E) = m^*(O \cap E') = m^*(E' \cap O)$

$\Rightarrow m^*(O-E) = m^*(E'-O') = m^*(E'-F) < \varepsilon$ from (2)

$m^*(O-E) < \varepsilon$ (Hence proved)

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

Topic #52

Theorem: Let $\phi \neq E \subset R$, then the following statements are equivalent

(i) There is a G_δ set "G" with $E \subset G$, $m^*(G-E) = 0$ → give

(ii) There is a F_σ set "F" with $F \subset E$, $m^*(E-F) = 0$ → prove

proof: (i) $\Rightarrow$ (ii) $\because \phi \neq E \subset R \Rightarrow \phi \neq E' \subseteq R$

$\therefore$ For $\phi \neq E'$ given $E' \subseteq G$, $m^*(G-E') = 0$

$\therefore \quad 0 = m^*(G-E') = m^*(G \cap (E')') = m^*(G \cap E)$

$= m^*(E \cap G) = m^*(E \cap (G')') = m^*(E-G')$

$= m^*(E-F)$, where $G_\delta = G = \bigcap_{n=1}^{\infty} G_n$

$\Rightarrow G_\delta' = G' = \left(\bigcap_{n=1}^{\infty} G_n\right)'$

$\Rightarrow G_\delta' = \bigcup_{n=1}^{\infty} G_n'$

$\Rightarrow G_\delta' = \bigcup_{n=1}^{\infty} F_n' = F = F_\sigma$

(ii) $\Rightarrow$ (i)

Conversely, again for $E' \subset R$, with $F \subset E'$

$m^*(E'-F) = 0$ (given).

$\therefore \quad 0 = m^*(E'-E) = m^*(E' \cap F') = m^*(F' \cap E')$

$= m^*(F'-E) = m^*(G-E)$

where $F_\sigma' = F' = \left(\bigcap_{k=1}^{\infty} F_k\right)' = \bigcup_{k=1}^{\infty} F_k' = \bigcup_{k=1}^{\infty} G_n = G = G_\delta$

(proved)

Topic #52

Lebesgue Measure:

Let "M" be the class of measurable sets, which is σ-algebra, Then the Lebesgue measure of any set $E \subseteq M$ is defined to be its lebesgue outer measure.

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

$$m^* \equiv m ; \left\{ A,\ m^*(A) = m^*(A \cap E) + m^*(A \cap E') \right.$$

Topic #53

Theorem: let $\{E_i\}_{i=1}^{n}$ be a sequence a in measurable space "M", then m is

(i) countably subadditive

(ii) finitely additive provided that $E_i \cap E_j = \phi, \forall\ i \neq j$.

(iii) countably additive if E_i's are pairwise disjoint.

(iv) "m" is monotonic.

proof: (i) All E_i's are measurable.

$\Rightarrow$ $\bigcup_{i=1}^{n} E_i$ is also measurable, $\therefore$ "M" is space of lebesgue measurable set i

$$\Rightarrow m\left(\bigcup_{i=1}^{n} E_i\right) = m^*\left(\bigcup_{i=1}^{\infty} E_i\right) \leq \sum_{i=1}^{\infty} m^*(E_i) = \sum_{i=1}^{\infty} m(E_i)$$

(ii) $\forall\ \phi \neq A \subseteq R$ and for the sequence $\{E_i\}_{n=1}^{n}$ of pairwise disjoint measurable sets ;

$$m^*\left(A \cap \left(\bigcup_{i=1}^{\infty} E_i\right)\right) = \sum_{i=1}^{\infty} m^*(A \cap E_i)$$

put $A = R \Rightarrow A\left(\bigcup_{i=1}^{\infty} E_i\right) = \bigcup_{i=1}^{\infty} E_i$ $\quad A \cap E_i = R \cap E_i$

$$\Rightarrow m^*\left(\bigcup_{i=1}^{\infty} E_i\right) = \sum_{i=1}^{\infty} m^*(E_i)$$

$$\Rightarrow m\left(\bigcup_{i=1}^{\infty} E_i\right) = \sum_{i=1}^{\infty} m(E_i)$$

(iii) $m^*\left(\bigcup_{i=1}^{n} E_i\right) = \sum_{i=1}^{n} m^*(E_i)$

$\Rightarrow m\left(\bigcup_{i=1}^{n} E_i\right) = \sum_{i=1}^{n} m(E_i)$ finitly additive.

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

iv) Monotonicity : Let $\phi \neq A, B \in M$, such that $B \subseteq A$.

$\Rightarrow A = B \cup (A - B)$

$\therefore m^*$ is finitely additive,

$\Rightarrow m^*(A) = m^*(B) + m^*(A-B)$

$\Rightarrow m^*(A) \geq m^*(B)$ $\because m^*(A-B) \geq 0$

$\Rightarrow m^*(B) \leq m^*(A)$



$\because B \cap (A-B) \neq \phi$

Topic # 54

Theorem: Lebesgue measure "m" is translation invariant.

proof: $\because$ Given that say $\phi \neq E \in M$ is measurable.

$\because$ "E" is measurable $\Leftrightarrow$ for given $\varepsilon > 0$, $\exists$ an open set $U \supseteq E$, such that $m^*(U-E) < \varepsilon$

"U" is open set in $R \Rightarrow$ "U+y" is also open in R, $\forall\ y \in R$

if $E \subseteq U \Rightarrow E+y \subseteq U+y$

$\Rightarrow (U+y)-(E+y) = (U-E)+y$

$\Rightarrow m^*[(U+y)-(E+y)] = m^*[(U-E)+y]$



$\because$ Lebesgue outer measure is translation invariant.

$\Rightarrow m^*(U-E) < \varepsilon$

But $m^* = m \Rightarrow m[(U+y)-(E+y)] = m(U-E) < \varepsilon$

$\Rightarrow$ "E+y" is also measurable $\Rightarrow$ "m" is translation invariant.

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

Topic # 55

Theorem: Suppose E_1 and E_2 are two measurable sets, then $m(E_1 \cup E_2) + m(E_1 \cap E_2) = m(E_1) + m(E_2)$

proof: If $m(E_1) = m(E_2) = \infty$, then the result is obvious.

Case I: If $m(E_1) < \infty$, $m(E_2) < \infty$

$\therefore$ if E_1 is measurable, then for any $\phi \neq A \subseteq R$

$$m^*(A) = m^*(A \cap E_1) + m^*(A \cap E_1') \longrightarrow (1)$$

For particularly if $A = E_2$

$$(1) \Rightarrow m^*(E_2) = m^*(E_2 \cap E_1) + m^*(E_2 \cap E_1') \longrightarrow (2)$$

Similarly $E_2 \longleftrightarrow E_1$

$$m^*(E_1) = m^*(E_1 \cap E_2) + m^*(E_1 \cap E_2') \longrightarrow (3)$$

Adding (2) and (3): $(m^* \equiv m)$

$$m(E_1) + m(E_2) = m(E_1 \cap E_2) + m(E_1 \cap E_2') + m(E_2 \cap E_1) + m(E_2 \cap E_1') \longrightarrow (A)$$

$$E_1 \cup E_2 = (E_1 - E_2) \cup (E_2 - E_1) \cup E_1 \cap E_2$$

$$= (E_1 \cap E_2') \cup (E_1 \cap E_1') \cup (E_1 \cap E_2)$$



$$\Rightarrow m(E_1 \cup E_2) = m[(E_1 \cap E_2') \cup (E_2 \cap E_1') \cup (E_1 \cap E_2)]$$

"m" is finitely addative:

$$m(E_1 \cup E_2) = m(E_1 \cap E_2') + m(E_2 \cap E_1') + m(E_1 \cap E_2)$$

$$(A) \Rightarrow m(E_1) + m(E_2) = m(E_1 \cup E_2) + m(E_1 \cap E_2)$$

proved

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

Topic # 56

$R = R^2 = R^3$

Cantor Set:



Cantor dust

# Step	# Dropped Interval	length	Interval
n = 1	$1 = m = 2^{1-1}$	$l(I_{11}) = \frac{1}{3}$	$(\frac{1}{3}, \frac{2}{3})$
n = 2	$2 = m = 2^{2-1}$	$l(I_{21}) = l(I_{22}) = \frac{1}{3^2}$	$(\frac{1}{9}, \frac{2}{9}), (\frac{7}{9}, \frac{8}{9})$
n = 3	$4 = m = 2^{3-1}$	$l(I_{31}) = ___ = \frac{1}{3^3}$	
n	$m = 2^{n-1}$	$l(I_{n.}) = ___ = \frac{1}{3^n}$	

Dropped Interval $= G_1 = \underbrace{I_{11}}_{n=1}, \underbrace{I_{21}, I_{22}}_{n=2}, \underbrace{I_{31}, I_{32}, I_{33}, I_{34}}_{n=3}, \dots$

$I_{n1}, I_{n2}, \dots I_{nm}, \dots$

$n \to \infty, \; m = 2^{n-1}$

$\therefore G_1 = \{I_{11}\}, \{I_{21}, I_{22}\}, \{I_{31}, I_{32}, I_{33}, I_{34}\}, \dots$

$n \to \infty, \; m = 2^{n-1}$

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

$$\Rightarrow G_1 = \bigcup_{n=1}^{\infty} \bigcup_{m=1}^{2^{n-1}} I_{nm}$$

$\therefore$ Cantor Set $= C = [0,1] - G_1 = G_1'$ ← universal set.

Topic #57

Measure of Cantor Set:

$\because$ "G_1" is open, being countable union of disjoint open intervals $\Rightarrow$ "G_1" is measurable.

$\because C = G_1'$ = closed and is also measurable.

$$m(G_1) = m\left(\bigcup_{n=1}^{\infty} \bigcup_{m=1}^{2^{n-1}} I_{nm}\right) \quad \because \text{"}m\text{" is countably additive}$$

$$= \sum_{n=1}^{\infty} \sum_{m=1}^{2^{n-1}} m(I_{nm}) = \sum_{n=1}^{\infty} \sum_{m=1}^{2^{n-1}} l(I_{nm})$$

expanding firstly by "n"

$$= \sum_{m=1}^{2^{1-1}} l(I_{1m}) + \sum_{m=1}^{2^{2-1}} l(I_{2m}) + \sum_{m=1}^{2^{3-1}} l(I_{3m}) + \cdots\cdots$$

$$= \underbrace{l(I_{11})} + \underbrace{l(I_{21}) + l(I_{22})} + \underbrace{l(I_{31}) + l(I_{32}) + l(I_{33}) + l(I_{34})} + \cdots\cdots$$

$$= \frac{1}{3} + \left(\frac{1}{3^2} + \frac{1}{3^2}\right) + \left(\frac{1}{3^3} + \frac{1}{3^3} + \frac{1}{3^3} + \frac{1}{3^3}\right) + \cdots\cdots$$

$$m(G_1) = \frac{1}{3} + \frac{2}{3^2} + \frac{4}{3^3} + \cdots\cdots \quad \begin{cases} a = \frac{1}{3}, & r = \frac{2}{3} \\ S_\infty = \frac{a}{1-r} = \frac{1/3}{1-2/3} = 1 \end{cases}$$

$\boxed{m(G_1) = 1}$ $\because [0,1] = G \cup C \Rightarrow m[0,1] = m(G_1) + m(C)$ (disjoint) $\Rightarrow \boxed{m(C) = 0}$

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

Topic #58

Generalized Cantor Set and its Measure:



$$m(G_1) = \frac{\alpha}{3} + \left(\frac{\alpha}{9} + \frac{\alpha}{9}\right) + \left(\frac{\alpha}{27} + \frac{\alpha}{27} + \frac{\alpha}{27} + \frac{\alpha}{27}\right) + \cdots$$

$$m(G_1) = \frac{\alpha}{3} + \frac{2\alpha}{3^2} + \frac{4\alpha}{3^3} + \cdots$$

$a = \alpha/3$, $r = 2/3$

$$S_\infty = \frac{a}{1-r} = \frac{\alpha/3}{1-2/3} = \frac{\alpha/3}{1/3} = \alpha$$

$$\boxed{m(G_1) = \alpha}$$

$\because [0,1] = G_1 \cup C$ (disjoint)

$$\Rightarrow m[0,1] = m(G_1) + m(C)$$

$$1 = \alpha + m(C)$$

$$\Rightarrow \boxed{m(C) = 1 - \alpha}$$

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

Topic # 59

Theorem: Let $\{E_n \uparrow\}_{n=1}^{\infty}$ be an increasing sequence of measurable sets. Then

$$m^*\left(\bigcup_{n=1}^{\infty} E_n\right) = m\left(\bigcup_{n=1}^{\infty} E_n\right) = \lim_{n\to\infty} m^*(E_n)$$

Proof: $\because$ Given that $(E_n, \uparrow)$, such that

$\Rightarrow E_1 \subseteq E_2 \subseteq E_3 \subseteq \cdots$ of measurable sets

Defining $F_1 = E_1$, $F_2 = E_2 - E_1$, $F_3 = E_3 - E_2$, $\cdots$

$F_n = E_n - E_{n-1}$, where $\{F_n\}_{n=1}^{\infty}$ is a seq. of disjoint measurable sets.

$$\Rightarrow \bigcup_{k=1}^{n} F_k = \bigcup_{k=1}^{n} E_k \Rightarrow \bigcup_{k=1}^{\infty} F_k = \bigcup_{k=1}^{\infty} E_k.$$

$$\therefore m^*\left(\bigcup_{k=1}^{\infty} E_k\right) = m\left(\bigcup_{k=1}^{\infty} E_k\right) = m\left(\bigcup_{k=1}^{\infty} F_k\right) \quad \text{(disjoint + measurable)}$$

$$= \sum_{k=1}^{\infty} m(F_k) = \lim_{n\to\infty} \sum_{k=1}^{\infty} m(F_k)$$

$$= \lim_{n\to\infty} m\left(\bigcup_{k=1}^{\infty} F_k\right) = \lim_{n\to\infty} m\left(\bigcup_{k=1}^{\infty} E_k\right)$$

$$= \lim_{n\to\infty} m^*(E_n) \qquad \text{(proved)}$$

Topic # 60

Theorem: Let $\{E_n \downarrow\}_{n=1}^{\infty}$ be a decreasing sequence of measurable sets such that $m(E_k) < \infty$ for some $k \in \mathbb{N}$, Then

$$m^*\left(\bigcap_{n=1}^{\infty} E_n\right) = m\left(\bigcap_{n=1}^{\infty} E_n\right) = \lim_{n\to\infty} m^*(E_n)$$

This PDF document was edited with Icecream PDF Editor. Upgrade to PRO to remove watermark.

Scanned with CamScanner

proof: Let $m(E_1) < \infty$, Say (without lose of generality)

$\therefore \langle E_n, \downarrow \rangle \Rightarrow E_1 \supseteq E_2 \supseteq E_3 \supseteq \cdots$

$\Rightarrow \phi = E_1 - E_1 \subseteq E_1 - E_2 \subseteq E_1 - E_3 \subseteq \cdots E_1 - E_n \subseteq \cdots$

$E_1 \subseteq E_n \subseteq \cdots$

$$\Rightarrow m^*\left(\bigcup_{n=1}^{\infty}(E_1 - E_n)\right) = \lim_{n\to\infty} m^*(E_1 - E_n)$$

$$\bigcup_{n=1}^{\infty}(E_1 - E_n) = \bigcup_{n=1}^{\infty}(E_1 \cap E_n')$$

$$= (E_1 \cap E_1') \cup (E_1 \cap E_2') \cup \cdots$$

$$= E_1 \cap [E_1' \cup E_2' \cup \cdots]$$

$$= E_1 \cap [E_1 \cap E_2 \cap \cdots]'$$

$$= E_1 \cap \left(\bigcap_{n=1}^{\infty} E_n\right)'$$

$$= E_1 - \left(\bigcap_{n=1}^{\infty} E_n\right)$$

Sets Relation ↑

$$m^*\left(E_1 - \bigcap_{n=1}^{\infty} E_n\right) = \lim_{n\to\infty} m^*(E_1 - E_n)$$

$$m^*(E_1) - m^*\left(\bigcap_{n=1}^{\infty} E_n\right) = \lim_{n\to\infty} m^*(E_1) - \lim_{n\to\infty} m^*(E_n)$$

$$m^*(E_1) - m^*\left(\bigcap_{n=1}^{\infty} E_n\right) = m^*(E_1) - \lim_{n\to\infty} m^*(E_n)$$

$$\Rightarrow m^*\left(\bigcap_{n=1}^{\infty} E_n\right) = \lim_{n\to\infty} m^*(E_n) \quad \text{(proved)}.$$

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

Topic # 61

Borel Set:

e.g. $G_\delta = \bigcap_{n=1}^{\infty} U_n$; $F_\sigma = \bigcup_{n=1}^{\infty} F_n$

Definition: A set "E" is said to be a Borel set if it can be expressed as countable union or intersection of open or closed sets.

e.g. $\{x\} = \bigcap_{n=1}^{\infty} \left(x - \frac{1}{n}, x + \frac{1}{n}\right)$ → G_δ Borel sets.

Topic # 62

Borel Measure Space:

Restriction of Lebesgue measure "m" to the class of Borel set "B" is said to be Borel measure space. $(\mathbb{R}, B, m)$

Theorem: Every Borel set is measurable.

proof: Every open or closed set is measurable. ⇒ Countable union or intersection of closed or open set is also measurable.

∴ Borel set is also measurable being union or intersection of closed or open sets

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

Topic #63

Theorem: Countable set are Borel sets with zero measure.

Proof: Case I: Singleton:

$$\forall\ a \in \mathbb{R}: \{a\} = \bigcap_{n=1}^{\infty} \left(a - \frac{1}{n}, a + \frac{1}{n}\right) \quad ; n \in \mathbb{N}$$

Countable ↓ finite ↔ infinite. $\mathbb{Z}, \mathbb{N}, \mathbb{R}, \mathbb{Q}$ ---

$\therefore \left\{\left\{a - \frac{1}{n}, a + \frac{1}{n}\right\}, \downarrow\right\}$ is decreasing sequence.

with $m(E_n) = \frac{2}{n} < \infty$

where $E_n = \left(a - \frac{1}{n}, a + \frac{1}{n}\right)$

$$\Rightarrow m\{a\} = m\left\{\bigcap_{n=1}^{\infty} E_n\right\} = \lim_{n \to \infty} m(E_n)$$

$$= \lim_{n \to \infty} l(E_n)$$

$$= \lim_{n \to \infty} \left(\frac{2}{n}\right) = \frac{2}{\infty}$$

$$= 0$$

Finite Set

$$\{x_1, x_2, x_3, \dots\dots x_n\} = \bigcup_{i=1}^{n} \{x_i\}$$

$$\Rightarrow m\left(\bigcup_{i=1}^{n} \{x_i\}\right) = \sum_{i=1}^{n} m\{x_i\} = 0$$

Infinite Set:

$$\{x_1, x_2, x_3, \dots\dots\dots\} = \bigcup_{i=1}^{\infty} \{x_i\}$$

$$\Rightarrow m\left(\bigcup_{i=1}^{\infty} \{x_i\}\right) = \sum_{i=1}^{\infty} m\{x_i\} = 0$$

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

Topic #64

Sum Modulo 1:

Let $x, y \in A = [0,1)$ and we define sum modulo "1" of x and y as

$$x \hat{+} y = \begin{cases} x+y & \text{if } x+y<1 \\ x+y-1 & \text{if } x+y \geq 1 \end{cases}$$

Translation Modulo 1:

Let $E \subseteq A = [0,1)$ and $y \in A$, we define a translation modulo "1" of "E" as.

$$E \hat{+} y = \{ x \hat{+} y, x \in E \}.$$

$\downarrow I$

Set of all "Sum Modulo 1"

Topic #65

Theorem: Let $E \subset A = [0,1)$ be a measurable set, Then for ~~any~~ each $y \in A$. The set $E \hat{+} y$ is measurable and $m(E) = m(E \hat{+} y)$.

proof: Suppose $E_1 = E \cap [0, 1-y)$ and

$E_2 = E \cap [1-y, 1)$

$\therefore E_1 [0, 1-y), [1-y, 0)$ are measurable

$\Rightarrow E_1$ and E_2 are also measurable being the intersection of two measurable sets.

Also $E_1 \cap E_2 = \{E \cap [1-y, 1)\} \cap \{E \cap [0, 1-y]\}$

$= E \cap \{[0, 1-y) \cap [1-y, 1)\}$

$= E \cap \phi = E$

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

and $E_1 \cup E_2 = E \cup \phi = E$

$\Rightarrow m(E) = m(E_1) + m(E_2) \longrightarrow (1)$

"m" is finitely additive.

Let $y_1 \in A$, if $y_1 \in E_1$

$\Rightarrow y_1 < 1-y \Rightarrow y_1 + y < 1 \Rightarrow y_1 \hat{+} y = y_1 + y$

$E_1 + y$

$\Rightarrow E_1 \hat{+} y = E_1 + y \longrightarrow (a)$

if $y_2 \in E_2 \Rightarrow y_2 \geq 1-y \Rightarrow y_1 + y \geq 1$

$\Rightarrow y_2 \hat{+} y = y_2 + y - 1$

$\Rightarrow E_2 \hat{+} y = E_2 + (y-1) \longrightarrow (b)$

$\because$ "m" is translation invariant.

$\Rightarrow E_1 + y$ and $E_2 + (y-1)$ are measurable.

$\Rightarrow E_1 \hat{+} y$ and $E_2 \hat{+} (y-1)$ are measurable.

$\Rightarrow m(E_1 \hat{+} y) = m(E_1 + y) = m(E_1)$

Similarly

$$m(E_2 \hat{+} y) = m(E_2 + y - 1) = m(E_2)$$

$\because E_1 \cup E_2 = E$, $E_1 \cap E_2 = \phi$

$\Rightarrow E \hat{+} y = (E_1 + y) \cup (E_2 + y)$ and $(E_1 + y) \cap (E_2 + y) = \phi$

"$E \hat{+} y$" being union of two disjoint measurables sets its measurable.

$\Rightarrow m(E \hat{+} y) = m(E_1 \hat{+} y) + m(E_2 \hat{+} y)$

$= m(E_1 + y) + m(E_2 + y)$

$= m(E_1) + m(E_2) = m(E)$

$\Rightarrow$ "m" is also invariant w.r.t translation modulo "1".

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

Topic # 66

Measurable Function:

Let "M" be the measurable space (σ-algebra of measurable sets). a function $f: E \to \mathbb{R}$ is Lebesgue measurable function on $\phi \neq E \in M$.

if $\{x \in E : f(x) > a\} \in M$ for every $a \in \mathbb{R}$.



For Example (1): Suppose "f" is real valued identically constant function $f(x) = \beta$, Then:

$$\{x : f(x) > a\} = \begin{cases} \mathbb{R} & \text{if } \beta > a \\ \phi \in M \end{cases}$$

$\therefore \mathbb{R}, \phi \in M$

$\Rightarrow$ "f" is measurable.

For Example (2):

$$f(x) = \begin{cases} 1 & \text{if } x \in A \\ 0 & \text{if } x \notin A \end{cases}$$

$$\{x : f(x) > a\} = \begin{cases} \phi \\ A \neq X \\ X = A \end{cases}$$

$\phi, A, X \in M \Rightarrow$ "f" is measurable.

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

Example (3): $(R, B, m) \to$ (Borel σ-algebra)

$f(x) = x$

$\Rightarrow \{x : f(x) > a\} = \begin{cases} (a, +\infty) \in B \\ \text{i.e always measurable.} \end{cases}$

$\Rightarrow$ "f" is measurable.

Topic # 67

Theorem: Let $f : E \to R \cup \{\pm\infty\}$, $\phi \neq E \in M$, Then.

(i) "f" is measurable $\Leftrightarrow$ (ii) $\{x \in E : f(x) \geq a\}$ is measurable for all $a \in R \Leftrightarrow$ (iii) $\{x \in E : f(x) < a\}$ is measurable $\forall\ a \in R \Leftrightarrow$ (iv) $\{x \in E : f(x) \leq a\}$ is measurable $\forall\ a \in R$.

Proof: (i) $\to$ (ii) "f" is measurable $\to \{x \in E : f(x) > a\} \in M$

$$\{x \in E : f(x) \geq a\} = \bigcap_{n=1}^{\infty} \left\{ x \in E : f(x) > a - \frac{1}{n} \right\}$$

Let $y \in \{x \in E : f(x) \geq a\} \Rightarrow f(y) \geq a, \forall\ a \in R$

$\Rightarrow f(y) > a - \frac{1}{n}, \forall\ n \in N$

$\Rightarrow y \in \{x \in E : f(x) > a - \frac{1}{n}\}, \forall\ n \in N$

$\Rightarrow y \in \bigcap_{n=1}^{\infty} \{x \in E : f(x) > a - \frac{1}{n}\}$

$$\{x \in E : f(x) \geq a\} \subset \bigcap_{n=1}^{\infty} \{x \in E : f(x) > a - \frac{1}{n}\} \quad \rightarrow (a)$$

Let conversely $t \in \bigcap_{n=1}^{\infty} \{x \in E : f(x) > a - \frac{1}{n}\}$

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

$\Rightarrow f(t) > a - \frac{1}{n}, \forall\ n \in \mathbb{N}$

$\Rightarrow f(t) \geq a, \forall\ a \in \mathbb{R}$ ~~$\Rightarrow t \in$~~

$\Rightarrow t \in \{x \in E : f(x) \geq a\}$

$\Rightarrow \bigcap_{n=1}^{\infty} \{x \in E : f(x) > a - \frac{1}{n}\} \subseteq \{x : f(x) \geq a\}$ → (b)

from (a) and (b)

$\Rightarrow \{x \in E : f(x) \geq a\} = \bigcap_{n=1}^{\infty} \{x \in E : f(x) > a - \frac{1}{n}\}$

$\therefore \{x \in E : f(x) > a - \frac{1}{n}\}$ is measurable.

$\Rightarrow \bigcap_{n=1}^{\infty} \{x \in E : f(x) > a - \frac{1}{n}\}$ is also measurable.

$\forall\ n \in \mathbb{N}$

$\Rightarrow \{x \in E : f(x) \geq a\}$ is also measurable.

Topic #68

(ii) $\Rightarrow$ (iii) $\{x \in E : f(x) \geq a\} \in M \Rightarrow \{x \in E : f(x) < a\} \in M$.

$\{x \in E : f(x) < a\} = \{x \in E : f(x) \geq a\}^c \quad \forall\ a \in \mathbb{R}$

$\therefore \{x \in E : f(x) \geq a\} \in M \Rightarrow \{x \in E : f(x) \geq a\}^c \in M$

$\Rightarrow \{x \in E : f(x) < a\} \in M$ i.e. is measurable.

Topic #69

(iii) $\Rightarrow$ (iv) $\underbrace{\{x \in E : f(x) < a\}}_{\in M} \Rightarrow \{x \in E : f(x) < a\} \in M$.

ple show that $\{x \in E : f(x) \leq a\} =$

$= \bigcap_{n=1}^{\infty} \{x \in E : f(x) < a + \frac{1}{n}\}$

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

Let $y \in \{x \in E : f(x) \leq a\}$ ——— (a)

$\Rightarrow f(y) \leq a$, $\forall\ a \in R$

$\Rightarrow f(y) < a + \frac{1}{n}$, $\forall\ n \in N$

$\Rightarrow y \in \{x \in E : f(x) < a + \frac{1}{n}\}$, $\forall\ n \in N$

$\Rightarrow y \in \bigcap_{n=1}^{\infty} \{x \in E : f(x) < a + \frac{1}{n}\}$ ——— (B)

From (a) and (B) $\Rightarrow \{x \in E : f(x) \leq a\} \subseteq \bigcap_{n=1}^{\infty} \{x \in E : f(x) < a + \frac{1}{n}\}$

Let $z \in \bigcap_{n=1}^{\infty} \{x \in E : f(x) < a + \frac{1}{n}\}$

$\Rightarrow z \in \{x \in E : f(x) < a + \frac{1}{n}\} \Rightarrow f(z) < a + \frac{1}{n}, \forall n \in N$

$\Rightarrow f(z) \leq a\ \forall\ a \in R \Rightarrow z \in \{x \in E : f(x) \leq a\}$

$\Rightarrow \bigcap_{n=1}^{\infty} \{x \in E : f(x) < a + \frac{1}{n}\} \subseteq \{x \in E : f(x) \leq a\}$

$\therefore \bigcap_{n=1}^{\infty} \{x \in E : f(x) < a + \frac{1}{n}\} = \{x \in E : f(x) \leq a\}$

Given that $\{x \in E : f(x) < a + \frac{1}{n}\} \in M$.

$\Rightarrow \bigcap_{n=1}^{\infty} \{x \in E : f(x) < a + \frac{1}{n}\} \in M$

$\Rightarrow \{x \in E : f(x) \leq a\} \in M$. i.e is measurable.

Topic # 70

(iv) $\Rightarrow$ (i) ; $\{x \in E : f(x) \leq a\} \in M \Rightarrow$ "f" is measurable.

$f : E \to R \cup \{+\infty\}$, $\{x \in E : f(x) < a\} \in M$,

$\{x \in E : f(x) \leq a\} \in M$, $\forall\ a \in R$

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

$= \{x \in E : f(x) \leq a\}^c = \{x \in E : f(x) > a\} \in M$

$\Rightarrow$ "f" is measurable.

Topic # 71

Theorem: Let $\phi \neq E \in M$, $f : E \to R \cup \{\pm\infty\}$, then $\{x \in E : f(x) = a\} \in M$, $\forall a \in R$.

Proof: Case I: $a \in R$

$\Rightarrow \{x \in E : f(x) = a\} = \{x \in E : f(x) \geq a\} \cap \{x \in E : f(x) \leq a\}$

$\therefore \{x \in E : f(x) \geq a\}$ and $\{x \in E : f(x) \leq a\} \in M$.

$\Rightarrow \{x \in E : f(x) \geq a\} \cap \{x \in E : f(x) \leq a\} \in M$.

$\Rightarrow \{x \in E : f(x) = a\} \in M$ (measurable)

Case II: If $a = \infty$, Then $\{x \in E : f(x) = \infty\} = \bigcap_{n=1}^{\infty} \{x \in E : f(x) > n\}$, $\forall n \in N$

Let $y \in \{x \in E : f(x) = \infty\} \Rightarrow y \in \{x \in E : f(x) > n\}$

$\Rightarrow y \in \bigcap_{n=1}^{\infty} \{x \in E : f(x) > n\}$, $\forall n \in N$

$\Rightarrow \{x \in E : f(x) = \infty\} \subseteq \bigcap_{n=1}^{\infty} \{x \in E : f(x) > n\}$ ——— (a)

Now let $t \in \bigcap_{n=1}^{\infty} \{x \in E : f(x) > n\}, \forall n \in \mathbb{N}$

$\Rightarrow t \in \{x \in E : f(x) > n\}, \forall n \in \mathbb{N}$

$\Rightarrow f(t) > n, \forall n \in \mathbb{N}$

$\Rightarrow t \in \{x \in E : f(x) = \infty\}$

$\Rightarrow \bigcap_{n=1}^{\infty} \{x \in E : f(x) > n\} \subseteq \{x \in E : f(x) = \infty\} \longrightarrow (b)$

From (a) and (b)

So, $\{x \in E; f(x) = \infty\} = \bigcap_{n=1}^{\infty} \{x \in E : f(x) > n\}$

Case III: If $a = -\infty$, Then $\{x \in E : f(x) = -\infty\} =$

$$\bigcap_{n=1}^{\infty} \{x \in E : f(x) < -n\}$$

Let $z \in \{x \in E : f(x) = -\infty\} \Rightarrow z \in \{x \in E : f(x) < -n\}$

$\Rightarrow z \in \bigcap_{n=1}^{\infty} \{x \in E : f(x) < -n\}, \forall n \in \mathbb{N}$

$\Rightarrow \{x \in E : f(x) = -\infty\} \subseteq \bigcap_{n=1}^{\infty} \{x \in E : f(x) < -n\} \longrightarrow (a)$

Now let $s \in \bigcap_{n=1}^{\infty} \{x \in E : f(x) < -n\}, \forall n \in \mathbb{N}$

$\Rightarrow s \in \{x \in E : f(x) < -n\}, \forall n \in \mathbb{N}$

$\Rightarrow f(s) < -n, \forall n \in \mathbb{N}$

$\Rightarrow s \in \{x \in E : f(x) = -\infty\}$

$\Rightarrow \bigcap_{n=1}^{\infty} \{x \in E : f(x) < -n\} \subseteq \{x \in E : f(x) = -\infty\} \longrightarrow (b)$

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

From (a) and (b)

So, $\{x \in E : f(x) = -\infty\} = \bigcap_{n=1}^{\infty} \{x \in E : f(x) < -n\}$

Topic # 72

Positive anel Negative parts of Extented Real Valued Function:

$$|f(x)| = \begin{cases} +f(x) & \text{if } f(x) \geq 0 \\ -f(x) & \text{if } f(x) < 0 \end{cases}$$

$$f^+ = \begin{cases} f(x) & \text{if } f(x) \geq 0 \\ 0 & \text{if } f(x) < 0 \end{cases}$$

$$= \max\{f(x), 0\} = f \vee 0$$

$$f^- = \begin{cases} 0 & \text{if } f(x) \geq 0 \\ -f(x) & \text{if } f(x) < 0 \end{cases}$$

$$= \max\{-f(x), 0\} = -f \vee 0$$

where $f : A \to \mathbb{R} \cup \{\pm\infty\}$, $A \neq \phi$, $\forall x \in A$

Topic # 73

Theorem: Show that

(i) $f^+ = \frac{f + |f|}{2}$, (ii) $f^- = \frac{|f| - f}{2}$, (iii) $f = f^+ - f^-$

(iv) $|f| = f^+ + f^-$, (v) $\max(f, g) = \frac{(f+g) + |f-g|}{2}$

(vi) $\min(f, g) = \frac{(f+g) - |f-g|}{2}$

Proof:

(i) If $f(x) \geq 0 \Rightarrow f^+(x) = f(x) \Rightarrow f^+ = f$

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

$$|f(x)| = |f|(x) = f(x)$$

$$\therefore \frac{f+|f|}{2}(x) = \frac{f(x)+|f|(x)}{2} = \frac{f(x)+f(x)}{2} = f(x) = f^+(x)$$

If $f(x) < 0 \Rightarrow f^+(x) = 0$, $|f|(x) = |f(x)| = -f(x)$

$$\therefore \frac{f+|f|}{2}(x) = \frac{f(x)+|f|(x)}{2} = \frac{f(x)-f(x)}{2} = 0 = f^+(x)$$

$$\Rightarrow \frac{f+|f|}{2} = f^+ \quad \text{(proved)}$$

(ii) If $f(x) > 0 \Rightarrow f^-(x) = 0$, $|f(x)| = |f|(x) = +f(x)$

$$\therefore \frac{|f|-f}{2}(x) = \frac{|f|(x)-f(x)}{2} = \frac{+f(x)-f(x)}{2} = 0 = f^-(x)$$

If $f(x) < 0 \Rightarrow f^-(x) = -f(x)$, $|f(x)| = |f|(x) = -f(x)$

$$\therefore \frac{|f|-f}{2}(x) = \frac{|f|(x)-f(x)}{2} = \frac{-f(x)-f(x)}{2} = -f(x) = f^-(x)$$

$$\Rightarrow \frac{|f|-f}{2} = f^- \quad \text{(proved)}$$

(iii) $f = f^+ - f^-$

If $f(x) \geq 0 \Rightarrow f^+(x) = f(x)$, $f^-(x) = 0$

$$\therefore (f^+ - f^-)(x) = f^+(x) - f^-(x) = f(x) - 0 = f(x) = f$$

$$\Rightarrow f^+ - f^- = f \quad \text{(proved)}.$$

(iv) $|f| = f^+ + f^-$

If $f(x) \geq 0 \Rightarrow f^+(x) = f(x)$, $f^-(x) = 0$

$$\therefore (f^+ + f^-)(x) = f^+(x) + f^-(x) = f(x) + 0 = f(x) = |f(x)|$$

$$\Rightarrow f^+ + f^- = |f| \quad \text{(proved)}.$$

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

(V) $\max(f,g) = \dfrac{(f+g)+|f-g|}{2}$

$$|f-g| = \begin{cases} +(f-g), & \text{if } \geq 0 \\ -(f-g), & \text{if } < 0 \end{cases}$$

$$|f-g|(x) = |f(x)-g(x)| = \begin{cases} f(x)-g(x), & f(x) \geq g(x) \\ g(x)-f(x), & f(x) < g(x) \end{cases}$$

Case I: If $f(x) \geq g(x) \Rightarrow f(x)-g(x) \geq 0$

$\Rightarrow |f(x)-g(x)| = f(x)-g(x)$

$\therefore \dfrac{(f+g)+|f-g|}{2}(x) = \dfrac{f(x)+g(x)+|f(x)-g(x)|}{2}$

$= \dfrac{f(x)+g(x)+f(x)-g(x)}{2} = f(x)$

Note: $f(x) \geq g(x)$ then $\max(f,g) = f(x)$

$\Rightarrow \dfrac{(f+g)+|f-g|}{2} = \max(f,g)$

Case II: If $f(x) < g(x) \Rightarrow f(x)-g(x) < 0$

$\Rightarrow |f(x)-g(x)| = -(f(x)-g(x)) = g(x)-f(x)$

and if $f(x) < g(x)$ then $\max(f,g) = g(x)$

$\therefore \dfrac{(f+g)+|f-g|}{2}(x) = \dfrac{f(x)+g(x)+|f(x)-g(x)|}{2}$

$= \dfrac{f(x)+g(x)+g(x)-f(x)}{2} = g(x)$

$\Rightarrow \dfrac{(f+g)+|f-g|}{2} = \max(f,g)$ (proved)

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

(vi) $\min(f,g) = \dfrac{(f+g)-|f-g|}{2}$

$$|f-g|(x) = |f(x)-g(x)| = \begin{cases} f(x)-g(x) & ; f(x) \geq g(x) \\ g(x)-f(x) & ; f(x) < g(x) \end{cases}$$

Case I: If $f(x) \geq g(x) \Rightarrow f(x)-g(x) \geq 0$

$\Rightarrow |f(x)-g(x)| = f(x)-g(x)$ and

If $f(x) \geq g(x) \Rightarrow \min(f,g) = g(x)$

$\therefore \dfrac{(f+g)-|f-g|}{2}(x) = \dfrac{f(x)+g(x)-|f(x)-g(x)|}{2}$

$= \dfrac{f(x)+g(x)-(f(x)-g(x))}{2}$

$= \dfrac{f(x)+g(x)-f(x)+g(x)}{2} = g(x)$

$\Rightarrow \dfrac{(f+g)-|f-g|}{2} = \min(f,g)$

Case II: If $f(x) < g(x) \Rightarrow f(x)-g(x) < 0$

$\Rightarrow |f(x)-g(x)| = g(x)-f(x)$ and

If $f(x) < g(x) \Rightarrow \min(f,g) = f(x)$

$\therefore \dfrac{(f+g)-|f-g|}{2}(x) = \dfrac{f(x)+g(x)-|f(x)-g(x)|}{2}$

$= \dfrac{f(x)+g(x)-g(x)+f(x)}{2} = f(x)$

$\Rightarrow \dfrac{(f+g)-|f-g|}{2} = \min(f,g)$ (Proved).

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

Topic # 74

Characteristic Function:

Let $\phi \neq X$ and $A \subseteq X$, then characteristic function $\chi_A : X \to R$ is defined and given by:

$$\chi_A(x) = \begin{cases} 1 & \text{if } x \in A \\ 0 & \text{if } x \notin A \end{cases}$$

If $A = \phi$, $\Rightarrow \chi_A(x) = 0 = 0(x)$, $\forall x \in X$ (0(x): zero function)

$\Rightarrow \chi_\phi = 0$

And if $A = X \Rightarrow \chi_A(x) = 1 = 1(x)$ $\forall x \in X$ (1(x): Identity function)

$\Rightarrow \chi_X = 1$

Topic # 75

Theorem: Prove that if $A \subseteq B$, then $\chi_A \leq \chi_B$.

Proof: Case I; If $x \in A$ and

$A \subseteq B \Rightarrow x \in B$

$\Rightarrow \chi_A = 1 = \chi_B \Rightarrow \chi_A \leq \chi_B$



Case II; If $x \in B$ but $x \notin A$

$\Rightarrow \chi_A = 0$ and $\chi_B = 1$

$\Rightarrow \chi_A \leq \chi_B$

Case III; If $x \notin A$ nor $x \notin B$

$\Rightarrow \chi_A = 0 = \chi_B \Rightarrow \chi_A \leq \chi_B$

(proved)



This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

Topic # 76

Theorem: Show that $\chi_{A\cap B} = \chi_A \cdot \chi_B$ 1

Proof: If $x \in A \cap B \Rightarrow x \in A$ and $x \in B$

$\Rightarrow \chi_{A\cap B} = 1 = \chi_A = \chi_B = 1 \cdot 1 = \chi_A \cdot \chi_B$

If $x \notin A\cap B \Rightarrow$ {
- $x \in A$ but $x \notin B \Rightarrow \chi_A = 1, \chi_B = 0$
- $x \in B$ but $x \notin A \Rightarrow \chi_B = 1, \chi_A = 0$
- $x \notin A$ and $x \notin B \Rightarrow \chi_A = \chi_B = 0$

$\therefore \chi_A \cdot \chi_B = 0 = O(x)$ }

$x \notin A\cap B \Rightarrow \chi_{A\cap B} = 0$

For all, $x \notin A\cap B$; $\chi_{A\cap B} = 0 = O(x) = \chi_A \cdot \chi_B$

$\chi_{A\cap B} = \chi_A \cdot \chi_B$ (proved)

Topic # 77

Theorem: Show that $\chi_{A-B} = \chi_A - \chi_{A\cap B}$.

Proof: Case I: If $x \in A - B$

$\Rightarrow x \in A$ but $x \notin B, x \notin A\cap B$

$\Rightarrow \chi_{A-B} = 1$



$\chi_A - \chi_{A\cap B} = 1 - 0 = 1 = \chi_{A-B}$

Case II If $x \notin A - B \Rightarrow$ {
- $x \notin A, x \notin B, x \notin A\cap B \longrightarrow$ (a)
- $x \notin A, x \in B, x \notin A\cap B \longrightarrow$ (b)
- $x \in A, x \in B, x \in A\cap B \longrightarrow$ (c)

$\Rightarrow \chi_{A-B} = 0 - 0(x)$

$\chi_A - \chi_{A\cap B} =$ {
- (a) $\Rightarrow 0 - 0 = 0$
- (b) $\Rightarrow 0 - 0 = 0$
- (c) $\Rightarrow 1 - 1 = 0$

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

$\therefore$ For all cases; $\chi_{A-B} = \chi_A - \chi_{A\cap B}$ (proved)

Topic #78

Theorem: Show that $\chi_{A\times B} = \chi_A \cdot \chi_B$

Proof: $A\times B = \{(x,y) : x\in A, y\in B\}$

Case I:

$(x,y) \in A\times B \Rightarrow \begin{cases} x\in A \text{ and } x\in B \\ \Rightarrow \chi_A = 1 = \chi_B \end{cases}$

$\Rightarrow \chi_{A\times B}(x,y) = 1$

$\therefore \chi_{A\times B} = \chi_A \cdot \chi_B$

Case II: $\begin{cases} x\in A \text{ but } y\notin B \rightarrow (a) \\ x\notin A \text{ but } y\in B \rightarrow (b) \\ x\notin A \text{ and } y\notin B \rightarrow (c) \end{cases}$

If $(x,y) \notin A\times B \Rightarrow$

$\Rightarrow \chi_{A\times B}(x,y) = 0$

Also: $\chi_A \cdot \chi_B = \begin{cases} (a) \Rightarrow 1\cdot 0 = 0 \\ (b) \Rightarrow 0\cdot 1 = 0 \\ (c) \Rightarrow 0\cdot 0 = 0 \end{cases}$

In all cases: $\chi_{A\times B} = \chi_A \cdot \chi_B$ (proved)

Topic #79

Theorem: If $A = \bigcup_{n=1}^{\infty} A_n$ and $\{A_n\}_{n=1}^{\infty}$ is a sequence of pairwise disjoint subsets of X. Then show that $\chi_A = \sum_{n=1}^{\infty} \chi_{A_n}$

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

Case I:

Proof: If $x \in A \Rightarrow x \in \bigcup_{n=1}^{\infty} A_n$

$\Rightarrow x \in A_k$ exactly for $n=k$



$x \notin \bigcup_{n=1}^{\infty} A_n - \{A_k\}$

$\Rightarrow \chi_{A_k}(x) = 1, \chi_{A_n}(x) = 0, \forall n \neq k$

$x \in A \Rightarrow \chi_A(x) = 1$

$\therefore \chi_A(x) = 1 = 0+0+\cdots+\underset{n=k}{1}+0+0+\cdots$

$\chi_A(x) = 1 = \sum_{n=1}^{\infty} \chi_{A_n}(x) \Rightarrow \chi_A = \sum_{n=1}^{\infty} \chi_{A_n}$

Case II: $x \notin A = \bigcup_{n=1}^{\infty} A_n \Rightarrow x \notin A_n, \forall n \in \mathbb{N}$

$\Rightarrow \chi_A(x) = 0$ $\left\{ \Rightarrow \chi_{A_n}(x) = 0, \forall n \in \mathbb{N} \right.$

$= 0+0+0+\cdots+0+\cdots = \sum_{n=1}^{\infty} \chi_{A_n}(x)$

$\Rightarrow \chi_A = \sum_{n=1}^{\infty} \chi_{A_n}$ (proved)

Topic # 80

Theorem: Let $\phi \neq A \subseteq X$ and for characteristic function $\chi_A : X \to \mathbb{R}$, χ_A is measurable $\Leftrightarrow$ "A" is measurable.

Proof: Suppose "χ_A" is measurable (function)

$\Rightarrow \forall a \in \mathbb{R}$ is set; $\{x : \chi_A(x) > a\}$ is measurable.

In particular for $a = 0$; $\{x : \chi_A(x) > 0\} = A$

By definition of χ_A.

$\therefore A \in M$ (measurable)

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

Conversely, let $A \in M$ (measurable) and we prove "χ_A" is measurable.

In other words we want to prove the $\{x : \chi_A(x) \geq a\} \in M, \forall\ a \in R$

$$\{x : \chi_A(x) \geq a\} = \begin{cases} \phi & \text{if } a \geq 1 \\ A & \text{if } 0 \leq a < 1 \\ A \cup A^c & \text{if } a < 0 \end{cases}$$

(i) $a \geq 1$, $\chi_A(x) \geq 1 \Rightarrow$ there does not exist $x \in A$ such that $\chi_A(x) \geq 1$

(ii) $0 \leq a < 1$, $\chi_A(x) > 0.9$ (say)

(iii) $a < 0$ (say -2), $\chi_A(x) > -2$

In all cases; if $A \in M$ (measurable)

$\Rightarrow A^c \in M$ (measurable)

$\Rightarrow A \cup A^c \in M$ (measurable)

$\phi \in M$ (measurable)

$\Rightarrow \{x : \chi_A(x) \geq a\} \in M$ (measurable)

So "χ_A" is measurable. (Proved)

Topic # 81

Algebra of Measurable Functions:

If "f" and "g" are two extended real valued and measurable functions defined on same domain $\phi \neq D$. then (i) $f + k$, (ii) kf, (iii) $f \pm g$, (iv) fg, (v) $\frac{f}{g}, g \neq 0$, (vi) $|f|$, (vii) $f \vee g$ (viii) $f \wedge g$ are measurable functions.

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

Topic # 82 & 83

Proof: is "$f+k$" is measurable, $k \in R$

$\because$ "f" is measurable $\Rightarrow \{x : f(x) > a\} \in M, \forall a \in R$

$\Rightarrow \{x : f(x) > a-k\} \in M$ (measurable)

$\Rightarrow \{x : f(x)+k > a\} \in M, \forall a \in R$

$\Rightarrow$ "$f+k$" is measurable.

83 (ii) "kf" is measurable, $\forall k \in R$ | $f \to kf$ Scaler compositon

proof: $\because$ "f" is measurable.

$\Rightarrow \{x : f(x) > q\} \in M, \forall a \in R$

(a) If $k > 0$: $\because f(x) > q, \forall x \in R$

$\Rightarrow f(x) > \frac{q}{k} \Rightarrow kf(x) > q, \forall a \in R$

$\therefore \{x : kf(x) > a\} \in M \ \forall a \in R$

$\Rightarrow$ "kf" is measurable.

(b) If $k < 0$; $f(x) > a$, $\forall a \in R$ (given)

$\Rightarrow f(x) > \frac{a}{k} \Rightarrow kf(x) < a \quad (\because k < 0)$

$\Rightarrow \{x : kf(x) < a\} \in M$

$\Rightarrow$ "kf" is measurable.

(c) If $k = 0 \Rightarrow \{x : kf(x) > a\}, \forall a \in R$

$= \{x : 0 f(x) > a\} \because \forall a \in R$

$= \{x : 0 > a\}, \forall a \in R$ ... (1)

$$\{x : 0 > a\} = \begin{cases} D & ; \ a < 0 \quad (\exists x : f(x) > a) \\ \phi & ; \ a \geq 0 \end{cases}$$

"D" is measurable (given) and ϕ is measurable.

This PDF document was edited with Icecream PDF Editor. Upgrade to PRO to remove watermark.

Scanned with CamScanner

$\therefore \{x : |kf(x)| > a\}$ is measurable.

$\Rightarrow$ "kf" is measurable.

Topic # 84

(iii) "$f+g$" is measurable.

proof: We have to proof "$f+g$" is measurable.

$\Rightarrow \{x : (f+g)(x) > a\} \in M$ (To prove)

$\{x : (f+g)(x) > a\} = \{x : f(x) + g(x) > a\}$

$= \{x : f(x) > a - g(x)\}$, $\forall\, a \in R$

$\therefore \{f(x), a - g(x) \in R \Rightarrow \exists\, p \in Q$ such that

$f(x) > p > a - g(x)\}$

because "Q" is dense in R : $\bar{Q} = R$

$\therefore \{x : (f+g)(x) > a\} = \bigcap_{p \in Q} \{x : f(x) > p\} \cap \{x : a - g(x) < p\}$

i.e. our claim.

Let $y \in \{x : (f+g)(x) > a\} \Rightarrow y \in \{x : f(x) + g(x) > a\}$

$\Rightarrow f(y) + g(y) > a \Rightarrow f(y) > a - g(y)$

$\Rightarrow$ there exist($\exists$) $p \in Q$ such that $f(y) > p > a - g(y)$

$\bar{Q} = R$

$\Rightarrow f(y) > p$ and $a - g(y) < p$.

$\Rightarrow y \in \{x : f(x) > p\}$ and $y \in \{x : a - g(x) < p\}$

$\Rightarrow y \in \{x : f(x) > p\} \cap \{x : a - g(x) < p\}$

$\Rightarrow y \in \bigcup_{p \in Q} [\{x : f(x) > p\} \cap \{x : a - g(x) < p\}]$

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

$$\Rightarrow \{x : (f+g)(x) > a\} \subseteq \bigcup_{p \in Q} \left[\{x : f(x) > p\} \cap \{x : a - g(x) < p\}\right] \longrightarrow (A)$$

Now Let $t \in \bigcup_{p \in Q} \left[\{x : f(x) > p\} \cap \{x : a - g(x) < p\}\right]$

$\Rightarrow t \in \{x : f(x) > p\} \cap \{x : a - g(x) < p\}$ for some $p \in Q$

$\Rightarrow t \in \{x : f(x) > p\}$ and $t \in \{x : a - g(x) < p\}$

$\Rightarrow f(t) > p$ and $a - g(t) < p \Rightarrow p > a - g(t)$

$\Rightarrow f(t) > p > a - g(t) \Rightarrow f(t) > a - g(t)$

$\Rightarrow f(t) + g(t) > a \rightarrow (f+g)(t) > a$

$\Rightarrow t \in \{x : (f+g)(x) > a\}$

$\Rightarrow \bigcup_{p \in Q} \left[\{x : f(x) > p\} \cap \{x : a - g(x) < p\}\right] \subseteq \{x : (f+g)(x) > a\} \longrightarrow (B)$

From (A) and (B) $\rightarrow \{x : (f+g)(x) > a\} =$

$$= \bigcup_{p \in Q} \left[\{x : f(x) > p\} \cap \{x : a - g(x) < p\}\right]$$

$\therefore$ $\begin{cases} \text{"}f\text{" is measurable} \Rightarrow \{x : f(x) > p\} \in M \\ \text{"}g\text{" is measurable} \Rightarrow \{x : a - g(x) < p\} \in M \end{cases}$

$\Rightarrow \{x : f(x) > p\} \cap \{x : a - g(x) < p\} \in M$, for $p \in Q$

$\Rightarrow \bigcup_{p \in Q} \left[\{x : f(x) > p\} \cap \{x : a - g(x) < p\}\right] \in M$

$\Rightarrow \{x : (f+g)(x) > a\} \in M$ (measurable)

$\rightarrow$ "$f+g$" is measurable function.

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

Now we have to prove "f-g" is measurable.

⇒ If g is measurable ⇒ (-1)g = -g is also M.

⇒ f + (-1)g = f - g is also measurable.

(proved)

Topic#85

(iv) "fg" is measurable.

proof: $fg = \frac{1}{4}[(f+g)^2 - (f-g)^2]$

Firstly we prove "f^2" is measurable.

⇒ we need to prove $\{x : f^2(x) > a\} \in M, \forall a \in \mathbb{R}$

$y = x^2$ (Say function)

$y = a$ (line)

$\Rightarrow y^2 = a^2$

$y = x^2$

$a^2 = x^2$

$\Rightarrow x = \pm a$



Case 2: $f^2(x) > a$

$\Rightarrow f(x) > a$, $f(x) < -a$

$\{x : f^2(x) > a\} = \begin{cases} \{x : f(x) > \sqrt{a}\} \cup \{x : f(x) < -\sqrt{a}\}, & a \geq 0 \\ \text{All Domain} \cdot (D) & , a < 0 \end{cases}$

$\because$ "f" is measurable $\Rightarrow \{x : f(x) > \sqrt{a}\} \in M$

and $\{x : f(x) < -\sqrt{a}\} \in M$

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

$\Rightarrow \{x: f(x) > \sqrt{q}\} \cup \{x: f(x) < -\sqrt{q}\} \in M$

and "Domain" is given to be measurable. i.e $D \in M$

$\therefore \{x: f^2(x) < a\} \in M \Rightarrow$ "f^2" is measurable.

$\because f, g$ is measurable $\Rightarrow (f \pm g)$ is also measurable.

$\Rightarrow (f \pm g)^2$ is measurable.

$\Rightarrow \frac{1}{4}[(f+g)^2 - (f-g)^2] = fg$ is also measurable.

($\because Kf$ is also measurable) $\forall K \in R$.

Topic # 86

prove that if "f" is measurable function, then "$\frac{1}{f}$" is also measurable.

Proof: "f" is measurable $\Rightarrow \{x: f(x) > a\} \in M, \forall a \in R$

To prove "$\frac{1}{f}$" is measurable, we show that

$\{x: \frac{1}{f}(x) > a\} \in M, \forall a \in R$.

$$\frac{1}{f}(x) = \frac{1}{f(x)} > a$$

$$\left\{x: \frac{1}{f}(x) > a\right\} = \begin{cases} \{x: f(x) < \frac{1}{a}\} \cap \{x: f(x) > 0\}; & a > 0 \\ \{x \in D: f(x) > 0\}; & a = 0 \\ \{x: f(x) > 0\} \cup [\{x: f(x) < \frac{1}{a}\} \cap \{x: f(x) < a\}]; & a < 0 \\ \phi \end{cases}$$

All the sets on R.H.S are measurable.

$\Rightarrow \{x: \frac{1}{f}(x) > a\}$ is also measurable, $\forall a \in R$.

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

$\Rightarrow$ $\frac{1}{f}$ is measurable.

(v) $\frac{f}{g}$, $g \neq 0$, is measurable.

proof: $\frac{f}{g}$ is also measurable because $\frac{f}{g} = \frac{1}{g} \cdot f$ is measurable being product of two measurable functions.

Topic # 87

(vi) $|f|$ is measurable.

proof: $\because$ "f" is measurable $\Rightarrow \{x : f(x) > a\} \in M, \forall a \in R$

To prove $|f|$ is measurable, we need to show:

$\{x : |f|(x) > a\} \in M, \forall a \in R$

$$\{x : |f|(x) > a\} = \{x : |f(x)| > a\}$$

$$= \begin{cases} \underbrace{\{x : f(x) > a\}}_{\in M} \cup \underbrace{\{x : f(x) < -a\}}_{\in M}, & \forall a > 0 \\ \{x : |f|(x) > a\} = \\ \emptyset \in M, & a < 0 \end{cases}$$

In all cases our R.H.S, we get a measurable set. $\Rightarrow \{x : |f|(x) > a\} \in M$ (measurable), $\forall a \in R$

$\Rightarrow |f|$ is also measurable.

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

(vii) $f \vee g = \max(f, g)$ is measurable.

proof: $\because f \vee g = \max(f, g) = \frac{1}{2}[(f+g) + |f-g|]$

So $f \vee g$ is measurable because $(f+g)$ and $|f-g|$ is measurable and "kf" is measurable, $\forall\ k \in \mathbb{R}$.

(viii) $f \wedge g = \min(f, g)$ is measurable.

proof: $\because f \wedge g = \min(f, g) = \frac{1}{2}[(f+g) - |f-g|]$

So $f \wedge g$ is measurable because $(f+g)$ and $|f-g|$ is measurable and "kf" is measurable, $\forall\ k \in \mathbb{R}$.

Topic # 88

Restriction of a function:

Let $f: X \to \mathbb{Y}$ be a function and $A \subset X$, Then $g: A \to \mathbb{Y}$ defined as $g(x) = f(x)$, $x \in A$ is called the restriction of "f" to $A \subset X$ and given by $f_A = g$.

Theorem: Let "f" be an extended real valued measurable function on "D". Let $A \subset D$, Then restriction of "f" to "A" is also measurable and $A \in \mathcal{M}$.

Proof: $\because$ "f" is measurable $\Rightarrow \{x \in D : f(x) > a\} \in \mathcal{M}$. $\forall\ a \in \mathbb{R}$

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

For any $a \in \mathbb{R}$, To prove "f_A" is measurable we have $\{x \in A : f_A(x) > a\} \in M$

Note: $\{x \in A : f_A(x) > a\} = \{x \in A : f(x) > a\} \cap A$

On R.H.S both the sets are measurable.

$\Rightarrow \{x \in A : f_A(x) > a\} \in M$

$\because$ Intersection of two measurable sets is measurable.

$\Rightarrow$ "f_A" is measurable.

Topic # 89

Theorem: An extended real valued function defined on a set of measure zero is measurable.

Proof: Let $\phi \neq E \subset \mathbb{R}$, such that $m^*(E) = 0$ and $f : E \to \mathbb{R} \cup \{\pm\infty\}$, To prove "$f$" is measurable, we need to show that:

$\{x \in E : f(x) > a\} \in M$, $\forall\ a \in \mathbb{R}$.

$\because \{x \in E : f(x) > a\} \subset E$

$\Rightarrow m^*\{x \in E : f(x) > a\} \leq m^*(E)$

$\because$ By monotonicity of m^*.

$\Rightarrow m^*\{x \in E : f(x) > a\} \leq 0$ ..... (given)

But by non-negativity of "m^*"

$\Rightarrow m^*\{x \in E : f(x) > a\} \geq 0$

$\Rightarrow m^*\{x \in E : f(x) > a\} = 0$; $\forall\ a \in \mathbb{R}$.

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

$\Rightarrow \{x \in E : f(x) > a\} \in M \; ; \forall \; a \in \mathbb{R}$

So "f" is measurable.

Topic # 90

Theorem: Let $\{f_n\}$ be a sequence of extended real valued measurable function on "D", then.

(i) $\sup_{n \in \mathbb{N}} f_n$, $\inf_{n \in \mathbb{N}} f_n$ are measurable.

(ii) $\max_{1 \le i \le n} (f_i)$, $\min_{1 \le i \le n} (f_i)$ are measurable.

(iii) $\overline{\lim_{n \in \mathbb{N}}} f_n$, $\underline{\lim_{n \in \mathbb{N}}} f_n$ are measurable.

(iv) If $\lim_{n \to \infty} f_n = f$, Then "f" is measurable.

Proof: Let $f = \inf_{n \in \mathbb{N}} f_n$ and To prove f is measurable Then we have to shows $\{x : f(x) > a\} \in M, \forall a \in \mathbb{R}$

We claim : $\{x : f(x) > a\} = \bigcup_{n=1}^{\infty} \{x : f_n(x) > a\}$

Let $y \in \{x : f(x) > a\} \Rightarrow f(y) > a$

$\because f = \inf_{n \in \mathbb{N}} f_n \Rightarrow f(y) \le f_n(y), \forall n \in \mathbb{N}$

$\because f(y) > a \Rightarrow a < f(y) \le f_n(y), \forall n \in \mathbb{N}$

$\Rightarrow f_n(y) > a \; \forall \; n \in \mathbb{N} \Rightarrow y \in \{x : f_n(x) > a\} \; \forall n \in \mathbb{N}$

$\Rightarrow y \in \bigcap_{n=1}^{\infty} \{x : f_n(x) > a\}$

$\Rightarrow \{x : f(x) > a\} \subseteq \bigcap_{n=1}^{\infty} \{x : f_n(x) > a\} \longrightarrow (A)$

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

Now let $t \in \bigcap_{n=1}^{\infty} \{x : f_n(x) > a\}$

$\Rightarrow t \in \{x : f_n(x) > a\}, \forall a \in R, \forall n \in N$

$\Rightarrow f_n(t) > a, \forall n \in N$

$\Rightarrow a < f_n(t), \forall n \in N$

$\Rightarrow$ "a" is lower bound $\{f_1(t), f_2(t), \cdots\}$

But $f = \inf_{n \in N} f_n$ is also its lower bound.

$\Rightarrow a < \inf_{n \in N} f_n, \forall n \in N$

$\Rightarrow f(t) > a \Rightarrow \inf_{n \in N} f_n > a$

$f(t) > a \Rightarrow t \in \{x : f(t) > a\} \longrightarrow (A)$

$\Rightarrow \bigcap_{n=1}^{\infty} \{x : f_n(x) > a\} \subset \{x : f(x) > a\} \longrightarrow (B)$

From (A) and (B) $\{x : f(x) > a\} = \bigcap_{n=1}^{\infty} \{x : f_n(x) > a\}$

So $f = \inf_{n \in N} f_n$ is measurable.

Now let $h = \sup_{n \in N} f_n$

... The set $\{x : h(x) > a\} = \bigcup_{n=1}^{\infty} \{x : f_n(x) > a\}$ is measurable, So $h = \sup_{n \in N} f_n$ is measurable.

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

Topic# 9

$\max_{1\le i\le n} f_i$, $\min_{1\le i\le n} f_i$ are measurable.

$\{f_n\}_{n=1}^{\infty}$ is seq of measurable functions.

Proof: Let $f = \max_{1\le i\le n} f_i$, To prove "f" is measurable, we need to show that $\{x : f(x) > a\} \in M$ and we claim

$$\{x : f(x) > a\} = \bigcup_{k=1}^{n} \{x : f_k(x) > a\} \quad \forall a \in \mathbb{R}.$$

let $y \in \{x : f(x) > a\} \Rightarrow f(y) > a$

$\therefore f(x) = \max_{1\le i\le n} f_i$ So there exist $J = \{1, 2, \dots, n\}$

$\Rightarrow f(x) = f_j(x) \Rightarrow f(y) = f_j(y) > a$

$\Rightarrow y \in \{x : f_j(x) > a\}$, for some $j = \{1, 2, \dots, n\}$

Hence $y \in \bigcup_{i=1}^{n} \{x : f_i(x) > a\}$

Now let $y \in \bigcup_{i=1}^{n} \{x : f_i(x) > a\}$

Then $y \in \{x : f_k(x) > a\}$ for some $k = \{1, 2, \dots, n\}$

$\Rightarrow f_k(y) > a$, By definition $f_i(y) \le \max f_i(y) = f(y)$ $\forall i = 1, 2, \dots, n.$

and $f_k(y) \le f(y) \Rightarrow h(y) > a$

$\Rightarrow y \in \{x : h(x) > a\}$.

The countable union of measurable sets is measurable. So $f = \max_{1\le i\le n} f_i$ is measurable.

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

$\overline{\lim}_{n\in\mathbb{N}}$ → Limit superior

$\underline{\lim}_{n\in\mathbb{N}}$ → Limit inferior

Let $g = \min_{1\le i\le n} f_i$, To prove "g" is measurable we need to show that $\{x : g(x) > a\} \in M \quad \forall a \in \mathbb{R}$

and we claim $\{x : g(x) > a\} = \bigcap_{i=1}^{n} \{x : f_i(x) > a\}$

The countable intersection of measurable sets is measurable. So $g = \min_{1\le i\le n} f_i$ is measurable.

Topic #92

Show that $\overline{\lim}_{n\in\mathbb{N}} f_n, \underline{\lim}_{n\in\mathbb{N}} f_n$ are measurables.

Proof:

$\overline{\lim}_{n\in\mathbb{N}} f_n = \inf_{n\in\mathbb{N}} \left(\sup_{k\ge n} f_k\right) \rightarrow ①$

$\sup_{k\ge n} f_k = \sup\{f_k, f_{k+1}, f_{k+2}, \cdots\}, k = 1, 2, 3, \cdots$

$= \begin{cases} \sup\{f_1, f_2, f_3, \cdots\} = F_1 \text{ is measurable being "sup" of measurable functions.} \\ \sup\{f_2, f_3, f_4, \cdots\} = F_2, \cdots \end{cases}$

$\Rightarrow \{F_n\}_{n=1}^{\infty}$ is a sequence of measurable functions

$\Rightarrow \inf_{n\in\mathbb{N}} \{F_n\}_{n=1}^{\infty}$ is also measurable

$\Rightarrow \overline{\lim}_{n\in\mathbb{N}} f_n = \inf_{n\in\mathbb{N}} \{f_n\} = \inf_{n\in\mathbb{N}} (\sup_{k\ge n} f_k)$ is also measurable

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner

Similarly $\underline{\lim}_{n\in\mathbb{N}} f_n = \sup_{n\in\mathbb{N}} (\inf_{k\geq n} f)$ is measurable.

Topic # 93

If $\lim_{n\to\infty} f_n = f$, then "f" is measurable.

Proof: $\because \lim_{n\to\infty} f_n = \overline{\lim}_{n\in\mathbb{N}} f_n = \underline{\lim}_{n\in\mathbb{N}} f_n = f$ $\{\{f_n\}_{n=1}^{\infty}$ ↓ Limit (given)

But $\overline{\lim}_{n\in\mathbb{N}} f_n$ and $\underline{\lim}_{n\in\mathbb{N}} f_n$ are measurable.

$\Rightarrow$ "f" is also measurable.

MTH644 – Measure Theory Handouts

Written by Mahar Afaq Safdar Muhammadi

REGARDS: VIRTUAL ALERTS

ہمیشہ دعاؤں میں یاد رکھیے گا۔ جزاک اللہ

This PDF document was edited with Icecream PDF Editor.
Upgrade to PRO to remove watermark.

Scanned with CamScanner