

# MTH-603 Numerical Analysis Update MCQS For Quiz-1 File Solve By Vu Topper RM

85% To 100% Marks



وَتَعَزُّ مِنْ تَشَاءٍ وَتَذِلُّ مِنْ تَشَاءٍ

PROFESSIONAL ONLINE ACADEMY



**We Offers**

LMS Handling

Important Notes

Online Classes

Assignments

Quiz & GDB's

Projects

**NOTHING  
IS  
IMPOSSIBLE**

**Join Us  
Now**


For More Info

Contact us at:

**Rizwan Manzoor**



**0322-4021365**



بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

For More Help Contact What's app 03224021365

While using the Gauss-Seidel Method for finding the solution of the following system  $2x+2y+z=3$ ,  $x+3y+z=2$ ,  $x+y+z=2$  with the initial guess  $(0,0,0)$ , the next iteration would be

- A.  $(32, -23, -2)$
- B.  $(32, 57, 19)$
- C.  $(32, 16, 13)$**
- D.  $(32, 23, 2)$

Which of the following system of equation is diagonally dominant?

- A.  $8x+y-z=5$ ,  $2x+13y+5z=7$ ,  $x+2y+9z=3$**
- B.  $8x+y-z=5$ ,  $x+2y+9z=3$ ,  $2x+13y+5z=7$
- C.  $2x+13y+5z=7$ ,  $8x+y-z=5$ ,  $x+2y+9z=3$
- D.  $x+2y+9z=3$ ,  $8x+y-z=5$ ,  $2x+13y+5z=7$

Which of the following systems of linear equations has a strictly diagonally dominant coefficient matrix?

- A.  $-2x_1+7x_2+2x_3=5$ ,  $6x_1-2x_2+3x_3=1$ ,  $x_1+x_2-5x_3=-13$
- B.  $-2x_1+7x_2+2x_3=5$ ,  $6x_1-2x_2+3x_3=1$ ,  $x_1+x_2-5x_3=-13$
- C.  $6x_1-2x_2+3x_3=1$ ,  $-2x_1+7x_2+2x_3=5$ ,  $x_1+x_2-5x_3=-13$
- D.  $x_1+x_2-5x_3=-13$ ,  $6x_1-2x_2+3x_3=1$ ,  $-2x_1+7x_2+2x_3=5$**

While solving a system of three linear equations in three variables by Jacobi's method, the first approximate solution is NECESSARY to be taken as  $(0,0,0)$ .

- A. True
- B. False**

While using the Gauss-Seidel Method for finding the solution of the system of equation, the following system

$x+2y+2z=3$ ,  $x+3y+3z=2$ ,  $x+y+5z=2$  can be rewritten as

- A.  $x=3-2y-2z$ ,  $y=2-3x-2z$ ,  $z=2-5x-5y$
- B.  $x=3-2y-2z$ ,  $y=2-3x-2z$ ,  $z=2-5x-5y$**
- C.  $x=1-23y-23z$ ,  $y=1-32x-32z$ ,  $z=1-y-2x$

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

For More Help Contact What's app 03224021365

D.  $x=3-2y-2zy=2-x-3zz=2-y-x$

While solving the system;  $x+4y = 4$ ,  $2x-y = 1$  by Gauss-Seidel method, which of the following ordering will lead more rapidly to the desired order of accuracy?

A.  $2x-y = 1$ ,  $x+4y = 4$

B. no need to reordering

**C.  $x+4y = 1$ ,  $2x-y = 4$**

D.  $x-2y = 1$ ,  $x+4y = 4$

In Gauss-Jacobi's method, the corresponding elements of  $x(r+1)$  replaces those of  $x(r)$  as soon as they become available.

A. True

**B. False**

An augmented matrix may also be used to find the inverse of a matrix by combining it with the ..... matrix.

A. Inverse

B. None of the given choices

**C. Identity**

D. Square

The system of linear equations;  $2x+3y = 6$ ,  $x+4y = 5$  is not strictly diagonal dominant as-----

A.  $|2| < |3|$

**B.  $|2| < |4|$**

C.  $|2| < |1|$

D.  $|3| < |4|$

While using the Gauss-Seidel Method for finding the solution of the following system  $3x+y+z=11$ ,  $2x+5y-z=16$ ,  $x+y+5z=4$  with initial guess  $(0,0,0)$ , the next iteration would be

A.  $(113,165,-45)$

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

- B. (113, -165, -45)
- C. (113, 2615, -725)
- D. (113, 511, 213)**

While solving a system of linear equations by Gauss Jordan Method, in which of the following case the partial pivoting is essential?

- A. When lower triangular entries are zero
- B. When pivots are zero
- C. When the upper triangular entries are zero
- D. When diagonal entries are zeros**

Which of the following system of equation is diagonally dominant

- A.  $3x+y+z=3$ ,  $x+y+5z=2$ ,  $2x+5y-z=4$
- B.  $x+y+5z=2$ ,  $3x+y+z=3$ ,  $2x+5y-z=4$**
- C.  $3x+y+z=3$ ,  $2x+5y-z=4$ ,  $x+y+5z=2$
- D.  $2x+5y-z=4$ ,  $3x+y+z=3$ ,  $x+y+5z=2$

In ..... method, matrix [A] of the system of equations is decomposed into the product of two matrices [L] and [U], where [L] is a lower-triangular matrix and [U] is an upper-triangular matrix with 1's on its main diagonal.

- A. Gauss-Seidel
- B. Crout's Reduction**
- C. None of the given choices
- D. Gaussian elimination

Which of the following systems of linear equations has a strictly diagonally dominant coefficient matrix?

- A.  $7x_1+5x_2-1x_3=9$ ,  $-x_1+3x_2+x_3=1$ ,  $x_1+2x_2+4x_3=1$
- B.  $4x_1+2x_2+3x_3=1$ ,  $7x_1+5x_2-8x_3=9$ ,  $-5x_1+3x_2+2x_3=1$**
- C.  $4x_1+2x_2+3x_3=1$ ,  $-5x_1+3x_2+2x_3=1$ ,  $7x_1+5x_2-8x_3=9$
- D.  $-5x_1+3x_2+2x_3=1$ ,  $7x_1+5x_2-8x_3=9$ ,  $4x_1+2x_2+3x_3=1$

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

Which of the following rearrangement make strictly diagonal dominant, the system of linear equations;  $x-3y+z = -2$ ,  $-6x+4y+11z=1$ ,  $5x-2y-2z=9$ ?

- A.  $5x-2y-2z=9$ ,  $x-3y+z = -2$ ,  $-6x+4y+11z=1$
- B.  $5x-2y-2z=9$ ,  $-6x+4y+11z=1$ ,  $x-3y+z = -2$
- C. No need to rearrange as system is already in diagonal dominant form.
- D.  $-6x+4y+11z=1$ ,  $x-3y+z = -2$ ,  $5x-2y-2z=9$**

How many Eigen values will exist corresponding to the function;  $\text{Exp}(ax) = e^{ax}$ , when the matrix operator is of differentiation?

- A. Infinite many
- B. Unique**
- C. None
- D. Finite Multiple

Jacobi's method is highly recommended for ..... matrix to compute all the Eigen values and the corresponding eigenvectors.

- A. None of the given choices
- B. non real symmetric
- C. real un-symmetric
- D. real symmetric**

For a function;  $y=f(x)$ , if  $y_0$ ,  $y_1$  and  $y_2$  are 2,3 and 5 respectively then which of the following will be 2nd order Leading difference at  $y_0 = 2$  ?

- A. 2
- B. 1**
- C. -2
- D. -1

Direct methods can be more rapid than iterative algorithms.

- A. true**
- B. false

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

If the product of two matrices is an identity matrices that is  $AB = I$ , then which of the following is true?

- A. B is singular
- B. A is singular
- C. A is inverse of B**
- D. A is transpose of B

A and its transpose matrix have \_\_\_\_\_ eigenvalues.

- A. same**
- B. different

For differences methods we require the initial approximation.

- A. true**
- B. false

**Question No:1**

**(Marks:1)**

**Vu-Topper RM**

In Jacobi's Method, We assume that the .....elements does not vanish.

- A. Diagonal**
- B. Off-diagonal
- C. Row
- D. Column

ok

**Question No:2**

**(Marks:1)**

**Vu-Topper RM**

While solving a system of linear equations, which of the following approach is economical for the computer memory?

- A. Direct
- B. Iterative**
- C. Analytical
- D. Graphical

Page 69

ok

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

**Question No:3**

**(Marks:1)**

**Vu-Topper RM**

Back substitution procedure is used in .....

**A. Gaussian Elimination Method**

B. Jacobi's method

C. Gauss-Seidel method

D. None of the given choices

**Question No:4**

**(Marks:1)**

**Vu-Topper RM**

If  $n \times n$  matrices A and B are similar, then they have the same eigenvalues (with the same multiplicities).

**A. True**      **ok**

B. False

C. Both

D. None

**Question No:5**

**(Marks:1)**

**Vu-Topper RM**

The basic idea of relaxation method is to reduce the largest residual to .....

A. One

B. Five

C. Two

**D. Zero**

**Page 83**

**ok**

**Question No:6**

**(Marks:1)**

**Vu-Topper RM**

The Jacobi's method is a method of solving a matrix equation on a matrix that has no zeros along its \_\_\_\_\_.

**A. Main diagonal**

**Page 104**

**ok**

B. last column

C. last row

D. first row

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

**Question No:7**

**(Marks:1)**

**Vu-Topper RM**

If A is a nxn triangular matrix (upper triangular, lower triangular) or diagonal matrix, the eigenvalues of A are the diagonal entries of A.

**A. TRUE**

B. FALSE

**Question No:8**

**(Marks:1)**

**Vu-Topper RM**

A 3 x 3 identity matrix have three and different eigen values.

A. TRUE

**B. FALSE**

**Question No:9**

**(Marks:1)**

**Vu-Topper RM**

Which of the following is a reason due to which the LU decomposition of the system of linear equations;  $x+y = 1$ ,  $x+y = 2$  is not possible?

A. Associated coefficient matrix is singular

B. All values of l's and u's can't be evaluated

C. Determinant of coefficient matrix is zero

**D. All are equivalent** ok

**Question No:10**

**(Marks:1)**

**Vu-Topper RM**

Gauss - Jordan Method is similar to .....

A. Gauss-Seidel method

B. Iteration's method

C. Relaxation Method

**D. Gaussian elimination method**

Page 95

ok

**Question No:11**

**(Marks:1)**

**Vu-Topper RM**

While using Relaxation method, which of the following is the largest Residual for 1st iteration on the system;  $2x+3y = 1$ ,  $3x + 2y = -4$  ?

A. -4

**B. 3** ok

C. 2

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

D. 1

**Question No:12**

**(Marks:1)**

**Vu-Topper RM**

Gauss–Seidel method is also known as method of .....

**A. Successive displacement**

**Page 263**

**ok**

B. Iterations

C. False position

D. None of the given choices

**Question No:13**

**(Marks:1)**

**Vu-Topper RM**

Jacobi's Method is a/an.....

**A. Iterative method**

**Page 69**

**ok**

B. Direct method

**Question No:14**

**(Marks:1)**

**Vu-Topper RM**

The characteristics polynomial of a  $3 \times 3$  identity matrix is \_\_\_\_\_, if  $x$  is the eigen values of the given  $3 \times 3$  identity matrix. where symbol  $^{\wedge}$  shows power.

A.  $(x-1)^3$

B.  $(x+1)^3$

**C.  $x^3-1$**

**ok**

D.  $x^3+1$

**Question No:15**

**(Marks:1)**

**Vu-Topper RM**

They can be used only to find the eigenvalue of  $A$  that is largest in absolute value—we call this eigenvalue the dominant eigenvalue of  $A$ .

**A. TRUE**

B. FALSE

**Question No:16**

**(Marks:1)**

**Vu-Topper RM**

In ..... method, a system is reduced to an equivalent diagonal form using elementary transformations.

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

- A. Jacobi's
- B. Gauss-Seidel
- C. Relaxation

**D. Gaussian elimination**

**Page 262**

**ok**

**Question No:17**

**(Marks:1)**

**Vu-Topper RM**

The linear equation:  $2x+0y-2=0$  has ----- solution/solutions.

**A. Unique**

**Page 48**

- B. no solution
- C. infinite many
- D. finite many

**Question No:18**

**(Marks:1)**

**Vu-Topper RM**

Under elimination methods, we consider, Gaussian elimination and ...methods.

- A. Gauss-Seidel
- B. Jacobi

**C. Gauss-Jordan elimination**

**Page 48**

D. None of the given choices

**Question No:19**

**(Marks:1)**

**Vu-Topper RM**

Which of the following method is not an iterative?

- A. Jacobi's method
- B. Gauss-Seidel method
- C. Relaxation methods

**D. Gauss-Jordan method**

**ok**

**Question No:20**

**(Marks:1)**

**Vu-Topper RM**

An eigenvector V is said to be normalized if the coordinate of largest magnitude is equal to zero.

A. TRUE

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

**B. FALSE**

**Page 97**

**ok**

**Question No:21**

**(Marks:1)**

**Vu-Topper RM**

Exact solution of  $2/3$  is not existing.

**A. TRUE**

B. FALSE

**Question No:22**

**(Marks:1)**

**Vu-Topper RM**

When the condition of diagonal dominance becomes true in Jacobi's Method. Then it means that the method is .....

A. Stable

B. Unstable

**C. Convergent**

**Page 70**

**ok**

D. Divergent

**Question No:23**

**(Marks:1)**

**Vu-Topper RM**

Gauss-Seidel method is similar to .....

A. Iteration's method

B. Regula-Falsi method

C. Jacobi's method

**D. None of the given choices**

**Page 263**

**ok**

**Question No:24**

**(Marks:1)**

**Vu-Topper RM**

Sparse matrices arise in computing the numerical solution of .....

A. Ordinary differential equations

**B. Partial differential equations**

**Page 69**

**ok**

C. Linear differential equations

D. Non-linear differential equations

**Question No:25**

**(Marks:1)**

**Vu-Topper RM**

While solving by Gauss-Seidel method, which of the following is the first

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

Iterative solution for the system;  $x-2y=1$ ,  $x+4y=4$  ?

**A. (1, 0.75)** ok

B. (0,0)

C. (1,0)

D. (0,1)

**Question No:26**

**(Marks:1)**

**Vu-Topper RM**

While solving a system of linear equations by Gauss Jordan Method, after all the elementary row operations if there lefts also zeros on the main diagonal then which of the is true about the system?

A. System may have unique solutions

B. System has no solution

C. System may have multiple numbers of finite solutions

**D. System may have infinite many solutions** ok

**Question No:27**

**(Marks:1)**

**Vu-Topper RM**

Numerical methods for finding the solution of the system of equations are classified as direct and ..... methods

A. Indirect

**B. Iterative** Page 48

C. Jacobi

D. None of the given choices

**Question No:28**

**(Marks:1)**

**Vu-Topper RM**

If the Relaxation method is applied on the system;  $2x+3y=1$ ,  $3x+2y=-4$ , then largest residual in 1st iteration will reduce to -----.

**A. Zero** ok

B. 4

C. -1

D. -1

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

**Question No:29**

**(Marks:1)**

**Vu-Topper RM**

While using Relaxation method, which of the following is the Residuals for 1st iteration on the system;  $2x+3y = 1$ ,  $3x + 2y = 4$  ?

- A. (2,3)
- B. (3,-2)
- C. (-2,3)
- D. (1,4)

ok

**Question No:30**

**(Marks:1)**

**Vu-Topper RM**

If the order of coefficient matrix corresponding to system of linear equations is 33 then which of the following will be the orders of its decomposed matrices; 'L' and 'U'?

- A. Order of 'L' = 31, Order of 'U' = 13
- B. Order of 'L' = 32, Order of 'U' = 2\*3
- C. Order of 'L' = 33, Order of 'U' = 33
- D. Order of 'L' = 34, Order of 'U' = 43

**Question No:31**

**(Marks:1)**

**Vu-Topper RM**

While solving the system;  $x-2y = 1$ ,  $x+4y = 4$  by Gauss-Seidel method, which of the following ordering is feasible to have good approximate solution?

- A.  $x+4y = 1$ ,  $x-2y = 4$
- B.  $x+2y = 1$ ,  $x-4y = 4$
- C.  $x+4y = 4$ ,  $x-2y = 1$
- D. No need to reordering

ok

**Question No:32**

**(Marks:1)**

**Vu-Topper RM**

While solving the system;  $x-2y = 1$ ,  $x+4y = 4$  by Gauss-Seidel method, which of the following ordering is feasible to have good approximate solution?

- A. Complicated
- B. Easiest

Page 51

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

**Question No:33**

**(Marks:1)**

**Vu-Topper RM**

Full pivoting, in fact, is more ...than the partial pivoting.

A. Easiest

**B. Complicated**

**Page 51**

**Question No:34**

**(Marks:1)**

**Vu-Topper RM**

For the equation  $X^3 + 3x - 1 = 0$  the root of the equation lies in the interval...

A. (1, 3)

B. (1, 2)

**C. (0, 1)**

D. (1, 2)

**Question No:35**

**(Marks:1)**

**Vu-Topper RM**

lies in the category of iterative method.

A. Bisection Method

B. Regula Falsi Method

C. Secant Method

D. **All**

**Page 8**

**Question No:36**

**(Marks:1)**

**Vu-Topper RM**

If  $n \times n$  matrices A and B are similar, then they have the different eigenvalues (with the same multiplicities).

A. TRUE

**B. FALSE**

**Question No:37**

**(Marks:1)**

**Vu-Topper RM**

The Jacobi's method is a method of solving a matrix equation on a matrix that has \_\_\_\_ zeros along its main diagonal.

**No**

**Page 892**

**ok**

At

least one

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

**Question No:38**

**(Marks:1)**

**Vu-Topper RM**

To apply Simpson's  $3/8$  rule, the number of intervals in the following must be

- A. 10
- B. 11
- C. **12**
- D. 13

**Question No:39**

**(Marks:1)**

**Vu-Topper RM**

The Gauss-Seidel method is applicable to strictly diagonally dominant or symmetric \_\_\_\_\_ definite matrices A.

- A. **Positive** ok
- B. negative

**Question No:40**

**(Marks:1)**

**Vu-Topper RM**

Differences methods find the \_\_\_\_\_ solution of the system.

- A. **Numerical** ok
- B. Analytical

**Question No:41**

**(Marks:1)**

**Vu-Topper RM**

To apply Simpson's  $1/3$  rule, the number of intervals in the following must be

- A. **2** Simpson rule
- B. 3
- C. 5
- D. 7

**Question No:42**

**(Marks:1)**

**Vu-Topper RM**

Bisection and false position methods are also known as bracketing method and are always

- A. Divergent

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

Question No:43

(Marks:1)

Vu-Topper RM

The Inverse of a matrix can only be found if the matrix is.

- A. Singular
- B. **None Singular**
- C. Scalar
- D. Diagonal

Question No:44

(Marks:1)

Vu-Topper RM

In interpolation is used to represent the  $\delta$  Forward difference  $\Delta$

- A. **Central difference** Page 117
- B. Backward difference

Question No:45

(Marks:1)

Vu-Topper RM

The base of the decimal system is \_\_\_\_\_

- A. **10**
- B. 0
- C. 2
- D. 8

Question No:46

(Marks:1)

Vu-Topper RM

Bisection method is ... method

- A. Open Method
- A. **Bracketing Method** Page 26

Question No:47

(Marks:1)

Vu-Topper RM

A 3 x 3 identity matrix have three and \_\_\_\_\_ eigen values.

- A. **Same** ok
- B. different

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

For More Help Contact What's app 03224021365

**Question No:48**

**(Marks:1)**

**Vu-Topper RM**

Eigenvalues of a symmetric matrix are all \_\_\_\_\_.

- A. complex
- B. zero
- C. Real
- D. positive

**Page 104**

**ok**

**Question No:49**

**(Marks:1)**

**Vu-Topper RM**

The Jacobi iteration converges, if A is strictly diagonally dominant.

- A. TRUE
- B. FALSE

**Page 69**

**ok**

**Question No:50**

**(Marks:1)**

**Vu-Topper RM**

Below are all the finite difference methods EXCEPT \_\_\_\_\_.

- A. jacobi's method
- B. Newton's backward difference method
- C. Stirling formula
- D. Forward difference method

**Question No:51**

**(Marks:1)**

**Vu-Topper RM**

Two matrices with the same characteristic polynomial need not be similar.

- A. TRUE
- B. FALSE

**ok**

**Question No:52**

**(Marks:1)**

**Vu-Topper RM**

The determinant of a diagonal matrix is the product of the diagonal elements.

- A. TRUE
- B. FALSE

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

**Question No:53**

**(Marks:1)**

**Vu-Topper RM**

The Gauss-Seidel method is applicable to strictly diagonally dominant or symmetric positive definite matrices A.

**A. TRUE**

B. FALSE

**Question No:54**

**(Marks:1)**

**Vu-Topper RM**

The determinant of a \_\_\_\_\_ matrix is the product of the diagonal elements.

**A. Diagonal**

**Page 70**

**ok**

B. Upper triangular

C. Lower triangular

D. Scalar

**Question No:55**

**(Marks:1)**

**Vu-Topper RM**

For differences methods we require the set of values.

**A. TRUE**

**ok**

B. FALSE

**Question No:56**

**(Marks:1)**

**Vu-Topper RM**

If  $x$  is an eigen value corresponding to eigen value of  $V$  of a matrix  $A$ . If  $a$  is any constant, then  $x - a$  is an eigen value corresponding to eigen vector  $V$  is an of the matrix  $A - aI$ .

**A. TRUE**

B. FALSE

**Question No:57**

**(Marks:1)**

**Vu-Topper RM**

Central difference method seems to be giving a better approximation; however, it requires more computations.

A. True

B. **False**

**Page 71**

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

**Question No:58**

**(Marks:1)**

**Vu-Topper RM**

Iterative algorithms can be more rapid than direct methods.

**A. TRUE** ok

B. FALSE

**Question No:59**

**(Marks:1)**

**Vu-Topper RM**

Central Difference method is the finite difference method.

**A. TRUE** ok

B. FALSE

**Question No:60**

**(Marks:1)**

**Vu-Topper RM**

The Jacobi's method is a method of solving a matrix equation on a matrix that has no zeros along its main diagonal.

**A. TRUE** Page 892

B. FALSE

**Question No:61**

**(Marks:1)**

**Vu-Topper RM**

Power method is applicable if the eigen vectors corresponding to eigen values are linearly independent.

**A. TRUE** Page 102 ok

B. FALSE

**Question No:62**

**(Marks:1)**

**Vu-Topper RM**

Power method is applicable if the eigen values are \_\_\_\_\_.

**A. Real and Distinct** Page 102

B. real and equal

C. positive and distinct

D. negative and distinct

**Question No:63**

**(Marks:1)**

**Vu-Topper RM**

Simpson's rule is a numerical method that approximates the value of a definite integral by using polynomials.

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

**A. Quadratic**

**Page 174**

B. Linear

C. Cubic

D. Quartic

**Question No:64**

**(Marks:1)**

**Vu-Topper RM**

In Simpson's Rule, we use parabolas to approximating each part of the curve. This proves to be very efficient as compared to Trapezoidal rule.

**A. TRUE**

B. FALSE

**Question No:65**

**(Marks:1)**

**Vu-Topper RM**

The predictor-corrector method an implicit method. (multi-step methods)

**A. TRUE**

**Page 212**

B. FALSE

**Question No:66**

**(Marks:1)**

**Vu-Topper RM**

Generally, Adams methods are superior if output at many points is needed.

**A. TRUE**

B. FALSE

**Question No:67**

**(Marks:1)**

**Vu-Topper RM**

The Trapezoidal rule is a numerical method that approximates the value of a \_\_\_\_\_.

A. Indefinite integral

**B. Definite integral**

**Page 176**

C. Improper integral

D. Function

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

**Question No:68**

**(Marks:1)**

**Vu-Topper RM**

The need of numerical integration arises for evaluating the definite integral of a function that has no explicit \_\_\_\_\_ or whose anti derivative is not easy to obtain.

**A. Antiderivative**

B. Derivatives

**Question No:69**

**(Marks:1)**

**Vu-Topper RM**

An indefinite integral may \_\_\_\_\_ in the sense that the limit defining it may not exist.

**A. Diverge**

B. Converge

**Question No:70**

**(Marks:1)**

**Vu-Topper RM**

An improper integral is the limit of a definite integral as an endpoint of the interval of integration approaches either a specified real number or  $\infty$  or  $-\infty$  or, in some cases, as both endpoints approach limits.

**A. TRUE**

B. FALSE

**Question No:71**

**(Marks:1)**

**Vu-Topper RM**

While using Relaxation method, which of the following is the Residuals for 1st iteration on the system;  $2x + 3y = 1$ ,  $3x + 2y = 4$  ?

**A. (2,3)**

B. (1,4)

C. (4,4)

D. (3,4)

**Question No:72**

**(Marks:1)**

**Vu-Topper RM**

For the system of equations;  $x = 2$ ,  $y = 3$ . The inverse of the matrix associated with its coefficients is-----.

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

**A. Singular**

B. Non-singular

C. Non identity

**Question No:73**

**(Marks:1)**

**Vu-Topper RM**

The number of significant digits in the number 608.030060 is:

A. 7

B. 8

**C. 9**

D. 1

**Question No:74**

**(Marks:1)**

**Vu-Topper RM**

For two matrices A and B, such that "A = Inverse of B", then which of the following is true?

A. B=Inverse of A

B. A and B are not-singular

C. AB=Identity Matrix

**D. All choice is true**

**Question No:75**

**(Marks:1)**

**Vu-Topper RM**

If a system of equations has a property that each of the equation possesses one large coefficient and the larger coefficients in the equations correspond to different unknowns in different equations, then which of the following iterative method is preferred to apply?

A. Gauss-Seidel method

B. Gauss-Jordon method

**C. Gauss elimination**

**ok**

D. Croat's method

**Question No:76**

**(Marks:1)**

**Vu-Topper RM**

Two matrices with the \_\_ characteristic polynomial need not be similar.

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

**A. Same**

B. different

**Question No:77**

**(Marks:1)**

**Vu-Topper RM**

Which of the following method is simplest one to integrate numerically a given tabular function but give more error?

A. Rectangular method

**B. Trapezoidal method**

C. Simpson's 1/3 Rule

D. Simpson's 3/8 Rule

**Question No:78**

**(Marks:1)**

**Vu-Topper RM**

At which of the following points the Minimum value of 2nd derivative of function  $f(x) = -(2/x)$  in the interval:  $[1,4]$  exists?

**A. At  $x=1$**

B. At  $x=2$

C. At  $x=3$

D. At  $x=4$

**Question No:79**

**(Marks:1)**

**Vu-Topper RM**

The Gaussian elimination method fails if any one of the pivot elements becomes.....?

**A. Zero**

B. One

C. Two

D. Five

**Question No:80**

**(Marks:1)**

**Vu-Topper RM**

Gauss elimination and Gauss-Jordan methods are popular among many methods for finding the .....of a matrix.

A. Identity

B. Transpose

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

**C. Inverse** ok

D. None of the given choices

**Question No:81**

**(Marks:1)**

**Vu-Topper RM**

In Gauss-Seidel method, each equation of the system is solved for the unknown with ----- coefficient, in terms of remaining unknowns.

**A. Largest** ok

B. Smallest

C. Any positive

D. any negative

**Question No:82**

**(Marks:1)**

**Vu-Topper RM**

Power method is applicable if the Eigen vectors corresponding the Eigen values are linearly

A. Dependent

**B. Independent**

**Page 6**

**Question No:83**

**(Marks:1)**

**Vu-Topper RM**

Power method is applicable if the Eigen values are real and distinct.

A. False

**B. True**

**Page 102**

**Question No:84**

**(Marks:1)**

**Vu-Topper RM**

The Power method can be used only to find the eigenvalue of A that is largest in absolute value—we call this eigenvalue the dominant eigenvalue of A.

A. False

**B. True**

**ok**

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

**Question No:85**

**(Marks:1)**

**Vu-Topper RM**

If  $x$  is an eigenvalue corresponding to eigenvalue of  $V$  of a matrix  $A$ . If  $a$  is any constant, then  $x - a$  is an eigen value corresponding to eigen vector  $V$  is an of the matrix  $A - a I$ .

A. False

**B. True**

**Question No:86**

**(Marks:1)**

**Vu-Topper RM**

The dominant or principal eigenvector of a matrix is an eigenvector corresponding to the Eigen value of largest magnitude (for real numbers, largest absolute value) of that matrix,

A. False

**B. True**

**Question No:87**

**(Marks:1)**

**Vu-Topper RM**

Eigen values of a \_\_\_\_\_ matrix are all real.

A. Antisymmetric

**B. Symmetric**

C. Rectangular

D. Triangular

**Question No:88**

**(Marks:1)**

**Vu-Topper RM**

In Trapezoidal rule, the integral is computed on each of the sub-intervals by using linear interpolating formula, ie. For  $n=1$  and then summing them up to obtain the desired integral.

A. False

**B. True**

**Question No:89**

**(Marks:1)**

**Vu-Topper RM**

In Runge – Kutta Method, we do not need to calculate higher order derivatives and find greater accuracy.

A. False

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

**B. True**

**Question No:90**

**(Marks:1)**

**Vu-Topper RM**

The Trapezoidal Rule is an improvement over using rectangles because we have much less "missing" from our calculations. We used \_\_\_\_\_ to model the curve in trapezoidal Rule.

**A. Straight lines**

B. curves

C. parabolas

D. constant

**Question No:91**

**(Marks:1)**

**Vu-Topper RM**

Euler's Method numerically computes the approximate derivative of a function.

**A. False**

B. True

**Question No:92**

**(Marks:1)**

**Vu-Topper RM**

Euler's Method numerically computes the approximate \_\_\_\_\_ of a function.

**A. Antiderivative**

B. Derivative

C. Error

D. Value

**Question No:93**

**(Marks:1)**

**Vu-Topper RM**

If we wanted to find the value of a definite integral with an infinite limit, we can instead replace the infinite limit with a variable, and then take the limit as this variable goes to \_\_\_\_\_.

A. Constant

B. Finite

**C. Infinity**

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

D. Zero

**Question No:94**

**(Marks:1)**

**Vu-Topper RM**

The Jacobi iteration \_\_\_\_\_, if A is strictly diagonally dominant.

**A. Converges**      **ok**

B. diverges

**Question No:95**

**(Marks:1)**

**Vu-Topper RM**

By using determinants, we can easily check that the solution of the given system of linear equation exists and it is unique.

A. False

**B. True**      **ok**

**Question No:96**

**(Marks:1)**

**Vu-Topper RM**

Direct method can more rapid than iterative algorithms

A. False

**B. True**

**Question No:97**

**(Marks:1)**

**Vu-Topper RM**

The dominant eigenvector of a matrix is an eigenvector corresponding to the eigenvalue of largest magnitude (for real numbers, smallest absolute value) of that matrix.

A. False

**B. True**

**Question No:98**

**(Marks:1)**

**Vu-Topper RM**

The central difference method is finite difference method.

A. False

**B. True**

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

**Question No:99**

**(Marks:1)**

**Vu-Topper RM**

The absolute value of a determinant ( $|\det A|$ ) is the product of the absolute values of the eigenvalues of matrix A

A. False

**B. True** ok

**Question No:100**

**(Marks:1)**

**Vu-Topper RM**

Eigenvectors of a symmetric matrix are orthogonal, but only for distinct eigenvalues.

A. False

**B. True**

**Question No:101**

**(Marks:1)**

**Vu-Topper RM**

6. Let A be an  $n \times n$  matrix. The number x is an eigenvalue of A if there exists a non-zero vector v such that  $Av = xv$

A.  $Av = xv$

**B.  $Ax = xv$**  ok

C.  $Av + xv = 0$

D.  $Av = Ax$

**Question No:102**

**(Marks:1)**

**Vu-Topper RM**

In Jacobi's Method, the rate of convergence is quite \_\_\_\_\_ compared with other methods.

**A. Slow** ok

B. Fast

**Question No:103**

**(Marks:1)**

**Vu-Topper RM**

Numerical solution of  $2/3$  up to four decimal places is \_\_\_\_\_.

A. 0.667

B. 0.6666

**C. 0.6667** ok

D. 0.66667

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

**Question No:104**

**(Marks:1)**

**Vu-Topper RM**

Euler's method is only useful for a few steps and small step sizes; however, Euler's method together with Richardson extrapolation may be used to increase the \_\_\_\_\_.

**A. Order and accuracy**

B. Divergence

**Question No:105**

**(Marks:1)**

**Vu-Topper RM**

The first langrage polynomial with equally spaced nodes produced the formula for \_\_\_\_\_.

A. Simpson's rule

**B. Trapezoidal rule**

C. Newton's method

D. Richardson's method

**Question No:106**

**(Marks:1)**

**Vu-Topper RM**

The need of numerical integration arises for evaluating the indefinite integral of a function that has no explicit antiderivative or whose antiderivative is not easy to obtain.

A. False

**B. True**

**Question No:107**

**(Marks:1)**

**Vu-Topper RM**

The Euler method is numerically unstable because of \_\_\_\_\_ convergence of error.

**A. Slow**

B. Fast

C. Moderate

D. No

**Question No:108**

**(Marks:1)**

**Vu-Topper RM**

Adams – Bash forth is a multistep method.

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

A. False

B. True

**Question No:109**

**(Marks:1)**

**Vu-Topper RM**

Multistep method does not improve the accuracy of the answer at each step.

A. False

B. True

**Question No:110**

**(Marks:1)**

**Vu-Topper RM**

Generally, Adams methods are superior if output at \_\_\_\_\_ points is needed.

A. Many

B. Two

C. Single

D. At most

**Question No:111**

**(Marks:1)**

**Vu-Topper RM**

Symbol used for forward differences is

A.  $\nabla$

B.  $\Delta$

C.  $\delta$

D.  $\mu$

**Page 12**

**Question No:112**

**(Marks:1)**

**Vu-Topper RM**

The relationship between central difference operator and the shift operator is given by

A.  $\delta = E - E^{-1}$

B.  $\delta = E + E^{-1}$

C.  $\delta = E^{1/2} + E^{-1/2}$

D.  $\delta = E^{1/2} - E^{-1/2}$

**Page 152**

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

**Question No:113**

**(Marks:1)**

**Vu-Topper RM**

Muller's method requires -----starting points

A. 1

**B. 3**      **Page 41**

C. 2

D. 4

**Question No:114**

**(Marks:1)**

**Vu-Topper RM**

If we retain  $r+1$  terms in Newton's forward difference formula, we obtain a polynomial of degree ---- agreeing with  $y_x$  at  $x_0, x_1, \dots, x_n$ .

A.  $r+2$

**B.  $r$**

C.  $r+1$

D.  $r-1$

**Question No:115**

**(Marks:1)**

**Vu-Topper RM**

Octal number system has the base -----

A. 2

B. 10

**C. 8**

D. 16

**Question No:116**

**(Marks:1)**

**Vu-Topper RM**

Rate of change of any quantity with respect to another can be modeled by

**A. An ordinary differential equation**

B. A partial differential equation

C. A polynomial equation

D. None of the given choices

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

**Question No:117**

**(Marks:1)**

**Vu-Topper RM**

Adam-Moulton P-C method is derived by employing

- A. Newton's backward difference interpolation formula**
- B. Newton's forward difference interpolation formula
- C. Newton's divided difference interpolation formula
- D. None of the given choices

**Question No:118**

**(Marks:1)**

**Vu-Topper RM**

Newton Raphson method is ..... method

- A. Bracketing Method
- B. Open**
- C. Random
- D. none

**Question No:119**

**(Marks:1)**

**Vu-Topper RM**

Eigenvalue is

- A. Real**
- B. Vector
- C. odd
- D. even

**Question No:120**

**(Marks:1)**

**Vu-Topper RM**

Bisection and false position methods are also known as

- A. bracketing method**
- B. open method
- C. random

**Question No:121**

**(Marks:1)**

**Vu-Topper RM**

If  $f(x)$  contains trigonometric, exponential or logarithmic functions then this equation is known as

- A. Transcendental equation**
- B. Algebraic

**Page 6**

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

- C. Polynomial
- D. Linear

**Question No:122**

**(Marks:1)**

**Vu-Topper RM**

In interpolation  $\delta$  is used to represent the

- A. Forward difference
- B. Central difference**
- C. Backward difference

**Question No:123**

**(Marks:1)**

**Vu-Topper RM**

P in Newton's forward difference formula is defined as

- A.  $P=(x-x_0)/h$**
- B.  $P=(x+x_0)/h$
- C.  $P=(x+x_n)/h$
- D.  $P=(x-x_n)/h$

**Question No:124**

**(Marks:1)**

**Vu-Topper RM**

Newton's divided difference interpolation formula is used when the values of the are

- A. Equally spaced**
- B. Not equally spaced
- C. Constant
- D. None of the above

**Question No:125**

**(Marks:1)**

**Vu-Topper RM**

Given the following data

|      |   |   |   |   |
|------|---|---|---|---|
| X    | 0 | 1 | 2 | 4 |
| F(x) | 1 | 1 | 2 | 5 |

The value of  $f(2,4)$  is

- A. 1.5

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

B. 2

**C. 3**

D. 1

**Question No:126**

**(Marks:1)**

**Vu-Topper RM**

They can be used only to find the Eigen value of A that is largest in absolute value—we call this Eigen value the dominant Eigen value of A.

A. False

**B. True**

**Question No:127**

**(Marks:1)**

**Vu-Topper RM**

If the order of coefficient matrix corresponding to system of linear equations is  $3 \times 3$  then which of the following will be the orders of its decomposed matrices; 'L' and 'U'?

A. Order of 'L' =  $3 \times 1$ , Order of 'U' =  $1 \times 3$

B. Order of 'L' =  $3 \times 2$ , Order of 'U' =  $2 \times 3$

**C. Order of 'L' =  $3 \times 3$ , Order of 'U' =  $3 \times 3$**  **ok**

D. Order of 'L' =  $3 \times 4$ , Order of 'U' =  $4 \times 3$

**Question No:128**

**(Marks:1)**

**Vu-Topper RM**

For the equation  $x + 3x - 1 = 0$ , the root of the equation lies in the interval.....

A. (1, 3)

B. (1, 2)

**C. (0, 1)**

D. (1, 2)

**Question No:129**

**(Marks:1)**

**Vu-Topper RM**

If the root of the given equation lies between a and b, then the first approximation to the root of the equation by bisection method is

**A.  $\frac{a+b}{2}$**

B.  $\frac{a-b}{2}$

C.  $\frac{b-a}{2}$

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

D. None of the given options

**Question No:130**

**(Marks:1)**

**Vu-Topper RM**

Let A be an  $n \times n$  matrix. The number  $x$  is an eigenvalue of A if there exists a non-zero vector  $v$  such that \_\_\_\_\_.

A.  $Av = xv$

B.  $Av + xv = 0$

C.  $Av = Ax1$

**D.  $Av = \lambda v$  ok**

**Question No:131**

**(Marks:1)**

**Vu-Topper RM**

By using determinants, we can easily check that the solution of the given system of linear equation \_\_\_\_\_ and it is \_\_\_\_\_.

**A. Exits, unique**

B. Exists, consistent

C. Trivial, unique

D. Non trivial, inconsistent

**Question No:132**

**(Marks:1)**

**Vu-Topper RM**

In ..... method, the elements above and below the diagonal are simultaneously made zero.

A. Jacobi's

B. Gauss-Seidel

**C. Gauss-Jordan Elimination**

**Page 59**

**ok**

D. Relaxation

**Question No:133**

**(Marks:1)**

**Vu-Topper RM**

Which of the following is equivalent form of the system of equations in matrix form;  $AX=B$  ?

A.  $XA = B$

**B.  $X = B$  (Inverse of A) ok**

C.  $X = (\text{Inverse of } A)B$

D.  $BX = A$

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

**Question No:134**

**(Marks:1)**

**Vu-Topper RM**

If the determinant of a matrix A is not equal to zero then the system of equations will have.....

- A. A unique solution**
- B. Many solutions
- C. Infinite many solutions
- D. None of the given choices

**Question No:135**

**(Marks:1)**

**Vu-Topper RM**

Sparse matrix is a matrix with .....

- A. Some elements are zero
- B. Many elements are zero**
- C. Some elements are one
- D. Many elements are one

**Page 69**

**ok**

**Question No:136**

**(Marks:1)**

**Vu-Topper RM**

How many Eigen vectors will exist corresponding to the function;  
 $\text{Exp}(ax) = e^{ax}$ , when the matrix operator is of differentiation?

- A. Infinite many**
- B. Finite Multiple
- C. None

**ok**

**Question No:137**

**(Marks:1)**

**Vu-Topper RM**

Which of the following is the meaning of partial pivoting while employing the row transformations?

- A. Making the largest element as pivot**
- B. Making the smallest element as pivot
- C. Making any element as pivot
- D. Making zero elements as pivot

**Page 50**

**Question No:138**

**(Marks:1)**

**Vu-Topper RM**

Differences methods are iterative methods.

- A. False

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

**B. True**

**Question No:139**

**(Marks:1)**

**Vu-Topper RM**

Eigenvalues of a \_\_\_\_\_ matrix are all real.

**A. Symmetric**

B. Anti-symmetric

C. Rectangular

D. Triangular

**Question No:140**

**(Marks:1)**

**Vu-Topper RM**

For a system of linear equations, the corresponding coefficient matrix has the value of determinant;  $|A| = 0$ , then which of the following is true?

A. The system has unique solution

B. The system has finite multiple solutions

C. The system has infinite may solutions

**D. The system has no solution**

**Page 48**

**Question No:141**

**(Marks:1)**

**Vu-Topper RM**

For the system;  $2x+3y = 1$ ,  $3x + 2y = -4$ , if the iterative solution is  $(0,0)$  and ' $dx_i = 2$ ' is the increment in ' $y$ ' then which of the following will be taken as next iterative solution?

A.  $(2,0)$

B.  $(0,3)$

C.  $(0,2)$

**D.  $(1,-4)$**

**ok**

**Question No:142**

**(Marks:1)**

**Vu-Topper RM**

While using Relaxation method, which of the following is increment ' $dx_i$ ' corresponding to the largest Residual for 1st iteration on the system;  $2x+3y = 1$ ,  $3x + 2y = -4$  ?

A. -2

B. 2

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

C. 4 ok  
D. 3

**Question No:143** (Marks:1) **Vu-Topper RM**

If system of equations is inconsistent then its means that it has .....

A. No Solutions ok

B. Many solutions Infinite

C. Many solutions

D. None of the given choices

**Question No:144** (Marks:1) **Vu-Topper RM**

Relaxation Method is a/an .....

A. Direct method

B. Iterative method ok

**Question No:145** (Marks:1) **Vu-Topper RM**

The eigenvectors of a square matrix are the non-zero vectors that, after being multiplied by the matrix, remain ..... to the original vector.

A. Perpendicular

B. Parallel ok

C. Diagonal

D. None of the given choices

**Question No:146** (Marks:1) **Vu-Topper RM**

In Jacobi's method after finding D1, the sum of the diagonal elements of D1 should be ..... to the sum of the diagonal elements of the original matrix A.

A. Greater than

B. Less than

C. Same Page 106

D. Different

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

**Question No:147**

**(Marks:1)**

**Vu-Topper RM**

In interpolation is used to represent the d Forward difference?

**A. Central difference**

B. Backward difference

**Question No:148**

**(Marks:1)**

**Vu-Topper RM**

Power method is applicable if the eigen vectors corresponding to eigen values are linearly ----.

**A. Independent**

B. Dependent

**Question No:149**

**(Marks:1)**

**Vu-Topper RM**

Power method is applicable if the Eigen vector corresponding to pigeon values Ara linearly independent

A. False

**B. True**

**Question No:150**

**(Marks:1)**

**Vu-Topper RM**

In Jacobi method we assume that the \_\_\_\_\_ element does not vanish

**A. Diagonal**

B. off – Diagonal

C. Row

D. Column

**Question No:151**

**(Marks:1)**

**Vu-Topper RM**

In Jacobi method that rat off convergence is quit slow compared with other method

A. False

**B. True**

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

**Question No:152**

**(Marks:1)**

**Vu-Topper RM**

White solving the system linear equation  $x-y=2$ ,  $-x+y=3$  By Jacobi method if  $(0, 0)$  be first approximate solution then which of that following is second approximate solution

**A. (2, 3)**

B.  $(0, 3)$

C.  $(2, 0)$

**Question No:153**

**(Marks:1)**

**Vu-Topper RM**

The liner equation:  $0x+0y = 2$  has \_\_\_\_\_ solution

A. unique

**B. No solution**

C. infinite many

D. finite many

**Question No:154**

**(Marks:1)**

**Vu-Topper RM**

When the condition of diagonal dominance become true in Jacobi method then it's means that the method is

A. stable

B. un- stable

**C. Convergent**

D. Divergent

**Question No:155**

**(Marks:1)**

**Vu-Topper RM**

A series  $16+8+4+2+1$  is replaced by the series  $16+8+4+2$  then it is called

A. inherent error

B. local round of error

**C. local truncation error**

**Question No:156**

**(Marks:1)**

**Vu-Topper RM**

If one root of the equal is  $3-7i$ , then the other root will be

A.  $-3-7i$

بري صحبت سے تنہائی بہتر ہے اور تنہائی سے نيك صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

**B.  $-3+7i$**

C.  $3-7i$

D.  $3+7i$

**Question No:157**

**(Marks:1)**

**Vu-Topper RM**

The number system that has a base 2 is called...system.

A. octal

**B. Binary**

C. Decimal

D. Hexadecimal

**Question No:158**

**(Marks:1)**

**Vu-Topper RM**

Which method required is derivative of that solution

A. Bisection method

B. Regular falsi method

C. Muller method

**D. Newton Raphson method**

**Question No:159**

**(Marks:1)**

**Vu-Topper RM**

If the Relaxation method is applied on the system  $2x+3y=1$   $3x+2y=-4$  then largest residual in 1<sup>st</sup> iteration will reduce to

Select the correct option

**A. Zero**

B. 4

C. -1

D. -1

**Question No:160**

**(Marks:1)**

**Vu-Topper RM**

The number system that has a base 8 is called \_ system

A. Decimal

**B. Octal**

C. Binary

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

**Question No:161**

**(Marks:1)**

**Vu-Topper RM**

In Gauss Seidel method each equation of the system is solved for the unknown with \_ coefficient in terms of remaining unknown

- A. smallest
- B. largest**
- C. any largest
- D. any negative

**Question No:162**

**(Marks:1)**

**Vu-Topper RM**

Which method is required a derivative of a solution?

- A. Regular falsi method
- B. Neuton Raphson method**
- C. Muller method
- D. Bisection method

**Question No:163**

**(Marks:1)**

**Vu-Topper RM**

If a system of equation has a property that each of the equation possesses one large coefficient in the equations correspond to different unknown in different equation then which of the following iterative method id preferred to apply?

- A. Crout s method
- B. Gauss Seidel method**
- C. Gauss – Jordan method
- D. Gauss elimination method

**Question No:164**

**(Marks:1)**

**Vu-Topper RM**

With the initial vector (0;0;0;) the residual would be

- A.  $R_1 = 2, R_2 = 1, R_3 = 1$
- B.  $R_1 = 1, R_2 = 3, R_3 = 2$
- C.  $R_1 = 5, R_2 = 9, R_3 = 2$**
- D.  $R_1 = 3, R_2 = 2, R_3 = 1$

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

**Question No:165**

**(Marks:1)**

**Vu-Topper RM**

In full providing we interchange rows and columns such that the element in the matrix of the variable also get charged

**A. largest**

B. Middle

C. smallest

D. none of the given choice

**Question No:166**

**(Marks:1)**

**Vu-Topper RM**

For the system  $2x+3y=1$   $3x + 2y = -4$  if the iterative solution is  $(0 ; 0,)$  and  $dx_i = 2$  is the increment in  $y$  then which of the following will be taken as next iterative so

A.  $(0;3)$

**B.  $(0;2)$**

C.  $(1;4)$

D.  $(2;0)$

**Question No:167**

**(Marks:1)**

**Vu-Topper RM**

If the determinant of the matrix  $A$  is equal to zero then the system of equation will have ....

**A. No solution or infinitely many solution**

B. unique solution

C. infinite many solution

D. no solution

The number system that has a base 16 is called...system.

**Hexadecimal**

بری صحبت سے تنہائی بہتر ہے اور تنہائی سے نیک صحبت بہتر ہے

**For More Help Contact What's app 03224021365**

**Visit My YouTube Channel**  
**For More Important Notes**  
**Channel Name = #VuTopperRM**

برى صحت سے تٲهائى بهٲر هے اور تٲهائى سے نيك صحت بهٲر هے

**For More Help Contact What's app 03224021365**