Q.1 With α = 0.05, how much of the distribution represents the critical region?

A. 10%
B. 5%
C. 1%
D. 50%

Q.2 Which of the formula is used to measure the mean of Binomial Distribution?

A. p+n
B. np
C. np (1-p)
D. n-p

Q.3 If we add a constant to each score in statistics then it will not change the _____

A. Sample
B. Standard deviation
C. None of the given options
D. Variable

Q.4 Sample means should be relatively close to the:

A. Population means
B. Population size
C. Critical region
D. Sample size

Q.5 When a variable is measured on a scale consisting of exactly two categories: the resulting data are called:

A. Distribution
B. Population
C. Binomial
D. Normal

Q.6 Why is the sample variance considered biased if divided by n?

A. It eliminates variability
B. It underestimates the population variance
C. It ignores the mean
D. It overestimates standard deviation

Q.7 Which of the following two main areas are discussed in Inferential Statistics?

A. Level of the significant degree of freedom
B. Point estimate and interval estimate
C. Estimation of parameter testing of hypothesis
D. Biased estimator and unbiased estimator

Q.8 In statistical models, ANOVA stands for:

A. Analysis of variable
B. Analysis of verification
C. Analysis of Variance
D. Analysis of variation

Q.9 Which of the following measure of dispersion provides the difference between the third and first quartile?

A. Inter Quartile Range

B. Range

C. Standard Deviation

D. Variance

Q.10 What does the null hypothesis (H0) state in hypothesis testing?

A. The treatment has an effect

B. The treatment has no effect

C. The data support the research hypothesis

D. The sample mean is extreme
