Q.1 Which assumption ensures that measurements from one participant are not influenced by another?

A. Equal variances
B. Random sampling
C. Normality
D. Independence of observations

Q.2 In research, Ho is used to state_____ statistically

A. Null hypothesis
B. Alternate hypothesis
C. No hypothesis
D. All hypothesis

Q.3 While using SPSS, in which of the following view, the researcher selects the level of measurement of each variable?

A. Data view
B. Comment box
C. Dialogue box
D. Variable view

Q.4 Higher variability can _____the chances of finding a significant treatment effect in a hypothesis test.

A. Deviate
B. Increase
C. Not change
D. Decrease

Q.5 Which is an assumption of independent-sample t-test?

A. Different variances
B. Skewed populations
C. Independent observations
D. Dependent observations

Q.6 In SPSS, the correlation coefficient ranges from –1 to 1. This value indicates the _____between two variables.

A. Coefficients of the variables
B. Significance level
C. Inverse relationship
D. Strength of the relationship

Q.7 In ANOVA, the dependent variable should be:

A. On ordinal Scale

B. Measured on interval or ratio scale

C. On categorical scale

D. On nominal Scale

Q.8 Why is the independent-samples t-test widely used?

A. It ignores population assumptions

B. Most studies compare two separate groups

C. It requires only one group

D. It compares more than two means

Q.9 In research, which of the following is a measure of the stability or consistency of test scores?

A. Reliability

B. Non-probability

C. Validity

D. Probability

Q.10 If an alpha level of .05 is used, that means the researcher is taking a _____ risk of committing a Type I error.

A. 5%

B. 7%

C. 10%

D. 6%
