

Virtual University of Pakistan

(bif401)

(bioinformatics 1)

MCQ's for final term

Email address

bif401@vu.edu.pk

Course Instructor

(Dr. Saffee Ullah)

Created By Sadaf Parvez

Group of VU Biologists

www.facebook.com/vubiologists

www.facebook.com/groups/vubiologists

VU Biologists

www.facebook.com/groups/vubiologists

----- provide mass of intact molecules and its fragments

MS1 and MS2

theoretical data is usually ----- as compared to exp. Data,

larger

Three scoring schemes can be applied to -----

score the match at each stage of protein search

Simply sum the scores up (a linear function)

$$\text{Score} = \text{Score}_{MW} + \text{Score}_{PST} + \text{Score}_{Exp} < \text{Thr}$$

In the above scoring method, MW and PST stands for ____ & ____ respectively

MS1 & MS2

_____ are assembled by Ribosomes in cellular cytoplasm

Proteins

Amino acids have characteristics like polarity, hydrophobicity, and charge states. these characteristics are governed by

Elemental composition of an amino acid's side chain (R group)

Hydrophobic amino acids are non-polar due to

very little dipole moments between H and C

Core of amino acids is _____ and surface is _____

Hydrophobic & hydrophilic

At pH ____ Five amino acids are charged, 2 negatively and 3 positively

7

pK values for an amino acid is the pH at which:

Exactly half of the chargeable group is charged

The pK of aspartic and glutamic acids are ____-& ____ respectively.

3.9 & 4.2

If $\text{pH} < \text{pK}$ for an amino acid, the amine side chains ____

Gain proton and hence basic

If $\text{pH} > \text{pK}$ for an amino acid, _____ loses a proton and hence acidic

The carboxyl side chain

Lysine and arginine have pK ____ & ____ respectively.

10.5 & 12.5

Glycine residues increase backbone flexibility because :

They have no R group (only an H)

_____ reduce the flexibility of polypeptide chains.

Proline residues

_____ Cement together by making _____ to stabilize 3-D protein structures

Cysteines, disulfide bonds

Chemically _____ remain in the core of protein.

Inactive amino acids

Backbone is highly polar (hydrophilic) due to polar ____ and ____ in each peptide unit

-NH & C=O

A-helices & B-sheets are stabilized by _____

H-bonds

The very few hydrophobic "patches" on protein surface are involved in:

Protein-protein interactions

Alpha Helix is Stabilized by H-bonds between every ____ in backbone

~ 4th residue

Proteins fold spontaneously to achieve:

Thermodynamic stability

If a protein is ____, then it can lead to a lack of function in the protein

Misfolded

[Download More Files from VUAnswer.com](#)

Given algorithms and procedures to fold a protein, we can:

Fold amino acid chains to form 3D proteins

Each amino acid can fold into ____, ____ & ____

Alpha Helices, Beta Sheets and Loops

If computations of protein folding take $1/10^{\text{th}}$ of a nano-second (10^{-10}), then to compute all the folding possibilities will take

1.6×10^{30} years

All the information required for folding a protein into its native structure is present within the:

Protein's amino acid sequence

(Memorize the table given below)

Protein Structures	
Energies of Various Bonds & Interactions	
Bond Type	kJ/mol
Covalent Bond	250
Electrostatic	5
van der Waals	5
Hydrogen bond	20

Primary sequence can also be referred to as ____

1' structure

Protein structures are organized into ____ modular conformations

1', 2', 3' and 4'

[Download More Files from VUAnswer.com](#)

VU Biologists

www.facebook.com/groups/vubiologists

Edman Degradation & Tandem Mass Spectrometry are methods for obtaining _____

1' structures

2' structures are also referred to as

Secondary structures

C- Terminus is _____ charged

Negatively

N-terminus is _____ charged

Positively

2' structures or secondary protein structures are formed as a result of H-Bond formation between _____ in a protein backbone

N and C termini

_____ connect helices and sheets

Loops

_____ are Secondary structures that are not helices, sheets, or recognizable turns

Coils

_____ are also secondary structure which form the first structures after folding of protein's amino acids

Loops and Coils

Combinations of Alpha helices, Beta sheets, coils and loops help form _____

3' structures [Download More Files from VUAnswer.com](http://www.vuanswer.com)

_____, _____ & _____ interactions enforce the 3' structure

Covalent bonds, Hydrogen bonds and hydrophobic

The stability of all the structures of protein is as:

4' > 3' > 2' > 1'

The protein folding into 3D structures leads to:

Reduction in bond angles

Peptide bond between two amino acids is _____

Planar & rigid

The angle between the 1st, overlapped and the 4th points forms a _____

Dihedral angle

_____ can be used to construct the backbone of a protein towards its visualization

C-Alphas

The _____ is used to express the size of atoms, molecules and extremely small biological structures, the lengths of chemical bonds, the arrangement of atoms in crystals.

Angstrom

Atoms of phosphorus, sulfur, and chlorine are _____ in covalent radius, while a hydrogen atom is _____ respectively.

~1 Å & 0.25 Å

Crystallography data gives _____ of atomic coordinates

Relative positions

_____ Proteins are used to determine protein structures.

Crystallized

(Memorize the below file format of protein data base to attempt additional mcq's in exam, for more, you can read ppt of lecture no. 179)

HEADER - Contains a brief description of the structure, the date and the PDB ID code.

TITLE - The title of the structure.

COMPND - Brief details of the structure.

SOURCE - Identifies which organism the structure came from.

KEYWDS - Lists a set of useful words/phrases that describe the structure.

AUTHOR - The scientists depositing the structure.

REVDAT - The date of the last revision.

_____ contains protein structure information

PDB

PDB has the coordinates of C-Alphas for over _____ proteins

50,000

To view a protein, we need to evaluate the:

physical location of its atoms

To trace the backbone of a protein, _____ trace can be used

CA atoms

Coordinates of CA atoms can be obtained from the _____

PDB

Protein structures are visualized using several online tools. These tools include:

Rasmol, CHIME, Swiss PDB Viewer and Cn3D.

CPK stands for:

Corey-Paulin-Koltun Diagrams

In CPK diagrams, each atom is represented by a _____

Solid sphere

Spheres are equal to atomic _____

Van der Waal radius

Ribbon Diagrams represent the protein secondary structures by using:

Simple cartoon figures

Balls & Stick (BS) Models have atoms as _____ and intermediate bonds as _____ respectively
colored balls, sticks

“Proteins fold for a unique, stable and minimum free kinetic energy structure”. The given statement is called:

Anfinsen's thermodynamic hypothesis

There is _____ free energy accessible to each atom for further interactions.

Lesser

The greater the number of bonds, the more _____ a protein becomes.

Stable

The basic idea of thermodynamic stability is to _____ bonding in order to _____ the free energy

Maximize, minimize

We can calculate energy of a folded protein based on the _____ of atomic interactions
number and types

To determine the number of each type of interaction within a protein, we need to find its

inter-atomic distances

Based on specific atomic distances in protein, we can guess the _____

type of atomic interaction

Energies of Interactions

www.ucdavis.edu

Force	Strength (kJ/mol)	Distance (nm)
Van der Waals	0.4-4.0	0.3-0.6
Hydrogen Bonds	12-30	0.3
Ionic Interactions	20	0.25
Hydrophobic Interactions	<40	varies

$$\text{Energy}_{\text{TOTAL}} = \text{Atoms}_{\text{VWF}} \times \text{Energy}_{\text{VWF}} + \text{Atoms}_{\text{HB}} \times \text{Energy}_{\text{HB}} + \text{Atoms}_{\text{IonicInteraction}} \times \text{Energy}_{\text{IonicInteraction}}$$

To determine the structure of protein, we use _____

X-Ray or NMR

If two atoms are participating in a covalent bond, their distance is:

~0.96Å

In case of hydrogen bond formation between atoms, the inter-atomic distance is :

~1.97Å

X-Ray data should have a minimum of _____ resolution

1.97Å

,"Helix Formers" are generally _____ amino acids (M, A, L...)

Hydrophobic

Alpha Helices are formed by hydrogen bonding (O-H) between _____ atoms in the protein backbone

C_i and N_{i+4}

_____ residues are needed to make a Beta Strand

5 to 10

Download More Files from VUAnswer.com

The sub-structures of beta-sheets are:

- **Beta Strand**
- **Beta Sheet**
- **Beta Barrel**
- **Beta Sandwiches**

_____ is made of a single beta sheet that twists and coils upon itself.

Beta-barrel

Beta Sandwiches are made of two beta sheets which are usually _____ so their strands are aligned

twisted and packed

Loops are formed by amino acids present in the middle of _____ in a protein backbone

the Alpha Helices and Beta Sheets

Variability in _____ allows loops to join Alpha Helices and Beta Sheets in a variety of ways.

length and conformation

Loops are mostly comprised of _____ amino acids

charged and polar

Hairpin loops are two amino acids long and join _____ Beta strands

anti-parallel

Essentially, a secondary structure which is not a helix, sheet or loop, is a _____

coil

Coils are apparently _____ regions

disordered

_____ are semi-independent functional structures in a protein

Domains

Domains have over _____ residues

~40

Download More Files from VUAnswer.com

_____ interact (H-bonds) more internally than externally

Locally Compact – Domains

Domains have a _____ core

hydrophobic

various types of domains are:

- **Alpha Domains**
- **Beta Domains**
- **Alpha/Beta Domains**
- **Alpha + Beta Domains**
- **Alpha & Beta Multi-Domains**
- **Membrane & cell-surface proteins**

(must see the structures of these domains in lecture 196)

Domains can be classified into structural classes. Classes can be further classified into

Architecture and Topologies

_____ classifies proteins by their structural similarity

CATH

FSSP stands for _____ based on the DALI algorithm

Family of Structurally Similar Proteins

SCOP stands for:–

Structural Classification of Proteins

So for two different proteins, sharing the same domain, we may want to compare:

only a portion of the overall structure i.e. a domain

Whole protein structures can be compared by calculating the _____ between their Alpha Carbons positions

root mean squared difference (RMSD)

Download More Files from VUAnswer.com

VU Biologists

www.facebook.com/groups/vubiologists

The _____ the RMSD, the similar are the proteins

Lower

Full protein structures can be compared and ranked by the overall _____ in positions between their Alpha Carbons

differences

_____ tells us about the quality of the matches

RMSD

_____ mostly found in Alpha Helices

Alanine

The first such algorithm to predict 2' given an amino acid sequence was the:

Chou-Fasman Algorithm

we only know _____ 3D protein structures, but 10 times more sequences

100,000

For a primary sequence, and a tentative 2' structure, _____ can help us compute the overall propensity

propensity table

___ amino acids are needed to start an Alpha Helix and ___ amino acids for Beta Sheet

4,5

Loops are small amino acids

~ 3-4

Alpha Helices are formed from 4 contiguous amino acids having an Alpha-Helix propensity over _____

1.0

Download More Files from VUAnswer.com

