

Question # 10 of 10 (Start time: 03:18:29 AM, 05 February 2021)

If the median of an arrangement of numbers is equal to the mean of its middle terms then the arrangement contains

Select the correct option



Odd number of terms



Even number of terms



Unlimited number of terms



Prime number of terms

Question # 9 of 10 (Start time: 03:17:43 AM, 05 February 2021)

The measure of central tendency which divides the data into two equal parts is called....

Select the correct option



Mode



Median



Mean



Geometric mean

Question # 8 of 10 (Start time: 03:17:13 AM, 05 February 2021)

Ogives are used to determine at different values.

Select the correct option



Frequency



Frequency distribution



The cumulative Frequency



None of these

Question # 7 of 10 (Start time: 03:17:00 AM, 05 February 2021)

..... is the mean of inverse values.

Select the correct option



Harmonic mean



Geometric mean



Arithmetic mean



Midhing

The most frequent value in the data is called

Select the correct option



Mean



Median



Mode



Midrange

Question # 5 of 10 (Start time: 03:15:13 AM, 05 February 2021)

Last year a small statistical consulting company paid each of its five statistical clerks \$22,000, two statistical analysts \$50,000 each, and the senior statistician/owner \$270,000. The number of employees earning less than the mean salary is:

Select the correct option



4



5



6



7



Question # 4 of 10 (Start time: 03:14:47 AM, 05 February 2021)

In Positively Skewed Distribution, which of the following is true

Select the correct option



Mean > median > mode



Mode > median > mean



Mean > mode > median



None of these

Question # 3 of 10 (Start time: 03:14:02 AM, 05 February 2021)

Mode of the data 25,35,21,56,25,56,65 is

Select the correct option



35



25 & 56



21



65



The frequency of the required class can be obtained from the cumulative frequency

Select the correct option



by subtracting the cumulative frequency of the relative class from the next cumulative frequency



by subtracting the cumulative frequency of the relative class from the total frequency



by subtracting the cumulative frequency of the relative class from the previous cumulative frequency



None of these

The frequency of the required class can be obtained from the cumulative frequency

Select the correct option



by subtracting the cumulative frequency of the relative class from the next cumulative frequency



by subtracting the cumulative frequency of the relative class from the total frequency



by subtracting the cumulative frequency of the relative class from the previous cumulative frequency



None of these

Question # 1 of 10 (Start time: 03:12:01 AM, 05 February 2021)

If the mean of the data is 50 and sum of the data is 450 then the number of observations in the data will be

Select the correct option



9



500



22500



None of these

Question # 10 of 10 (Start time: 01:42:11 AM, 05 February 2021)

The formula for mean is

Select the correct option



The sum of values /2



The sum of values/total no of values



The sum of value



none of these

Question # 9 of 10 (Start time: 01:41:12 AM, 05 February 2021)

Geometric mean of 130, 140, 160 is -----

Select the correct option



130.5



135.5



142.8



145.7

View

Question # 8 of 10 (Start time: 01:39:58 AM, 05 February 2021)

In stem and leaf display stem shows

Select the correct option



Decimal number



Unit number



Thousandth number



None of these

Which of these numbers be a probability?

Select the correct option



0



1.001



1.002



1.003

Probability of occurrence of an event A = ...

(Where X=Number of outcomes favorable to event A and n=total number of outcomes)

Select the correct option

☐ nX

☒ n/X

☐ X/n

☐ None of these

AM Knowledge

The following formula for probability

$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$ is called

Select the correct option



The law of addition



The law of subtraction



The law of multiplication



None of these

There are 5 white balls and 4 black balls. In how many ways can we select 3 white and 2 black balls?

Select the correct option



30



40



50



60

AM Knowledge Official

The moving averages can be used for

Select the correct option

- ☐ better graph
- ☒ forecasting purpose
- ☐ middle value
- ☐ median



When two events are mutually exclusive, the probability of either one of those occurring is the sum of individual probabilities. This is called the

Select the correct option



OR Rule



AND Rule



OR-AND Rule



AND-OR Rule

AM Knowledge Official

Question # 7 of 10 (Start time: 01:38:31 AM, 05 February 2021)

Fifth decile for 10.1, 10.3, 10.5, 10.6, 10.7, 10.9 is approximately _____

Select the correct option



10.1



10.3



10.6



10.9

Question # 6 of 10 (Start time: 01:37:52 AM, 05 February 2021)

In the arrangement of numbers,by following stem and leaf format a number say '27' is located by taking

Select the correct option



2 as its leaf and 7 as its stem



2 as its stem and 7 as its leaf



27 as its stem with nil leaf



Nil as its stem 27 as a leaf

Question # 5 of 10 (Start time: 01:37:08 AM, 05 February 2021)

Trimmed mean is the mean of

Select the correct option



The remaining data



The middle data



The data from last



None of these

Truncated mean is also called

Select the correct option



Arithmetic mean



Trimmed mean



Harmonic Mean



None of these

Question # 3 of 10 (Start time: 01:35:41 AM, 05 February 2021)

For the class 165 - 184 the mid point will be



Select the correct option



172.5



174.5



176.5



To find mid point at least two classes are required.

Question # 2 of 10 (Start time: 01:35:06 AM, 05 February 2021)

Which of the following statements is true for Positively Skewed Distribution?

Select the correct option



Mean > median > mode



Mean = median = mode



Mode > median = mean



None of above

Central tendency is tendency of data values to ----- in the central part of data

Select the correct option



Cluster



Disperse



both of these



None of these

Class width of the given data is defined as ratio of

Select the correct option



Range and number of values in the given data



Range and the number of classes



Range and mid point/value



None of these

The average of smallest and largest values in the data is called....

Select the correct option



Range



Midrange



Interquartile range



Midhinge

Question # 8 of 10 (Start time: 01:20:51 AM, 05 February 2021)

The average of the first and third quartiles is called...

Select the correct option



Range



Midrange



Interquartile range



Midhinge



Question # 7 of 10 (Start time: 01:19:48 AM, 05 February 2021)

If the mean of the data is 50 and sum of the data is 450 then the number of observations in the data will be

Select the correct option



9



500



22500



None of these

Average is calculated as

Select the correct option



Sum of observation/number of observations



Number of observations /Sum of observation



Sum of observation-number Of observations



Sum of observation+number of observations

.....is used in compounding such as investments that are accumulated over a period of time

Select the correct option



Harmonic mean



Geometric mean



Arithmetic mean



Mid range

Question # 4 of 10 (Start time: 01:17:54 AM, 05 February 2021)

The formula for mean is

Select the correct option



The sum of values /2



The sum of values/total no of values



The sum of value



none of these



Bivariate categorical data can be organized as

Select the correct option



The Summary Table



Contingency Tables



Side by Side Bar charts



Only option 2 & 3



Question # 2 of 10 (Start time: 01:15:17 AM, 05 February 2021)

Mode of a set of values can be

Select the correct option



Exactly one



Two



Three



More than one



In an ordered array of data having even number of values, median is the . . .

Select the correct option



Sum of 2 middle values.



average of 2 middle values.



average of 4 middle values.



difference of 2 middle values.

When a coin is tossed once, the probability of getting head is

Select the correct option



0.50



0.51



0.52



0.53

The number of terms in the expansion of the binomial $(p+q)^n$ is

Select the correct option



n



$n-1$



$n+1$



$2n$



AM Kno

Which of these numbers cannot be a probability?

Select the correct option



0.5



0



1



1.001



A worker out of 500 gets a prize by lotter. What is the chance of any one individual say Zain being selected?

Select the correct option



1/500



-1/500

AM Knowledge Official



1/600



-1/600