MIC501P – MICROBIOLOGY

Practical Exam Notes

Prepared by: *Sir Mughal*

Main Group link: https://chat.whatsapp.com/DWXwJpXahbxFVQutey8JXN

Subject Group link: https://chat.whatsapp.com/KpornC5HiMQ5P4aLcuS9ai

M.C.Qs (Multiple Choice Questions)

1. Color of Gram Positive Bacteria is **PURPLE** or **VIOLET**.
2. Color of Gram Negative Bacteria is **PINK** or **RED**.
3. L.B is a standard media used to grow **E. COLI**.
4. Molds are identified on the basis of **MORPHOLOGY**.
5. Optimum temperature of incubation for bacteria is **37°C**.
6. Generation time of bacteria is **SEVERAL HOURS**.
7. Particular type of bacteria separate in **SOLID MEDIUM**.
8. Which reagent is used in catalase test? **CATALASE REAGENT (H_2O_2 3%)**.
9. Magnifying power of Oil immersion lens is **100x**.
10. Which of these is used for removing microbes and their spores? **AUTOCLAVING** or **STEAM STERILIZATION**.
11. For open manipulation of bacteria, **LAMINAR AIR FLOW CABINET** is used.
12. How many biosafety levels are there? **4**
13. Amino Acids, Proteins and Urea provide **NITROGEN**.
14. Disk Diffusion method is used for **ANITMICROBIAL SUSCEPTIBILITY**.
15. The phase in which growth rate is exactly equal to the death rate is called **STATIONARY PHASE**.
16. Lease dangerous biosafety level to human health is **BIOSAFETY LEVEL 1 (BSL-1)**.
17. **GREEN PIGMENTS** are produced by Psuedomonas aeruginosa.
18. **BSL-4** have the strictest methods because dealing with agents that are most dangerous to human health.
19. The **CENTRIFUGE** is used to separate fractions.
20. **SPECTROPHOTOMETER** is used to measure optical density.
21. STERILIZATION refers to total destruction of all microorganisms.
22. Gram staining is categorized as **DIFFERENTIAL STAINING**.
23. In **LOG PHASE / EXPONENTIAL PHASE**, the bacterial biomass increase linearly with time.
24. **SPORES** cannot be killed by alcohol.
25. Temperature of autoclave is **121°C**.
26. In gram staining, **IODINE** is used as a mordant.
27. Lauria broth was first made by **BERTANI**.
28. Reagent of indole test is called **KOVAC's** or **EHRLICH's**.

Compiled by: Sir Mughal

Get in touch with us:
Sir Mughal: https://wa.me/923132310940
Sir Jawad: https://wa.me/923036100859

SHORT QUESTIONS

1. What is indole test?

Indole test is a biochemical test used in microbiology to determine whether a bacterium can break down the amino acid tryptophan to produce indole using the enzyme tryptophanase.

In this test, bacteria are grown in a medium containing tryptophan (commonly peptone water or tryptone broth). After incubation, a few drops of Kovac's reagent are added. If indole is present, it reacts with the reagent to form a red or cherry-red layer on the top of the medium, indicating a positive result.

A positive indole test means the organism produces tryptophanase and can convert tryptophan into indole, pyruvate, and ammonia. A negative result shows no color change, meaning the organism lacks this enzyme.

This test is commonly used to differentiate members of the *Enterobacteriaceae* family, for example, *Escherichia coli* (indole positive) from *Klebsiella pneumoniae* (indole negative).

2. Define Agar plate, Agar stab and broth culture?

Agar plate: An agar plate is a sterile Petri dish containing a solid growth medium made by mixing nutrients with agar. It is used to grow, isolate, and study microorganisms on a solid surface where colonies can be observed.

Agar stab: An agar stab is a technique where microorganisms are introduced into a semi-solid agar medium by inserting a straight inoculating needle deep into the agar. It is used to study bacterial growth patterns, oxygen requirements, and motility.

Broth culture: A broth culture is a liquid nutrient medium in which microorganisms are grown. The bacteria are inoculated into the liquid and incubated, allowing them to multiply freely throughout the medium, often resulting in turbidity (cloudiness).

3. What are components of L.B broth?

Components of L.B. (Luria-Bertani) broth:

LB broth is a commonly used nutrient-rich liquid medium for bacterial growth. Its main components are:

- ***Tryptone (peptone):*** Provides amino acids and peptides as a nitrogen and carbon source.
- ***Yeast extract:*** Supplies vitamins (especially B-complex), growth factors, and additional nutrients.
- ***Sodium chloride (NaCl):*** Maintains osmotic balance and helps bacterial cell stability.
- ***Distilled water:*** Acts as the solvent for all components.
- ***pH (usually adjusted to ~7.0):*** Ensures optimal bacterial growth conditions.

4. What are modes of action of chemical sterilization?

Chemical sterilizing agents kill or inhibit microorganisms through several mechanisms:

- ***Protein denaturation:*** Chemicals disrupt hydrogen bonds and other interactions in proteins, causing enzymes and structural proteins to lose function (e.g., alcohols, phenols).
- ***Cell membrane disruption:*** Some chemicals damage the lipid bilayer, increasing permeability and causing leakage of cellular contents (e.g., detergents, alcohols).
- ***Oxidation of cellular components:*** Oxidizing agents destroy essential cellular molecules like proteins, lipids, and DNA by adding oxygen or removing electrons (e.g., hydrogen peroxide, chlorine compounds).
- ***Alkylation of proteins and nucleic acids:*** Alkylating agents add alkyl groups to DNA and proteins, preventing replication and enzyme activity (e.g., formaldehyde, ethylene oxide).
- ***Enzyme inactivation:*** By binding to active sites or altering structure, chemicals stop metabolic activity essential for survival.

Compiled by SIR MUGHAL

Get in touch with us:
Sir Mughal: https://wa.me/923132310940
Sir Jawad: https://wa.me/923036100859

5. Describe Antimicrobial Susceptibility Test.

Antimicrobial Susceptibility Test (AST):

Antimicrobial Susceptibility Testing is a laboratory method used to determine the effectiveness of different antimicrobial drugs against a specific microorganism. It helps to identify which antibiotics can inhibit or kill the bacteria and which are resistant.

In this test, a pure culture of bacteria is first prepared and then spread evenly on the surface of a suitable agar medium (commonly Mueller-Hinton agar). Antibiotic-impregnated discs are placed on the agar surface. The plate is then incubated to allow bacterial growth.

After incubation, clear zones around the antibiotic discs, called zones of inhibition, are observed. These zones indicate that bacterial growth has been prevented by the antibiotic. The size of the zone is measured in millimeters and compared with standard charts to classify the organism as sensitive, intermediate, or resistant.

Common methods include the Kirby-Bauer disc diffusion method, broth dilution, and automated systems. This test is widely used in clinical laboratories to guide appropriate antibiotic therapy and prevent antibiotic resistance.

6. Define Light Microscope. Name three types.

Light Microscope:

A light microscope is an optical instrument that uses visible light and a system of lenses to magnify small objects, allowing detailed observation of cells and microorganisms that cannot be seen with the naked eye.

Three types of light microscopes:

- ***Simple microscope*** – uses a single lens for low magnification.
- ***Compound microscope*** – uses two or more lenses (objective and eyepiece) for higher magnification and better resolution.
- ***Stereo (dissecting) microscope*** – provides a three-dimensional view of larger specimens at low magnification.

7. Explain dry heat method.

Dry heat method (Dry heat sterilization):

Dry heat sterilization is a method of killing microorganisms and their spores using high temperature without moisture. It destroys microbes mainly through oxidation of cell components, protein denaturation, and dehydration of cells.

In this method, sterilization is carried out using devices like a hot air oven or direct flame. The hot air oven is commonly used in laboratories and operates at temperatures such as 160°C for 2 hours or 170°C for 1 hour, depending on the material being sterilized.

Dry heat is especially suitable for materials that can withstand high temperatures but may be damaged by moisture, such as glassware, metal instruments, powders, oils, and fats.

Compared to moist heat, dry heat requires higher temperatures and longer exposure time to achieve effective sterilization.

Compiled by: SIR MUGHAL

Get in touch with us:
Sir Mughal: https://wa.me/923132310940
Sir Jawad: https://wa.me/923036100859

8. Write down the procedure of oxidase test.

Procedure of Oxidase Test:

The oxidase test is used to detect the presence of cytochrome c oxidase enzyme in bacteria.

Procedure:

- Take a clean glass slide or filter paper strip.
- Place a small amount of oxidase reagent (commonly tetramethyl-p-phenylenediamine dihydrochloride**)** on it.
- Using a sterile wooden stick or glass rod (not a metal loop), pick a fresh colony of the test organism.
- Smear the colony onto the reagent-impregnated area.
- Observe the color change within 10–30 seconds**.**

Result interpretation:

- ***Positive test:*** Development of a dark purple/blue color indicates presence of cytochrome c oxidase.
- ***Negative test:*** No color change or a very slight change after 30 seconds indicates absence of the enzyme.

9. What is CFU?

CFU (Colony Forming Unit):

CFU is a measure used in microbiology to estimate the number of viable (living) microorganisms in a sample that are capable of forming colonies on a solid growth medium. Each CFU represents either a single microbial cell or a group of cells that grow together to form one visible colony on an agar plate. CFU is commonly expressed as CFU/mL (for liquid samples) or CFU/g (for solid samples) and is calculated after performing serial dilution and plate counting. It is widely used in food microbiology, clinical diagnostics, and water quality testing to determine microbial load.

10. What is diffusion sterilization method?

Diffusion sterilization method:

Diffusion sterilization is a chemical sterilization technique in which a sterilizing agent spreads in the form of vapors or gases and penetrates all areas of the material or enclosed space to destroy microorganisms, including spores.

In this method, volatile chemical agents such as ethylene oxide gas, formaldehyde vapor, or hydrogen peroxide vapor diffuse through air or within a closed chamber and come in contact with microorganisms. These agents kill microbes by alkylating proteins and nucleic acids, denaturing enzymes, or oxidizing cellular components.

The process is usually carried out in a sealed sterilization chamber under controlled conditions of temperature, humidity, and exposure time to ensure complete sterilization.

It is commonly used for sterilizing heat-sensitive medical instruments, plastics, surgical equipment, and laboratory materials that cannot withstand high temperatures.

11. What are the effects of Gas sterilization? Highlight one disadvantage.

Effects of Gas Sterilization:

Gas sterilization (commonly using ethylene oxide or formaldehyde gas) affects microorganisms in the following ways:

- ***DNA damage:*** The gas alkylates nucleic acids, preventing DNA replication and transcription.
- ***Protein inactivation:*** It reacts with functional groups in proteins and enzymes, leading to loss of metabolic activity.
- ***Spore destruction:*** Highly resistant bacterial spores are also destroyed due to disruption of essential cellular components.
- ***Cell membrane disruption:*** Alters permeability and damages membrane structure, leading to cell death.

Compiled by: SIR MUGHAL

Get in touch with us:
Sir Mughal: https://wa.me/923132310940
Sir Jawad: https://wa.me/923036100859

Disadvantage:

- Gas sterilization is toxic and potentially harmful to humans, so strict handling, aeration, and safety precautions are required after sterilization.

12. What is significance of serial dilution in plate cell counted?

Significance of serial dilution in plate cell counting:

Serial dilution is important in plate cell counting because it helps to obtain a countable number of microbial colonies from a very concentrated sample. Its main significance includes:

- ***Reduces microbial concentration:*** It gradually decreases the number of microorganisms step by step, making it easier to plate an appropriate dilution.
- ***Obtains countable plates:*** Ensures that colonies fall within the ideal range (usually 30–300 CFU per plate**)** for accurate counting.
- ***Improves accuracy:*** Helps in estimating the actual number of viable cells in the original sample using CFU calculations.
- ***Avoids overcrowding:*** Prevents formation of confluent growth where colonies merge and cannot be counted individually.
- ***Facilitates quantitative analysis:*** Allows calculation of microbial load in terms of CFU/mL or CFU/g.

13. Write use of microscope, cover slip and slide.

Uses of microscope:

- To observe microorganisms such as bacteria, fungi, and protozoa that are not visible to the naked eye.
- To study cell structure, shape, and arrangement.
- To examine tissues, blood cells, and biological samples in laboratories.
- To support diagnosis in medical microbiology and pathology.

Uses of cover slip:

- To cover the specimen on the slide and protect it from contamination and drying.
- To flatten the specimen into a single thin layer for clearer viewing.
- To prevent damage to the microscope lens by avoiding direct contact with the sample.

Uses of slide:

- To hold the specimen or sample for microscopic observation.
- To provide a flat, transparent surface for placing biological materials.
- To support staining and preparation of samples for detailed study.

14. Which type of media is appropriate if we study the nutrient requirements from newly born bacteria and why?

Appropriate media: *Defined (synthetic) medium*

Reason:

A defined medium is the most suitable when studying the nutritional requirements of newly born bacteria because its exact chemical composition is known.

- It contains precisely measured nutrients (specific carbon, nitrogen, salts, vitamins, etc.).
- Since all components are known, we can systematically add or remove one nutrient at a time.
- This helps to identify which nutrients are essential for bacterial growth and development.
- It allows accurate study of the metabolic and nutritional needs of newly formed bacterial cells without interference from unknown ingredients (unlike complex media like nutrient broth or LB broth).

Compiled by: SIR MUGHAL

Get in touch with us:
Sir Mughal: https://wa.me/923132310940
Sir Jawad: https://wa.me/923036100859

15. Write significance of biosafety levels.

Significance of Biosafety Levels (BSL):

Biosafety levels are important laboratory containment guidelines that ensure safe handling of microorganisms based on their risk of infection and hazard level**.**

- ***Protect laboratory personnel:*** They reduce the risk of infection by using appropriate safety practices, equipment, and facility design.
- ***Prevent environmental contamination:*** They ensure that harmful microorganisms do not escape into the surrounding environment.
- ***Control exposure risk:*** Different biosafety levels (BSL-1 to BSL-4) match the danger level of pathogens being handled.
- ***Standardize laboratory safety:*** They provide internationally accepted procedures for working safely with infectious agents.
- ***Enable safe research:*** Allow scientists to study dangerous pathogens under controlled and secure conditions.
- ***Protect public health:*** Help prevent accidental outbreaks or spread of infectious diseases from laboratories.

16. Define media and its types.

Media (Culture Media):

Media are nutrient preparations used in the laboratory to support the growth, multiplication, and maintenance of microorganisms under controlled conditions.

Types of media:

- ***Based on physical state:***
  - ***Liquid (broth) media –*** e.g., nutrient broth, used for mass growth of bacteria.
  - ***Solid media –*** contains agar, used for colony isolation (e.g., nutrient agar).
  - ***Semi-solid media –*** low agar concentration, used for motility studies.

- ***Based on chemical composition:***
  - ***Defined (synthetic) media –*** exact chemical composition is known.
  - ***Complex (undefined) media –*** contains extracts like yeast extract or peptone with unknown exact composition.

- ***Based on function:***
  - ***Selective media –*** allows growth of specific organisms while inhibiting others.
  - ***Differential media –*** distinguishes organisms based on visible changes.
  - ***Enriched media –*** contains extra nutrients for fastidious organisms.
  - ***Transport media –*** used to preserve and transport samples without growth.

17. What is the primary objective of streaking bacteria in a solid medium by a quadrant streaking method?

The main objective of streaking bacteria using the quadrant streaking method is to obtain isolated pure colonies from a mixed or concentrated bacterial sample. By gradually diluting the number of bacteria across four quadrants of a solid agar plate, individual cells become separated and grow into distinct colonies. Each isolated colony is assumed to originate from a single bacterial cell, which allows:

- Isolation of pure cultures
- Separation of different bacterial species from a mixed sample
- Easy study of colony characteristics and identification of microorganisms

Compiled by: SIR MUGHAL

Get in touch with us:
Sir Mughal: https://wa.me/923132310940
Sir Jawad: https://wa.me/923036100859

18. What are membrane filters?

Membrane filters:

Membrane filters are thin, porous filtration membranes made of materials such as cellulose acetate, nitrocellulose, or polycarbonate that are used to separate microorganisms and particles from liquids or gases. They work by physically trapping microorganisms on the surface of the filter while allowing the liquid or air to pass through. The pore size is usually very small (commonly 0.22 µm or 0.45 µm) which is effective for removing bacteria. Uses include:

- Sterilization of heat-sensitive liquids (e.g., antibiotics, serum)
- Microbiological testing of water for bacterial contamination
- Air filtration in sterile environments
- Separation and concentration of microorganisms for study

19. How to prepare LB broth?

Preparation of LB (Luria-Bertani) broth:

Materials required:

- Tryptone – 10 g/L
- Yeast extract – 5 g/L
- Sodium chloride (NaCl) – 10 g/L
- Distilled water – 1 liter
- pH meter or pH paper
- Conical flask, balance, and autoclave

Procedure:

- Measure 1 liter of distilled water in a conical flask.
- Weigh and add 10 g tryptone**,** 5 g yeast extract, and 10 g NaCl into the water.
- Stir well until all components dissolve completely.
- Check and adjust the pH to around 7.0 if needed.
- Cover the flask with cotton plug or cap loosely.
- Sterilize the medium in an autoclave at 121°C for 15 minutes (15 psi pressure)**.**
- After sterilization, allow it to cool before use.

20. What is generation time of bacteria?

Generation time of bacteria:

Generation time is the time required for a bacterial cell to divide and form two daughter cells (i.e., the time needed for one complete cell division cycle). In simpler terms, it is the time taken for a bacterial population to double in number under ideal growth conditions.Generation time varies among bacteria depending on species and environmental conditions such as temperature, nutrients, and pH. For example, *Escherichia coli* has a generation time of about 20 minutes under optimal conditions, while some bacteria may take several hours or even days.

Compiled by SIR MUGHAL

Get in touch with us:
Sir Mughal: https://wa.me/923132310940
Sir Jawad: https://wa.me/923036100859

21. Write difference between bright field microscopy, fluorescent field microscopy and dark field microscopy in microbiological sciences.

Differences between Bright Field, Fluorescence, and Dark Field Microscopy

Feature	Bright Field Microscopy	Fluorescence Microscopy	Dark Field Microscopy
Light source	Visible light passes directly through specimen	UV or high-energy light excites fluorophores	Oblique light; only scattered light enters objective
Background	Bright	Dark	Dark
Appearance of specimen	Dark or stained specimen	Brightly glowing colored specimen	Bright, glowing specimen
Need for staining	Usually required	Fluorescent dyes required	Not required
Contrast	Low to moderate	Very high and specific	High
Type of observation	General morphology of cells	Specific molecules, proteins, DNA	Live, thin, unstained organisms
Common use	Routine lab observation	Molecular and diagnostic studies	Observation of spirochetes and small live microbes

22. How AST (Antimicrobial Susceptibility Testing) results are interpreted?

Interpretation of AST (Antimicrobial Susceptibility Testing) results:

AST results are interpreted by measuring the zone of inhibition around antibiotic discs (Kirby-Bauer method) or by determining MIC values in dilution methods.

1. Disc diffusion method (Zone of inhibition):

- After incubation, the clear area around each antibiotic disc is measured in millimeters (mm).
- The size of the zone is compared with standard reference charts (CLSI/EUCAST guidelines).
- Based on this comparison, the organism is classified as:

Category	Meaning
Sensitive (S)	Antibiotic effectively inhibits bacterial growth; likely to work in treatment
Intermediate (I)	Antibiotic has limited effectiveness; may work at higher dose or in specific conditions
Resistant (R)	Antibiotic does not inhibit growth; treatment is likely ineffective

2. MIC method (Minimum Inhibitory Concentration):

- MIC is the lowest concentration of an antibiotic that prevents visible bacterial growth.
- The MIC value is compared with standard breakpoints to classify bacteria as S, I, or R.

Compiled by: SIR MUGHAL

Get in touch with us:
Sir Mughal: https://wa.me/923132310940
Sir Jawad: https://wa.me/923036100859

23. Write a short note on bacterial morphology and arrangements.

Bacterial morphology refers to the shape, size, and structural appearance of bacterial cells. Bacteria show different basic shapes such as:

- ***Cocci (spherical):*** Round-shaped bacteria.
- ***Bacilli (rod-shaped):*** Cylindrical or elongated bacteria.
- ***Spirilla (spiral-shaped):*** Rigid helical forms.
- ***Vibrios (comma-shaped):*** Curved rod-shaped bacteria.

Some bacteria may also appear as coccobacilli (intermediate between cocci and bacilli) or filamentous forms. Bacterial arrangement refers to how bacterial cells are organized after cell division, depending on the plane of division and whether cells remain attached. Common arrangements include:

- ***Single cells:*** Independent bacteria.
- ***Diplococci / diplobacilli:*** Pairs of cells.
- ***Streptococci / streptobacilli:*** Chains of cells.
- ***Staphylococci:*** Irregular grape-like clusters.
- ***Tetrads:*** Groups of four cells.
- ***Sarcinae:*** Cubical packets of eight or more cells.

24. Differentiate between graduated pipettes and Pasteur pipettes.

Difference between Graduated pipettes and Pasteur pipettes:

Feature	Graduated Pipettes	Pasteur Pipettes
Structure	Long glass/plastic tube with calibrated markings	Simple glass/plastic tube with a narrow tip and rubber bulb
Measurement	Allows accurate measurement of specific volumes	No accurate volume measurement
Precision	High precision and accuracy	Low precision, approximate transfer only
Use	Used in quantitative laboratory work	Used for transfer of liquids in microbiology
Control of volume	Can deliver different measured volumes	Delivers drops or small amounts of liquid
Calibration	Graduated markings present	No calibration
Typical application	Chemical analysis, microbiology quantification	Culture handling, reagent transfer, staining procedures

25. Why it is crucial to flame inoculation loop in process of streaking?

Flaming the inoculation loop is crucial in the streaking process to ensure aseptic conditions and prevent contamination.

- ***Sterilization of loop:*** Flaming kills any microorganisms present on the loop from previous use.
- ***Prevents cross-contamination:*** It avoids transfer of unwanted microbes between samples or streaking areas.
- ***Ensures pure culture isolation:*** Helps obtain isolated colonies from a single organism.
- ***Maintains laboratory safety:*** Reduces risk of spreading infectious or environmental microbes.

Compiled by: SIR MUGHAL

Get in touch with us:
Sir Mughal: https://wa.me/923132310940
Sir Jawad: https://wa.me/923036100859

26. Why IMVIC test is used? What results are expected?

IMViC test: purpose and expected results

Why IMViC test is used:

The IMViC test is a group of biochemical tests used in microbiology to differentiate and identify members of the *Enterobacteriaceae* family, especially to distinguish coliform bacteria (like *Escherichia coli*) from non-coliforms (like *Enterobacter* and *Klebsiella*).

IMViC stands for:

- **I** – Indole test
- **M** – Methyl Red test
- **V** – Voges–Proskauer test
- **C** – Citrate utilization test

Expected results:

Organism	Indole	Methyl Red (MR)	Voges-Proskauer (VP)	Citrate
Escherichia coli	+	+	–	–
Enterobacter aerogenes	–	–	+	+
Klebsiella pneumoniae	–	–	+	+

27. Why streaking of bacteria is necessary?

Importance of streaking bacteria:

Streaking of bacteria is necessary in microbiology to isolate and separate individual bacterial cells from a mixed population so that they can grow into pure colonies.

- ***Isolation of pure culture:*** It helps obtain a single type of microorganism from a mixed sample.
- ***Separation of colonies:*** Dilutes bacteria across the agar surface so individual colonies can be seen clearly.
- ***Identification:*** Pure colonies allow study of colony morphology and biochemical characteristics.
- ***Avoids contamination:*** Ensures that only one species is studied for accurate results.
- ***Further testing:*** Pure cultures are required for antibiotic sensitivity tests and biochemical analysis.

28. Write advantages and disadvantages of ethylene oxide.

Advantages:

- ***Highly effective sterilant:*** Kills bacteria, viruses, fungi, and even resistant spores.
- ***Good penetration ability:*** Can sterilize packaged and complex instruments (e.g., catheters, syringes).
- ***Low temperature process:*** Suitable for heat- and moisture-sensitive materials like plastics and rubber.
- ***Wide application:*** Used in medical devices, surgical instruments, and laboratory equipment.

Disadvantages:

- ***Toxic and carcinogenic:*** Exposure can be harmful to human health and requires strict safety measures.
- ***Long processing time:*** Sterilization and aeration take several hours to remove toxic residues.
- ***Highly flammable and explosive:*** Requires controlled handling and special equipment.
- ***Residue risk:*** May leave chemical residues on materials if not properly aerated.
- ***Expensive setup:*** Requires specialized sterilization chambers and monitoring systems.

Compiled by SIR MUGHAL

Get in touch with us:
Sir Mughal: https://wa.me/923132310940
Sir Jawad: https://wa.me/923036100859

LONG QUESTIONS

1. Explain Biosafety levels.

Biosafety Levels (BSL):

Biosafety levels are a system of laboratory containment practices designed to protect laboratory personnel, the environment, and the public from exposure to infectious biological agents. They classify laboratories into four levels (BSL-1 to BSL-4) based on the risk group of microorganisms, their infectivity, severity of disease, and route of transmission.

BSL-1 (Basic level):

BSL-1 is used for well-characterized organisms that are not known to consistently cause disease in healthy adults. These organisms present minimal potential hazard. Examples include non-pathogenic laboratory strains such as *Escherichia coli* K-12. Work is performed on open laboratory benches using standard microbiological practices. Basic personal protective equipment such as lab coats and gloves is used, and hand hygiene is strictly followed. No special containment equipment is required, and the laboratory design is similar to a standard teaching laboratory with a sink and easy-to-clean surfaces.

BSL-2 (Moderate risk level):

BSL-2 is used for organisms that pose moderate hazards to personnel and the environment. These may cause human disease but are generally treatable. Examples include *Staphylococcus aureus*, *Salmonella* species, and Hepatitis B virus. Laboratory access is restricted to trained personnel. Procedures that may generate aerosols are performed in a Class II biosafety cabinet. Autoclaving facilities are required for decontamination of waste. Additional precautions include biohazard signage, sharps safety, and controlled laboratory entry.

BSL-3 (High containment level):

BSL-3 laboratories are designed for pathogens that can cause serious or potentially lethal diseases through inhalation. Examples include *Mycobacterium tuberculosis* and SARS-associated coronaviruses. Strict access control is implemented, and only authorized personnel are allowed inside. Laboratory personnel are required to use respiratory protection. The facility is designed with negative air pressure to prevent contaminated air from escaping. All manipulations involving infectious material are conducted inside biosafety cabinets to minimize aerosol exposure.

BSL-4 (Maximum containment level):

BSL-4 is the highest biosafety level and is used for extremely dangerous and exotic pathogens that have a high risk of life-threatening disease and no available treatment or vaccine. Examples include Ebola virus and Marburg virus. Personnel must wear full-body, air-supplied positive pressure suits. The laboratory is completely isolated from other areas and has its own dedicated air supply and waste decontamination systems. Entry and exit involve multiple airlocks and decontamination showers to prevent contamination release.

2. What are basic constituents of media required for microbial growth?

Basic constituents of media required for microbial growth:

Culture media are specially prepared nutrient formulations that provide all essential requirements for the growth, multiplication, and maintenance of microorganisms under laboratory conditions. The basic constituents of microbial growth media include several chemical components that fulfill nutritional and physiological needs of microorganisms.

Compiled by SIR MUGHAL

Get in touch with us:
Sir Mughal: https://wa.me/923132310940
Sir Jawad: https://wa.me/923036100859

Water: Water is the most essential component of all culture media. It acts as a universal solvent in which all nutrients are dissolved and also serves as a medium for biochemical reactions inside microbial cells. Without water, microbial metabolism and growth cannot occur.

Carbon source: Carbon is required for the synthesis of cellular components such as proteins, carbohydrates, lipids, and nucleic acids. It also provides energy to microorganisms. Common carbon sources include glucose, lactose, and other carbohydrates. Autotrophic microorganisms may use carbon dioxide as their carbon source.

Nitrogen source: Nitrogen is essential for the synthesis of amino acids, proteins, enzymes, and nucleic acids. It is supplied in media through compounds such as peptone, amino acids, ammonium salts, or nitrates.

Mineral salts: Minerals are required in small amounts for enzyme activation, osmotic balance, and structural functions. Important minerals include sodium, potassium, magnesium, calcium, iron, and phosphate ions. These help in maintaining cellular stability and metabolic activity.

Growth factors: Some microorganisms require specific organic compounds for growth because they cannot synthesize them themselves. These include vitamins, amino acids, purines, and pyrimidines. Yeast extract is a common source of growth factors in culture media.

Energy source: Microorganisms require an energy source for metabolic activities. This is usually obtained from carbohydrates or light in phototrophic organisms. The energy source supports ATP production and cellular processes.

Buffering agents: Buffers are added to maintain a stable pH in the medium. Microbial growth often produces acidic or alkaline by-products, and buffering agents prevent harmful pH changes that could inhibit growth.

3. Write components of bacterial colony shape.

Components of bacterial colony shape:

Bacterial colony shape refers to the visible characteristics of a colony formed on solid culture media. It is an important criterion for microbial identification and characterization. The morphology of a colony is described using several components.

***Size*:** Size refers to the diameter of the colony and is usually described as small, medium, or large. It depends on the growth rate and nutrient utilization of the organism. Some bacteria produce pinpoint colonies, while others form large spreading colonies.

***Shape (Form)*:** Shape describes the overall outline of the colony when viewed from above. Common shapes include circular, irregular, filamentous, rhizoid, and punctiform. This characteristic helps in distinguishing different bacterial species.

***Margin (Edge)*:** Margin refers to the border or edge of the colony. It may be smooth (entire), wavy (undulate), lobate, serrated, filamentous, or curled. The nature of the margin reflects the pattern of bacterial growth at the colony boundary.

***Elevation*:** Elevation describes how the colony rises above the agar surface. It may be flat, raised, convex, umbonate, or crateriform. This feature helps in understanding the vertical growth pattern of bacteria.

***Surface appearance*:** This refers to the texture and appearance of the colony surface. It may be smooth, rough, glistening, dull, or wrinkled depending on the bacterial species and capsule production.

Get in touch with us:
Sir Mughal: https://wa.me/923132310940
Sir Jawad: https://wa.me/923036100859

Compiled by SIR MUGHAL

***Opacity*:** Opacity describes how much light passes through the colony. Colonies may be transparent, translucent, or opaque. This property is useful in differentiating species based on pigment and density.

Color (Pigmentation): Some bacteria produce pigments that give colonies distinct colors such as white, yellow, red, or green. Pigmentation may be useful for identification and may also have biological roles like protection from UV radiation.

***Consistency*:** Consistency refers to the physical nature of the colony when touched with a loop. It may be creamy, dry, mucoid, or brittle, depending on extracellular product formation such as polysaccharide capsules.

4. What is Agar Diffusion method?

Agar Diffusion Method:

The agar diffusion method is a microbiological technique used to evaluate the antimicrobial activity of a substance (usually antibiotics or disinfectants) by observing its ability to inhibit the growth of microorganisms on a solid agar medium.

In this method, a uniform lawn of bacteria is first prepared on the surface of a sterile agar plate, commonly Mueller-Hinton agar. After this, small wells are made in the agar or antibiotic-impregnated discs are placed on the surface. The antimicrobial agent then diffuses through the agar medium in all directions, creating a concentration gradient.

As the compound diffuses outward, it inhibits bacterial growth in the surrounding area if the organism is sensitive. After incubation, this inhibition appears as a clear zone around the disc or well, known as the zone of inhibition. The size of this zone depends on the effectiveness of the antimicrobial agent, its diffusion rate, and the susceptibility of the microorganism.

The most common example of this method is the Kirby-Bauer disc diffusion technique, which is widely used in clinical laboratories to determine antibiotic susceptibility patterns and guide appropriate treatment choices.

This method is simple, cost-effective, and widely used for screening antimicrobial activity and comparing the effectiveness of different drugs.

5. Describe streak plate method.

Streak Plate Method:

The streak plate method is a fundamental microbiological technique used to isolate pure colonies of bacteria from a mixed microbial population. It works on the principle of mechanical dilution of microorganisms across the surface of a solid agar medium.

In this method, a sterile inoculating loop is first dipped into a microbial sample containing bacteria. The loop is then streaked over a small section of the agar plate in a back-and-forth motion. This first area contains a high concentration of bacteria. The loop is then sterilized by flaming and cooled, and a portion of bacteria from the first streaked area is dragged into a second quadrant. This process is repeated into third and fourth quadrants, progressively diluting the number of bacterial cells.

As the streaking continues, fewer and fewer bacterial cells are deposited in each successive area. After incubation, isolated single colonies appear in the later streaked sections where individual bacterial cells were separated from the original mixture. Each colony is assumed to arise from a single bacterial cell, forming a pure culture.

The streak plate method is widely used in microbiology for isolation, identification, and purification of bacterial species, as well as for further biochemical and antibiotic sensitivity testing.

Compiled by: SIR MUGHAL

Get in touch with us:
Sir Mughal: https://wa.me/923132310940
Sir Jawad: https://wa.me/923036100859

6. Define Solid medium, Liquid medium and semi solid medium.

Solid medium:

A solid medium is a type of culture medium that contains a solidifying agent, usually agar (1.5–2%), which makes the medium firm and stable. It is used for the isolation of microorganisms and formation of distinct colonies. Bacteria grow as visible colonies on the surface of the medium, allowing morphological studies such as shape, size, and color. Example: Nutrient agar.

Liquid medium:

A liquid medium, also called broth, is a culture medium that does not contain any solidifying agent. It is used for the mass cultivation of microorganisms. In this medium, bacteria grow uniformly throughout the liquid, often causing turbidity, sediment formation, or surface pellicle formation. Example: Nutrient broth, LB broth.

Semi-solid medium:

A semi-solid medium contains a low concentration of agar (0.2–0.5%), making it soft and jelly-like. It is used mainly to study bacterial motility, oxygen requirements, and growth patterns. Motile bacteria spread through the medium, while non-motile bacteria grow only at the inoculation site. Example: Motility test medium.

7. How a surface can be sterilized with 70% ethanol?

Sterilization of surfaces using 70% ethanol:

70% ethanol is commonly used as a chemical disinfectant for surface sterilization in microbiology and laboratory settings because it is highly effective in killing a wide range of microorganisms, including bacteria, fungi, and some viruses.

To sterilize a surface, the area is first cleaned to remove visible dirt or organic matter, as these can reduce the effectiveness of ethanol. Then, 70% ethanol is applied directly to the surface using a sterile cotton swab, wipe, or spray. The surface is thoroughly wetted to ensure complete contact with the disinfectant.

The ethanol works by denaturing proteins and disrupting lipid membranes of microorganisms. The presence of 30% water is important because it slows evaporation and allows better penetration into microbial cells, increasing its effectiveness compared to absolute alcohol.

After application, the ethanol is left on the surface to air dry. During this contact time, microorganisms are destroyed, resulting in a sterile or disinfected surface suitable for laboratory work such as inoculation, streaking, or aseptic handling.

This method is widely used for disinfecting benches, instruments, work areas, and hands (in some cases) before and after microbiological procedures.

8. Describe the procedure of gram staining.

Gram Staining Procedure

1. Preparation of smear: A clean, grease-free glass slide is taken. A small drop of sterile water (if required) is placed, and a thin bacterial smear is prepared using a sterile inoculating loop. The smear is spread evenly to form a thin film.

2. Air drying and heat fixing: The smear is allowed to air dry completely. After drying, the slide is heat-fixed by passing it 2–3 times over a flame. This kills the bacteria and fixes them firmly onto the slide.

3. Primary staining (Crystal violet): The smear is flooded with crystal violet stain and left for about 1 minute. Then it is gently washed with distilled water. All bacterial cells appear purple at this stage.

Compiled by SIR MUGHAL

4. Mordant application (Gram's iodine): Gram's iodine is added to the smear and left for about 1 minute. It forms a crystal violet–iodine (CV-I) complex inside the cell wall, making the dye less soluble.

5. Decolorization step: The slide is washed with alcohol or acetone-alcohol mixture for a few seconds. This is the most critical step. Gram-positive bacteria retain the CV-I complex and remain purple due to thick peptidoglycan layer, while Gram-negative bacteria lose the stain and become colorless.

6. Counterstaining (Safranin): Safranin is applied for about 1 minute and then rinsed with water. Gram-negative bacteria take up the counterstain and appear pink/red, while Gram-positive bacteria remain purple.

7. Microscopic observation: The slide is air-dried and observed under oil immersion (100× objective lens). Bacteria are identified based on their color reaction:
Gram-positive = purple
Gram-negative = pink/red

9. Differentiate between moist heat method and dry heat method. Write their disadvantages and applications with examples.

Difference between Moist Heat Method and Dry Heat Method (Sterilization)

1. Basic principle: Moist heat sterilization works by coagulation and denaturation of proteins in the presence of water. Moisture increases heat penetration and makes it more effective at lower temperatures. Dry heat sterilization works by oxidation, dehydration, and protein denaturation in the absence of moisture, requiring higher temperatures and longer exposure time.

2. Temperature and time: Moist heat requires comparatively lower temperature and shorter time (e.g., autoclaving at 121°C for 15 minutes). Dry heat requires higher temperature and longer time (e.g., hot air oven at 160°C for 2 hours).

3. Mechanism of killing microorganisms: Moist heat kills microbes mainly by coagulating cellular proteins and enzymes. Dry heat kills microbes mainly by oxidative damage and dehydration of cell components.

4. Penetration power: Moist heat has high penetration power due to steam. Dry heat has low penetration power compared to moist heat.

5. Efficiency: Moist heat is generally more efficient and faster. Dry heat is slower and less efficient but suitable for certain materials.

Disadvantages of Moist Heat Method

- Not suitable for heat-sensitive materials like oils, powders, and some plastics.
- May damage moisture-sensitive instruments.
- Requires proper pressure control in autoclaves.
- Cannot be used for materials that are easily corroded by moisture.

Disadvantages of Dry Heat Method

- Requires high temperature and long exposure time.
- Poor penetration compared to steam.
- Not suitable for many heat-labile materials.
- Energy-consuming process.

Compiled by: SIR MUGHAL

Get in touch with us:
Sir Mughal: https://wa.me/923132310940
Sir Jawad: https://wa.me/923036100859

Applications of Moist Heat Method

- Sterilization of culture media and laboratory solutions using autoclave.
- Sterilization of surgical instruments and dressings.
- Pasteurization of milk and beverages.
- Sterilization of wound dressings and medical waste.
- Examples**:** Autoclaving nutrient agar, sterilizing surgical tools, pasteurization of milk.

Applications of Dry Heat Method

- Sterilization of glassware (Petri dishes, test tubes, pipettes).
- Sterilization of metal instruments.
- Sterilization of powders, oils, and fats.
- Incineration of contaminated waste.
- Examples**:** Hot air oven sterilization of glass Petri plates, flaming of inoculating loop, incineration of infected material.

10. How to grow and preserve bacteria in LB broth?

Growth of bacteria in LB broth:

LB (Luria-Bertani) broth is a nutrient-rich liquid medium used for cultivation of bacteria such as *Escherichia coli*. To grow bacteria, a sterile LB broth is first prepared and cooled to room temperature. A small amount of bacterial inoculum (from a colony or glycerol stock) is introduced into the broth under aseptic conditions near a flame or in a laminar flow hood to avoid contamination.

The inoculated broth is then incubated at an appropriate temperature, usually 37°C, with shaking (100–200 rpm) to ensure proper aeration and uniform distribution of nutrients. During incubation, bacteria multiply rapidly and the culture becomes turbid, indicating active growth. Growth can be monitored by measuring optical density using a spectrophotometer at 600 nm (OD600).

Preservation of bacteria grown in LB broth:

Bacteria grown in LB broth can be preserved using short-term or long-term methods. For short-term storage, cultures can be kept at 4°C in a refrigerator for a few days to weeks, slowing down metabolic activity.

For long-term preservation, bacteria from LB culture are mixed with 20–50% glycerol and stored at -20°C or -80°C. Glycerol acts as a cryoprotectant and prevents ice crystal formation that can damage cells. Alternatively, bacteria can be sub-cultured periodically onto fresh LB agar plates to maintain viability.

In some cases, bacteria are also preserved by lyophilization (freeze-drying) for long-term storage, which keeps them viable for years.

Proper aseptic techniques must be followed throughout to prevent contamination and maintain pure cultures.

11. How to view a specimen under light microscope?

1. Preparation of slide: A clean glass slide is taken and the specimen is placed on it. A drop of water (for wet mount) or stain (if required) is added. A cover slip is gently placed to avoid air bubbles.

2. Placement of slide on stage: The prepared slide is placed on the microscope stage and secured with stage clips to prevent movement.

3. Starting with low power objective: The microscope is first adjusted to the low power objective lens (4× or 10×). Coarse adjustment knob is used to bring the specimen into focus. This step helps in locating the specimen.

4. Adjustment of light and focus: The diaphragm is adjusted to control light intensity, and mirror or light source is aligned properly. Fine adjustment knob is then used for clearer focus.

Compiled by: SIR MUGHAL

Get in touch with us:
Sir Mughal: https://wa.me/923132310940
Sir Jawad: https://wa.me/923036100859

5. Switching to higher magnification: After locating the specimen under low power, the objective lens is switched to higher magnifications (40× or 100× oil immersion lens) for detailed observation. Only the fine adjustment knob is used at higher power.

6. Oil immersion technique (for 100× lens): A drop of immersion oil is placed on the slide, and the 100× objective is carefully focused. Oil increases resolution by reducing light refraction.

7. Observation and recording: The specimen is observed carefully, noting structure, shape, and arrangement. Observations are recorded for analysis.

12. Differentiate between enriched and differential media. Write components and primary objectives.

Difference between Enriched and Differential Media

Feature	Enriched Media	Differential Media
Purpose	Promote growth of fastidious organisms	Differentiate microorganisms
Function	Provides extra nutrients	Shows visible biochemical differences
Indicators	Usually absent	Present (pH dyes, indicators)
Selectivity	Non-selective but nutrient-rich	Not necessarily selective but differentiating
Example	Blood agar, chocolate agar	MacConkey agar, EMB agar

Components

Enriched media components:

- Basic nutrients (peptone, agar, salts)
- Enrichment substances like: Blood (sheep blood agar), Serum, Egg yolk, Yeast extract

These components provide additional growth factors like hemin, NAD, amino acids, and vitamins.

Differential media components:

- Basic nutrient medium (peptone, agar, salts)
- Specific substrates (lactose, sucrose, etc.)
- Indicators (neutral red, eosin, methylene blue, phenol red)

Primary Objectives

Enriched media objectives:

- To support growth of fastidious microorganisms
- To recover bacteria from clinical samples
- To enhance growth of nutritionally demanding organisms
- Example: isolation of *Streptococcus* using blood agar

Differential media objectives:

- To distinguish between different bacterial species
- To identify metabolic properties such as lactose fermentation
- To aid in preliminary identification of pathogens
- Example: *E. coli* forms pink colonies on MacConkey agar, while non-lactose fermenters remain colorless

Compiled by: SIR MUGHAL

Get in touch with us:
Sir Mughal: https://wa.me/923132310940
Sir Jawad: https://wa.me/923036100859

13. What is growth curve that is obtained in bacterial population?

Bacterial Growth Curve (in a population):

The bacterial growth curve is a graphical representation that shows the change in the number of bacterial cells over time when bacteria are grown in a closed (batch) culture system. When bacterial population is plotted against time, a characteristic sigmoid (S-shaped) curve is obtained. The growth curve consists of four distinct phases:

1. Lag phase: This is the initial phase where bacteria are adapted to the new environment. There is little or no increase in cell number. Metabolically, cells are active and synthesizing enzymes required for growth.

2. Log (Exponential) phase: In this phase, bacteria divide at a constant and maximum rate. Cell number increases exponentially, and the population doubles at regular intervals (generation time). This is the phase of highest metabolic activity.

3. Stationary phase: In this phase, growth rate slows down and the number of new cells equals the number of dying cells. This occurs due to nutrient depletion, oxygen limitation, and waste accumulation. The total population remains constant.

4. Death (Decline) phase: In this final phase, the number of dying cells exceeds the number of new cells. This leads to a decrease in viable cell count due to unfavorable conditions.

14. Write a detailed note on Biosafety Level 4.

Biosafety Level 4 (BSL-4):

Biosafety Level 4 (BSL-4) is the highest level of biological containment used in microbiology laboratories. It is designed for the handling of extremely dangerous and exotic infectious agents that pose a high risk of life-threatening disease, are easily transmitted, and for which there are usually no available vaccines or effective treatments.

BSL-4 laboratories are used for pathogens that can cause severe or fatal diseases in humans and may also spread through aerosols. Examples include Ebola virus, Marburg virus, Lassa virus, and other highly virulent hemorrhagic fever viruses. Because of the extreme risk, strict safety measures are implemented to prevent any exposure or accidental release.

In BSL-4 facilities, laboratory access is highly restricted and controlled. Only specially trained and authorized personnel are allowed to enter. Workers must undergo extensive training in biosafety and emergency procedures before handling any pathogens.

A defining feature of BSL-4 is the use of full-body, air-supplied positive pressure suits, also known as “space suits.” These suits provide complete isolation from the laboratory environment and are connected to a dedicated air supply system to prevent inhalation of infectious aerosols.

The laboratory itself is designed as a completely isolated unit. It operates under negative air pressure, ensuring that air flows into the lab but cannot escape. All air leaving the facility passes through HEPA (High-Efficiency Particulate Air) filters to remove any infectious particles.

Entry and exit procedures are highly controlled and include multiple security barriers such as airlocks, chemical showers, and decontamination systems. All materials leaving the laboratory must be sterilized, usually by autoclaving or chemical disinfection.

Waste management is strictly regulated. All liquid and solid waste is decontaminated before disposal to ensure no viable pathogens are released into the environment.

BSL-4 laboratories are rare and are typically located in specialized government or high-security research centers. Their main purpose is to study emerging infectious diseases, develop vaccines, and understand high-risk pathogens under maximum containment conditions.

Compiled by SIR MUGHAL

Get in touch with us:
Sir Mughal: https://wa.me/923132310940
Sir Jawad: https://wa.me/923036100859

15. Write a detailed note on Biosafety Level 3.

Biosafety Level 3 (BSL-3)

Biosafety Level 3 (BSL-3) is an intermediate-high containment level used in microbiology laboratories for the handling of infectious agents that can cause serious or potentially life-threatening diseases through inhalation exposure. These pathogens are usually indigenous or exotic and may have the potential for aerosol transmission, making strict laboratory control essential.

BSL-3 laboratories are designed for work with microorganisms such as Mycobacterium tuberculosis, Bacillus anthracis, SARS-associated coronaviruses, and other airborne infectious agents. These organisms are highly pathogenic and require enhanced safety precautions compared to BSL-1 and BSL-2 levels.

A key feature of BSL-3 is strict access control. Entry into the laboratory is restricted to trained and authorized personnel only. Laboratory doors are kept closed during work, and access is carefully monitored to prevent unauthorized entry.

All procedures involving infectious materials are performed in a Class II or Class III biosafety cabinet to prevent the release of aerosols and protect laboratory workers from inhalation exposure. Personnel are required to wear appropriate protective clothing, including laboratory coats, gloves, and sometimes respiratory protection depending on the procedure.

The facility design of a BSL-3 laboratory includes negative air pressure systems, which ensure that air flows into the laboratory but does not escape to surrounding areas. This prevents contaminated air from leaving the containment zone. The air leaving the laboratory is passed through HEPA filters to remove infectious particles before release.

BSL-3 laboratories also include specialized engineering controls such as sealed windows, self-closing doors, and directional airflow systems. These features help maintain containment and reduce the risk of accidental exposure.

Decontamination procedures are strictly followed. All biological waste, culture materials, and contaminated equipment are sterilized using autoclaving or chemical disinfection before disposal. Laboratory surfaces are regularly disinfected to maintain a sterile working environment.

Personnel working in BSL-3 laboratories undergo specialized biosafety training and medical surveillance to ensure early detection of possible infections. Emergency protocols are also in place in case of accidental exposure or spills.

BSL-3 laboratories are commonly used in clinical diagnostics, research on tuberculosis, and studies of emerging infectious diseases. They play a critical role in public health protection, vaccine development, and epidemiological research involving high-risk pathogens.

16. What is Laminar Air Flow Cabinet?

Laminar Air Flow (LAF) Cabinet:

A Laminar Air Flow Cabinet is a specialized laboratory equipment used to provide a sterile working environment by supplying air that has been filtered through HEPA (High Efficiency Particulate Air) filters in a smooth, uniform, and unidirectional flow.

The main principle of LAF is that air moves in a laminar (non-turbulent) flow pattern, either horizontally or vertically, which continuously sweeps away dust particles, microorganisms, and contaminants from the work area. This ensures that the workspace remains free from contamination during microbiological and aseptic procedures.

The cabinet consists of a blower fan, pre-filter, HEPA filter, working bench, and UV light. The HEPA filter removes particles as small as 0.3 microns with very high efficiency, ensuring that only sterile air reaches the working surface. UV light is often used before work to sterilize the internal chamber by killing microorganisms.

There are two main types of laminar flow cabinets: horizontal laminar flow cabinets, where air flows directly towards the user, and vertical laminar flow cabinets, where air flows downward onto the work surface. Vertical flow cabinets are commonly preferred in microbiology because they provide better protection against contamination.

Compiled by: SIR MUGHAL

Get in touch with us:
Sir Mughal: https://wa.me/923132310940
Sir Jawad: https://wa.me/923036100859

Laminar air flow cabinets are widely used in microbiology laboratories for aseptic inoculation, culture handling, media preparation, and tissue culture work. They are essential for maintaining sterile conditions and preventing contamination of samples and cultures.

17. How autoclave does performs sterilization?

Working of Autoclave for Sterilization:

An autoclave is a laboratory device used for sterilization by moist heat under pressure. It works on the principle that increased pressure raises the boiling point of water, allowing steam to reach higher temperatures than 100°C, which effectively kills all microorganisms including bacterial spores.

The material to be sterilized (such as culture media, glassware, surgical instruments, or contaminated waste) is placed inside the autoclave chamber. The chamber is then sealed tightly to make it airtight.

Water inside the autoclave is heated to produce saturated steam. As steam builds up, pressure inside the chamber increases, typically to 15 psi (pounds per square inch). At this pressure, the temperature rises to 121°C.

The steam penetrates deeply into materials and microbial cells. The main mechanism of sterilization is protein denaturation and coagulation of enzymes and cellular structures, which leads to irreversible microbial death.

The standard sterilization cycle is usually carried out at 121°C for 15–20 minutes at 15 psi pressure, although time may vary depending on the load.

After the sterilization cycle is complete, the pressure is gradually released and the materials are allowed to cool before removal. Proper drying is also important to avoid contamination.

Autoclaves are widely used in microbiology, hospitals, and laboratories for sterilizing media, instruments, dressings, and biohazard waste, ensuring complete destruction of all forms of microbial life including spores.

Compiled by: SIR MUGHAL

Get in touch with us:
Sir Mughal: https://wa.me/923132310940
Sir Jawad: https://wa.me/923036100859

LMS Handling

by Sir Mughal

Reliable • Professional • Result-Oriented



Our Key Features

- Timely Activities
- Expert Team Members
- Maximum Quiz Scores
- Assignment Solutions by *Subject-Oriented Experts*
- Paid Project Handling
- Guaranteed Results
- Market Competitive Rates

Contact Us

0313-2310940

shahzeb.zoologist04@gmail.com