Quiz 2 Bio302 (Molecular Biology)

Solved By Sami Ur Rehman (BS Biotech 3rd semester)

Leader sequence of CRISPR array is A-T rich and _____ in lenght

370

190

200

500

Which enzyme is used to add a formyl group rapidly to the methionine amino group?

Arg-rRNA transformylase

Met-tRNA transformylase

Met-tRNA transformase

Met-rRNA transformylase

Enhancers are the _____ elements that can act at a distance to modelate transcription.

RNA

Nucleotide

Protein

DNA

Which molecular technique was used to find the direction of RNA editing by Kenneth Stuart and colleagues?

Fast PCR

PCR

RT-PCR

Hot Start PCR

How many enzymatic steps are required by all aminoacyl-tRNA synthetases to attach an amino acid to a tRNA?

2

1

4

3

The Amino acids polymerizing machinery is composed of at least _____ RNA molecules.

3

1

4

2

The transcription machinery and the translation machinery in _____ are located in the same compartment.

Prokaryotes

Fungi

Viruses

Eukaryotes

Paul C. Zamecnik and Mahlon B. Hoagland (1957) showed that:

Before their incorporation into DNA, amino acids are attached to a class of RNA molecules

Before their incorporation into proteins, amino acids are attached to a class of RNA molecules

Before their incorporation into proteins, nucleotides are attached to a class of RNA molecules

Before their incorporation into proteins, amino acids are attached to a class of DNA molecules

In eukaroytes, translation is initiated by binding of ribosomes to the _____

TATA box

3' cap

Poly A tail

5' cap

The three nucleotide long units of protein coding region of the mRNA is called _____

Anticodon

Tricodon

Codon

Pre-codon

The anti-codon loop contain ……

A three nucleotide anticodon

Basic tools of genetic regulation are the ability of some proteins to bind to specific ……

Regulatory DNA sequences (not confirmed)

……. Product of the following gene is only found in brain.

ADAR2

The diameter of exit tunnel of bacterial ribosome is about …….. At its narrow point.

10 Aº

During translation, proteins are synthesized …….

By ribosomes using the information on mRNA

New amino acids are added to the ….. of the growing polypeptide chain during protein synthesis.

Carboxyl terminus