Objective questions:

Sn RNA are significantly involved in rRNAand mRNA processing and _____gene regulation
3-D arm contains the base _____Dihydrouridine
Secondary structure involves _____of the polyribonucleotide chain coil formation

Lineweaver-Burke plot, a *Double Reciprocal Plot*, that is useful in the practical determination of ___ Km and Vmax

V_o is plotted against [S], it is not always possible to determine when ____has been Achieved Vmax

____ reflects the affinity of the enzyme for the substrate. Km

Enzyme is not rigid, changes shape with _____ substrate

If *K*.eq of a reaction is greater than 1.0, □***GO*** is ____Negative
The precursors for DNA and RNA synthesis are nucleoside triphosphates

the mitochondrial membrane has no transporter fo262r oxaloacetate
Dna disruotion can accur in lab by change in_____
pH, salt, conc heating up 80 C ,all given
Dna is polymer of ribonucleotides
2ndary structure involves _____formAtion of polyribonucleotides chain
Hairpin loop ring coil
The sn RNA are significantly involes rRNA and mRNA processing
Dna replication
Transcription
Translation
Gene regulatiom
Lqyeses are the enzynes which di not catalyze the reaction
Hydrolysis
Oxidation
Both
The ionuzation of water can be meared by
Electrical conductivity

The solution is said to be ____ph then there is equal con of positive and negative ion
Nutral
Lineweaver burk plot accurate determination of
Km
Vmax
Kw
vmax and km
In michals menton v nod us the___€ reaction velocity
Initial
Total
Final
Accurate
____reflects the affinity of enzyme for substrate
Vmax
Km
Kw
None
Enzyme not rigid change the shape with
Coenzyme
Cofactor
Substrate
Phosphothetic group
The binding of enzymes to substrate moulds active site
Enzyme
Substrate
Ligand
Both a and h
Fatty acid catabolsim causes the oxidation beta carbon to form
Carboxyl
Keton
glycrol
The acetate formed from dietry fats binds ti coenzyme enter
Citric acid
ETC

Glyoxylate
In eukaryotes the citric acid take
Mitochondria
Cytosol
A buffer system consist of a _____its proyon accepter
Weak acid
Salt
Base
Buffer
Pepsin pH 2
Upon a warming ammunium molybdate in nitric acid
Yellow
Red
Voilet
Purple
SGOT level increased
Heart and liver
Kidny and liver
Heart and kidny
Keq is less 1.0 dalda Gnod is
Nutral
Pos
Neg
Infinity
PEP has one form that
Dehydride
Keto
Aldehyde
Enol
Precursor for DNA and RNA aynthesis
Monophosphate
Diphosphate
Tri
Tetra
Electron tend to flow toward half cell_____value of E

Neg
Pos
Nutral
Mass no. Is number of
Atomic molecules
Electron
Proton
Nutron
Pyruvate kinase rnzyme requires ion
Mn2+
K+
Mg+2
All
Reduction of pyruvate to lactate is catalyzed by lactate_____
Dehydrogenase
Aldolase
Carboxy
Hydrogenase
Stationary phase no. 3rd
Mitochondial membrane has no transport for
PEP
malte
Pyruvat
oxaloacetate
Oxidation phase of pentose phosphate pathway____
Glucose
NAD+
Pentose phosphate and NADPH
Function of sucrose emzyme is___
Break sucrose
Synthesis sucrose
Synthesis fructose
All
Alpha ketoglatarate complex resemble
ADP

PDH complex
Phosphate structure
Coenzyme b12 is the____enzyme
Hormone
Aditional factor
Cofactor
Brain prefer to use _____fuoe
Glucose
Ketone boddies
Protien
Fatty acid
Chilomicrone transports dietry____cholestrol___to tissues

SGOT is also called _____AST

Differnce between NAD and NADP_____phosphate

Subjective

TCA cycle (10)

E2 function (5)

Buret test theory procedure results?(5)

What is snRNA (5)

Glycolysis(3)

What are ketoacidiosis how it accur(3)

Internal loopin RNA(3)

Name 2 molecules produced by Propeonibacterium>>(2)

How ribonuleotides held together(2)

How many molecules of NADH are used during glycolysis(2)

biotechnologists
biotechnologists
vu
medical
students
(bio_
technologist)
biotechnologists