MTH101 LECTURES WISE MCQS 1 TO 22

BY VU BSCS

LEC 1

1. What are natural numbers?

- A) Numbers that include both positive and negative numbers
- B) Numbers that include zero and all positive integers
- C) Numbers that include all real numbers
- D) Numbers we count with, starting from 1

Correct Answer: D) Numbers we count with, starting from 1

2. Which of the following is a rational number?

- A) π
- B) √2
- C) 3/4
- D) e

Correct Answer: C) 3/4

3. What is the name for numbers that cannot be written as a ratio of two integers?

- A) Rational numbers
- B) Irrational numbers
- C) Real numbers
- D) Imaginary numbers

Correct Answer: B) Irrational numbers

4. The solution to the equation $x + 2 = 0$ is:

- A) 2
- B) -2
- C) 0
- D) 1

Correct Answer: B) -2

5. Which of the following represents a subset?

- A) $\{1, 2\} \subseteq \{1, 2, 3\}$
- B) $\{1, 2, 3\} \subseteq \{1, 2\}$
- C) $\{2, 4, 6\} \subseteq \{1, 3, 5\}$
- D) $\{5, 7\} \subseteq \{8, 9\}$

  Correct Answer: A) $\{1, 2\} \subseteq \{1, 2, 3\}$

6. Which of the following intervals is infinite?

- A) $(2, 5)$
- B) $(-\infty, 3]$
- C) $[1, 2]$
- D) $[0, 1]$

  Correct Answer: B) $(-\infty, 3]$

7. What is the order property of real numbers?

- A) If a + b = 0, a and b are positive.
- B) If b - a is positive, then a < b.
- C) If a > b, then b > a.
- D) If a < b, then a = b.

  Correct Answer: B) If b - a is positive, then a < b.

8. What does the notation $A \cup B$ represent?

- A) The intersection of sets A and B
- B) The union of sets A and B
- C) The complement of set A
- D) The difference of sets A and B

  Correct Answer: B) The union of sets A and B

9. In set theory, the symbol ∅ represents:

- A) A non-empty set
- B) An empty set
- C) A universal set
- D) A singleton set

  Correct Answer: B) An empty set

10. Which of the following numbers is irrational?

- A) 2/3
- B) -7
- C) √2
- D) 5

  Correct Answer: C) √2

11. Which of the following is an example of an inequality?

- A) $2 + 2 = 4$
- B) $5 < 7$
- C) $x + y = 0$
- D) $4 - 3 = 1$

Correct Answer: B) $5 < 7$

12. What is the coordinate of the origin on the real number line?

- A) 1
- B) -1
- C) 0
- D) ∞

Correct Answer: C) 0

13. What does the set-builder notation {x | x is a real number and 2 < x < 5} represent?

- A) All real numbers greater than 5
- B) All real numbers between 2 and 5, excluding 2 and 5
- C) All real numbers less than 2
- D) All real numbers greater than or equal to 2 and less than or equal to 5

  Correct Answer: B) All real numbers between 2 and 5, excluding 2 and 5

14. The inequality $a \leq b$ means:

- A) a is greater than b
- B) a is less than or equal to b
- C) a is less than b
- D) a is equal to b

  Correct Answ(↓) a is less than or equal to b

15. The interval (0, 1] represents:

- A) All numbers between 0 and 1, excluding 1
- B) All numbers between 0 and 1, including 1
- C) All numbers between 0 and 1, excluding both 0 and 1
- D) All numbers between 0 and 1, including 0

  Correct Answer: B) All numbers between 0 and 1, including 1

16. The solution set of the inequality $x < 5$ is:

- A) (-∞, 5)
- B) [5, ∞)
- C) (5, ∞)
- D) (-∞, 5]

  Correct Answer: A) (-∞, 5)

17. Which statement is true for $x^2 > 0$?

- A) x is positive
- B) x is negative
- C) x is non-zero
- D) x equals 0

  Correct Answer: C) x is non-zero

18. In which case will the inequality $a/c < b/c$ hold?

- A) When c is positive
- B) When c is negative
- C) When c equals zero
- D) Always true

  Correct Answer: A) When c is positive

19. What does the inequality $x \geq 2$ mean?

- A) x is greater than 2
- B) x is less than 2
- C) x is greater than or equal to 2
- D) x is less than or equal to 2

Correct Answer: C) x is greater than or equal to 2

20. The intersection $A \cap B$ of two sets A and B means:

- A) Elements that are in either A or B
- B) Elements that are only in A
- C) Elements that are in both A and B
- D) Elements that are in A but not in B

Correct Answer: C) Elements that are in both A and B

1. What is the absolute value of -7?

a) -7

b) 0

c) 7

d) -1

Answer: c) 7

2. What is the absolute value of 0?

a) -1

b) 0

c) 1

d) Undefined

Answer: b) 0

3. Solve the equation $|x| = 5$. What are the possible values of x?

a) 5

b) -5

c) 5 and -5

d) None

Answer: c) 5 and -5

4. Which of the following is true about the absolute value?

a) $|a|$ is always negative

b) $|a|$ is always non-negative

c) $|a|$ can be either positive or negative

d) $|a|$ is always positive unless $a = 0$

Answer: b) $|a|$ is always non-negative

5. Solve $|x - 3| = 7$. What are the possible values of x?

a) 10 and -4

b) 10 and 4

c) 7 and -7

d) 0 and 7

Answer: a) 10 and -4

6. What is the result of $|2 - 5|$?

a) -3

b) 3

c) 7

d) -7

Answer: b) 3

7. What is the absolute value of the product of -3 and 4?

a) -12

b) 12

c) -7

d) 7

Answer: b) 12

8. The solution to the inequality $|x - 4| < 3$ is:

a) $1 < x < 7$

b) $-3 < x < 3$

c) $x > 4$

d) $x < 3$

Answer: a) $1 < x < 7$

9. Solve $|x + 2| = 5$. What are the values of x?

a) 3 and -7

b) 5 and 2

c) -5 and 7

d) -3 and 5

Answer: a) 3 and -7

10. Which of the following satisfies the equation $|x - 1| = 2$?

a) $x = 1$

b) $x = -1$

c) $x = 3$ or $x = -1$

d) $x = 0$

Answer: c) $x = 3$ or $x = -1$

11. What is the absolute value of the square root of 16?

a) 4

b) -4

c) 0

d) 8

Answer: a) 4

12. The expression |a - b| represents:

a) The sum of a and b

b) The difference between a and b

c) The distance between a and b on a coordinate line

d) The product of a and b

Answer: c) The distance between a and b on a coordinate line

13. Which of the following is true about the triangle inequality theorem?

a) $|a + b| \leq |a| + |b|$

b) $|a + b| \geq |a| + |b|$

c) $|a + b| = |a| + |b|$

d) $|a + b| < |a| + |b|$

Answer: a) $|a + b| \leq |a| + |b|$

14. Solve $|x - 2| \geq 5$. What are the possible values of x?

a) $x \leq -3$ or $x \geq 7$

b) $x = -3$ or $x = 7$

c) $x > 2$

d) $x < 2$

Answer: a) $x \leq -3$ or $x \geq 7$

15. What is the result of |4/3|?

a) 4

b) -4/3

c) 4/3

d) 1

Answer: c) 4/3

16. What is the absolute value of -|x| if x = -6?

a) 6

b) -6

c) 12

d) 0

Answer: b) -6

17. What is the geometric interpretation of |x| on a coordinate line?

a) The distance between x and -x

b) The distance between x and 0

c) The distance between x and y

d) The distance between x and 1

Answer: b) The distance between x and 0

18. Solve |2x - 3| = 7. What are the possible values of x?

a) 5 and -2

b) 5 and 2

c) 4 and -4

d) 2 and -2

Answer: a) 5 and -2

19. Which of the following expressions is always true?

a) $|a - b| = -(a - b)$

b) $|a| = a$ if $a < 0$

c) $|a| \geq 0$ for all real numbers a

d) $|a + b| > |a| + |b|$

Answer: c) $|a| \geq 0$ for all real numbers a

20. The distance between points A and B with coordinates a = -2 and b = 5 is:

a) 3

b) -7

c) 7

d) 5

Answer: c) 7

1. **What is a coordinate plane?**

   A) The intersection of two coordinate lines at 45 degrees

   B) The intersection of two coordinate lines at 90 degrees

   C) A single line in space

   D) A cube in 3D space

   Answer: B) The intersection of two coordinate lines at 90 degrees

2. **What is an ordered pair?**

   A) Two numbers where order doesn't matter

   B) Two real numbers in an assigned order

   C) A combination of letters and numbers

   D) Any pair of coordinates on a line

   Answer: B) Two real numbers in an assigned order

3. **Which of the following is true about a point (x, y) in the plane?**

   A) It must always lie on the x-axis

   B) It is defined by its x- and y-coordinates

   C) It has no relationship to the axes

   D) It always lies on the origin

   Answer: B) It is defined by its x- and y-coordinates

4. **What are the labels commonly used for the coordinate axes?**

   A) m and n

   B) x and y

   C) a and b

   D) u and v

   Answer: B) x and y

5. **Which quadrant contains the point $(-3, 4)$?**

   A) Quadrant I

   B) Quadrant II

   C) Quadrant III

   D) Quadrant IV

   Answer: B) Quadrant II

6. **What is the x-intercept of a graph?**

   A) The point where the graph crosses the x-axis

   B) The point where the graph crosses the y-axis

   C) The point where the graph crosses the origin

   D) The highest point on the graph

   Answer: A) The point where the graph crosses the x-axis

7. **What is the y-intercept of a graph?**

   A) The point where the graph crosses the x-axis

   B) The point where the graph crosses the y-axis

   C) The point where the graph touches the origin

   D) The farthest point of the graph

   Answer: B) The point where the graph crosses the y-axis

8. **If the ordered pair $(3, 2)$ satisfies the equation $6x - 4y = 10$, what is the result when substituted?**

   A) 10

   B) 5

   C) 0

   D) 2

   Answer: A) 10

9. **Which equation is satisfied by the ordered pair $(2, 0)$?**

   A) $6x - 4y = 18$

   B) $6x - 4y = 10$

   C) $6x + 4y = 12$

   D) None of the above

   Answer: D) None of the above

10. **What is the graph of $y = x^2$?**

    A) A straight line

    B) A parabola opening upwards

    C) A hyperbola

    D) A parabola opening downwards

    Answer: B) A parabola opening upwards

11. **What is symmetry about the y-axis?**

    A) Points (x, y) and $(-x, y)$ are symmetric

    B) Points (x, y) and $(x, -y)$ are symmetric

    C) Points (x, y) and $(-x, -y)$ are symmetric

    D) Points $(-x, -y)$ and (y, x) are symmetric

    Answer: A) Points (x, y) and $(-x, y)$ are symmetric

12. **What is symmetry about the x-axis?**

    A) Points (x, y) and $(x, -y)$ are symmetric

    B) Points $(-x, y)$ and $(-x, -y)$ are symmetric

    C) Points (x, y) and $(-x, y)$ are symmetric

    D) Points (y, x) and (x, y) are symmetric

    Answer: A) Points (x, y) and $(x, -y)$ are

13. **What is symmetry about the origin?**

    A) Points (x, y) and $(x, -y)$ are symmetric

    B) Points (x, y) and $(-x, -y)$ are symmetric

    C) Points (x, y) and $(-x, y)$ are symmetric

    D) Points $(-x, y)$ and $(x, -y)$ are symmetric

    Answer: B) Points (x, y) and $(-x, -y)$ are symmetric

14. **What is the benefit of using symmetry in graphing?**

    A) It reduces the number of points to calculate

    B) It makes the graph more colorful

    C) It makes the graph change shape

    D) It allows one to ignore the axes

    Answer: A) It reduces the number of points to calculate

15. **What is the equation of a graph that is symmetric about the y-axis?**

    A) $y = x^2$

    B) $y = x^3$

    C) $y = x$

    D) $y = 2x + 1$

    Answer: A) $y = x^2$

16. **What happens when you substitute $-x$ into a function symmetric about the y-axis?**

    A) The equation changes completely

    B) The equation remains the same

    C) The equation becomes negative

    D) The equation vanishes

    Answer: B) The equation remains the same

17. **The graph of the equation $x = y^4$ is symmetric about which axis?**

    A) x-axis

    B) y-axis

    C) Origin

    D) None

    Answer: A) x-axis

18. **How do you check symmetry about the x-axis?**

    A) Replace x with $-x$

    B) Replace y with $-y$

    C) Replace both x and y with their negatives

    D) Add x and y together

    Answer: B) Replace y with $-y$

19. **What does it mean for a graph to be symmetric about the origin?**

A) Points are mirrored across both axes

B) Points are mirrored across the x-axis only

C) Points are mirrored across the y-axis only

D) Points remain unchanged

Answer: A) Points are mirrored across both axes

20. **If $y = x^2$, which of the following is true?**

A) The graph is symmetric about the x-axis

B) The graph is symmetric about the y-axis

C) The graph is symmetric about the origin

D) The graph is not symmetric

Answer: B) The graph is symmetric about the y-axis

1. What is the formula for the slope of a line passing through two points (x_1, y_1) and (x_2, y_2)?

a) $\frac{y_1 - y_2}{x_1 - x_2}$
b) $\frac{y_2 - y_1}{x_2 - x_1}$
c) $\frac{x_2 - x_1}{y_2 - y_1}$
d) $y_2 - y_1$

Answer: b

---

2. What is the slope of a vertical line?

a) 0
b) Undefined
c) 1
d) ∞

Answer: b

3. If two lines are parallel, their slopes are:

a) Negative reciprocals

b) Equal

c) Opposite

d) Undefined

Answer: b

---

4. If the slope of a line is $m = 2$, what does it mean?

a) The line rises 2 units for every 1 unit of run

b) The line falls 2 units for every 1 unit of run

c) The line is vertical

d) The line is horizontal

Answer: a

5. What is the slope of the line passing through the points $(6, 2)$ and $(9, 8)$?

a) 2

b) 3

c) 6

d) $\frac{2}{3}$

Answer: b

6. For a non-vertical line, the slope m is related to the angle of inclination θ by:

a) $m = \sin\theta$

b) $m = \cos\theta$

c) $m = \tan\theta$

d) $m = \cot\theta$

Answer: c

---

7. A line with an undefined slope is:

a) Horizontal

b) Vertical

c) Diagonal

d) Parallel to the x-axis

Answer: b

8. If the slope of one line is 3, what is the slope of a line perpendicular to it?

a) -3
b) $\frac{1}{3}$
c) $-\frac{1}{3}$
d) 3

Answer: c

---

9. If the slope of a line is negative, the line:

a) Slopes upward to the right
b) Slopes downward to the right
c) Is horizontal
d) Is vertical

Answer: b

10. What is the equation of a vertical line passing through the point $(a, 0)$?

a) $x = a$

b) $y = a$

c) $y = 0$

d) $x = 0$

Answer: a

---

11. What is the slope of the line parallel to the y-axis?

a) 0

b) 1

c) Undefined

d) -1

Answer: c

12. What is the slope of the line passing through points $(2, 9)$ and $(4, 3)$?

a) 3

b) -3

c) $\frac{3}{2}$

d) $-\frac{3}{2}$

Answer: b

---

13. What is the angle of inclination for a line with slope $m = 1$?

a) $30°$

b) $45°$

c) $60°$

d) $90°$

Answer: b

14. The slope of a horizontal line is:

a) Undefined

b) 0

c) 1

d) ∞

Answer: b

---

15. If two lines are perpendicular, the product of their slopes is:

a) 0

b) 1

c) -1

d) Undefined

Answer: c

16. What is the equation of a line that passes through $(2, 3)$ with a slope of $-\frac{3}{2}$?

a) $y - 3 = -\frac{3}{2}(x - 2)$

b) $y = -\frac{3}{2}x + 6$

c) $y = 3x + 2$

d) Both a and b

Answer: d

---

17. The equation of a line in point-slope form is:

a) $y = mx + b$

b) $y - y_1 = m(x - x_1)$

c) $Ax + By + C = 0$

d) $y = m(x - x_1)$

Answer: b

18. The general form of a line equation is:

a) $y = mx + b$
b) $Ax + By + C = 0$
c) $y = mx + C$
d) $x = y + b$

Answer: b

---

19. The slope of the line $8x + 5y = 20$ is:

a) $\frac{8}{5}$
b) $-\frac{8}{5}$
c) $\frac{5}{8}$
d) $-\frac{5}{8}$

Answer: b

20. What is the y-intercept of the line $8x + 5y = 20$?

a) 5

b) $-\frac{8}{5}$

c) 4

d) -4

Answer: c

---

21. The slope of the line that passes through $(-2, 7)$ and $(5, 7)$ is:

a) 0

b) 1

c) Undefined

d) ∞

Answer: a

22. In the equation $y = mx + b$, the variable b represents:

a) Slope

b) x-intercept

c) y-intercept

d) Angle of inclination

Answer: c

---

23. A line that crosses the y-axis at $b = -4$ with slope $m = -9$ is written as:

a) $y = -9x - 4$

b) $y = -9x + 4$

c) $y = 9x - 4$

d) $y = 9x + 4$

Answer: a

24. The equation of a line parallel to the x-axis is:

a) $x = 0$

b) $x = a$

c) $y = b$

d) $y = 0$

Answer: c

---

25. If a line passes through the points $(3, 4)$ and $(-2, -1)$, its slope is:

a) 1

b) -1

c) 5

d) 0

Answer: a

26. The equation of a line passing through the origin with slope 2 is:

a) $y = 2x + 1$

b) $y = 2x$

c) $y = x + 2$

d) $y = -2x$

Answer: b

---

27. If $m = \tan(\theta)$, and $\theta = 45°$, what is the slope m?

a) 0

b) 1

c) -1

d) Undefined

Answer: b

28. What is the angle of inclination for a line with slope $m = -1$?

a) 90°

b) 45°

c) 135°

d) 180°

Answer: c

---

29. The slope-intercept form of a line equation is:

a) $y = mx + b$

b) $y - y_1 = m(x - x_1)$

c) $Ax + By + C = 0$

d) $y = m + x + b$

Answer: a

30. What is the product of the slopes of two perpendicular lines?

a) 1

b) 0

c) -1

d) Undefined

Answer: c

LEC 5

Distance Between Two Points

1. **What is the formula for the distance between two points (x_1, y_1) and (x_2, y_2) in a coordinate plane?**

   a) $\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$

   b) $|x_2 - x_1| + |y_2 - y_1|$

   c) $(x_2 - x_1)^2 + (y_2 - y_1)^2$

   d) None of the above

   Answer: a

2. **What is the distance between the points $(-2, 3)$ and $(1, 7)$?**

   a) 3

   b) 5

   c) 7

   d) 10

   Answer: b

3. **If the distance between two points (x_1, y_1) and (x_2, y_2) is denoted by $d(P_1, P_2)$, then**

   a) $d(P_1, P_2) = d(P_2, P_1)$

   b) $d(P_1, P_2) \neq d(P_2, P_1)$

   c) $d(P_1, P_2) = 0$

   d) None of the above

   Answer: a

4. **The points A(4, 6), B(1, -3), and C(7, 5) form a right triangle because**

   a) $AB^2 + AC^2 = BC^2$

   b) $AB^2 + BC^2 = AC^2$

   c) $AB = AC = BC$

   d) None of the above

   Answer: a

The Midpoint Formula

5. **What is the formula for the midpoint of a line segment joining two points (x_1, y_1) and (x_2, y_2)?**

   a) $(x_1 + x_2, y_1 + y_2)$

   b) $\left(\frac{x_1+x_2}{2}, \frac{y_1+y_2}{2}\right)$

   c) $\left(\frac{x_2-x_1}{2}, \frac{y_2-y_1}{2}\right)$

   d) $(x_1 - x_2, y_1 - y_2)$

   Answer: b

6. **What is the midpoint of the line segment joining the points $(3, -4)$ and $(7, 2)$?**

   a) (5, -1)

   b) (10, -2)

   c) (4, 1)

   d) (3, 0)

   Answer: a

7. **What is the equation of a circle centered at $(-5, 3)$ with radius 4?**

   a) $(x+5)^2 + (y-3)^2 = 16$

   b) $(x-5)^2 + (y+3)^2 = 16$

   c) $(x+5)^2 + (y-3)^2 = 4$

   d) $(x+5)^2 + (y-3)^2 = 9$

   Answer: a

8. **What is the standard form equation of a circle with center $(1, -2)$ and passing through the point $(4, 2)$?**

   a) $(x-1)^2 + (y+2)^2 = 25$

   b) $(x-1)^2 + (y+2)^2 = 16$

   c) $(x+1)^2 + (y-2)^2 = 25$

   d) None of the above

   Answer: a

9. **Which of the following represents the equation of a unit circle?**

   a) $(x+1)^2 + (y-1)^2 = 1$

   b) $(x-1)^2 + (y+1)^2 = 0$

   c) $x^2 + y^2 = 1$

   d) $x^2 + y^2 = 0$

   Answer: c

10. **Which of the following values of k results in no graph for the equation $(x - x_0)^2 + (y - y_0)^2 = k$?**

    a) $k > 0$

    b) $k = 0$

    c) $k < 0$

    d) None of the above

    Answer: c

Quadratic Equations (y = ax² + bx + c)

11. **The graph of $y = ax^2 + bx + c$ is a**

    a) Line

    b) Circle

    c) Parabola

    d) Ellipse

    Answer: c

12. **The x-coordinate of the vertex of a parabola represented by $y = ax^2 + bx + c$ is given by**

    a) $x = \frac{-b}{2a}$

    b) $x = \frac{-a}{2b}$

    c) $x = \frac{c}{a}$

    d) $x = \frac{-b}{a}$

    Answer: a

13. **For $y = x^2 - 2x - 2$, the x-coordinate of the vertex is**

    a) 0

    b) 1

    c) -1

    d) 2

    Answer: b

14. **If $y = -x^2 + 4x - 5$, then the parabola opens**

    a) Upwards

    b) Downwards

    c) Left

    d) Right

    Answer: b

15. **The y-intercept of $y = ax^2 + bx + c$ is found by**

    a) Setting $y = 0$

    b) Setting $x = 0$

    c) Setting $b = 0$

    d) Solving the quadratic formula

    Answer: b

16. **The solution to the quadratic equation $x^2 - 2x - 2 = 0$ is**

    a) $x = 1$

    b) $x = 2$

    c) $x = 1 \pm \sqrt{3}$

    d) $x = -1 \pm \sqrt{2}$

    Answer: c

Inequalities

17. **The solution set of the inequality $x^2 - 2x - 2 > 0$ is**

    a) $(-\infty, 1)$

    b) $(-\infty, 1-\sqrt{3}) \cup (1+\sqrt{3}, \infty)$

    c) $(1-\sqrt{3}, 1+\sqrt{3})$

    d) None of the above

    Answer: b

18. **Which of the following is the quadratic formula?**

    a) $\frac{-b \pm \sqrt{b^2-4ac}}{2a}$

    b) $\frac{-a \pm \sqrt{b^2-4ac}}{2b}$

    c) $\frac{-b}{2a}$

    d) None of the above

    Answer: a

19. **If a ball is thrown straight up with initial velocity $24.5\,\mathrm{m/s}$, the maximum height reached by the ball is**

    a) 20 meters

    b) 25 meters

    c) 30.625 meters

    d) 40 meters

    Answer: c

20. **In the equation $s(t) = -4.9t^2 + 24.5t$, the time at which the ball reaches its maximum height is**

    a) 2.5 seconds

    b) 5 seconds

    c) 1 second

    d) 3.5 seconds

    Answer: a

1. **What is a function?**
   - A) A relationship where one variable depends on another
   - B) A static value
   - C) A geometric shape
   - D) A type of equation

     Answer: A) A relationship where one variable depends on another
2. **In the function $f(x) = 2x + 3$, which variable is independent?**
   - A) f
   - B) x
   - C) 2
   - D) 3

     Answer: B) x

3. **Which of the following is a valid function?**

- A) $y^2 = x + 4$
- B) $y = 3x + 2$
- C) $y = \pm 4$
- D) $x = y^2 - 5$

Answer: B) $y = 3x + 2$

4. **What does the notation $f(x)$ represent?**

- A) The product of f and x
- B) A function of x
- C) A constant
- D) The sum of f and x

Answer: B) A function of x

5. **Which of the following statements is true about the domain of a function?**
   - A) It includes all real numbers.
   - B) It can include imaginary numbers.
   - C) It is the set of all allowable input values.
   - D) It is always a finite set.

   Answer: C) It is the set of all allowable input values.

6. **For the function $g(x) = \frac{1}{x-3}$, what value must be excluded from the domain?**
   - A) 0
   - B) 1
   - C) 3
   - D) -3

   Answer: C) 3

7. **What is the range of the function $f(x) = x^2$?**
   - A) All real numbers
   - B) All non-negative real numbers
   - C) All integers
   - D) All positive real numbers

   Answer: B) All non-negative real numbers

8. **If $h(x) = \sqrt{x}$, what is the domain of $h(x)$?**
   - A) $(-\infty, \infty)$
   - B) $[0, \infty)$
   - C) $(-\infty, 0)$
   - D) $(0, \infty)$

   Answer: B) $[0, \infty)$

9. **Which of the following is an example of a piecewise function?**

- A) $f(x) = x^2$
- B)

$$f(x) = \begin{cases} x+2 & \text{if } x < 0 \\ x^2 & \text{if } x \geq 0 \end{cases}$$

- C) $f(x) = \sin(x)$
- D) $f(x) = e^x$

Answer: B)

$$f(x) = \begin{cases} x+2 & \text{if } x < 0 \\ x^2 & \text{if } x \geq 0 \end{cases}$$

10. **What is the natural domain of $f(x) = \frac{1}{x^2-4}$?**

    - A) $(-\infty, -2) \cup (-2, 2) \cup (2, \infty)$
    - B) $(-\infty, 2)$
    - C) $(-2, 2)$
    - D) $(-\infty, \infty)$

    Answer: A) $(-\infty, -2) \cup (-2, 2) \cup (2, \infty)$

11. **If $y = f(x) = 3x - 5$, what is $f(2)$?**

    - A) 1
    - B) 6
    - C) 7
    - D) 9

    Answer: C) 1

12. **What is the range of the function $f(x) = -x^2 + 4$?**

- A) $(-\infty, 4]$
- B) $[4, \infty)$
- C) $(-\infty, \infty)$
- D) $[0, 4]$

Answer: A) $(-\infty, 4]$

13. **In the function $f(x) = \frac{x+1}{x-1}$, what value is excluded from the domain?**

- A) 0
- B) 1
- C) -1
- D) 2

Answer: B) 1

14. **For the function defined as $f(x) = 5$ for all x, what is the range?**
    - A) All real numbers
    - B) {5}
    - C) $(-\infty, \infty)$
    - D) $[5, 5]$

    Answer: B) {5}

15. **What does the range of a function indicate?**
    - A) The set of all possible input values
    - B) The set of all possible output values
    - C) The values of the independent variable
    - D) The values of the dependent variable

    Answer: B) The set of all possible output values

16. **If $f(t) = 2t^2 + 3t - 1$, which of the following values is part of its natural domain?**

    - A) 0
    - B) -2
    - C) 3
    - D) All of the above

    Answer: D) All of the above

17. **If $f(x) = 2$ for $x \leq 3$ and $f(x) = x + 1$ for $x > 3$, what is $f(4)$?**

    - A) 2
    - B) 3
    - C) 4
    - D) 5

    Answer: D) 5

18. **Which of the following is NOT a function?**

- A) $y = x^3 - 3x$
- B) $y = \sqrt{x}$
- C) $y^2 = x$
- D) $y = 2x + 1$

Answer: C) $y^2 = x$

19. **What is the value of $f(-1)$ if $f(x) = x^2 - 4$?**

- A) -3
- B) -5
- C) 3
- D) 5

Answer: C) 3

19. **What is the value of $f(-1)$ if $f(x) = x^2 - 4$?**

- A) -3
- B) -5
- C) 3
- D) 5

Answer: C) 3

20. **If $g(x) = \frac{2}{x^2-1}$, what is the value of $g(0)$?**

- A) 0
- B) 1
- C) 2
- D) Undefined

Answer: C) 2

Multiple Choice Questions (MCQs)

1. **What is the definition of the sum of two functions f and g?**
   - A) $f(x) + g(x)$
   - B) $f(g(x))$
   - C) $f(x) \cdot g(x)$
   - D) $f(x) - g(x)$
   - **Correct Answer:** A
2. **If $f(x) = 2x$ and $g(x) = x^2$, what is $(f + g)(x)$?**
   - A) $x^2 + 2x$
   - B) $2x^2$
   - C) $2x + 2$
   - D) $2x^2 + 2x$
   - **Correct Answer:** A

3. **The domain of $f + g$ is defined as:**
    - A) The union of the domains of f and g
    - B) The intersection of the domains of f and g
    - C) The complement of the domains of f and g
    - D) The sum of the ranges of f and g
    - **Correct Answer:** B
4. **What is the product of two functions f and g?**
    - A) $f(g(x))$
    - B) $f(x) + g(x)$
    - C) $f(x) \cdot g(x)$
    - D) $f(x) - g(x)$
    - **Correct Answer:** C

5. **If $f(x) = x + 1$ and $g(x) = 2x$, what is $(f \cdot g)(x)$?**
   - A) $x^2 + 2x + 1$
   - B) $2x^2 + 2x$
   - C) $2x + 1$
   - D) $x^2 + 1$
   - **Correct Answer:** B
6. **Which of the following represents the quotient of functions f and g?**
   - A) $\frac{f(x)}{g(x)}$
   - B) $f(x) + g(x)$
   - C) $f(g(x))$
   - D) $f(x) - g(x)$
   - **Correct Answer:** A

7. **If $f(x) = 2x + 3$ and $g(x) = x - 1$, what is $(f - g)(x)$?**
   - A) $x + 4$
   - B) $x + 2$
   - C) $2x + 2$
   - D) $2x + 1$
   - **Correct Answer:** A
8. **Which operation allows you to evaluate $f(g(x))$?**
   - A) Addition
   - B) Subtraction
   - C) Multiplication
   - D) Composition
   - **Correct Answer:** D

9. **What is the natural domain of $f(x) = \sqrt{x}$?**
   - A) $(-\infty, \infty)$
   - B) $(0, \infty)$
   - C) $[0, \infty)$
   - D) $(-\infty, 0)$
   - **Correct Answer:** C
10. **For $f(x) = 3$ and $g(x) = 4x$, what is $(f \cdot g)(x)$?**
   - A) $12x$
   - B) $3 + 4x$
   - C) $3 + 12$
   - D) $3x + 4$
   - **Correct Answer:** A

11. **If $h(x) = x^2 + 1$, what is the decomposition of $h(x)$ using $f(x) = x$ and $g(x) = x^2$?**
    - A) $h(x) = f(g(x))$
    - B) $h(x) = g(f(x))$
    - C) $h(x) = g(x) + f(x)$
    - D) $h(x) = f(x) + g(x)$
    - **Correct Answer:** A
12. **Which of the following is true for $(f \circ g)(x)$ and $(g \circ f)(x)$?**
    - A) They are always equal.
    - B) They are never equal.
    - C) They can be equal in some cases.
    - D) They represent the same operation.
    - **Correct Answer:** C

13. **For the functions $f(x) = x^2$ and $g(x) = 2x + 1$, what is $(f \circ g)(x)$?**
    - A) $2x^2 + 1$
    - B) $(2x + 1)^2$
    - C) $x^2 + 1$
    - D) $x + 2$
    - **Correct Answer:** B
14. **A function that assigns the same value to every input is called:**
    - A) Monomial
    - B) Polynomial
    - C) Constant Function
    - D) Variable Function
    - **Correct Answer:** C

15. **What type of function is $f(x) = 3x^2 - 5x + 2$?**
    - A) Constant Function
    - B) Linear Function
    - C) Quadratic Function
    - D) Monomial
    - **Correct Answer:** C
16. **Which of the following functions is NOT a polynomial?**
    - A) $x^3 + 2x^2 + x$
    - B) $5x^2$
    - C) $x^{-1} + 3$
    - D) $7x + 1$
    - **Correct Answer:** C

17. **What is the result of $(f+g)(2)$ if $f(x)=x^2$ and $g(x)=3x$?**
    - A) 7
    - B) 10
    - C) 12
    - D) 8
    - **Correct Answer:** A
18. **For the functions $f(x)=x+3$ and $g(x)=x^2$, what is the domain of $(f \circ g)(x)$?**
    - A) $(-\infty, \infty)$
    - B) $[0, \infty)$
    - C) $[3, \infty)$
    - D) $(-3, \infty)$
    - **Correct Answer:** A

19. **If $h(x) = f(g(x))$ where f and g are defined, then $h(x)$ is an example of:**

- A) Linear Function
- B) Decomposition
- C) Composition of Functions
- D) Arithmetic Operation
- **Correct Answer:** C

20. **Which of the following is true for the composition of functions?**

- A) $f \circ g = g \circ f$
- B) $f(g(x)) = g(f(x))$
- C) Domains of f and g do not affect composition.
- D) The composition has a defined domain based on the domains of f and g.
- **Correct Answer:** D

Graphs of Functions MCQs

1. **What is the graph of the function $f(x) = x + 2$?**
    - A) A horizontal line
    - B) A vertical line
    - C) A line with slope 1 and y-intercept 2
    - D) A parabola
    - **Answer: C**
2. **Which of the following is true about the absolute value function $f(x) = |x|$?**
    - A) It is a straight line with a slope of 1.
    - B) It is piecewise defined.
    - C) It has a hole at $x = 0$.
    - D) It is always decreasing.
    - **Answer: B**

3. **What happens to the graph of $f(x)$ when we apply the transformation $g(x) = f(x) + 3$?**
    - A) The graph shifts down 3 units.
    - B) The graph shifts up 3 units.
    - C) The graph shifts left 3 units.
    - D) The graph shifts right 3 units.
    - **Answer: B**
4. **If $f(x) = x^2$, what is the graph of $g(x) = f(-x)$?**
    - A) It reflects $f(x)$ about the x-axis.
    - B) It reflects $f(x)$ about the y-axis.
    - C) It shifts $f(x)$ to the left.
    - D) It shifts $f(x)$ to the right.
    - **Answer: B**

5. **The function $t(x) = \frac{x^{-}-4}{x-2}$ has a hole at which x-value?**
   - A) $x = 2$
   - B) $x = 0$
   - C) $x = 4$
   - D) $x = -2$
   - **Answer: A**
6. **The graph of the function $g(x) = \begin{cases} 1 & \text{if } x \leq 2 \\ x+2 & \text{if } x > 2 \end{cases}$ is characterized by:**
   - A) A jump at $x = 2$
   - B) A hole at $x = 2$
   - C) A straight line with a slope of 1 for all x
   - D) A constant value for all x
   - **Answer: A**

7. **If $f(x)$ is multiplied by a constant $c > 1$, what happens to the graph?**
    - A) It compresses vertically.
    - B) It stretches vertically.
    - C) It shifts left.
    - D) It shifts right.
    - **Answer: B**
8. **The equation $y^2 + x^2 = 25$ describes which of the following?**
    - A) A line
    - B) A parabola
    - C) A circle
    - D) An ellipse
    - **Answer: C**

9. **Which transformation will result in a graph being translated left by 3 units?**
   - A) $g(x) = f(x + 3)$
   - B) $g(x) = f(x - 3)$
   - C) $g(x) = f(-x)$
   - D) $g(x) = f(x) + 3$
   - **Answer: A**
10. **What geometric transformation occurs when $f(x)$ is changed to $-f(x)$?**
   - A) Reflection about the x-axis
   - B) Reflection about the y-axis
   - C) Translation up
   - D) Translation down
   - **Answer: A**

11. **If a vertical line crosses the graph of a relation in two or more points, what can be concluded about that relation?**
    - A) It is a function.
    - B) It is not a function.
    - C) It is a linear function.
    - D) It is a quadratic function.
    - **Answer: B**
12. **For which function does the vertical line test fail?**
    - A) $y = x^2$
    - B) $y = |x|$
    - C) $x^2 + y^2 = 25$
    - D) $y = 2x + 1$
    - **Answer: C**

13. **If $f(x) = 3x - 1$, what is the slope of the graph?**
    - A) 3
    - B) -3
    - C) 1
    - D) 2
    - **Answer: A**
14. **How does the graph of $f(x) = \sin(x)$ change when transformed to $g(x) = 2\sin(x)$?**
    - A) It is compressed vertically.
    - B) It is stretched vertically.
    - C) It is reflected.
    - D) It shifts horizontally.
    - **Answer: B**

15. **Which of the following represents a function?**
    - A) $y^2 = x$
    - B) $x^2 + y^2 = 1$
    - C) $y = x^2 - 4$
    - D) $y = \sqrt{x}$ for $x < 0$
    - **Answer: C**
16. **If the function is defined as $g(x) = \begin{cases} 3 & x < 1 \\ x^2 & x \geq 1 \end{cases}$, what is $g(1)$?**
    - A) 3
    - B) 1
    - C) 0
    - D) 1
    - **Answer: B**

17. **The graph of $y = f(x - 4)$ is obtained by:**
    - A) Shifting $f(x)$ right by 4 units.
    - B) Shifting $f(x)$ left by 4 units.
    - C) Reflecting $f(x)$ about the y-axis.
    - D) Stretching $f(x)$.
    - **Answer: B**
18. **Which of the following statements about the graph of a function is true?**
    - A) It can intersect the x-axis at multiple points.
    - B) It can intersect the y-axis at multipl points.
    - C) It can have a vertical tangent line.
    - D) It cannot cross a vertical line more than once.
    - **Answer: D**

19. **If $g(x) = f(x) - 5$, how does the graph of $g(x)$ compare to the graph of $f(x)$?**
    - A) Shifted up 5 units.
    - B) Shifted down 5 units.
    - C) Reflected about the x-axis.
    - D) Stretched vertically.
    - **Answer: B**
20. **What does the notation $f(x)$ typically represent?**
    - A) The x-value of a function
    - B) The y-value of a function
    - C) The entire function
    - D) The slope of a function
    - **Answer: C**

21. **If $f(x)$ has a maximum point, what can be said about the slope of $f(x)$ at that point?**
    - A) It is positive.
    - B) It is negative.
    - C) It is zero.
    - D) It is undefined.
    - **Answer: C**
22. **Which of the following represents a transformation that compresses the graph vertically?**
    - A) $g(x) = 3f(x)$
    - B) $g(x) = \frac{1}{2}f(x)$
    - C) $g(x) = -f(x)$
    - D) $g(x) = f(x) + 2$
    - **Answer: B**

23. **The function $f(x) = \sqrt{x}$ is defined for which domain?**
    - A) All real numbers
    - B) $x \geq 0$
    - C) $x < 0$
    - D) $x \in \mathbb{R}$
    - **Answer: B**
24. **If a function's graph is symmetric with respect to the origin, what can be inferred about the function?**
    - A) It is an even function.
    - B) It is an odd function.
    - C) It has a maximum point.
    - D) It has a minimum point.
    - **Answer: B**

1. What does the notation $\lim_{x \to 0^+} f(x)$ represent?
   - A) The value of $f(x)$ at $x = 0$
   - B) The limit of $f(x)$ as x approaches 0 from the left
   - C) The limit of $f(x)$ as x approaches 0 from the right
   - D) The derivative of $f(x)$
   - **Correct Answer:** C
2. If $\lim_{x \to 2} f(x) = 3$ and $\lim_{x \to 2} g(x) = 4$, what is $\lim_{x \to 2} (f(x) + g(x))$?
   - A) 7
   - B) 12
   - C) 1
   - D) 8
   - **Correct Answer:** A

3. What is the value of $\lim_{x \to 0} \frac{\sin x}{x}$?
    - A) 0
    - B) 1
    - C) Infinity
    - D) Undefined
    - **Correct Answer:** B
4. Which of the following statements is true about $\lim_{x \to 0} \frac{\sin x}{x}$?
    - A) The limit does not exist.
    - B) The limit approaches negative infinity.
    - C) The limit approaches positive infinity.
    - D) The limit equals 1.
    - **Correct Answer:** D

5. If $f(x)$ oscillates between -1 and 1 as $x \to 0$, what can be said about $\lim_{x \to 0} f(x)$?
    - A) It exists and equals 0.
    - B) It does not exist.
    - C) It equals 1.
    - D) It equals -1.
    - **Correct Answer:** B
6. If $\lim_{x \to a} f(x) = \infty$, which of the following describes the behavior of $f(x)$?
    - A) It is bounded.
    - B) It approaches a finite value.
    - C) It increases without bound.
    - D) It oscillates indefinitely.
    - **Correct Answer:** C

7. What is the correct notation for the limit of $f(x)$ as x approaches 2 from the left?
   - A) $\lim_{x \to 2} f(x)$
   - B) $\lim_{x \to 2^+} f(x)$
   - C) $\lim_{x \to 2^-} f(x)$
   - D) $\lim_{x \to 0} f(x)$
   - **Correct Answer:** C
8. Which of the following is a condition for a limit to exist at a point?
   - A) The function must be continuous at that point.
   - B) The left-hand and right-hand limits must be equal.
   - C) The function must be defined at that point.
   - D) All of the above.
   - **Correct A ↓ er:** B

Questions on Tangents

9. The slope of the tangent line to a curve at a given point is given by:
    - A) The value of the function at that point.
    - B) The derivative of the function at tha point.
    - C) The area under the curve.
    - D) The integral of the function.
    - **Correct Answer:** B
10. The tangent line to the curve $y = f(x)$ at the point $(a, f(a))$ can be expressed as:
    - A) $y = f(a) + f'(a)(x - a)$
    - B) $y = f(a)(x - a) + f'(a)$
    - C) $y = f'(a)(x - a) + a$
    - D) $y = f(a) - f'(a)(x - a)$
    - **Correct A er:** A

11. If $f(x) = x^2$, what is the equation of the tangent line at the point (1, 1)?
    - A) $y = 2x - 1$
    - B) $y = 1 + 2(x - 1)$
    - C) $y = 2x + 1$
    - D) $y = 1 + x^2$
    - **Correct Answer:** B

Questions on Area

12. The area under a curve can be approximated by:
    - A) Finding the midpoint of the curve.
    - B) Using trapezoids.
    - C) Subdividing the area into rectangles.
    - D) All of the above.
    - **Correct Answer:** D

13. If a region cannot be filled with rectangles, we use:
    - A) The average value theorem.
    - B) Numerical methods to approximate the area.
    - C) Only circles to fill the area.
    - D) Exact integration.
    - **Correct Answer:** B
14. The process of finding the area under a curve as the number of rectangles approaches infinity is called:
    - A) Integral approximation.
    - B) The Fundamental Theorem of Calculus.
    - C) Riemann sums.
    - D) The limit process.
    - **Correct Answer:** D

15. The area under the curve $f(x)$ from a to b is represented as:
    - A) $\int_a^b f(x)\,dx$
    - B) $\lim_{n\to\infty} \sum f(x_i)\Delta x$
    - C) $A = b - a$
    - D) Both A and B.
    - **Correct Answer:** D
16. If a function $f(x)$ is continuous on the interval $[a, b]$, then the area under the curve is:
    - A) Always finite.
    - B) Undefined.
    - C) Infinite.
    - D) Always negative.
    - **Correct Answer:** A

General Questions on Limits

17. If $f(x) = \frac{1}{x}$, what is $\lim_{x \to 0^+} f(x)$?
    - A) 0
    - B) 1
    - C) Positive Infinity
    - D) Negative Infinity
    - **Correct Answer:** C
18. The limit $\lim_{x \to -\infty} (3x^2 + 2x)$ is:
    - A) 0
    - B) -Infinity
    - C) Positive Infinity
    - D) 0
    - **Correct Answer:** B

19. Which of the following functions does not have a limit as $x \to 0$?
    - A) $\sin(x)$
    - B) $\frac{\sin x}{x}$
    - C) $\tan(x)$
    - D) $\sin\left(\frac{1}{x}\right)$
    - **Correct Answer:** D
20. If $\lim_{x \to 2} f(x) = 5$, what can we conclude about $f(2)$?
    - A) $f(2) = 5$
    - B) $f(2)$ could be anything.
    - C) $f(2)$ is undefined.
    - D) $f(2) = 0$
    - **Correct Answer:** B

21. The left-hand limit and right-hand limit at a point are equal; thus, the overall limit:
    - A) Does not exist.
    - B) Must exist.
    - C) Is negative.
    - D) Is positive.
    - **Correct Answer:** B
22. If $\lim_{x \to c} f(x) = L$ and $f(c)$ is not equal to L, we can say:
    - A) The limit does not exist.
    - B) The limit is equal to c.
    - C) The function is continuous at c.
    - D) The function is discontinuous at c.
    - **Correct Answer:** D

Conceptual Questions

23. Which of the following could cause a limit to not exist?
    - A) Oscillation.
    - B) Jump discontinuity.
    - C) Unbounded behavior.
    - D) All of the above.
    - **Correct Answer:** D
24. What does $\lim_{x \to \infty} \frac{1}{x}$ approach?
    - A) 1
    - B) 0
    - C) Infinity
    - D) Undefined
    - **Correct Answer:** B

1. **What does the symbol Lim x->a f(x) represent?**
    - A) The value of f(x) at x = a
    - B) The behavior of f(x) as x approaches a
    - C) The value of x as f(x) approaches a
    - D) The derivative of f(x) at x = a
      Correct Answer: B
2. **Which of the following limits is equal to a constant k?**
    - A) $\lim_{x \to a} k$
    - B) $\lim_{x \to a} x$
    - C) $\lim_{x \to a} (x + k)$
    - D) $\lim_{x \to a} (k + x)$ **Correct Answer: A**

3. **Which theorem helps compute limits of the sum of two functions?**
    - A) Theorem 2.5.1
    - B) Theorem of Derivatives
    - C) Mean Value Theorem
    - D) Fundamental Theorem of Calculu

    Correct Answer: A
4. **If $f(x) = x$ and $g(x) = 5$, what is $\lim_{x \to 3} [f(x) + g(x)]$?**
    - A) 3
    - B) 5
    - C) 8
    - D) 15

    Correct Answer: C

5. **What is the value of $\lim_{x \to 2} (3x + 1)$?**

   - A) 5
   - B) 6
   - C) 7
   - D) 8

   Correct Answer: D

6. **What is $\lim_{x \to 3} (2x - 4)$?**

   - A) 2
   - B) 4
   - C) 6
   - D) 7

   Correct Answer: A

7. **What is lim $x \to a$ [k * g(x)] if k is a constant?**
    - A) k + g(a)
    - B) k * g(a)
    - C) g(a)
    - D) k * a

    Correct Answer: B
8. **Which of the following is true about limits of polynomials?**
    - A) Limits of polynomials do not exist
    - B) Limits of polynomials only exist if the polynomial is of degree 2
    - C) Limits of polynomials are determined by the highest power of x
    - D) Limits of polynomials depend on the leading coefficient only

    Correct Answer: C

9. **What is the value of $\lim_{x \to 5} (2x^2 - 4x + 3)$?**
   - A) 23
   - B) 53
   - C) 70
   - D) 78

   Correct Answer: C

10. **For the function $f(x) = 1/x$, what is $\lim_{x \to 0^+} f(x)$?**
   - A) 0
   - B) ∞
   - C) $-\infty$
   - D) 1

   Correct Answer: B

11. **For the same function $f(x) = 1/x$, what is $\lim_{x \to 0^-} f(x)$?**

    - A) 0
    - B) ∞
    - C) -∞
    - D) 1

    Correct Answer: C

12. **What is $\lim_{x \to +\infty} 1/x$?**

    - A) 0
    - B) ∞
    - C) -∞
    - D) 1

    Correct Answer: A

13. **What is the result of lim $x \to -\infty$ (x^2)?**

- A) 0
- B) $-\infty$
- C) ∞
- D) -1

Correct Answer: C

14. **For a rational function, if the numerator approaches a nonzero constant and the denominator approaches zero, the limit is:**

- A) 0
- B) ∞
- C) $-\infty$
- D) Undefined

Correct Answer: B

15. **What is $\lim_{x \to 4^+} (x^2 - 16) / (x - 4)$?**
    - A) 0
    - B) ∞
    - C) -∞
    - D) Undefined

    Correct Answer: B

16. **What happens when both numerator and denominator of a rational function approach zero as x approaches a?**
    - A) Limit does not exist
    - B) Limit is zero
    - C) Limit is infinity
    - D) Limit may exist if the factors can be canceled

    Correct Answer: D

17. **What is the limit of a constant function as x approaches any number?**
    - A) The constant itself
    - B) 0
    - C) ∞
    - D) Undefined

    Correct Answer: A
18. **What is the limit of a polynomial function as x approaches +∞?**
    - A) Determined by the lowest degree term
    - B) Determined by the highest degree term
    - C) 0
    - D) ∞

    Correct Answer: B

19. **What is $\lim_{x \to 3} (2x^2 - 5x + 4)$?**

- A) 7
- B) 13
- C) 16
- D) 19

Correct Answer: C

20. **What is $\lim_{x \to 0} (1/x)$?**

- A) 0
- B) ∞
- C) -∞
- D) Undefined

Correct Answer: D

21. **For the function $f(x) = x^2$, what is $\lim_{x \to +\infty} f(x)$?**

- A) 0
- B) ∞
- C) -∞
- D) Undefined

Correct Answer: B

22. **For the function $f(x) = 1/x^2$, what is $\lim_{x \to 0} f(x)$?**

- A) 0
- B) ∞
- C) -∞
- D) Undefined

Correct Answer: B

23. **For the function $f(x) = -1/x$, what is $\lim_{x \to 0^+} f(x)$?**

- A) 0
- B) ∞
- C) $-\infty$
- D) Undefined

Correct Answer: C

24. **What is the value of $\lim_{x \to -\infty} (3x^2 - 4x + 7)$?**

- A) ∞
- B) $-\infty$
- C) 7
- D) Undefined

Correct Answer: A

25. **What happens when the denominator of a rational function approaches zero and the numerator does not?**
    - A) The limit is zero
    - B) The limit is ∞ or -∞
    - C) The limit does not exist
    - D) The function is undefined

    Correct Answer: B

26. **For a function $f(x) = 1/x$, what is $\lim_{x \to +\infty} f(x)$?**
    - A) 0
    - B) ∞
    - C) -∞
    - D) 1

    Correct Answer: A

27. **What is lim $x \to +\infty$ $(2x^3 - 5x^2 + 3)$?**

- A) 0
- B) ∞
- C) -∞
- D) Undefined

Correct Answer: B

28. **What is lim $x \to -\infty$ $(3x^2 - x + 7)$?**

- A) 0
- B) ∞
- C) -∞
- D) Undefined

Correct Answer: B

29. **What is lim $x \to +\infty$ (4x - 2)?**

- A) 2
- B) 4
- C) ∞
- D) Undefined

Correct Answer: C

30. **What is the limit of $(x^2 + 5) / x^2$ as x approaches +∞?**

- A) 0
- B) 1
- C) 5
- D) ∞

Correct Answer: B

1. What does the formal definition of a limit include?

- a) Left-hand and right-hand limits
- b) Derivatives
- c) Continuity
- d) Integrals
- **Correct Answer: a)** Left-hand and right-hand limits

2. What does $\lim_{x \to a} f(x) = L$ mean intuitively?

- a) f(x) equals L at x = a
- b) As x approaches a, f(x) approaches L
- c) f(x) is undefined at x = a
- d) As x approaches infinity, f(x) approaches L
- **Correct Answer: b)** As x approaches a, f(x) approaches L

3. In the formal definition of limits, what does the symbol ε represent?

- a) An error
- b) A very small positive number
- c) The value of f(x) at x = a
- d) The slope of the tangent line
- **Correct Answer: b)** A very small positive number

---

4. What is δ used to signify in the formal definition of limits?

- a) The maximum distance from f(x) to L
- b) The distance of x from a
- c) The value of f(x) at a
- d) The derivative of the function
- **Correct Answer: b)** The distance of x from

5. Which of the following is true for the limit $\lim_{x \to a} f(x) = L$?

- a) $f(x)$ must equal L at $x = a$
- b) $f(x)$ can be undefined at $x = a$
- c) The limit only applies if $f(x)$ is continuous
- d) $f(x)$ must be continuous at $x = a$
- **Correct Answer: b)** $f(x)$ can be undefined at $x = a$

6. In the limit definition, which condition needs to be satisfied for all x in the interval around a?

- a) $|f(x) - L| < \varepsilon$
- b) $|x - a| < \varepsilon$
- c) $|f(x) + L| < \varepsilon$
- d) $|x + a| < \varepsilon$
- **Correct Answer: a)** $|f(x) - L| < \varepsilon$

---

7. What does it mean when we say f(x) approaches L as x approaches a?

- a) f(x) gets arbitrarily close to L as x gets close to a
- b) f(x) equals L at x = a
- c) x must always be equal to a
- d) f(x) cannot approach L if $x \neq a$
- **Correct Answer: a)** f(x) gets arbitrarily

8. If $\lim_{x \to a} f(x)$ exists, what must hold true?

- a) The left-hand and right-hand limits must be equal
- b) f(x) must be continuous
- c) f(x) cannot be undefined at x = a
- d) The limit only holds for polynomial functions
- **Correct Answer: a)** The left-hand and right-hand limits must be equal

9. What happens if the denominator of a rational function approaches zero while the numerator does not?

- a) The limit does not exist
- b) The limit is zero
- c) The limit is infinity or negative infinity
- d) The limit is the numerator divided by the denominator
- **Correct Answer: c)** The limit is infinity or negative infinity

10. What is the result of $\lim (x \to \infty) (4x^2 + 3x) / (2x^2 - 5)$?

- a) 2
- b) ∞
- c) 0
- d) 4
- **Correct Answer: a)** 2

---

11. What does the statement "for any $\varepsilon > 0$" in the formal limit definition mean?

- a) The value of ε must be exactly zero
- b) ε is a small positive number, but it can be as small as desired
- c) The limit holds only for $\varepsilon = 1$
- d) ε must be la[illegible]r than δ
- **Correct Answer: b)** ε is a small positive

12. Which of the following limits does not exist?

- a) $\lim (x \to 2) (3x - 5)$
- b) $\lim (x \to 0) 1/x$
- c) $\lim (x \to 1) (x^2 + 2x)$
- d) $\lim (x \to \infty) (1/x)$
- **Correct Answer: b)** $\lim (x \to 0) 1/x$

13. In the epsilon-delta definition of a limit, which of the following is true?

- a) δ depends on ε
- b) ε depends on δ
- c) ε and δ are always equal
- d) ε is always larger than δ
- **Correct Answer: a)** δ depends on ε

14. What happens if both the numerator and denominator of a rational function approach zero as x approaches a?

- a) The limit is zero
- b) The limit may need further simplification
- c) The limit does not exist
- d) The limit is the derivative of the numerator divided by the derivative of the denominator
- **Correct Answer: b)** The limit may need further simplification

15. What is the correct limit for $\lim (x \to \infty)$ $(3/x)$?

- a) ∞
- b) 3
- c) 0
- d) 1
- **Correct Answer: c)** 0

---

16. What is the limit of a constant function $f(x) = k$ as x approaches any value a?

- a) k
- b) 0
- c) a
- d) Undefined
- **Correct Answer: a)** k

17. What is the limit $\lim (x \to 2) (x^2 - 4) / (x - 2)$?

- a) 4
- b) 2
- c) Undefined
- d) Does not exist
- **Correct Answer: a)** 4

---

18. For the function $f(x) = 1/x$, what is $\lim (x \to 0^+)$?

- a) ∞
- b) -∞
- c) 0
- d) Does not exist
- **Correct Answer: a)** ∞

19. In the epsilon-delta definition, if $\varepsilon = 0.01$, what is δ used for?

- a) To define the interval around $x = a$
- b) To define the distance from $f(x)$ to L
- c) To calculate the derivative
- d) To calculate the integral
- **Correct Answer: a)** To define the interval around $x = a$

---

20. What does $\lim (x \to 0^-) f(x) = \infty$ mean?

- a) $f(x)$ is undefined at $x = 0$
- b) As x approaches 0 from the left, $f(x)$ becomes infinitely large
- c) $f(x)$ equals 0 at $x = 0$
- d) The limit do ot exist
- **Correct Answer: b)** As x approaches 0 from

1. **Which of the following is a condition for a function f to be continuous at a point c?**
   - A) $f(c)$ is defined
   - B) $\lim_{x \to c} f(x)$ exists
   - C) $\lim_{x \to c} f(x) = f(c)$
   - D) All of the above

     Answer: D) All of the above
2. **If a function f is a polynomial, what can we say about its continuity?**
   - A) It is continuous only at certain points
   - B) It is continuous everywhere
   - C) It is discontinuous everywhere
   - D) It is continuous only at rational points

     Answer: B) It is continuous everywhere

3. **For the function $f(x) = x^2 - 1$, which of the following is true?**
   - A) It is not continuous at $c = 2$
   - B) It is continuous at all real numbers
   - C) It is discontinuous at all points
   - D) It is continuous only at integers

   Answer: B) It is continuous at all real numbers
4. **The function $f(x) = \frac{2x^2+3}{x-1}$ is continuous at c provided:**
   - A) $c = 1$
   - B) $c \neq 1$
   - C) $c = 0$
   - D) c is any integer

   Answer: B) $c \neq 1$

5. **According to the Intermediate Value Theorem, if $f(a)$ and $f(b)$ have opposite signs, what can be concluded?**

   - A) There is no solution in (a, b)
   - B) There is exactly one solution in (a, b)
   - C) There are two solutions in (a, b)
   - D) There can be more than one solution

   Answer: B) There is exactly one solution in (a, b)

6. **Theorem 2.7.5 states that if $\lim_{x \to c} g(x) = L$ and f is continuous at L, then:**
   - A) $\lim_{x \to c}(f(g(x))) = f(c)$
   - B) $\lim_{x \to c}(f(g(x))) = L$
   - C) $\lim_{x \to c}(f(g(x))) = f(L)$
   - D) $\lim_{x \to c}(f(g(x)))$ does not exist

   Answer: C) $\lim_{x \to c}(f(g(x))) = f(L)$

7. **If $f(x) = \begin{cases} x^2 & \text{if } x < 0 \\ 2x + 1 & \text{if } x \geq 0 \end{cases}$, what can we say about the continuity of f at $x = 0$?**
   - A) Continuous
   - B) Discontinuous
   - C) Continuous from the left only
   - D) Continuous from the right only

   Answer: A) Continuous

8. **If $g(x)$ is a continuous function at c, what is true about $f(g(c))$ if f is also continuous at $g(c)$?**
   - A) $f(g(c))$ is undefined
   - B) $f(g(c))$ is discontinuous
   - C) $f(g(c))$ is continuous
   - D) Cannot be determined

   Answer: C) $f(g(c))$ is continuous
9. **What is the condition for a function f to be continuous from the right at point c?**
   - A) $f(c)$ is defined
   - B) $\lim_{x \to c^-} f(x)$ exists
   - C) $\lim_{x \to c} f(x) = f(c)$
   - D) All of the above

   Answer: D) All of the above

10. **For the function $f(x) = \frac{x-4}{x-2}$, what is the condition for its continuity?**
    - A) Continuous at $x = 2$
    - B) Discontinuous at $x = 2$
    - C) Continuous for all x
    - D) Continuous only at even integers

    Answer: B) Discontinuous at $x = 2$
11. **A function is continuous on a closed interval $[a, b]$ if:**
    - A) It is continuous on (a, b)
    - B) It is continuous from the right at a
    - C) It is continuous from the left at b
    - D) All of the above

    Answer: D) All of the above

12. **In the proof of continuity for polynomials, which theorem is referenced?**
    - A) The Mean Value Theorem
    - B) The Intermediate Value Theorem
    - C) The limit theorem for polynomials
    - D) The continuity theorem for rational functions

      Answer: C) The limit theorem for polynomials
13. **What is a point of discontinuity?**
    - A) A point where the function is defined
    - B) A point where $f(x)$ approaches infinity
    - C) A point where $f(x)$ is not continuous
    - D) A point where the limit exists

      Answer: C) A point where $f(x)$ is not

14. **Which of the following functions is continuous at $x = 1$?**

- A) $f(x) = \frac{1}{x-1}$
- B) $g(x) = x^2 + 2x + 1$
- C) $h(x) = \sqrt{x}$
- D) $k(x) = |x|$

Answer: B) $g(x) = x^2 + 2x + 1$

15. **For which of the following values is the function $f(x) = \frac{x^2-4}{x-2}$ continuous?**

- A) $x = 2$
- B) $x = -2$
- C) $x = 0$
- D) $x = 3$

Answer: B) $x = -2$ and D) $x = 3$ (but not A)

16. **The limit of a function $f(x)$ as $x \to c$ exists if:**

    - A) The left-hand limit and right-hand limit are equal
    - B) The function is defined at c
    - C) The function is continuous at c
    - D) Both A and B

      Answer: A) The left-hand limit and right-hand limit are equal

17. **Which of the following statements is true about rational functions?**

    - A) They are discontinuous everywhere
    - B) They are continuous everywhere except where the denominator is zero
    - C) They are continuous only at integers
    - D) They are continuous only at rational numbers

      Answer: B) They are continuous everywhere except where the denominator is zero

18. **The function** $f(x) = \begin{cases} 1 & \text{if } x \leq 0 \\ x & \text{if } x > 0 \end{cases}$ **is:**

- A) Continuous at $x = 0$
- B) Discontinuous at $x = 0$
- C) Continuous everywhere
- D) Continuous only at negative integers

  Answer: A) Continuous at $x = 0$

19. **What type of function is guaranteed to be continuous?**

- A) Rational functions
- B) Trigonometric functions
- C) Exponential functions
- D) All of the above

  Answer: D) All of the above

1. **What is the limit of $\sin(x)$ as x approaches 0?**
   - a) 1
   - b) 0
   - c) -1
   - d) Does not exist

   Answer: b) 0
2. **What is the limit of $\cos(x)$ as x approaches 0?**
   - a) 1
   - b) 0
   - c) -1
   - d) Does not exist

   Answer: a) 1

3. **What is the value of $\sin(0)$?**

   - a) 1
   - b) 0
   - c) -1
   - d) Undefined

   Answer: b) 0

4. **What is the value of $\cos(0)$?**

   - a) 1
   - b) 0
   - c) -1
   - d) Undefined

   Answer: a) 1

5. **Which of the following conditions must be met for a function $f(x)$ to be continuous at $x = c$?**
   - a) $f(c)$ is defined
   - b) $\lim_{x \to c} f(x)$ exists
   - c) $\lim_{x \to c} f(x) = f(c)$
   - d) All of the above

   Answer: d) All of the above
6. **For $f(x)$ to be continuous at $x = c$, which of the following must hold?**
   - a) $f(c)$ is not defined
   - b) $\lim_{x \to c} f(x) = 0$
   - c) $f(c) = \lim_{x \to c} f(x)$
   - d) None of the above

   Answer: c) $f(c) = \lim_{x \to c} f(x)$

7. **What is the relationship between h and x in the definition of continuity when $h \to 0$?**
   - a) $x = h - c$
   - b) $x = h + c$
   - c) $h = x - c$
   - d) $h = x + c$

   Answer: b) $x = h + c$

8. **What is the limit of $\sin(h + c)$ as $h \to 0$ if $c = 0$?**
   - a) 0
   - b) 1
   - c) Undefined
   - d) Does not exist

   Answer: a) 0

9. **What is the limit of $\cos(h + c)$ as $h \to 0$ if $c = 0$?**
   - a) 1
   - b) 0
   - c) Undefined
   - d) Does not exist

   Answer: a) 1

10. **Which of the following functions is continuous at $x = 0$?**
    - a) $\sin(x)$
    - b) $\cos(x)$
    - c) Both a and b
    - d) None of the above

    Answer: c) Both a and b

11. **What is the definition of continuity based on the ϵ-δ definition?**
    - a) $\lim_{h \to 0} f(h + c) = f(c)$
    - b) $\lim_{h \to 0} f(h)$ does not exist
    - c) $\lim_{x \to c} f(x) = 0$
    - d) None of the above

    Answer: a) $\lim_{h \to 0} f(h + c) = f(c)$

12. **Which of the following is a condition of continuity for the function $\cos(x)$ at $x = 0$?**
    - a) $\cos(0)$ is not defined
    - b) $\lim_{x \to 0} \cos(x) = 0$
    - c) $\lim_{x \to 0} \cos(x) = 1$
    - d) $\lim_{x \to 0} \cos(x)$ does not exist

    Answer: c) $\lim_{x \to 0} \cos(x) = 1$

13. **According to the ϵ-δ definition, which part ensures that the function value at a point matches the limit?**
    - a) $\lim_{x \to c} f(x)$
    - b) $f(c)$
    - c) $\lim_{h \to 0} f(h + c) = f(c)$
    - d) None of the above

    Answer: c) $\lim_{h \to 0} f(h + c) = f(c)$

14. **What must the limit of $\sin(x)$ approach as $x \to 0$ for $\sin(x)$ to be continuous?**
    - a) 1
    - b) 0
    - c) -1
    - d) Undefined

    Answer: b) 0

14. **What must the limit of $\sin(x)$ approach as $x \to 0$ for $\sin(x)$ to be continuous?**
    - a) 1
    - b) 0
    - c) -1
    - d) Undefined

    Answer: b) 0

15. **What must the limit of $\cos(x)$ approach as $x \to 0$ for $\cos(x)$ to be continuous?**
    - a) 0
    - b) -1
    - c) 1
    - d) Undefined

    Answer: c) 1

16. **Continuity of $f(x)$ at a point c requires that $f(c)$ be:**
    - a) Defined
    - b) Undefined
    - c) Equal to 0
    - d) Undefined at $x = 0$

    Answer: a) Defined

17. **If $f(x)$ is continuous at $x = c$, which of the following must be true?**
    - a) $\lim_{x \to c} f(x)$ does not exist
    - b) $\lim_{x \to c} f(x) = f(c)$
    - c) $f(x)$ is undefined at c
    - d) $\lim_{x \to c} f(x) = 0$

    Answer: b) $\lim_{x \to c} f(x) = f(c)$

18. **In the definition of continuity, $\lim_{x \to c} f(x) = f(c)$ means:**
    - a) The function jumps at $x = c$
    - b) The function is undefined at $x = c$
    - c) The function behaves smoothly at $x = c$
    - d) None of the above

    Answer: c) The function behaves smoothly at $x = c$

19. **Which theorem proves that $\sin(x)$ and $\cos(x)$ are continuous functions?**
    - a) Theorem 2.8.1
    - b) Theorem 2.8.2
    - c) Fundamental Theorem of Calculus
    - d) Mean Value Theorem

    Answer: a) Theorem 2.8.1

19. **Which theorem proves that $\sin(x)$ and $\cos(x)$ are continuous functions?**
    - a) Theorem 2.8.1
    - b) Theorem 2.8.2
    - c) Fundamental Theorem of Calculus
    - d) Mean Value Theorem

    Answer: a) Theorem 2.8.1

20. **What happens to $\sin(x)$ and $\cos(x)$ at $x = 0$ according to Theorem 2.8.1?**
    - a) They become undefined
    - b) They are continuous
    - c) They have a discontinuity
    - d) None of the above

    Answer: b) They are continuous

1. What is the slope of a secant line between two points $P(x_0, y_0)$ and $Q(x_1, y_1)$ on the graph of a function $y = f(x)$?

a) $\frac{f(x_1)-f(x_0)}{x_1-x_0}$

b) $\frac{x_1-x_0}{f(x_1)-f(x_0)}$

c) $f(x_1) - f(x_0)$

d) $x_1 - x_0$

Answer: a

2. What happens to the secant line as $x_1 \to x_0$?

a) The slope decreases

b) The secant line approaches the tangent line

c) The secant line disappears

d) The secant line becomes vertical

Answer: b

3. What is the geometric interpretation of the derivative of a function at a point?

a) The slope of the secant line

b) The slope of the tangent line

c) The area under the curve

d) The intercept of the tangent line

Answer: b

4. How is average velocity defined?

a) Distance traveled divided by time elapsed

b) Instantaneous velocity at a given point

c) Slope of the tangent line

d) The area under the curve of velocity

Answer: a

5. What is the instantaneous velocity of an object?

a) The average speed over a given time interval

b) The velocity at a given instant

c) The slope of the secant line

d) The total distance traveled

Answer: b

6. In terms of rates of change, what is velocity?

a) The rate of change of time with respect to distance

b) The rate of change of distance with respect to time

c) The slope of a vertical line

d) The area under the velocity-time graph

Answer: b

7. If $y = f(x)$, what represents the instantaneous rate of change of y with respect to x at a point x_0?

a) The slope of the secant line

b) The slope of the tangent line at x_0

c) The difference $f(x_1) - f(x_0)$

d) The average rate of change over an interval

Answer: b

8. If $d = f(t)$, what is the average velocity over a time interval $[t_0, t_1]$?

a) $\frac{f(t_1) - f(t_0)}{t_1 - t_0}$

b) $\frac{f(t_1)}{t_1}$

c) $f(t_1) - f(t_0)$

d) $f'(t_0)$

Answer: a

9. What is the instantaneous rate of change of $y = 2x + 1$ at any point x ?

a) 1

b) 0

c) 2

d) 4

Answer: c

10. Find the average rate of change of $y = 2x + 1$ over the interval $[3, 5]$.

a) 3

b) 1

c) 2

d) 8

Answer: c

11. For the function $y = 2x + 1$, what is the instantaneous rate of change at $x = -4$?

a) 4

b) -4

c) 2

d) 0

Answer: c

12. Which of the following is not an example of a rate of change?

a) The rate of change of bacteria population over time

b) The rate of change of distance traveled over time

c) The rate of change of quantity produced over cost

d) The rate of change of a metal rod's length over temperature

Answer: c

13. What does the derivative $f'(x)$ represent geometrically?

a) The y-intercept of the graph of $f(x)$

b) The slope of the tangent line to $f(x)$ at x

c) The total area under the graph of $f(x)$

d) The average rate of change of $f(x)$

Answer: b

14. Find the instantaneous rate of change of $y = 2x + 1$ at $x = 0$.

a) 1

b) 2

c) 0

d) -1

Answer: b

15. The slope of the tangent line at a point x_0 is given by which limit?

a) $\lim_{x_1 \to x_0} \frac{f(x_0)-f(x_1)}{x_0-x_1}$

b) $\lim_{x_1 \to x_0} \frac{f(x_1)-f(x_0)}{x_1-x_0}$

c) $\lim_{x_1 \to x_0} \frac{x_1-x_0}{f(x_1)-f(x_0)}$

d) $\lim_{x_0 \to x_1} f(x_0)$

Answer: b

16. Which of the following represents the average rate of change of a function over an interval $[x_0, x_1]$?

a) $\lim_{x_1 \to x_0} f'(x_0)$

b) $\frac{f(x_1)-f(x_0)}{x_1-x_0}$

c) $\frac{x_1-x_0}{f(x_1)-f(x_0)}$

d) $f(x_1) - f(x_0)$

Answer: b

17. The slope of a secant line gives the:

a) Instantaneous rate of change

b) Average rate of change

c) Derivative

d) Y-intercept

Answer: b

18. Which of the following is true about the instantaneous rate of change?

a) It is always positive

b) It is always negative

c) It can be positive, negative, or zero

d) It is the same as the average rate of change

Answer: c

19. For the function $y = 2x + 1$, what is the slope of the tangent line at any point?

a) 1

b) 2

c) 3

d) 0

Answer: b

20. What is the correct formula for the average rate of change of a function $y = f(x)$ between points x_0 and x_1?

a) $\frac{f(x_1)+f(x_0)}{x_1+x_0}$

b) $\frac{f(x_1)-f(x_0)}{x_1-x_0}$

c) $f(x_1) - f(x_0)$

d) $f'(x_0)$

Answer: b

1. The slope of the tangent line to the graph of $y = f(x)$ is given by:

- a) $\frac{f(x+h)-f(x)}{h}$
- b) $\lim_{h \to 0} \frac{f(x+h)-f(x)}{h}$
- c) $\frac{f(x_1)-f(x_0)}{x_1-x_0}$
- d) $f'(x_0)$
- **Answer**: b) $\lim_{h \to 0} \frac{f(x+h)-f(x)}{h}$

---

2. The derivative of the function $f(x) = x^2 + 1$ is:

- a) $2x + 1$
- b) x^2
- c) $2x$
- d) $2x^2$
- **Answer**: c) $2x$

3. The equation of the tangent line to $f(x) = x^2$ at the point $P(3, 9)$ is:

- a) $y = 6x - 9$
- b) $y = 6x - 27$
- c) $y = 9x - 27$
- d) $y = 2x + 1$
- **Answer**: b) $y = 6x - 27$

---

4. The process of finding a derivative is called:

- a) Integration
- b) Differentiation
- c) Substitution
- d) Approximation
- **Answer**: b) Differentiation

5. The derivative of $f(x) = x$ is:

- a) 1
- b) x
- c) 0
- d) x^2
- **Answer**: a) 1

---

6. If a function has a vertical tangent at $x = 0$, the derivative at $x = 0$ is:

- a) Undefined
- b) Zero
- c) Infinity
- d) Negative
- **Answer**: a) Undefined

7. The derivative of $f(x) = \sqrt{x}$ is:

- a) $\frac{1}{2\sqrt{x}}$
- b) $2x$
- c) $\frac{1}{x^2}$
- d) $\sqrt{x}$
- **Answer**: a) $\frac{1}{2\sqrt{x}}$

---

8. The derivative of $f(x) = e^x$ is:

- a) $e^x + x$
- b) e^x
- c) xe^{x-1}
- d) e^{x-1}
- **Answer**: b) e^x

9. The derivative of a constant function is:

- a) Zero
- b) Undefined
- c) The constant itself
- d) The constant multiplied by x
- **Answer**: a) Zero

---

10. The derivative of $f(x) = x^3$ is:

- a) $2x^2$
- b) $3x^2$
- c) $3x^3$
- d) x^2
- **Answer**: b) $3x^2$

11. A function is differentiable at x_0 if:

- a) It is continuous at x_0
- b) The limit defining the derivative exists at x_0
- c) Both a and b
- d) None of the above
- **Answer**: c) Both a and b

---

12. The derivative of $f(x) = \sin(x)$ is:

- a) $\cos(x)$
- b) $\sin(x)$
- c) $-\cos(x)$
- d) $-\sin(x)$
- **Answer**: a) $\cos(x)$

13. The derivative of $f(x) = \cos(x)$ is:

- a) $-\sin(x)$
- b) $\cos(x)$
- c) $\sin(x)$
- d) $-\cos(x)$
- **Answer**: a) $-\sin(x)$

---

14. A function is not differentiable at a point where the graph has a:

- a) Horizontal tangent
- b) Corner
- c) Continuous curve
- d) Positive slope
- **Answer**: b) Corner

15. The geometric interpretation of the derivative is:

- a) The area under the curve
- b) The slope of the tangent line at a point
- c) The value of the function
- d) The distance from the origin
- **Answer**: b) The slope of the tangent line at a point

---

16. The derivative of $f(x) = \ln(x)$ is:

- a) $\frac{1}{x^2}$
- b) $\frac{1}{x}$
- c) $\ln(x)$
- d) $x \ln(x)$
- **Answer**: b) $\frac{1}{x}$

17. The derivative of $f(x) = x^n$ (where n is a constant) is:

- a) nx^{n-1}
- b) n^x
- c) $n \cdot x^{n+1}$
- d) x^n
- **Answer**: a) nx^{n-1}

---

18. A function is not differentiable at a point of:

- a) Continuity
- b) Discontinuity
- c) Maximum
- d) Minimum
- **Answer**: b) Discontinuity

19. The derivative of $f(x) = \tan(x)$ is:

- a) $\sec^2(x)$
- b) $\sin(x)$
- c) $\cos^2(x)$
- d) $\tan(x)$
- **Answer**: a) $\sec^2(x)$

---

20. The derivative of $f(x) = \cot(x)$ is:

- a) $-\csc^2(x)$
- b) $\sec^2(x)$
- c) $\sin(x)$
- d) $\cos(x)$
- **Answer**: a) $-\csc^2(x)$

21. If $f'(x) = 0$ for all x, then $f(x)$ is:

- a) A linear function
- b) A constant function
- c) An exponential function
- d) Undefined
- **Answer**: b) A constant function

---

22. The slope of the tangent line to the graph o $f(x) = x^2$ at $x = 0$ is:

- a) 0
- b) 1
- c) 2
- d) Undefined
- **Answer**: a) 0

23. The derivative of $f(x) = x^{-1}$ is:

- a) $-x^{-2}$
- b) x^{-2}
- c) $-\frac{1}{x^2}$
- d) $\frac{1}{x^2}$
- **Answer**: c) $-\frac{1}{x^2}$

---

24. If a function $f(x)$ has a vertical tangent at $x = 0$, then:

- a) $f'(0) = \infty$
- b) $f'(0) = 0$
- c) $f'(0)$ does not exist
- d) $f'(0) = -\infty$
- **Answer**: c) $f'(0)$ does not exist

1. What is the derivative of a constant function $f(x) = c$?

- A) c
- B) 0
- C) c^2
- D) x

Answer: B) 0

2. The derivative of $f(x) = 5$ is:

- A) 5
- B) 1
- C) 0
- D) x^5

Answer: C) 0

3. According to the Power Rule, the derivative of $f(x) = x^n$ is:

- A) $n \cdot x^{n-1}$
- B) $n \cdot x^n$
- C) x^{n+1}
- D) $n \cdot x^{-n}$

Answer: A) $n \cdot x^{n-1}$

4. What is the derivative of $f(x) = x^3$?

- A) $3x$
- B) $3x^2$
- C) x^2
- D) $2x$

Answer: B) $3x^2$

5. The derivative of $f(x) = x^5$ using the Power Rule is:

- A) $5x^4$
- B) $5x^5$
- C) $4x^4$
- D) $6x^5$

Answer: A) $5x^4$

6. If $f(x) = c$ is a constant function, its derivative is geometrically interpreted as:

- A) The slope of a vertical line
- B) The slope of a horizontal line
- C) The slope of a curved line
- D) Undefined

Answer: B) The slo of a horizontal line

7. The derivative of $f(x) = 10x^2$ is:

- A) $10x^3$
- B) $20x$
- C) $5x^2$
- D) $2x$

Answer: B) 20x

8. What is the result of differentiating $f(x) = x^n$, where n is a positive integer?

- A) x^{n-1}
- B) $n \cdot x^{n-1}$
- C) $n \cdot x^n$
- D) x^n

Answer: B) $n \cdot x^{n-1}$

9. The quotient rule for the derivative of $f(x)/g(x)$ states:

- A) $\frac{g'(x)f(x)-f'(x)g(x)}{g(x)^2}$
- B) $\frac{f'(x)g'(x)}{g(x)^2}$
- C) $\frac{f(x)}{g'(x)}$
- D) $\frac{g'(x)}{f(x)^2}$

Answer: A) $\frac{g'(x)f(x)-f'(x)g(x)}{g(x)^2}$

10. If $f(x) = x^2 + 3x$ and $g(x) = x - 1$, what is the derivative of $\frac{f(x)}{g(x)}$?

- A) $\frac{(2x+3)(x-1)-(x^2+3x)(1)}{(x-1)^2}$
- B) $\frac{2x+3}{x-1}$
- C) $\frac{(2x+3)(x)}{x^2-1}$
- D) $\frac{x^2+3x}{(x-1)^2}$

Answer: A) $\frac{(2x+3)(x-1)-(x^2+3x)(1)}{(x-1)^2}$

11. The derivative of $f(x) = \frac{1}{g(x)}$ is given by:

- A) $-\frac{g'(x)}{g(x)^2}$
- B) $\frac{g(x)}{g'(x)^2}$
- C) $\frac{g'(x)}{g(x)^2}$
- D) $-\frac{g(x)}{g'(x)^2}$

Answer: A) $-\frac{g'(x)}{g(x)^2}$

12. The reciprocal rule for differentiation is derived using:

- A) The power rule
- B) The product rule
- C) The chain rule
- D) The quotient rule

Answer: D) The quotient rule

13. What is the derivative of $f(x) = 1/x^2$?

- A) $-2x^{-3}$
- B) $2x^{-2}$
- C) $-x^2$
- D) $2x$

Answer: A) $-2x^{-3}$

14. According to the generalized power rule, the derivative of x^{-n} is:

- A) $-n \cdot x^{-n-1}$
- B) $n \cdot x^{n-1}$
- C) $n \cdot x^{-n+1}$
- D) $-n \cdot x^{n-1}$

Answer: A) $-n \cdot x^{-n-1}$

15. What is the derivative of $f(x) = \frac{1}{x}$?

- A) $-x^{-2}$
- B) x^{-1}
- C) $1/x^2$
- D) $-1/x^2$

Answer: D) $-1/x^2$

16. The Power Rule applies to functions of the form $f(x) = x^n$. For which values of n does this rule hold?

- A) Positive integers only
- B) All integers
- C) Positive and negative integers
- D) Rational numbers only

Answer: B) All integers

17. The derivative of $f(x) = x^{-3}$ is:

- A) $3x^{-4}$
- B) $-3x^{-4}$
- C) $3x^{-3}$
- D) $-x^3$

Answer: B) $-3x^{-4}$

18. What is the derivative of $f(x) = \frac{x^2+1}{x}$?

- A) $\frac{x^2}{x^2}$
- B) $\frac{2x-1}{x^2}$
- C) 1
- D) $2x - x^{-2}$

Answer: B) $\frac{2x-1}{x^2}$

19. Which theorem allows us to compute the derivative of $f(x) = x^{-n}$?

- A) Quotient rule
- B) Chain rule
- C) Generalized power rule
- D) Constant rule

Answer: C) Generalized power rule

20. Using the power rule, what is the derivative of $f(x) = x^{\frac{1}{2}}$?

- A) $\frac{1}{2}x^{-\frac{1}{2}}$
- B) $x^{\frac{1}{2}}$
- C) $\frac{1}{2}x^{\frac{1}{2}}$
- D) $x^{-\frac{1}{2}}$

Answer: A) $\frac{1}{2}x^{-\frac{1}{2}}$

1. What is the derivative of a constant function $f(x) = c$?

- A) c
- B) 0
- C) c^2
- D) x

Answer: B) 0

2. The derivative of $f(x) = 5$ is:

- A) 5
- B) 1
- C) 0
- D) x^5

Answer: C) 0

3. According to the Power Rule, the derivative of $f(x) = x^n$ is:

- A) $n \cdot x^{n-1}$
- B) $n \cdot x^n$
- C) x^{n+1}
- D) $n \cdot x^{-n}$

Answer: A) $n \cdot x^{n-1}$

4. What is the derivative of $f(x) = x^3$?

- A) $3x$
- B) $3x^2$
- C) x^2
- D) $2x$

Answer: B) $3x^2$

5. The derivative of $f(x) = x^5$ using the Power Rule is:

- A) $5x^4$
- B) $5x^5$
- C) $4x^4$
- D) $6x^5$

Answer: A) $5x^4$

6. If $f(x) = c$ is a constant function, its derivative is geometrically interpreted as:

- A) The slope of a vertical line
- B) The slope of a horizontal line
- C) The slope of a curved line
- D) Undefined

Answer: B) The slo of a horizontal line

7. The derivative of $f(x) = 10x^2$ is:

- A) $10x^3$
- B) $20x$
- C) $5x^2$
- D) $2x$

Answer: B) 20x

8. What is the result of differentiating $f(x) = x^n$, where n is a positive integer?

- A) x^{n-1}
- B) $n \cdot x^{n-1}$
- C) $n \cdot x^n$
- D) x^n

Answer: B) $n \cdot x^{n-1}$

9. The quotient rule for the derivative of $f(x)/g(x)$ states:

- A) $\frac{g'(x)f(x)-f'(x)g(x)}{g(x)^2}$
- B) $\frac{f'(x)g'(x)}{g(x)^2}$
- C) $\frac{f(x)}{g'(x)}$
- D) $\frac{g'(x)}{f(x)^2}$

Answer: A) $\frac{g'(x)f(x)-f'(x)g(x)}{g(x)^2}$

10. If $f(x) = x^2 + 3x$ and $g(x) = x - 1$, what is the derivative of $\frac{f(x)}{g(x)}$?

- A) $\frac{(2x+3)(x-1)-(x^2+3x)(1)}{(x-1)^2}$
- B) $\frac{2x+3}{x-1}$
- C) $\frac{(2x+3)(x)}{x^2-1}$
- D) $\frac{x^2+3x}{(x-1)^2}$

Answer: A) $\frac{(2x+3)(x-1)-(x^2+3x)(1)}{(x-1)^2}$

11. The derivative of $f(x) = \frac{1}{g(x)}$ is given by:

- A) $-\frac{g'(x)}{g(x)^2}$
- B) $\frac{g(x)}{g'(x)^2}$
- C) $\frac{g'(x)}{g(x)^2}$
- D) $-\frac{g(x)}{g'(x)^2}$

Answer: A) $-\frac{g'(x)}{g(x)^2}$

12. The reciprocal rule for differentiation is derived using:

- A) The power rule
- B) The product rule
- C) The chain rule
- D) The quotient rule

Answer: D) The quotient rule

13. What is the derivative of $f(x) = 1/x^2$?

- A) $-2x^{-3}$
- B) $2x^{-2}$
- C) $-x^2$
- D) $2x$

Answer: A) $-2x^{-3}$

14. According to the generalized power rule, the derivative of x^{-n} is:

- A) $-n \cdot x^{-n-1}$
- B) $n \cdot x^{n-1}$
- C) $n \cdot x^{-n+1}$
- D) $-n \cdot x^{n-1}$

Answer: A) $-n \cdot x^{-n-1}$

15. What is the derivative of $f(x) = \frac{1}{x}$?

- A) $-x^{-2}$
- B) x^{-1}
- C) $1/x^2$
- D) $-1/x^2$

Answer: D) $-1/x^2$

16. The Power Rule applies to functions of the form $f(x) = x^n$. For which values of n does this rule hold?

- A) Positive integers only
- B) All integers
- C) Positive and negative integers
- D) Rational numbers only

Answer: B) All inte[illegible]s

17. The derivative of $f(x) = x^{-3}$ is:

- A) $3x^{-4}$
- B) $-3x^{-4}$
- C) $3x^{-3}$
- D) $-x^3$

Answer: B) $-3x^{-4}$

18. What is the derivative of $f(x) = \frac{x^2+1}{x}$?

- A) $\frac{x^2}{x^2}$
- B) $\frac{2x-1}{x^2}$
- C) 1
- D) $2x - x^{-2}$

Answer: B) $\frac{2x-1}{x^2}$

19. Which theorem allows us to compute the derivative of $f(x) = x^{-n}$?

- A) Quotient rule
- B) Chain rule
- C) Generalized power rule
- D) Constant rule

Answer: C) Generalized power rule

20. Using the power rule, what is the derivative of $f(x) = x^{\frac{1}{2}}$?

- A) $\frac{1}{2}x^{-\frac{1}{2}}$
- B) $x^{\frac{1}{2}}$
- C) $\frac{1}{2}x^{\frac{1}{2}}$
- D) $x^{-\frac{1}{2}}$

1. What does the Chain Rule help to find?

- a) Derivative of the sum of two functions
- b) Derivative of the product of two functions
- c) Derivative of the composition of two functions
- d) Integral of a function

Answer: c) Derivative of the composition of two functions

2. In the Chain Rule, if $y = f(g(x))$, what is $\frac{dy}{dx}$?

- a) $f'(g(x)) \cdot g'(x)$
- b) $g'(x) \cdot f(x)$
- c) $g'(x) + f(x)$
- d) $f'(x) \cdot g'(x)$ **Answer**: a) $f'(g(x)) \cdot g'(x)$

---

3. What is the derivative of $y = \sin(3x)$ using the Chain Rule?

- a) $3\cos(3x)$
- b) $\cos(3x)$
- c) $-3\cos(3x)$
- d) $3\sin(3x)$ **Answer**: a) $3\cos(3x)$

4. Given $y = \cos(5x^2)$, what is $\frac{dy}{dx}$?

- a) $-10x \sin(5x^2)$
- b) $5x^2 \cos(5x^2)$
- c) $5x \sin(5x^2)$
- d) $-5x^2 \sin(5x)$ **Answer**: a) $-10x \sin(5x^2)$

5. Which of the following is an alternative expression of the Chain Rule?

- a) Derivative of the outer function times the derivative of the inner function
- b) Derivative of the sum of two functions
- c) Derivative of the difference of two functions
- d) Derivative of the inner function squared

Answer: a) Derivative of the outer function times the derivative of the inner function

---

6. If $y = e^{2x+1}$, what is $\frac{dy}{dx}$?

- a) $e^{2x+1} \cdot 2$
- b) e^x
- c) $2e^x$
- d) $2e^{2x+1}$

Answer: a) $e^{2x+1} \cdot 2$

7. What is $\frac{d}{dx}(\tan(4x))$?

- a) $4\sec^2(4x)$
- b) $2\sec^2(x)$
- c) $\sec^2(4x)$
- d) $4\sin(4x)$ **Answer**: a) $4\sec^2(4x)$

---

8. For the function $y = \ln(3x + 1)$, what is $\frac{dy}{dx}$?

- a) $\frac{3}{3x+1}$
- b) $\ln(x)$
- c) $\frac{1}{3x+1}$
- d) $\frac{3x}{x+1}$ **Answer**: a) $\frac{3}{3x+1}$

9. If $y = (2x^2 + 1)^4$, the derivative $\frac{dy}{dx}$ is:

- a) $8x(2x^2 + 1)^3$
- b) $4(2x^2 + 1)^3$
- c) $8(2x^2 + 1)^4$
- d) $4x(2x^2 + 1)^2$ **Answer**: a) $8x(2x^2 + 1)^3$

---

10. In the Chain Rule, if $u = g(x)$, then $\frac{dy}{dx}$ is expressed as:

- a) $\frac{dy}{du} \cdot \frac{du}{dx}$
- b) $\frac{du}{dx} \cdot \frac{dx}{dy}$
- c) $\frac{dy}{dx} \cdot \frac{du}{dx}$
- d) $\frac{dy}{dx}$ **Answer**: a) $\frac{dy}{du} \cdot \frac{du}{dx}$

11. What is the derivative of $y = \cos(3x + 2)$?

- a) $-3\sin(3x + 2)$
- b) $3\cos(3x + 2)$
- c) $\sin(3x + 2)$
- d) $-\cos(3x + 2)$ **Answer**: a) $-3\sin(3x + 2)$

---

12. If $y = (5x^3 + 2)^6$, the derivative $\frac{dy}{dx}$ is:

- a) $90x^2(5x^3 + 2)^5$
- b) $6(5x^3 + 2)^5$
- c) $6x^2(5x^3 + 2)$
- d) $30x^2(5x + $ ^{6} **Answer**: a) $90x^2(5x^3 + 2)^5$

13. What is $\frac{d}{dx}\left(e^{x^2}\right)$?

- a) $2xe^{x^2}$
- b) e^{x^2}
- c) xe^{x^2}
- d) $2e^{x}$ **Answer**: a) $2xe^{x^2}$

15. If $y = \sin(2x)$, then $\frac{dy}{dx}$ is:

- a) $2\cos(2x)$
- b) $\cos(2x)$
- c) $-2\cos(2x)$
- d) $\sin(2x)$ **Answer**: a) $2\cos(2x)$

---

16. What is the derivative of $y = 3e^{4x}$?

- a) $12e^{4x}$
- b) $4e^{4x}$
- c) $3e^{x}$
- d) $12e^{x}$ **Answer**: a) $12e^{4x}$

17. In $\frac{dy}{dx} = f'(g(x)) \cdot g'(x)$, which part represents the derivative of the outer function?

- a) $f'(g(x))$
- b) $g'(x)$
- c) $f(g(x))$
- d) $f'(x)$ **Answer**: a) $f'(g(x))$

---

18. Which of the following functions requires the use of the Chain Rule for differentiation?

- a) $x^2 + 3x$
- b) $\cos(4x^2)$
- c) $2x + 5$
- d) x^2 **Answer**: b) $\cos(4x^2)$

19. For the function $y = \ln(7x + 3)$, what is $\frac{dy}{dx}$?

- a) $\frac{7}{7x+3}$
- b) $\frac{1}{7x+3}$
- c) $7\ln(7x + 3)$
- d) $\frac{7x}{x+3}$ Answer: a) $\frac{7}{7x+3}$

LEC 19

Implicit Differentiation MCQs

1. **What is the derivative of $x^2 + y^2 = 1$ using implicit differentiation?**
   - a) $2x + 2y\frac{dy}{dx} = 0$
   - b) $2x + \frac{dy}{dx} = 0$
   - c) $\frac{dy}{dx} = -\frac{x}{y}$
   - d) $2x - 2y\frac{dy}{dx} = 0$
   - **Answer:** a) $2x + 2y\frac{dy}{dx} = 0$
2. **Find $\frac{dy}{dx}$ if $x^2y + y^2x = 1$.**
   - a) $\frac{dy}{dx} = \frac{2x+y^2}{y+2xy}$
   - b) $\frac{dy}{dx} = \frac{-x^2}{y^2}$
   - c) $\frac{dy}{dx} = -\frac{2xy+y^2}{x^2+2xy}$
   - d) $\frac{dy}{dx} = -\frac{y}{x}$
   - **Answer:** c) $\frac{dy}{dx} = -\frac{2xy+y^2}{x^2+2xy}$

3. **Implicit differentiation is typically used when:**
   - a) y can easily be isolated.
   - b) y cannot be easily solved for in terms of x.
   - c) The equation is linear.
   - d) The equation is quadratic in x.
   - **Answer:** b) y cannot be easily solved for in terms of x.
4. **Find $\frac{dy}{dx}$ for $e^{xy} = 5$.**
   - a) $\frac{dy}{dx} = -\frac{y}{x}$
   - b) $\frac{dy}{dx} = \frac{-e^{xy}(x+y)}{x}$
   - c) $\frac{dy}{dx} = \frac{-e^{xy}}{y^2}$
   - d) $\frac{dy}{dx} = -\frac{y}{x+5}$
   - **Answer:** b) $\frac{dy}{dx} = \frac{-e^{xy}(x+y)}{x}$

5. **Which rule is used in implicit differentiation to differentiate terms with y ?**
   - a) Product rule
   - b) Chain rule
   - c) Quotient rule
   - d) Power rule
   - **Answer:** b) Chain rule
6. **Find $\frac{dy}{dx}$ for $y^3 + x^2y = 7$.**
   - a) $\frac{dy}{dx} = \frac{2xy}{3y^2 - x^2}$
   - b) $\frac{dy}{dx} = -\frac{2xy}{3y^2 + x^2}$
   - c) $\frac{dy}{dx} = \frac{3y^2 - 2x}{x}$
   - d) $\frac{dy}{dx} = -\frac{2xy}{y + x}$
   - **Answer:** b) $\frac{dy}{dx} = -\frac{2xy}{3y^2 + x^2}$

7. **In the equation $x^2 + xy + y^2 = 5$, find $\frac{dy}{dx}$.**
   - a) $\frac{dy}{dx} = \frac{-2x-y}{x+2y}$
   - b) $\frac{dy}{dx} = \frac{-x-y}{y+2x}$
   - c) $\frac{dy}{dx} = \frac{2x+y}{x+2y}$
   - d) $\frac{dy}{dx} = \frac{-x+y}{x+y}$
   - **Answer:** a) $\frac{dy}{dx} = \frac{-2x-y}{x+2y}$
8. **Find the derivative $\frac{dy}{dx}$ for $\sin(xy) = x + y$.**
   - a) $\frac{dy}{dx} = -\frac{\cos(xy)(x+y)}{1+\cos(xy)}$
   - b) $\frac{dy}{dx} = \frac{1-y}{x+\cos(xy)}$
   - c) $\frac{dy}{dx} = \frac{y\cos(xy)-1}{\cos(xy)-x}$
   - d) $\frac{dy}{dx} = \frac{\cos(xy)(x+y)-1}{\cos(xy)-y}$
   - **Answer:** c) $\frac{dy}{dx} = \frac{y\cos(xy)-1}{\cos(xy)-x}$

Derivatives of Rational Powers of x MCQs

9. **Find the derivative of $x^{3/2}$.**
   - a) $\frac{3}{2}x^{1/2}$
   - b) $\frac{2}{3}x^{-1/2}$
   - c) $\frac{3}{2}x^{1/2} + 1$
   - d) $\frac{1}{2}x^{3/2}$
   - **Answer:** a) $\frac{3}{2}x^{1/2}$

10. **The derivative of $x^{-4/3}$ is:**
   - a) $\frac{-4}{3}x^{-7/3}$
   - b) $\frac{4}{3}x^{-1/3}$
   - c) $-\frac{3}{4}x^{-7/3}$
   - d) $\frac{-4}{3}x^{-1/3}$
   - **Answer:** a) $\frac{-4}{3}x^{-7/3}$

11. **What is the derivative of $x^{2/5}$?**

- a) $\frac{2}{5}x^{-3/5}$
- b) $\frac{2}{5}x^{-1/5}$
- c) $\frac{5}{2}x^{-3/5}$
- d) $\frac{5}{2}x^{-7/5}$
- **Answer:** a) $\frac{2}{5}x^{-3/5}$

12. **Find the derivative of $\frac{1}{x^{1/2}}$.**

- a) $-\frac{1}{2}x^{-3/2}$
- b) $\frac{1}{2}x^{-1/2}$
- c) $\frac{-2}{x^{1/2}}$
- d) $\frac{-1}{2x^{3/2}}$
- **Answer:** a) $-\frac{1}{2}x^{-3/2}$

14. Find the derivative of $x^{5/3}$.

- a) $\frac{5}{3}x^{2/3}$
- b) $5x^{-2/3}$
- c) $\frac{3}{5}x^{2/3}$
- d) $5x^{4/3}$
- Answer: a) $\frac{5}{3}x^{2/3}$

LEC 20

1. What is the derivative of $\log_b(x)$ with respect to x?

a) $\frac{1}{x \ln(b)}$

b) $\ln(b) \cdot x$

c) $b^x \ln(b)$

d) $\frac{1}{x^2 \ln(b)}$

Answer: a) $\frac{1}{x \ln(b)}$

---

2. What is the derivative of $\ln(x)$?

a) $\frac{1}{x^2}$

b) $\ln(x^2)$

c) $\frac{1}{x}$

d) $x \ln(x)$

Answer: c) $\frac{1}{x}$

3. Which of the following is true for logarithmic differentiation?

a) It is used to differentiate exponential functions.

b) It is used to differentiate irrational powers.

c) It is used to simplify differentiation of complicated products or quotients.

d) It is used to differentiate trigonometric functions.

Answer: c) It is used to simplify differentiation of complicated products or quotients.

4. What is the derivative of $x^{\sqrt{2}}$?

a) $\sqrt{2}x^{\sqrt{2}-1}$

b) $\frac{1}{\sqrt{2}}x^{\sqrt{2}}$

c) $\sqrt{2}\ln(x)$

d) $x^{\sqrt{2}-2}$

Answer: a) $\sqrt{2}x^{\sqrt{2}-1}$

---

5. What is the derivative of a^x, where a is a constant?

a) $a^x \ln(a)$

b) $a^x \ln(x)$

c) $\ln(x)a^x$

d) $a^x \ln(a^x)$

Answer: a) $a^x \ln(a)$

6. If $y = e^x$, what is $\frac{dy}{dx}$?

a) $e^x + C$

b) $e^x \ln(e)$

c) e^x

d) xe^x

Answer: c) e^x

---

7. Which of the following functions has an inverse?

a) $f(x) = x^2$

b) $f(x) = e^x$

c) $f(x) = \sin(x)$

d) $f(x) = x^3 + x$

Answer: b) $f(x) = e^x$

8. What condition must a function satisfy to have an inverse?

a) It must be continuous.

b) It must be differentiable.

c) It must be one-to-one.

d) It must be even.

Answer: c) It must be one-to-one.

---

9. If $f(x) = \ln(x)$, what is the inverse of $f(x)$?

a) e^x

b) $\log_b(x)$

c) x^2

d) $\frac{1}{x}$

Answer: a) e^x

10. What is the derivative of the inverse function $f^{-1}(x)$ in terms of $f(x)$?

a) $\frac{1}{f'(x)}$
b) $f'(f^{-1}(x))$
c) $\frac{1}{f'(f^{-1}(x))}$
d) $-f'(x)$

Answer: c) $\frac{1}{f'(f^{-1}(x))}$

---

11. If $f(x) = 2^x$, what is the derivative of $f^{-1}(x)$?

a) $\frac{1}{x \ln(2)}$
b) $\frac{1}{2^x \ln(2)}$
c) $\frac{1}{\ln(2)}$
d) $2^x \ln(2)$

Answer: b) $\frac{1}{2^x \ln(2)}$

12. What is the derivative of $\ln(f(x))$ using the chain rule?

a) $\frac{1}{\ln(f(x))} \cdot f'(x)$

b) $\ln(f'(x))$

c) $\frac{1}{f(x)} \cdot f'(x)$

d) $\ln(f'(x)) \cdot f(x)$

Answer: c) $\frac{1}{f(x)} \cdot f'(x)$

---

13. If $f(x) = x^3$, what is the derivative of $\ln(f(x))$?

a) $\frac{3x^2}{x^3}$

b) $3\ln(x)$

c) $\frac{1}{x^3}$

d) $\frac{3}{x}$

Answer: d) $\frac{3}{x}$

14. What is the derivative of $\ln(x^2 + 1)$?

a) $\frac{1}{x^2+1}$
b) $\frac{2x}{x^2+1}$
c) $2\ln(x)$
d) $2x\ln(x)$

Answer: b) $\frac{2x}{x^2+1}$

---

15. If $f(x) = e^x + x^2$, what is $f'(x)$?

a) $e^x + 2x$
b) $e^x \ln(e) + 2x$
c) $xe^x + 2x$
d) $e^x + x^2$

Answer: a) $e^x + 2x$

16. What is the derivative of $\ln(7x)$?

a) $\frac{1}{x}$
b) $\frac{7}{x}$
c) $\frac{1}{x}\ln(7)$
d) $\frac{1}{x}$

Answer: d) $\frac{1}{x}$

---

17. The power rule for any real number r, $\frac{d}{dx}x^r$, gives:

a) rx^{r-1}
b) rx^{r+1}
c) $\frac{1}{r}x^{r-1}$
d) $r\ln(x)$

Answer: a) rx^{r-1}

18. If $f(x) = \sin(x)$, what is the inverse function $f^{-1}(x)$?

a) $\frac{1}{\cos(x)}$

b) $\arcsin(x)$

c) $\cos(x)$

d) $\sec(x)$

Answer: b) $\arcsin(x)$

---

19. What is the derivative of $\arcsin(x)$?

a) $\frac{1}{\sqrt{1-x^2}}$

b) $\frac{1}{\cos(x)}$

c) $\sqrt{1-x^2}$

d) $\frac{1}{x^2}$

Answer: a) $\frac{1}{\sqrt{1-x^2}}$

20. If $f(x) = \log_3(x)$, then the derivative of $f(x)$ is:

a) $\frac{1}{x \ln(3)}$

b) $\frac{1}{3 \ln(x)}$

c) $\frac{\ln(3)}{x}$

d) $\frac{3}{x}$

Answer: a) $\frac{1}{x \ln(3)}$

1. **Which of the following steps is NOT helpful in solving related rate problems?**
    - A) Draw a figure and label quantities.
    - B) Identify known and unknown rates of change.
    - C) Ignore the equation relating the quantities.
    - D) Differentiate the equation with respect to time.
    - **Answer: C) Ignore the equation relating the quantities.**

2. **If A is the area of a circle with radius r, what is the correct expression for the rate of change of area with respect to time?**
   - A) $\frac{dA}{dt} = 2r\pi$
   - B) $\frac{dA}{dt} = 2\pi r \frac{dr}{dt}$
   - C) $\frac{dA}{dt} = \pi r^2 \frac{dr}{dt}$
   - D) $\frac{dA}{dt} = \pi r^2$
   - **Answer: B)** $\frac{dA}{dt} = 2\pi r \frac{dr}{dt}$
3. **In the problem involving the oil spill, how fast is the area of the spill increasing when the radius is 60 ft?**
   - A) 120π square feet per second
   - B) 240π square feet per second
   - C) 360π square feet per second
   - D) 480π square feet per second
   - **Answer: B)** 240π **square feet per second**

4. **If a ladder of 5 feet is leaning against a wall and the base moves at 2 ft/sec, how fast is the top moving down when the base is 4 ft away?**
   - A) $-\frac{4}{3}$ ft/sec
   - B) $-\frac{3}{4}$ ft/sec
   - C) -2 ft/sec
   - D) $-\frac{8}{3}$ ft/sec
   - **Answer: A) $-\frac{4}{3}$ ft/sec**
5. **The Pythagorean theorem is useful in related rates problems involving which of the following shapes?**
   - A) Square
   - B) Triangle
   - C) Circle
   - D) Rectangle
   - **Answer: B) Triangle**

6. **What is the critical value of the function $f(x) = (x-2)^2$ where the function transitions from decreasing to increasing?**
   - A) $x = 1$
   - B) $x = 2$
   - C) $x = 0$
   - D) $x = -2$
   - **Answer: B)** $x = 2$
7. **In which interval is the function $f(x) = (x-2)^2$ increasing?**
   - A) $(-\infty, 2)$
   - B) $(2, \infty)$
   - C) $(-\infty, \infty)$
   - D) $(0, 2)$
   - **Answer: B)** $(2, \infty)$

8. **The slope of the tangent line to the graph of a decreasing function is:**
    - A) Positive
    - B) Negative
    - C) Zero
    - D) Undefined
    - **Answer: B) Negative**
9. **If $f'(x) > 0$ for all x in an interval, the function is:**
    - A) Decreasing
    - B) Increasing
    - C) Constant
    - D) Undefined
    - **Answer: B) Increasing**

10. **If $f'(x) = 0$ at a point, this point is known as:**
    - A) Maximum point
    - B) Critical point
    - C) Minimum point
    - D) Inflection point
    - **Answer: B) Critical point**
11. **In which of the following intervals is a function decreasing if $f'(x) < 0$ for all x in that interval?**
    - A) $(-\infty, 0)$
    - B) $(0, \infty)$
    - C) $(-\infty, \infty)$
    - D) None of these
    - **Answer: A) $(-\infty, 0)$**

12. **Which of the following represents the rate of change of one quantity with respect to another?**
    - A) Integration
    - B) Derivative
    - C) Algebra
    - D) Geometry
    - **Answer: B) Derivative**
13. **When the base of the ladder is moving away from the wall, the top of the ladder moves:**
    - A) Up the wall
    - B) Down the wall
    - C) Stays stationary
    - D) Moves in both directions
    - **Answer: B) Down the wall**

14. **Which of the following is TRUE for a concave-up function?**
    - A) The second derivative is negative.
    - B) The second derivative is positive.
    - C) The first derivative is positive.
    - D) The first derivative is zero.
    - **Answer: B) The second derivative is positive.**
15. **A function is said to be concave down if:**
    - A) Its graph is sloping upwards.
    - B) Its second derivative is positive.
    - C) Its second derivative is negative.
    - D) Its first derivative is zero.
    - **Answer: C) Its second derivative is negative.**

15. **A function is said to be concave down if:**
    - A) Its graph is sloping upwards.
    - B) Its second derivative is positive.
    - C) Its second derivative is negative.
    - D) Its first derivative is zero.
    - **Answer: C) Its second derivative is negative.**
16. **For a function $f(x)$, if $f'(x) = 0$ and $f''(x) > 0$, the point is:**
    - A) Maximum
    - B) Minimum
    - C) Inflection point
    - D) None of the above
    - **Answer: B) Minimum**

1. What is a relative maximum in a graph?

A) The highest value of the function

B) The highest point in a certain interval

C) The lowest point in the graph

D) The point where the derivative is undefined

Answer: B

2. A function has a relative minimum at a point if:

A) The function is decreasing in the interval

B) The function value at that point is less than or equal to other values in some interval

C) The function has a zero derivative at that point

D) The function is not differentiable at that point

Answer: B

3. At which of the following points is a relative extrema most likely to occur?

A) Where the function is undefined

B) At a critical point

C) Where the function has a vertical asymptote

D) At a point of inflection

Answer: B

4. What does it mean when a function has a critical point?

A) The function is continuous at that point

B) The derivative of the function is zero or undefined at that point

C) The function reaches its highest point at that point

D) The function is increasing at that point

Answer: B

5. A point where the tangent line to the graph is horizontal corresponds to:

A) A critical point

B) A point of inflection

C) A vertical asymptote

D) A discontinuity

Answer: A

6. Which of the following is NOT a requirement for a point to be a critical point?

A) The function is continuous at that point

B) The derivative is zero or undefined at that point

C) The function is differentiable at that point

D) The point exists in the domain of the function

Answer: C

7. The First Derivative Test is used to:

A) Determine where the function is increasing or decreasing

B) Find the concavity of the function

C) Determine if a function has vertical asymptotes

D) Identify horizontal asymptotes

Answer: A

8. A function has a relative maximum at a point if:

A) Its derivative changes from positive to negative at that point

B) Its second derivative is positive at that point

C) Its derivative changes from negative to positive at that point

D) Its second derivative is negative at that point

Answer: A

9. The Second Derivative Test is used to determine:

A) Where the function has a relative extremum

B) The concavity of the function at a critical point

C) Whether a function is increasing or decreasing

D) Where the function has asymptotes

Answer: B

10. What does it mean if the second derivative is positive at a point?

A) The graph is concave down

B) The graph is concave up

C) The function has a relative maximum

D) The function is decreasing

Answer: B

11. What type of function has vertical asymptotes?

A) Polynomial functions

B) Exponential functions

C) Rational functions

D) Linear functions

Answer: C

12. The horizontal asymptote of a rational function is determined by:

A) The degrees of the numerator and denominator

B) The critical points

C) The derivative of the function

D) The roots of the denominator

Answer: A

13. What does it mean if the first derivative of a function is positive in an interval?

A) The function is increasing in that interval

B) The function is decreasing in that interval

C) The function has a relative minimum in that interval

D) The function is concave down in that interval

Answer: A

14. What happens at an inflection point?

A) The function reaches a maximum

B) The concavity of the graph changes

C) The function is not differentiable

D) The function has a vertical asymptote

Answer: B

15. At which of the following points is a horizontal tangent found?

A) Where the derivative is 1

B) Where the derivative is 0

C) Where the second derivative is 0

D) Where the function is discontinuous

Answer: B

16. What is the primary purpose of the First Derivative Test?

A) To find inflection points

B) To determine where the function is concave

C) To identify relative extrema and intervals of increase/decrease

D) To determine the critical points of the function

Answer: C

17. What is the Second Derivative Test used to verify?

A) Whether a critical point is a relative extremum

B) Where the function has discontinuities

C) The exact value of the derivative

D) Where the function is continuous

Answer: A

18. In a polynomial function, what is the effect of the degree on the graph?

A) Determines the number of vertical asymptotes

B) Affects the number of turning points

C) Determines the x-intercepts

D) Affects the concavity of the graph

Answer: B

19. What is the general shape of the graph for a rational function at a vertical asymptote?

A) Flat line

B) Infinite slope

C) It tends to positive or negative infinity

D) It has a horizontal tangent

Answer: C

20. Which of the following is true about concavity in a graph?

A) If the second derivative is positive, the graph is concave down

B) If the second derivative is negative, the graph is concave up

C) Concavity changes at inflection points

D) Concavity is related only to horizontal asymptotes

Answer: C

EFFORT BY VU BSCS (M.TAQI)