بِسْمِ اللَّهِ الرَّحْمَٰنِ الرَّحِيمِ

Zoo_506

TODAY CURRENT MID TERM PAPERS

Fall 2022_23

ZOOLOGY
LEGENDS

"Don't Stop When you Tired,
Stop When you Done"


0304-4756496

MUHAMMAD ZAMAN

MCQS:

Prepare Quiz 1 Fall 2022 File

10+ Mcqs & Mostly from Past files



(1) In arThroPods The body is composed of a series of more or less Similar Pieces or Segments.

(2) Often Some of The bug; body Segments are fused with The Head to form a Crustacean CePhaloThorax

(3) In Archnida Generally have Segments of the body grouPed into tws - Sections.

(4) In The body of The adult insect There are apparently form 7 to 15 Segments.

(5) These ~~Sta~~ Sutures and The joint lines divide The Cuticula into areas called Plates or Sclerites

(6) Sense organ are Present in Antennae

(7) On The Second, or mesoThorax, and on Third, or metaThorax, both wings and legs occur in The majority of insects.

(8) The wings are Chitinous outgrowths form The body which very much size and form in different insects.

(9) The arrangement and number of Chief veins and Their branches are of importance in identifying insects.

Fareed

Date: / /

(10) The Alimentary canal extends form The mouth Through about The center of The body to The anus at The hinder end

(11) But They also receive much of The Carbondioxide gas Produced by The activities of The cell and Permit it to escaPe Through The sPiracles From The body.


(12) The cell by Their activities Produce not only carbon dioxide gas but also wast material Nitrogenous in nature which must be removed like All wastes from The body.

(13) The nervous system of insects is located along The m middle line. of The body quite near its under Surface.

(14) Below or behind The esoPhagus is a second ganglion, also in The head

(15) For The PurPose of together making hde in some object in which dePosit the eggs. oviPositor.

(16) Those which lay eggs hatching later are Termed oviParous


(17) In The Hemimetabola (or Paurometabola) The young insect on escaping from The egg, through resembling its Parents to some extent.

(18) The Feeding Periods between The molts (or ecdyses) are called Instal's

(19) In some flies The larva - shrinks within its larval skin and transforms, This skin, now called a Puparium functioning like a cocoon.

(20) The Nymphs of The Hemimetabola appear not to have attached sufficient attention to receive any special common ~~name~~ name.

Short Questions:

(Repeated Questions)

1. Define disease with example

Any harmful deviation from the normal structural or functional state of an organism is called Disease.

Example:

- Yellow Fever
- Malaria

2. What are tracheal gills?

Insects which in their early stage of live in water cannot, of course, breathe air into their bodies through spiracles during that period of their live.

There are enclosed in such cases and the animal obtains and usually through special structures called tracheal gills

3. How allergy is caused?

Allergy (hay fever, asthma, etc.) are frequently caused by exposure of susceptible persons to dried insect scales and hairs

4. Describe nematode disease and mention the name of pathogen?

Lymphatic filariasis (nematode worm): Culex, Anopheles, Aedes species of mosquito.

5. Physical transmission

• In case of Physical transmission the pathogen is simply carried externally or internally by the insect .Mostly Caused insects that have biting/chewing mouthparts.

6. Name the parts of legs in insect

- Coxa
- trochanter (two in a few cases)
- femur
- tibia
- tarsus

ZOOLOGY
LEGENDS

7. Contact insecticide

Certain materials called contact insecticides, which kill insects by contact with their bodies

Caused death by entering the spiracles and closing them up, thus producing suffocation.

Long question:

(Repeated Questions)

1. Explain and Name the types of four stages insect larvae.

Egg

Eggs are laid on plants by the adult female butterfly. These plants will then become the food for the hatching caterpillars.

Eggs can be laid from spring, summer or fall. This depends on the species of butterfly. Females lay a lot of eggs at once so that at least some of them survive. Butterfly eggs can be very small.

Caterpillar:

The Feeding Stage The next stage is the larva. This is also called a caterpillar if the insect is a butterfly or a moth. The job of the caterpillar is to eat and eat and eat. As the caterpillar grows it splits its skin and sheds it about 4 or 5 times.

Food eaten at this time is stored and used later as an adult. Caterpillars can grow 100 times their size during this stage. For example, a monarch butterfly egg is the size of a pinhead and the caterpillar that hatches from this tiny egg isn't much bigger. But it will grow up to 2 inches long in several weeks.

Pupa:

The Transition Stage When the caterpillar is full grown and stops eating, it becomes a pupa. The pupa of butterflies is also called a chrysalis.

Depending on the species, the pupa may suspended under a branch, hidden in leaves or buried underground. The pupa of many moths is protected inside a coccoon of silk. This stage can last from a few weeks, a month or even longer. Some species have a pupal stage that lasts for two years. It may look like nothing is going on but big changes are happening inside.

Special cells that were present in the larva are now growing rapidly. They will become the legs, wings, eyes and other parts of the adult butterfly. Many of the larva original cells will provide energy for these growing adult cells.

Adult:

ZOOLOGY LEGENDS

The Reproductive Stage The adult stage is what most people think of when they think of butterflies. They look very different from the larva. The caterpillar has a few tiny eyes, stubby legs and very short antennae. The adults have long legs, long antennae, and compound eyes. They can also fly by using their large and colorful wings.

The one thing they can't do is grow. Most adult butterflies live only one or two weeks, but some species hibernate during the winter and may live several months.

2. Name the parts of thorax and abdomen of insect.

• The thorax has its three segments usually quite clearly marked

• 1-Prothorax (have no wings)

• 2-Mesothorax (have fore wings)

• 3-Metathorax (hind wings)

• The first abdominal segment, when seemingly more apart of the thorax than of the abdomen, is called the median segment or ropodeum.

On the abdomen are various organs such as

◆ the ovipositor, ◆ sting ◆ cerci ◆ and styli, present in some cases, absent in others.

Different characteristics to identify the insect

- Bilaterally symmetrical.
- Consisting of a series of segments. (a) Head (b) thorax (c) abdomen.
- The animal is covered by a Shell like skeleton present outside the body, horny in nature.
- Three pairs of jointed legs attached to the thorax.
- Pair of antenna.
- Mouth parts.
- Two pair of wings.
- Breathing through Air tubes.
- Reproductive organs present near the hinder end of the body

3. Insect-borne diseases

Insect-borne diseases are viral and bacterial illnesses from insect (bug) bites.

• mosquitoes are known for spreading the Zika virus, Yellow Fever, and Malaria.

• Malaria (protozoan): Anopheles species of mosquito.

• Lymphatic filariasis (nematode worm): Culex, Anopheles, Aedes species of mosquito.

• Dengue (virus): Aedes species of mosquito.

ZOOLOGY LEGENDS

• Leishmaniasis (protozoan): mainly Phlebotomus species of sandfly.

4. Differentiate between pulivillus, pulvilli, and epodium ?

At the end of the last a pair of claws is generally found, and between them a sort of pad or cushion, the pulvillus Sometimes there are three of these, in which case the outer ones are called the pulvilli and the middle one the empodium

5. Nervous system of insects

The nervous system of insects is located along the middle line of the body quite near its under surface As in animals generally, it is composed of cells and fibers Ganglia: former are for the most part gathered together in clusters which are called ganglia And from each of the cells in a ganglion one or more nerve fibers pass out, to connect either with some other nerve cell or with some structure of the body

The larger nerves are really bundles of the fibers running side by side like the wires of a telephone cable Apparently each segment of the insect body once had but with the fusion of the segments tube of a fly Greatly many of these have also fused, reducing the separate ganglia in adult insects to a smaller number, which varies central in different kinds This fusion has been produced by the hinder ganglia moving forward until in some cases none is found in the abdomen

Each ganglion is connected to the one in front and the one behind by one or two bundles of nerve fibers which are called commissures Each consists of numerous fibers which taken together form the means of communication between the different parts of the system

6. Name head positions in insect

Heads with the mouth beneath are called hypognathous those with it in front are prognathous. Structures found on the head are

- A pair of antennae
- Two compound eyes (ocelli)
- The mouth parts

Four types of larvae.....

Types of larvae:

1. Larvae of butterflies and moths are commonly called **caterpillars**
2. Those of beetles are usually called **grubs**
3. Those of flies are called **maggots**.
4. Larvae found boring in wood, however, whether they will become moths, beetles or other insects, are uniformly called **borers**.

Zoology legends:

https://chat.whatsapp.com/HMS1CGrZW4h7QWa1D53roe

Zoo506:

https://chat.whatsapp.com/DszoDQiNCkDJzxt7drsjx2



Muhammad Zaman

03044756496

Jazak AllaH