



Sta 301 Quiz no - Microsoft Word
File
Home
Insert
Page Layout
References
Mailings
Review
View
Contact For Quiz
0343-8509619
1. The joint probability function, also called the
bivariate probability function
2. E(XY) is Equal to
E(X) E(Y)
3. The number of success, X in a hypergeometric experiment is called a
hyper geometric random variable
4. The probability distribution is called the
hyper geometric distribution
Activate Windows
Page: 1 of 4
Words: 425
160%
Type here to search
29°C
11:43 AM
8/14/2022


Sta 301 Quiz no - Microsoft Word
File
Home
Insert
Page Layout
References
Mailings
Review
View
Calibri (Body)
26
Paste
Clipboard
Font
Paragraph
Styles
Editing
Find
Replace
Select
Change Styles
Normal
No Spaci...
Heading 1
Heading 2
5. The outcomes of each trial may be classified into one of two categories,
success and failure
6. A normal distribution whose mean is zero and whose standard deviation is 1 is known as the
standard normal distribution.
7. Point estimation of a population parameter provides as an estimate a_____ value calculated
from the sample.
Single
8. An estimate is a numerical value of the unknown parameter obtained by applying a rule or a
formula, called
an estimator
9. sample is only a_____ of the population.
Activate Windows
Go to Settings to activate Windows.
Page: 1 of 4
Words: 425
160%
Type here to search
29°C
11:44 AM
8/14/2022


Sta 301 Quiz no - Microsoft Word
File
Home
Insert
Page Layout
References
Mailings
Review
View
Calibri (Body)
11
Paste
Clipboard
Font
Paragraph
Styles
Editing
Find
Replace
Select
Change Styles
¶ Normal
¶ No Spaci...
Heading 1
Heading 2
Part
10. The expected value of the sum of any two random variables is equal to the___of their expected values
Sum
11. The expected value of the sum of any two random variables is equal to the sum of their expected values.
E(X + Y) = E(X) + E(Y)

12. The probability of success changes on _____
each trial

13. The Poisson distribution is named after the French mathematician Sime'on Denis Poisson
(1781-1840)

14. The Poisson published its derivation in the year
1837.

15. The Poisson distribution has only _____ $\mu > 0$
one parameter

16. The mean and variance are equal in ___
THE POISSON DISTRIBUTION

17. The binomial distribution is a very important
discrete probability distribution

18. It was discovered by James Bernoulli about the
year 1700

19. E(4X + 5) = __

4 E(X) + 5

Activate Windows

20. The marginal p.d.f. of any of the_____ is obtained by integrating out the other variable from the joint p.d.f.


Variables

21. $E(3X + 5) = 3\,E(X) + 5$

22. he point to be understood here is that, from the joint probability function for (X, Y), we can obtain the INDIVIDUAL probability function of X and Y. Such individual probability functions are called

Activate Windows
Go to Settings to activate Windows.

Type here to search
29°C 11:47 AM 8/14/2022


Sta 301 Quiz no - Microsoft Word
File
Home
Insert
Page Layout
References
Mailings
Review
View
Paste
Clipboard
Calibri (Body)
11
Font
Paragraph
AaBbCcDc
¶ Normal
AaBbCcDc
¶ No Spaci...
AaBbC
Heading 1
AaBbCc
Heading 2
Change Styles
Styles
Find
Replace
Select
Editing
MARGINAL probability functions.
23. The expected value of the product of two independent r.v.'s is equal to the_____ of their expected values.
Product
24. The successive trials are not
independent.
Activate Windows

25. The hyper geometric probability distribution has _____N, n and k.

three parameters

26. _____may be defined as a physical process governed at least in part by some random mechanism

POISSON PROCESS

27. Poisson distribution can never be

negatively skewed

Activate Windows
Go to Settings to activate Windows.



Type here to search
29°C
11:49 AM
8/14/2022

In hyper geometric distribution formula, N= number of units in the population, n = number of units in the sample, and k is defined as

Select the correct option

- number of failures in the sample
- number of successes in sample
- number of failures in population
- number of successes in population

STA301 - Statistics and Probability (Quiz#3)

Question # 5 of 10 (Start time: 03:13:37 PM, 15 August 2022)

In hyper geometric distribution formula, N= number of units in the population, n = number of units in the sample, and k is defined as:

Select the correct option

- number of failures in the sample
- number of successes in sample
- number of failures in population
- number of successes in population

A random experiment has five outcomes in its sample space {s1, s2, s3, s4, s5}.
If P(s1)=0.2,P(s2)=0.3, P(s3)=0.1 and P(s4)=0.2
then P(s5)=?

Select the correct option

- ○ 0.8
- ○ 1
- ○ 0.2
- ○ 0.5

The variance of the Hypergeometric distribution is:

Select the correct option

○	{nk(N-k)(N-n)}/N*N(N-1) ✓
○	pq
○	npq
○	{nk(N-n)}/N(N-1)

The function F(x) gives the probability of the event that X takes a value a value...

lect the correct option

- $P(X<x)$
- $P(X\leq x)$
- $P(X>x)$
- $P(X=x)$

The standard normal distribution has it mean and variance equal to:

Select the correct option

- ○ 0 and 0 respectively
- ○ 0 and 1 respectively
- ○ 1 and 1 respectively
- ○ 1 and 0 respectively

not Sure

BC150402399: Muhammad Asad Latif

STA301 - Statistics and Probability (Quiz#3)

Question # 5 of 10 (Start time: 03:35:58 PM, 15 August 2022)

In sampling distribution of the sample proportion p is equal to:

Select the correct option

- ○ n/x
- ○ x
- ○ n
- ○ x/n

According to emperical rule, how much data lies between $\mu - 3\sigma$ and $\mu + 3\sigma$?

ct the correct option

- 95.44%
- 99.73%
- 68.26%
- 50%

Question # 4 of 10 (Start time: 03:34:09 PM, 15 August 2022)

The hyper geometric probability distribution is appropriate when a random sample of size n is drawn withou
population of N units:

Select the correct option

- total
- All
- finite
- Infinite

Binomial distribution is skewed to the right if:

elect the correct option

- p=q
- P<q
- p>q
- p=n

uestion # 10 of 10 (Start time: 03:43:48 PM, 15 August 2022)

Let X be a random variable with binomial distribution, that is (X=0,1,..., n). The expected value E[X] is

elect the correct option

- p
- Xnp
- np
- np(1-p)

Question # 7 of 10 (Start time: 03:40:07 PM, 15 August 2022)

If the first moment ratio is greater than 0, then the distribution will be:

Select the correct option

- ○ Negatively Skewed
- ○ Positively Skewed
- ○ Symmetrical
- ○ None of these

In sampling distribution of the sample proportion p is equal to:

Select the correct option

○	n/x
○	x
○	n
○	x/n

The joint density function f(x,y) will be a pdf if

Select the correct option

- ○ Both of integrals f(x,y) dx dy =0
- ○ Both of integrals f(x,y) dx dy =1 ✓
- ○ Both of integrals f(x,y) dx dy <1
- ○ Both of integrals f(x,y) dx dy >1

Question # 7 of 10 (Start time: 06:19:28 AM, 15 August 2022) Total Marks: 1

In which of the following distributions is the probability of a success usually small?

Select the correct option

- Binomial
- Hypergeometric
- All distribution
- Poisson

P[(X+Y) < 2] =? , where x=0, 1, 2 and y=0, 1, 2

Select the correct option

- f(0,0) + f(1,1) + f(1,0)
- f(0,0) + f(1,1) + f(2,0)
- f(0,1) + f(0,2) + f(2,0)
- f(0,0) + f(0,1) + f(1,0)

Question # 9 of 10 (Start time: 06:21:33 AM, 15 August 2022) **Total Marks:** 1

Which of the following is NOT CORRECT about a standard normal distribution?

Select the correct option

- P (Z = -1.0) = .03413
- P (Z = -2.5) = .4938
- P(Z = 2.0) = .0228
- P (0 = Z = 1.50) = .4332

Binomial distribution has two parameters; which are:

Subscribe for more Solutions

Select the correct option

- None
- p and n ✓
- p and q
- X and n

Uniform distribution is a:

Select the correct option

- ○ None of the above
- ○ Mixed
- ◉ Discrete distribution
- ○ Continuous Distribution

Question # 6 of 10 (Start time: 12:49:27 PM, 15 August 2022) Total Marks: 1

The number of parameters in normal distribution is(are):

Select the correct option

- 1
- 3
- 4
- 2

sec(s)

STA301 - Statistics and Probability (Quiz#3)

Quiz Start Time: 12:43 PM, 15 August 2022

Question # 7 of 10 (Start time: 12:50:42 PM, 15 August 2022)

Total Marks: 1

For the Poisson distribution $P(X = 1) = \frac{e^{-0.135}0.135^1}{1!}$ the mean value is:

Select the correct option

Reload Math Equations

- $e^{-0.135}$
- 0.135
- 1
- -0.135

Vu Study Center

Question # 8 of 10 (Start time: 12:51:53 PM, 15 August 2022) Total Marks: 1

The function F(x) gives the probability of the event that X takes a value a value.....

Select the correct option

- $P(X<x)$
- $P(X\leq x)$
- $P(X>x)$
- $P(X=x)$

Which of the following is true for the Poisson distribution:

Select the correct option

- ○ mean = standard deviation
- ◉ mean = variance
- ○ mean > variance
- ○ mean > variance

The shape parameter of normal distribution is:

Select the correct option

- Mean
- Mode
- Median
- Standard Deviation

Quiz

tps://vulms.vu.edu.pk/Quiz/QuizQuestion.aspx?ver=d15e2dac-b785-409b-9947-73e7a03f9025

Q: ... 06:09:16 AM, 15 August 2022)

Total Marks: 1

In a binomial experiment:

Select the correct option

www.thundershare.net

- Number of trials "n" is a fixed
- Number of trials "n" is a variable
- Number of trials "n" is not specified
- None of the above

Type here to search

New folder (2)

Quiz - Cent Brow...

STA301 Handouts...

ENG

6:09 am

15/08/2022