

Question # 1 of 10 ( Start time: 03:24:13 PM, 23 July 2023 )

Total Marks: 1

If a statistic used as an estimator, has its expected value equal to the true value of the population parameter being estimated then it is called

Select the correct option

|                                  |            |
|----------------------------------|------------|
| <input checked="" type="radio"/> | Unbiased   |
| <input type="radio"/>            | Sufficient |
| <input type="radio"/>            | Consistent |
| <input type="radio"/>            | Efficient  |

Click to Save Answer & Move to Next Question

Question # 5 of 10 (Start time: 03:27:36 PM, 23 July 2023)

Total Marks: 1

In a binomial probability distribution, relation between mean and variance is:

Select the correct option

|                                  |                   |
|----------------------------------|-------------------|
| <input type="radio"/>            | Mean > Variance   |
| <input type="radio"/>            | Mean < Variance   |
| <input checked="" type="radio"/> | Mean = Variance   |
| <input type="radio"/>            | Difficult to tell |

[Click to Save Answer & Move to Next Question](#)

Question # 3 of 10 ( Start time: 03:25:49 PM, 23 July 2023 )

Total Marks: 1

Which of the following is the property of hypergeometric experiment?

Select the correct option

|                                  |                                           |
|----------------------------------|-------------------------------------------|
| <input type="radio"/>            | p remains constant from trial to trial    |
| <input type="radio"/>            | Successive trials are independent         |
| <input type="radio"/>            | n is not fixed                            |
| <input checked="" type="radio"/> | Sampling is performed without replacement |

Click to Save Answer & Move to Next Question

Question # 7 of 10 ( Start time: 03:29:18 PM, 23 July 2023 )

Total Marks: 1

Which of the following is not true statement for the hypergeometric distribution?

Select the correct option

|                                  |                                   |
|----------------------------------|-----------------------------------|
| <input checked="" type="radio"/> | Trials are dependent              |
| <input type="radio"/>            | A constant probability of success |
| <input type="radio"/>            | A fixed number of trials          |
| <input type="radio"/>            | Only two possible outcomes        |

Click to Save Answer & Move to Next Question



Question # 2 of 10 ( Start time: 03:25:03 PM, 23 July 2023 )

Total Marks: 1

If we have the following information,  
 $P = 0.16$ ,  $q = 0.84$ ,  $Z = 1.96$  and  $e = 0.05$ , then sample size will be:

Select the correct option

|                                  |     |
|----------------------------------|-----|
| <input type="radio"/>            | 199 |
| <input type="radio"/>            | 205 |
| <input checked="" type="radio"/> | 207 |
| <input type="radio"/>            | 200 |

[Click to Save Answer & Move to Next Question](#)

Question # 10 of 10 ( Start time: 03:32:00 PM, 23 July 2023 )

Total Marks: 1

The method of Maximum Likelihood (ML) is used to find out:

Select the correct option

|                                  |                         |
|----------------------------------|-------------------------|
| <input type="radio"/>            | Correlation coefficient |
| <input type="radio"/>            | Random numbers          |
| <input checked="" type="radio"/> | Point estimates         |
| <input type="radio"/>            | Sample values           |

[Click to Save Answer & Move to Next Question](#)

Question # 4 of 10 ( Start time: 03:26:43 PM, 23 July 2023 )

Total Marks: 1

An urn contains 4 red balls and 6 green balls. A sample of 4 balls is selected from the urn without replacement. It is the example of:

Select the correct option

|                                  |                            |
|----------------------------------|----------------------------|
| <input type="radio"/>            | Exponential distribution   |
| <input type="radio"/>            | Binomial distribution      |
| <input type="radio"/>            | Poisson distribution       |
| <input checked="" type="radio"/> | Hypergometric distribution |

[Click to Save Answer & Move to Next Question](#)

Question # 6 of 10 ( Start time: 03:28:28 PM, 23 July 2023 )

Total Marks: 1

..... is used when we are dealing with proportions to find the probabilities:

Select the correct option

|                                  |                         |
|----------------------------------|-------------------------|
| <input checked="" type="radio"/> | Normal distribution     |
| <input type="radio"/>            | F distribution          |
| <input type="radio"/>            | t- distribution         |
| <input type="radio"/>            | Both Z & t distribution |

[Click to Save Answer & Move to Next Question](#)



Question # 8 of 10 ( Start time: 03:30:21 PM, 23 July 2023 )

Total Marks: 1

Which of the following is most important and most widely used method in point estimation?

Select the correct option

|                                  |                                  |
|----------------------------------|----------------------------------|
| <input type="radio"/>            | The method of fractional moments |
| <input type="radio"/>            | The method of moments            |
| <input checked="" type="radio"/> | The method of maximum likelihood |
| <input type="radio"/>            | The method of least square       |

Click to Save Answer & Move to Next Question

Question # 9 of 10 ( Start time: 03:31:10 PM, 23 July 2023 )

Total Marks: 1

Consistency is a ..... property.

Select the correct option

|                                  |                   |
|----------------------------------|-------------------|
| <input type="radio"/>            | Unique sample     |
| <input checked="" type="radio"/> | Large sample      |
| <input type="radio"/>            | Small sample      |
| <input type="radio"/>            | Non-random sample |

[Click to Save Answer & Move to Next Question](#)