8- Irrational numbers are
Ans- Can't be written as fraction

9:- The null set is designated with the symbol.
Ans- ϕ

10- Range of $A = [(0,2)(2,4)(3,4)(4,5^-)]$ is
Ans-
Rang of $A = (0,2,3,4)$ not confirm

11- If A and B are matrices, then which from the following is true.
Ans:-
$AB \neq BA$

12- Graph is symmetric with respect to the origin is the graph of an.
Ans:-
odd function

13- An unordered collection of distinct objects.
Ans:-
Set

14- Sets are generally represented by.
Ans:-
Capital letters

Quiz STA 100

(8) Irrational number are
Can't be written as friction

(9) The Null is designated with Symbol
$\emptyset$

(10) Range of A = $[(0,2), (2,4), (3,4), (4,5)]$ is
2, 4, 5

(11) If A and B are Matrices Then which form the following is true
$AB \neq BA$

(12) Graph is Symmetric with respect to the origin is the graph of
odd Function

(13) An unordered Collection of distinct objects.
Set

(14) Set is generally represented by.
Capital letter

(15) If $F(-x) = F(x)$ for every number x is domain of f is
even Function

Created by Amaan Khan

My status
Just now

Quiz STA100

1. If we solve ax2+bx+c=0. By completing square it gives us.
Ans. quadratic formulae.
2. Set of rational number is denoted by (Q)
3. 0.3333 is (rational number)
4. If f(-x)= f(x) for every value of x then it is (even function)
5. An unordered collection of distinct object is (set)

Questions same arahee hay only 3 questions different hoo gee

0

Created by Amaan Khan

Quiz STA 100

(16) The Set Consist of member of pre-image or input of function Called
Domain

(17) Two Set A and B are equal if (all element of Set A are in Set B).
$A \subseteq B$ and $B \subseteq A$

(18) Cardinatity of Set A is denoted by.
$|A|$

(19) $\{ x / x < 6 \quad x \text{ is Counting number} \}$ is an example of.
Set builder.

(20) If $b^2 - 4ac < 0$
There will be no real Solution.

(21) $(a+bi) + (c+di)$
$(a+c) + (b+d)i$

(22) $i(3i+2) = 2$

Created by Amaan Khan

15- If $f(-x) = f(x)$ for every number x in domain of f, then f is

Ans:-

Even function

16- The set consisting of members of the pre-image or inputs of a function is called.

Ans:-

Domain

17- Two sets A and B are equal if.

Ans:- $A \subseteq B$ and $B \subseteq A$.

if all elements of set A are in set B

18- Cardinality of Set A is denoted by.

Ans:-

$|A|$

Created by Amaan Khan

① Number in form $z = a + bi$ where $a, b \in R$ and $i = \sqrt{-1}$ is called a **Complex number.**

② If a function is to have an inverse which is also a function then it must be **One to one.**

$(A \cap B) \cap C = B \cap (A \cap C)$ **Associative of intersectio**

The null set designated with the symbols ———

Irrational numbers are **Can't be written as fraction.**

If every element of set A is an element of set B is denoted by ———

0.333333 is **Rational numbers.**

Rang of $A = [\{(0,2),(2,4)(3,4)(4,5)\}]$ is ———

Math type MTEF12!f+!.

feaagaail lev 2aal CVAUfe BSiUv2L———

Created by Amaan Khan

Stat 100 Quiz #1

1- Axis of symmetry in form of vertical line separates parabola into

Ans- Two equal halves.

2- %Mth type!MTEF!2!1!+-%
feaaaaaadI ev2aaatCvAUfeBSiuvZL2vd9azLbvvNv2CaerbuLwBLn%

Ans-

3- 0.333333 is

Ans:- Rational number

4- Number in form $z=a+bi$ where $a,b \in R$ and $i=\sqrt{-1}$ is called a

Ans:- Complex number

5- $(A \cap B) \cap C = B \cap (A \cap C)$

Ans- Associative property of intersection

6- If a function is to have an inverse which is also a function then it must be.

Ans- One to One function

7- If every element of set A is an element of set B is denoted by.

Ans-

$A \subseteq B$

Created by Amaan Khan

=M.Sc Zoology 2nd Semester= NASIR Ali.

Quiz.1 STA100.

① Axis of Symmetry in form of Vertical Separates parabola into
Two equal Halves

② % Math Type ! MTEF ! 2! 1! ± 7.
Feaaaaart 1ev2aaat CvAUfeBsiuv2L2vd9a2
LbVVNv2CaerbuLwBLn%.

③ 0.3333 is = Rational Number

④ Number in Form $z = a+bi$ where $a.b \in R$ and $i = \sqrt{-1}$ is Called
Complex Number.

⑤ $(A \cap B) \cap C = B \cap (A \cap C)$.
Associative Property of intersection

⑥ If every element of Set A is the element of Set B is denoted by
$A \subseteq B$

⑦ If a function is to have one inverse which is also a function Then it must be?
one to one function.

Axis of symmetry in form of vertical line separates parabola into **Two equal halves**?

39. The graph of quadratic function will form parabola.
40. The solution of quadratic equation are also roots.

Created by Amaan Khan

(36) Object in Set are Called.

element.

(37) Given Complex No $Z = a+ib$ its Conjugate is

$\bar{Z} = a-ib$

(38) Quadritic function also called.

Co-efficient.

Quiz STA 100

(23) Mapping where two or more Number of Pre-image mapped to exactly one no of image.

Many one relation

(24) The Quadratic formula for $ax^2+bx+c=0$

$$x = \frac{b \pm \sqrt{b^2-4ac}}{2a}$$

(25) We Can add two matrices having real number A and B if their

Order is Same.

(26) Root of $x^2+8x+12$

$-2, -6$

(27) If order of Matrix A is $m*p$ and the order of B is $p*n$ Then order of matrix AB is

$m \times n$

(28) Set of rational Number is denoted by

Q.

(29) The Solution of quadritic equat also called it

Roots

Created by Amaan Khan

1. Axis of symtry in form of vertical spirated parabola into Two equal halves
2. 0.3333 is a Rational nmbrs > Tehreem Fatima
3. Nmbr in form $z = a+bi$ where a, b ∈ R & $i = \sqrt{-1}$ called complx nmbrs
4. $(A \cap B) \cap C = B \cap (A \cap C)$. Associate property of intersection
5. If function is to have an inverse which also function one to one fncn
6. If every element of set A is an element of set B is denoted $A \subseteq B$.
7. Irrational numbers are cant be writen as fraction
8. The null set designated with symbol $\emptyset$
9. If A & B are matrix, then which form is true $AB \neq BA$.
10. If 2 set A & B are given, set consisting of all element which either A or B or in both Union of A & B.
11. In quadratic equation, when a is positive then graphs concaves downward
12. Graph is symtric with respect to origin is graph of an Odd function
13. An unordered collection of distinct object Set
14. Sets are generally represented by Capital letters
15. If we solve $ax^2+bx+c=0$ by comping square it gives us Quadratic formula
16. Set of rational number is denoted by Q
17. If $f(-x) = f(x)$ for every value of x then it is Even function
18. If $b^2-4ac < 0$, there will be NO real solution
19. Cardinality of set A $|A|$
20. The set consisig of member of function is called Domain
21. Two sets A & B are equal if $A \subseteq B$ and $B \subseteq A$
22. $(a+bi)+(c+di)$ $(a+c)+(b+d)i$
23. $\{x \mid x < 6$ and is counting numbers$\}$ is exple of Set builder method.
24. we can add two matrix having real numbers A and B if their order is same.
25. The Quadratic formula for $ax^2+bx+c=0$ $x = \frac{b \pm \sqrt{b^2-4ac}}{2a}$
26. Maping where two or more member of pair range is mapped to exactly one nmbr is many-one relation
27. $f(3i+2)$ 2
28. If order matrix A is $m \times p$ & order of B is $p \times n$, then order of matrix m x n
29. Range of $A = [(0,2)(2,4)(3,4)(4,5)]$ is Range of A (0, 2, 3, 4).
30. Roots of $x^2+8x+12$ are $-2, -6$
31. If matrix has equal number of column & rows then its Square matrix
32. which of folwing property of matrix multiplication coorect All mentioned.
33. object in set are called Elements
34. Two set A & B contaig these element which are of A or B A∪B
35. Given

Created by Amaan Khan

Quiz 1

Q determinant of null matrix

ans 0

Q if the order of 3×2 then what should be the order of matrix B , Solution is possible.

ans 3×1

Q Two matrix same order can be

ans added and subtract

Q- determinant is process of allocating a real number to every

ans square.

Q If the order of matrix A is 2×3 then what should be the order of matrix B such that solution of matrix A and B possible

2×3

Created by Amaan Khan

Q1

determinant of Identity matrix?

ans 1

Q2 If a scalar λ is multiplied with the matrix then which one following statement is correct?

ans Two λ can be multiplied with all elements of the matrix

Q3 Given a square matrix A, if there exist a matrix B such that $AB = BA = I$, then A is called the ___ of B

ans multiple inverse

Q2 which of the following matrix have no inverse.

ans Non-Singular matrix

Q1 To add two matrix order must be

ans

MTH 100

10. A square matrix which one diagonal (1s) and zero else where is called a/an ... matrix.

ans :- Identity matrix

09. A minor of an element in a third order determinant is a .....

ans Second order

08. If B is a matrix whose determinant zero then it inverse. . . .

(a) will be zero (b) cannot determine
(c) will be negative (d) will be positive.

07. if matrix have A an order 3×5 matrix B order 5×3, order of A×B is

ans => 3×3

06. A and B are any two matrices of order m and n respectively and m>n and p<q for the product condition m n p q AB hold.

(a) n=p (b) n>p (c) n<p

Created by Amaan Khan

Quiz 07

1) Identify the sequence Arithmetics of 8, 13, 18, 23

(a) Arithmetic (b) Geometric

Ans (c)

2) Explicit formula for the sequence with term 2, 7, 12, 17 is

(a) $5n+3$ (b) $5n-3$ (c) $n+2$

Ans (d) $n-3$

3) For a geometric sequence if 2nd term is equal to 4th term then value of common ratio is.

(a) -1 only (b) 1 only (c) 0

Ans (d) $+1$ or -1

4) If the sum of first n natural number is $n(n+1)/2$ the first average will be.

(a) $n/2$ (b) $(n-1)/2$ (c) $(n+1)/2$

Ans (d) $n(n+1)/2$

5) If the first term $a=1$ common ratio $r=1/2$ then sixth term of geometric sequence is

(a) 5 (b) 6 (c) 3

Ans (d) 4

Created by Amaan Khan

Q3 07

5. Sum of how many term of an arithmetic series will be zero provided that 1st term = 20, common difference = 20?

(a) 20 (b) 10 (c) 3

Ans (d) 2

7. The 9999th term of the sequence $(-1, 1, -1, 1, -1, 1 \ldots)$ is

(a) -1 (b) 9999

Ans (c) -9999 (d) 1

9. For a geometric progression if first term = 7 and common ratio = 3 then the 4th term is ...

(a) 121 (b) 144 (c) 189

Ans (d) 16

9. For an arithmetic series, if $a_1 = -3$, $d = 2$ and $S_n = 21$ then the number of term "n" in the series are

Ans (a) 7 (b) -7 (c) 10

Created by Amaan Khan

Quiz 02

10) The explicit formula when $a_1 = 6$ and $d = 2$ is ___

(a) $a_n = 6 + 2(n+1)$ (b) $a_n = 6 + 2n$

Ans (c) $a_n = 6 - 2(n-1)$ ✓ (d) $a_n = 6 + 2(n-1)$

11) For the sequence 5, 10, 20, 40 ... eight term will be ___

✓ (a) 640 (b) 19 (c) 70

Ans (d) 90

12) If first term of geometric sequence is 243 and sixth term is $\frac{1}{3}$, then the common ratio will be.

(a) 3 (b) 1 ✓ $\frac{1}{3}$

Ans (d) -1

13) For a sequence whose n^{th} term is $(-200/n)$ then its 200th term is

Ans (a) 200 (b) 1

✓ (c) -1 (d) -4000

Created by Amaan Khan

Quiz 02

14) If $C_n = n(n+1)$ for all integers $n \geq 1$ then which of the following is the first three term of a sequence.

(a) 0, 2, 6 ... (b) 2, 6, 12 ...

Ans (c) 1, 2, 3 ... (d) 1, 2, 6

15) Which of the following is an arithmetic sequence.

Ans (a) 1, 4, 7 ... (b) 1, 4, 9, ...

(c) 1/2, 1/4, 1/6 (d) 2, 4, 8, ...

16) Sum of first 888 term of sequence $(-1, 1, -1, 1 \ldots)$ is?

(a) 0 , (b) 1 (c) -1 , (d) Undefined

17) Sum of first 2000 term of the sequence $(-1, 1, -1, 1 \ldots)$ is

Ans (a) -1 (b) 1 (c) 0 (d) Undefined

Created by Amaan Khan

02

18) If the common ratio = 3, for the following geometric sequence $(x-1), (x-1)(x-1), (x-1)(x-1)(x-1)$ --- then the value of x is.

Ans (a) 3 (b) 4 (c) 2 (d) 1

19) For a geometric sequence if nth term = 32, first term = common ratio = 2 then the value of 'n' is —

Ans (a) 2 also! (b) 3 (c) 5

(d) 1

20) In first term = a = 2 common difference = d = 4 and term number = n = 10 then what is value of 10th term in arithmetic sequence?

Ans (a) 40 (b) 39 (c) 38

(d) 37

Created by Amaan Khan

Quiz 02

Q1/ For the sequence of natural numbers, its sum of 'n' natural numbers = summation $(n) = 21$ then value of 'n' is ---

Ans (a) -7 (b) -6 (c) 6 (d) 7

Quiz 02

22) The first term of sequence whose nth term = $2n > 2$ is ---

Ans (a) 2 (b) 4 (c) 6 (d) 1

23) Sum of the first 20 term of the following series: $1+2+3+4 \ldots$ is

Ans (a) 55 (b) 200 (c) 220 (d) 210

24) Sum of the geometric series: $1+(1/2)+(1/4)+(1/8)+\ldots$

Ans (a) 2 (b) 3/2 (c) 1 (d) 5/2

25) [illegible]

Created by Amaan Khan

Quiz 02

a) If the difference b/w successive term of a series is a constant, then the series is said to be a/an --- series?

Ans (a) Arithmetic (b) Geometric
(c) Harmonic (d) none

b) For a geometric sequence $(1, b, b, b, bbb \ldots)$ if its common ratio is 3 then value of b is

Ans (a) -3 (b) 3
(c) 1 or -1 (d) 1

c) Identify the sequence as Arithmetic, Geometric or neither
8, 10, 20, 40

Ans (a) Arithmetic (b) Geometric

Created by Amaan Khan

MTH 100 Quiz 03

1) $\sin^2\theta + \cos^2\theta$ = ----

(a) -1 (b) 0 (c) 1/2 (d) 1 ✓

2) $\cos(-\theta)$ = ----

(a) $\sec(\theta)$ ~~(b) $\cos(\theta)$~~ (b) $\cos\theta$

(c) $-\cos\theta$ (d) $-\cos(-\theta)$

3) $\sin(-\theta)$ = ----

(a) $\sin\theta$ (b) $-\sin\theta$ (c) $\cos\theta$

(d) $\cos(-\theta)$

4) The sum formula for $\tan(A+B)$ is -----

$$\frac{\tan A + \tan B}{1 - \tan A \tan B}$$

5) All the trigonometric function have positive value in ---- quadrant

(a) 1st (b) 2nd

Created by Amaan Khan

Quiz 03

6) The function $Y = \sin 4x$ has

period

(a) $\frac{\pi}{2}$ (b) $\frac{\pi}{3}$ (c) $\frac{\pi}{4}$ (d) ~~2π~~

7) The domain of inverse tangent function is - - - -

(-infinity, infinity)

8) If i adde) -5 to each value coordinate point (-23), what would be new coordinate?

(a) (7,2) (b) (-7,-2) (c) (3,2)

(d) (10, -15)

9) The smallest interval function repeat it self is called

(a) Slope (b) Period (c) Range

(d) Graph

10) The point where the x- and y- axis cross (0,0)

(a) intercept (b) Slop

Created by Amaan Khan

03

11) Using the distance formula find the distance b/w the two points A(1,2), B(4,6)

(a) -5 (b) 5

(c) 13 (d) 7

12) Smallest interval ~~function~~ for which the trigonometric function repeat it- self is called

(a) amplitude (b) Period

(c) angle (d) range

13) Power reducing formula $\cos^2 A$ is ---.

(a) $\frac{2 \tan A}{1 + \tan^2 A}$ (b) $1 - \cos^2 A$

(c) $\frac{\cos 2A + 1}{2}$ (d) $\frac{1 - \cos 2A}{2}$

14) Sin θ is positive in the first- and the --- quadrant

Second (Homework pg 37)

Created by Amaan Khan

33

15 ⇒ The ---- form of a trigonometric function is the height from the mean (or the rest) value of the trigonometric function to the maximum or minimum value

Amplitude (Howard pg 40)

16 ⇒ The difference formula for Sin(A-B) is

(Howard pg 48) Sin(A-B) = SinAcosB - CosASinB

17 ⇒ Equation that is not identity is called a(an) ---- equation

(a) Identical (b) linear
(c) Conditional (d) transcendental

18 ⇒ Cos θ is positive in the ---- and the third quadrant

First (Howard pg 38)

Created by Amaan Khan

Q no 3

19) The range of cosine function is b/w -1 and 1

(a) True (b) False

20) Cos θ is negative in the second and ---- quadrant

Fourth (Now see pg 38)

21) The amplitude of the function is the height from the rest value to the extreme value

(a) True (b) False

22) Equation of circle centered at (0, -5) and radius 5 is

(a) $S(x^2) \cdot (y+5)^2 = 8$ (b) $S\{x^2 + (y+5)^2\} = 64$

(c) $S\{x^2 + y + (y+5)^2\} = 645$ (d) $S(x^2) \cdot (y+5)^2$

Created by Amaan Khan

Qns 23

23

The center of circle having equation $(x-2)^2 - (y-5)^2 - 36 = 0$ is at-

(a) $(-2, -5)$ (b) $(2, -5)$

(c) $(-2, 5)$ (d) $(2, 5)$

24) Using the distance formula find the distance b/w the two point- A(4, 7), B(3, -2)

(a) 7 (b) 19

(c) $\sqrt{82}$ (d) 20

25) $\cos(180 - \theta) = \ldots$

$= -\cos\theta$ (Identity Pg 82)

26) $1 + \cot^2\theta = \ldots$

$\csc^2\theta$ (Identity Pg 92)

27) The range of $y = \arccos x$ is

$[-1, 1]$ $[0, \pi]$

Created by Amaan Khan

Quiz 03

28) The mid point of the line segment with the end point $(-4, 4), (-2, 2)$ is

(a) $(-6, 6)$ (b) $(1, 2)$

(c) $(-3, 3)$ (d) $(-2, -2)$

29) Equation of circle centered at $(0, -5)$ and radius $\sqrt{8}$ is

$$x^2 + (y+5)^2 = 8$$

30) Tangent function is periodic but its periodic is ___ degree

(a) 60 (b) 90

(c) 180 (d) 360

31) The domain of $y = \arcsin x$ is

(a) $[-1, 1]$ (b) $(-\pi, \pi)$

(c) $[-\infty, \infty]$ (d) $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right]$

Created by Amaan Khan

Quiz 03

2) $\cos 2A =$

(a) $2\cos^2 A - 1$ (b) $1 - \cos^2 A$

(c) $\cos^2 A + \sin^2 A$ (d) $2 \sin A \cos A$

3) The sin function is periodic with periodic 360 degree and range b/w - --

(a) 0 and -1 (b) 0 and 1

(c) -1 and 0.5 (d) -1 and 1

4) $\sin\left(-\frac{\pi}{4}\right) =$ - --

(a) $-\sin\left(\frac{\pi}{4}\right)$ (b) $\sin\left(\frac{\pi}{4}\right)$

(c) $\text{cosec}\left(\frac{\pi}{4}\right)$ (d) $-\sec\left(\frac{\pi}{4}\right)$

Created by Amaan Khan

05

MTH 100

18) For individual observation, reciprocal of arithmetic mean is called Harmonic mean

Created by Amaan Khan

(04) MTH 100

14) Budget of a household can be represented effectively through which of following chart.
(a) Bar (b) Frequency
(c) Pie (d) Histogram

15) if the mean, median mode of two data sets are equal then
(a) both data set should be equal
(b) both data set not necessarily equal
(c) both data set should not be equal
(d) none of them

16) when data is arranged, middle value in set of observation is identified as
(a) median (b) mean
(c) variance (d) standard deviation

17) median of the following data
4, 5, 6, 7, 8, 9, 10
(a) 6 (b) 6.5
(c) 7 (d) 7.5

Created by Amaan Khan

(3)

MTH 100

10) The cumulative frequencies are plotted against upper class boundaries of the corresponding class
(a) True (b) False

11) Arithmetic means 10 and number of observation are 20 then sum of all value is.
(a) ~~20~~ 20 (b) 100
(c) 200 (d) None of them

12) A quantative variable which can take any in a given range is called
(a) Simple (b) discrete
(c) Continous (d) none of them

13) If a variable cannot take numerical value then variable is called
(a) Constant (b) discreat
(c) Continous (d) Quantative

Created by Amaan Khan

2) MTH 100

4) Arithmetic mean 2, 4 and 6
(a) 3 (b) 4
(c) 5 (d) 6

5) The central tendency is measured by
(a) range (b) average
(c) dispersion (d) deviation

7) pieces of numerical and non-numerical information are called data.
(a) True (b) False

8) The mode of the list (1, 2, 2, 3, 3, 3, 4) is
(a) 2 (b) 2.5
(c) 3 (d) 4

9) Relative frequency calculated by dividing the frequency of each interval. (a) length of interval
(b) sum of all interval
(c) total number of observation
(d) None of them

Created by Amaan Khan

(1)

MTH 100 Quiz 04

1) Frequency Density =
(a) Cumulative Frequency / Relative
(b) Cumulative / Class width
(c) Frequency / Class width
(d) Class width / Frequency

2) Which of the following is not used to measure the central tendency.
(a) Arithmetic (b) Mode
(c) Median (d) Standard Deviation

3) A bar chart which represent continuous data is called
(a) Histogram (b) Pie chart
(c) Discrete chart (d) None of them

4) Method used to compute average of central value of collected data is consider.
(a) measure of central tendency
(b) measure of negative skewness
(c) negative variation
(d) Positive variation.

Created by Amaan Khan

Created by Amaan Khan


MTH100:Quiz # 2

Question # 8 of 10 (Start time: 04:07:09 PM, 23 January 2017)

$\cos(\theta - 180) =$ _____.

Select the correct option

- ○ $\sin\theta$
- ○ $-\sin\theta$
- ● $-\cos(\theta)$
- ○ $\cos(\theta)$



...aspx?ver=da847507-5c31-48f6-97c9-35a92b771211

MC160402583: Muhammad Muslim

MTH100:Quiz # 2

Question # 9 of 10 (Start time: 04:08:00 PM, 23 January 2017)


The amplitude of the function is the height from the rest value to the extreme value

Select the correct option

- ● True
- ○ False

Created by Amaan Khan

Created by Amaan Khan



Question # 8 of 10 (Start time: 03:57:19 PM, 23 January 2017)

The sine function is periodic, with period 360 degrees, and range between _____.

Select the correct option

- 0 and -1
- 0 and 1
- -1 and 0.5
- -1 and 1

Question # 5 of 10 (Start time: 08:27:19 PM, 20 January 2017) Total Marks: 1

In the cosine function, the maximum value of _____ is taken when $\theta = 0^\circ$, $\pm 360^\circ$, $\pm 720^\circ$ and so on.

Select the correct option

Reload Math Equations

- -1
- 0
- 0.5
- 1

Facebook

Click to Save Answer & Move to Next Question

Created by Amaan Khan

Created by Amaan Khan

Question # 9 of 10 (Start time: 03:57:53 PM, 23 January 2017)

$\tan\theta =$

Select the correct option

- ○ $\frac{\cos\theta}{\sin\theta}$
- ● $\frac{\sin\theta}{\cos\theta}$
- ○ $\frac{\cot\theta}{\sin\theta}$
- ○ $\frac{\cot\theta}{\cos\theta}$

MC160402583: Muhammad Muslim

MTH100:Quiz # 2

Question # 7 of 10 (Start time: 04:06:15 PM, 23 January 2017)


$sin\theta$ is positive in the first and the _____ quadrant.

Select the correct option

- ○ first
- ● second
- ○ third
- ○ fourth

Created by Amaan Khan


Created by Amaan Khan





Created by Amaan Khan

Created by Amaan Khan



MTH100:Quiz # 2

Question # 9 of 10 (Start time: 07:37:59 PM, 19 January 2017)

$\tan 2A =$ _____.


Select the correct option

- $\frac{\tan A + \tan B}{1 - \tan A \tan B}$
- $\frac{2\tan A}{1 - \tan^2 A}$
- $\frac{2\tan A}{1 + \tan^2 A}$
- $\frac{\sin^2 A}{\cos^2 A}$



Created by Amaan Khan

Created by Amaan Khan



MC160402583: Muhammad Muslim

MTH100:Quiz # 2

Question # 6 of 10 (Start time: 04:05:26 PM, 23 January 2017)

$\tan 2A = ____.$

Select the correct option

- ○ $\frac{\tan A + \tan B}{1 - \tan A \tan B}$
- ● $\frac{2\tan A}{1 - \tan^2 A}$
- ○ $\frac{2\tan A}{1 + \tan^2 A}$
- ○ $\frac{\sin^2 A}{\cos^2 A}$

MTH100:Quiz # 2 — Quiz Start Time: 08:24 PM, 20 January 2017

Question # 3 of 10 (Start time: 08:25:55 PM, 20 January 2017) — Total Marks: 1

$\sin 2A = ____.$

Select the correct option


Reload Math Equations

- ○ $1 - 2\sin^2 A$
- ○ $\cos^2 A - \sin^2 A$
- ○ $\cos^2 A + \sin^2 A$
- ● $2\sin A \cos A$

Click to Save Answer & Move to Next Question

Created by Amaan Khan

Created by Amaan Khan



MC160401314: Qamar Zaman

MTH100:Quiz # 2

Question # 2 of 10 (Start time: 07:32:42 PM, 19 January 2017)

$\cos 2A =$ _____.


Select the correct option

●	$2\cos^2 A - 1$
○	$1 - \cos^2 A$
○	$\cos^2 A + \sin^2 A$
○	$2 \sin A \ \cos A$




Created by Amaan Khan

Created by Amaan Khan



Created by Amaan Khan

Created by Amaan Khan



Created by Amaan Khan

Created by Amaan Khan

Question # 8 of 10 (Start time: 08:29:26 PM, 20 January 2017) Total Marks: 1


$\tan\theta =$

Select the correct option

Reload Math Equations

- $\frac{\cos\theta}{\sin\theta}$
- $\frac{\sin\theta}{\cos\theta}$
- $\frac{\cot\theta}{\sin\theta}$
- $\frac{\cot\theta}{\cos\theta}$

Click to Save Answer & Move to Next Question



Question # 1 of 10 (Start time: 08:24:34 PM, 20 January 2017) Total Marks: 1

The domain of *[Math Processing Error]* is .....

Select the correct option

Reload Math Equations


- [-1,1]
- *[Math Processing Error]*
- *[Math Processing Error]*



Click to Save Answer & Move to Next Question

Created by Amaan Khan

Created by Amaan Khan



MC160401314: Qamar Zaman

MTH100:Quiz # 2

Question # 7 of 10 (Start time: 07:36:20 PM, 19 January 2017)


The domain of $y = \arccos x$ is.....

Select the correct option

- [x] $[-1, 1]$
- [ ] $[0, 1]$
- [ ] $[0, \infty]$
- [ ] $[-\infty, 0]$

Created by Amaan Khan

Created by Amaan Khan



Question # 7 of 10 (Start time: 08:28:32 PM, 20 January 2017) Total Marks: 1

The trigonometric functions with the property that they keep repeating themselves are called _____

Select the correct option

- sine functions
- cosine functions
- periodic functions
- tangent functions

Click to Save Answer & Move to Next Question

Created by Amaan Khan

Created by Amaan Khan

MTH100:Quiz # 2

Question # 10 of 10 (Start time: 04:09:12 PM, 23 January 2017)

The midpoint of the line segment with the endpoints (−1, 1), (7, −5) is

Select the correct option

- ○ (3,2)
- ○ (-4,2)
- ● (3,-2)
- ○ (4,-2)

Muhammad Muslim

MTH100:Quiz # 2

Question # 4 of 10 (Start time: 04:03:29 PM, 23 January 2017)


The midpoint of the line segment with the endpoints (−4, 4), (−2, 2) is

Select the correct option

- ○ (-6, 6)
- ○ (2,2)
- ● (-3,3)
- ○ (-2,-2)

Created by Amaan Khan

Created by Amaan Khan



MC160401314: Qamar Zaman

MTH100:Quiz # 2

Question # 8 of 10 (Start time: 07:37:20 PM, 19 January 2017)

The difference formula for $sin (A - B)$ is _____.

Select the correct option

○	\[\cos A\,\cos B + \sin A\,\sin B\]
●	\[\sin A\,\cos B - \cos A\,\sin B\]
○	\[\sin A\,\cos B + \cos A\,\sin B\]
○	\[\cos A\,\cos B - \sin A\,\sin B\]

Created by Amaan Khan

Created by Amaan Khan

Question # 10 of 10 (Start time: 03:58:21 PM, 23 January 2017)

The range of the cosine function is between _____.

Select the correct option

- 0 and -1
- 0 and 1
- -1 and 0.5
- -1 and 1

MTH100:Quiz # 2

Question # 2 of 10 (Start time: 04:01:20 PM, 23 January 2017)


The domain of $y = \arccos x$ is.....

Select the correct option

- $[-1, 1]$
- $[0, 1]$
- $[0, \infty]$
- $[-\infty, 0]$

Created by Amaan Khan

Created by Amaan Khan



MC160401314: Qamar Zaman

MTH100:Quiz # 2

Question # 5 of 10 (Start time: 07:34:28 PM, 19 January 2017)

The cosine function has "period" 360° as it repeats itself after each revolution of _____.

Select the correct option

○	60°
○	90°
○	180°
●	360°

Created by Amaan Khan

Created by Amaan Khan

Question # 10 of 10 (Start time: 08:30:40 PM, 20 January 2017) Total Marks: 1

$\sin(-\theta) = ____.$

Select the correct option

Reload Math Equations

- ○ $\sin\theta$
- ● $-\sin\theta$
- ○ $\cos\theta$
- ○ $\cos(-\theta)$

Click to Save Answer & Move to Next Question

Question # 9 of 10 (Start time: 08:30:10 PM, 20 January 2017) Total Marks: 1

The domain of inverse sine function and inverse cosine function is same.


Select the correct option

- ● True
- ○ False

Click to Save Answer & Move to Next Question


Created by Amaan Khan

Created by Amaan Khan



Created by Amaan Khan

Created by Amaan Khan



MC160401314: Qamar Zaman

MTH100:Quiz # 2

Question # 3 of 10 (Start time: 07:33:14 PM, 19 January 2017)

What is the slope of the line that passes through (8, 4) and (6, 7)?


Select the correct option

- 2/3
- 3/2
- -2/3
- -3/2




Created by Amaan Khan

Created by Amaan Khan



Created by Amaan Khan

Created by Amaan Khan



MC160401314: Qamar Zaman

MTH100:Quiz # 2

Question # 1 of 10 (Start time: 07:32:18 PM, 19 January 2017)


$\tan\theta =$

Select the correct option

- ○ $\frac{\cos\theta}{\sin\theta}$
- ● $\frac{\sin\theta}{\cos\theta}$
- ○ $\frac{\cot\theta}{\sin\theta}$
- ○ $\frac{\cot\theta}{\cos\theta}$

Created by Amaan Khan


Created by Amaan Khan



Created by Amaan Khan

Created by Amaan Khan





Created by Amaan Khan