Description: When a carboxylic acid is treated with an alcohol and an acid catalyst, an ester is formed (along with

Scanned with CamScanner

- This flow of phosphoryl groups is catalyzed by kinases, and proceeds with an overall loss of free energy under intracellular conditions.

Scanned with CamScanner

Question # 8 of 10 (Start time: 10:42:46 AM, 30 July 2021)
Total Marks: 1

Pyruvate is oxidized, with loss of its carboxyl group to yield _____.

Select the correct option

o	Acetyl-coenzyme A correct
o	Starch
o	Malate
o	Succinyl CoA

Scanned with CamScanner

Question # 6 of 10 (Start time: 10:40:42 AM, 30 July 2021)
Total Marks: 1

The process in which the enolate of an aldehyde or a ketone reacts at the α-carbon with the carbonyl of another molecule to obtain β-hydroxy aldehyde or β-hydroxy ketone, is called as _____.

Select the correct option

O	Decarboxylation
O	Hydrolysis
O	Aldol condensation correct
O	Claisen condensation

Scanned with CamScanner

Question # 5 of 10 (Start time: 10:38:58 AM, 30 July 2021) Total Marks: 1

The payoff phase of glycolysis includes the energy-conserving steps in which _____ takes place.

Select the correct option

O	Methylation
O	Nitration
O	Phosphorylation
O	All of given

correct

Click to Save Answer & Move to Next Question

Scanned with CamScanner

BIO303:Quiz

Question # 4 of 10 (Start time: 10:38:24 AM, 30 July 2021)

During fermentsation, material rich in _____ is supplied to a pure culture of a specific microorganism

Select the correct option

- ○ Carbohydrates

Correct

- ○ Lipids
- ○ Nucleic Acids
- ○ Proteins

Scanned with CamScanner

Question # 3 of 10 (Start time: 10:37:41 AM, 30 July 2021) Total Marks:

The reduction of pyruvate to lactate is catalyzed by lactate_____, which forms lactate at pH 7.

Select the correct option

- Phosphatase
- Decarboxylation
- Dehydrogenase
- Kinase

correct

Scanned with CamScanner

Question # 7 of 10 (Start time: 10:41:33 AM, 30 July 2021)
Total Marks: 1

The loss of ____ from an alcohol results in the introduction of a C=C bond between two adjacent carbon atoms.

Select the correct option

o	H2O
o	OH-
o	H+
o	ATP

correct

Scanned with CamScanner

1. 'High-energy' compounds have a $\Delta G'^{0}$ of hydrolysis more negative than -25 kj/mol

Scanned with CamScanner