Question No : 1 of 34

The graph of an odd function is symmetrical about ----------

Answer (Please select your correct option)

- x-axis
- y-axis
- origin
- None of these

Start Time: 9:41 AM

87:00
Time Left

Question Summary : (Attempted Question)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17

Conduct Instructions

Done

Start | VU Examination Syste...

Click here to FINIS

Address http://server/vutes/client/Paper.aspx

MTH633 Group Theory

Question No : 26 of 41

If H has 6 elements, then all cosets of H must have _____ elements.

Answer (Please select your correct option)

- 3
- 9
- 12
- 6

Start Time 8:05 AM

110:00 Time Left

Question Summary : (Attempted Question)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21

Conduct Instructions

Done

Start | VU Examination Syste... | MathType (Lite mode) - ...

MTH631 Real Analysis II

Question No : 3 of 34

_____ is the only region in R

Answer (Please select your correct option)

- Set of finite points of R
- Interval
- Any finite sequence of R
- None of these

Start Time 9:41 AM

84:00
Time Left

Question Summary : (Attempted Question)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18

Conduct Instructions

Done

Start | VU Examination Syste...

Click here to FINISH

MTH631 Real Analysis II

Question No : 10 of 34

The rate of change of $f(x,y)$ in the direction of a unit vector is called the ---------- of $f(x,y)$.

Answer (Please select your correct option)

- differential
- directional derivative
- mixed derivative
- None of these

Start Time: 9:41 AM

74:00
Time Left

10

Question Summary : (Attempted Question)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24

Conduct Instructions

Click here to FINISH Your Exam

Done

Start | VU Examination Syste...

Question No : 11 of 34 Marks:

A function of several variables may have first order partial derivatives at a point X_0 but ---------- continuous at X_0.

Answer (Please select your correct option)

- fail to be
- must be
- always not
- None of these

Start Time: 9:41 AM

74:00
Time Left

11

Question Summary : (Attempted Question)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26

Conduct Instructions

Click here to FINISH Your Exam

Done

Start | VU Examination Syste...



Question No : 12 of 34

A function of the form $L(X) = f_{x_1}(X_0)x_1 + f_{x_2}(X_0)x_2 + \ldots + f_{x_n}(X_0)x_n$ is called ———

Answer (Please select your correct option)

- quadractic function
- differential of f at X_0
- constant function
- None of these

Start Time: 9:41 AM

74:00 Time Left

12

Question Summary : (Attempted Question)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22

Conduct Instructions

Click here to FINISH Your Exam

Done

Start | VU Examination Syste...

If $h = f(x,y), x = f(r,\theta)$ and $y = f(r,\theta)$, then using the chain rule, $\frac{\partial h}{\partial \theta}$ = ...

Answer (Please select your correct option)

- C $\frac{\partial f}{\partial r}\frac{\partial r}{\partial \theta} + \frac{\partial f}{\partial r}\frac{\partial r}{\partial \theta}$
- C $\frac{\partial f}{\partial x}\frac{\partial x}{\partial \theta} + \frac{\partial f}{\partial y}\frac{\partial y}{\partial \theta}$
- C $\frac{\partial f}{\partial x} + \frac{\partial x}{\partial \theta} + \frac{\partial f}{\partial y} + \frac{\partial y}{\partial \theta}$
- C None of these

Start Time: 9:41 AM

74:00
Time Left

13

Question Summary : (Attempted Question)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22

Conduct Instructions

Done

Click here to FINISH Your Exam

Start | VU Examination Syste...

HP LI750

Question No : 14 of 34

If $h = f(x,y), x = r\cos\theta$ and $y = r\sin\theta$, where then using the chain rule, $\frac{\partial h}{\partial r}$ = ...

Answer (Please select your correct option)

- $\cos\theta\frac{\partial f}{\partial \theta} + \sin\theta\frac{\partial f}{\partial r}$
- $\cos\theta\frac{\partial f}{\partial x} + \sin\theta\frac{\partial f}{\partial y}$
- $\cos\theta\frac{\partial f}{\partial x} - \sin\theta\frac{\partial f}{\partial y}$
- None of these

Start Time: 9:41 AM

74:00 Time Left

14

Question Summary : (Attempted Question)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24

Conduct Instructions

Click here to FINISH Your Exam

Done

Start | VU Examination Syste...

Address http://server/vutes/client/login.aspx

MTH631 Real Analysis II

Question No : 9 of 34

A point on the graph of a function at which the tangent line is parallel to x-axis is called the .....

Answer (Please select your correct option)

- critical point
- point of inflection
- point of relative extrema
- None of these

Start Time 9:41 AM

74:00 Time Left

Question Summary : (Attempted Question)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23

Conduct Instruction

Click here to FINISH Your Exam

Done

Start | VU Examination Syste...

Address http://server/vutes/client/login.aspx

MTH631 Real Analysis II

Question No : 5 of 34 Mark

If f and g are continuous functions at the point X_0 in R^n, then $\frac{f}{g}$ is continuous at X_0 provided that _____

Answer (Please select your correct option)

- $g(X_0) = 0$
- $g(X_0) \neq 0$
- $f(X_0) = g(X_0) = 0$
- None of these

Start Time: 9:41 AM

78:00 Time Left

Question Summary : (Attempted Question)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25

Conduct Instructions

Click here to FINISH Your Exam

Done

Start VU Examination Syste...

Question No : 3 of 50

_____ is the only region in R.

Answer (Please select your correct option)

- Set of finite points of R
- Interval
- Any finite sequence of R
- None of these

Start Time: 9:41 AM

81:00 Time Left

3

Question Summary : (Attempted Question)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18

Conduct Instructions

Done

Start | VU Examination Syste...

Click here to FINISH Your

Question No : 4 of 34

For the function $f(x,y)=\frac{xy}{x^2+y^2}$, the limit of $f(x,y)$ as $(x,y)\to(0,0)$ along the line $y=x$ is ----

Answer (Please select your correct option)

- $\frac{1}{2}$
- $-\frac{1}{2}$
- 0
- 1

Start Time: 9:41 AM

80:00 Time Left

4

Question Summary : (Attempted Question)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22

Conduct Instructions

Done

Click here to FINISH Your Exam

Start | VU Examination Syste...

Address http://server/vutes/client/login.aspx

MTH631 Real Analysis II

Question No : 2 of 34

If H is an open covering of a compact subset S then, S can be covered by ----------

Answer (Please select your correct option)

- infinitely many sets from H
- finitely many sets from H
- atleast one set from H
- None of these

Start Time: 9:41 AM

83:00 Time Left

2

Question Summary : (Attempted Question)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19

Conduct Instructions

Done

Click here to **FINISH** Your Exam

Start VU Examination Syste...

Address http://server/vutes/client/login.aspx

MTH631 Real Analysis II

Question No : 15 of 34

If $p(X)$ is a homogeneous polynomial of degree r in $X - X_0$. If $p(X) \leq 0$ for all $X \neq X_0$. Then p is called _____

Answer (Please select your correct option)

- positive semidefinite
- positive definite
- negative semidefinite
- negative definite

Start Time: 9:41 AM

71:00 Time Left

15

Question Summary : (Attempted Question)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25

Conduct Instructions

Click here to FINISH Your Exam

Done

Start | VU Examination Syste...

Question No : 12 of 34

A function of the form $L(X) = f_{x_1}(X_0)x_1 + f_{x_2}(X_0)x_2 + \ldots + f_{x_n}(X_0)x_n$ is called ———

Answer (Please select your correct option)

- quadractic function
- differential of f at X_0
- constant function
- None of these

Start Time: 9:41 AM

74:00 Time Left

12

Question Summary : (Attempted Question)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22

Conduct Instructions

Click here to FINISH Your Exam

Done

Start | VU Examination Syste...

Address http://server/vutes/client/login.aspx

MTH631 Real Analysis II

Question No : 15 of 34

If $p(X)$ is a homogeneous polynomial of degree r in $X - X_0$. If $p(X) \leq 0$ for all $X \neq X_0$. Then p is called _____

Answer (Please select your correct option)

- positive semidefinite
- positive definite
- negative semidefinite
- negative definite

Start Time: 9:41 AM

71:00 Time Left

15

Question Summary : (Attempted Question)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25

Conduct Instructions

Click here to FINISH Your Exam

Done

Start | VU Examination Syste...

If $h = f(x,y), x = f(r,\theta)$ and $y = f(r,\theta)$, then using the chain rule, $\frac{\partial h}{\partial \theta}$ = ...

Answer (Please select your correct option)

- C $\frac{\partial f}{\partial r}\frac{\partial r}{\partial \theta} + \frac{\partial f}{\partial r}\frac{\partial r}{\partial \theta}$
- C $\frac{\partial f}{\partial x}\frac{\partial x}{\partial \theta} + \frac{\partial f}{\partial y}\frac{\partial y}{\partial \theta}$
- C $\frac{\partial f}{\partial x} + \frac{\partial x}{\partial \theta} + \frac{\partial f}{\partial y} + \frac{\partial y}{\partial \theta}$
- C None of these

Start Time: 9:41 AM

74:00
Time Left

13

Question Summary : (Attempted Question)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22

Conduct Instructions

Done

Click here to FINISH Your Exam

Start | VU Examination Syste...

HP LI750

Question No : 14 of 34

If $h = f(x,y), x = r\cos\theta$ and $y = r\sin\theta$, where then using the chain rule, $\frac{\partial h}{\partial r}$ = ...

Answer (Please select your correct option)

- $\cos\theta\frac{\partial f}{\partial \theta} + \sin\theta\frac{\partial f}{\partial r}$
- $\cos\theta\frac{\partial f}{\partial x} + \sin\theta\frac{\partial f}{\partial y}$
- $\cos\theta\frac{\partial f}{\partial x} - \sin\theta\frac{\partial f}{\partial y}$
- None of these

Start Time: 9:41 AM

74:00 Time Left

14

Question Summary : (Attempted Question)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24

Conduct Instructions

Click here to FINISH Your Exam

Done

Start | VU Examination Syste...

Address http://server/vutes/client/login.aspx

MTH631 Real Analysis II

Question No : 18 of 34 **Marks: 1**

A function f is said to be absolutely integrable on $[a,b)$ if f is locally integrable on $[a,b)$ and ----------

Answer (Please select your correct option)

- $\int_a^b |f(x)|dx < \infty$
- $\int_a^b |f(x)|dx = \infty$
- $\int_a^b |f(x)|dx \geq \infty$
- None of these

Start Time: 9:41 AM

65:00 Time Left

18

Question Summary : (Attempted Question)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26

Conduct Instructions

Click here to **FINISH** Your Exam

Done

Start VU Examination Syste...

Address http://server/vutes/client/login.aspx

MTH631 Real Analysis II

Question No : 18 of 34 **Marks: 1**

A function f is said to be absolutely integrable on $[a,b)$ if f is locally integrable on $[a,b)$ and ----------

Answer (Please select your correct option)

- $\int_a^b |f(x)|dx < \infty$
- $\int_a^b |f(x)|dx = \infty$
- $\int_a^b |f(x)|dx \geq \infty$
- None of these

Start Time: 9:41 AM

65:00 Time Left

18

Question Summary : (Attempted Question)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26

Conduct Instructions

Click here to **FINISH** Your Exam

Done

Start VU Examination Syste...

MTH631 Real Analysis II

Question No : 19 of 34

An absolutely convergent integral is always ----------

Answer (Please select your correct option)

- divergent
- convergent
- conditionally convergent
- None of these

Start Time: 9:41 AM

62:00
Time Left

19

Question Summary : (Attempted Question)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22

Conduct Instructions

Click here to FINISH Your Exam

Done

Start | VU Examination Syste...

Address http://server/vutes/client/login.aspx

MTH631 Real Analysis II

Question No : 20 of 34

If f is locally integrable on $[a,b)$ and $\int_a^b |f(x)|dx < \infty$, then $\int_a^b f(x)dx$ is _____

Answer (Please select your correct option)

- divergent
- conditionally convergent
- convergent
- None of these

Start Time: 9:41 AM

62:00 Time Left

20

Question Summary : (Attempted Question)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23

Conduct Instructions

Done

Click here to FINISH Your Exam

Start | VU Examination Syste...

http://server/vutes/client/login.aspx

MTH631 Real Analysis II

Question No : 34 of 34 **Marks: 10 (Budgeted Time Min)**

Suppose $U_0 \in T$ is a limit point of T, g is continuous at U_0 and f is continuous at $X_0 = G(U_0)$. Then, show that $f \circ g$ is continuous at U_0.

Answer (Please click here to Add Answer)

Start Time: 9:41 AM
62:00
Time Left
Question Summary : (Attempted Question)
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30
Conduct Instructions
Click here to FINISH Your Exam
Done
Start
VU Examination Syste...

Address http://server/vutes/client/login.aspx

MTH631 Real Analysis II

Question No : 33 of 34 **Marks: 5 (Budgeted Time Mi**

Show that $(0,0)$ is a critical point of the function $p(x,y) = x^2 - 2xy + y^2 - x^4 - y^4$ and it has positive semidefinite second differentials at $(0,0)$

Answer (Please click here to Add Answer)

100%

Start Time: 9:41 AM

62:00

Time Left

33

Question Summary : (Attempted Question)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30

Conduct Instructions

Click here to FINISH Your Exam

Done

Start VU Examination Syste...

File Edit View Tools Help

Address http://server/vutes/client/login.aspx

MTH631 Real Analysis II

Question No : 32 of 34 **Marks:**

Define a locally integrable function. Also show that the function $f(x) = \log x$ is locally integrable on $[0, 1)$.

Answer (Please click here to Add Answer)

Normal Arial 12 100%

Start Time 9:41 AM

61:00 Time Left

32

Question Summary : (Attempted Question)

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25

Conduct Instructions

Click here to **FINISH** Your Exam

Done

Start VU Examination Syste...

MTH631 FINAL PAPER


7:46
33%
Details
MTH631
Question No : 15 of 23
The values of x in a specific interval for which a given power series converges is called
Answers : Please select your correct option :
interval of continuity
radius of convergence
interval of convergence
None of these
50:00


33%
Details
MTH631
page 46
Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536
49:00


7:46
33%
Details
MTH631
page 75
49:00


Details
page 48
49:00


33%
Details
MTH9631
Question No : 12 of 23
For the series $f(x) = \sum_{n=0}^{\infty} (-1)^n x^n = \frac{1}{1+x}$, the $\lim_{x \to 1^-} f(x) =$
Answer: (Please select your correct option)
$\frac{1}{2}$
$-\frac{1}{2}$
$\pm\frac{1}{2}$
None of these
49:00


7:46
33%
Details
page 50
Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536
50:00


page 27
Mubashar Ali
Diplomaticsmile13@gamil.com
Whatsup num 0314-7654536
51:00


7:46
33%
Details
MTH631 Real Analysis II
Question No : 13 of 23
In the definition of uniform convergence the natural number N always depends on
Answer (Please select your correct option)
Mubashar Ali
Diplomaticsmile13@gamil.com
Whatsup num 0314-7654536
51:00


7:46
33%
Details
page 30
Mubashar Ali
Diplomaticsmile13@gamil.com
Whatsup num 0314-7654536
51:00


Details
MTH631
Question No : 11 of 23
The limit of a uniformly convergent series of integrable functions is always
page 25
Answer (Please select your correct option)
integrable
not integrable
differentiable
None of these
Mubashar Ali
Diplomaticsmile13@gamil.com
Whatsup num 0314-7654536
51:00


7:46
33%
Details
MTH631 Real Analysis II
Question No : 10 of 23
If a series converges absolutely uniformly on S, then
page 20
Answers (Please select your correct option)
it converges uniformly on S
it diverges
it is differentiable on S
None of these
Mubashar Ali
Diplomaticsmile13@gamil.com
Whatsup num 0314 7654536
52:00
Time Left


7:46
33%
MTH631 Real Analysis II
Details
Question No : 9 of 23
Which of the following is true about the expression $\int_a^b f(x)dx = \lim_{n\to\infty} \int_a^b f_n(x)dx$?
page 11
Answer: (Please select your correct option)
It is always true
It is not true generally
It is always false
None of these
Mubashar Ali
Diplomaticsmile13@gamil.com
Whatsup num 0314 7654536
54:00
Time Left


7:46
33%
Details
MTH631 Real Analysis II
Question No : 8 of 23
Suppose that a sequence of function (f_n) is pointwise convergent on the set S. Then
Answer (Please select your correct option)
(f_n) always converges uniformly
(f_n) may converge uniformly
(f_n) always diverges
None of these
Mubashar Ali
Diplomaticsmile13@gamil.com
Whatsup num 0314-7654536
55:00


7:46
33%
Details
MTH6311 Real Analysis II
Question No : 7 of 23
Which of the following is true about a uniformly continuous function, f, on [a, b]?
Answer : (Please select your correct option)
The function is bounded.
The function achieves its maximum on the set [a, b].
The derivative of f is bounded.
None of these
55:00
Time Left


7:46
Details
Question No : 6 of 23
Answer: (Please select your correct option)
has no conclusion
None of these
diverges
converges
Mubashar Ali
Diplomaticsmile13@gamil.com
Whatsup num 0314-7654536
56:00
Time Left


7:46
33%
MTH631 Real Analysis II
Details
Question No : 5 of 23
Which property is not satisfied by norm defined on a set S.
page 5
Answers: (Please select your correct option)
None of these
Mubashar Ali
Diplomaticsmile13@gamil.com
Whatsup num 0314-7654536
57:00


7:45
33%
Details
MTH631 Real Analysis II
Question No : 4 of 21
page 3
Answer : (Please select your correct option)
Mubashar Ali
Diplomaticsmile13@gamil.com
Whatsup num 0314-7654536
57:00


7:45
33%
Details
MTH631 Real Analysis II
Question No : 3 of 23
By Stone and Weierstrass theorem, the trigonometric polynomials are dense in C[a,b] provide
page 56 handout
Answer (Please select your correct option)
Mubashar Ali
Diplomaticsmile13@gamil.com
Whatsup num 0314-7654536
58:00
Time Left


7:45
33%
Details
MTH631 Real Analysis II
Question No : 2 of 23
sin(cx) is a periodic function having period equal to
page 54
Answer (Please select your correct option)
π/c
2c/π
2π/c
c/π
Start Time: 10:26 AM
58:00
Time Left

2019


Press Esc to exit full screen
Question No : 1 of 23
An infinite series of the form $\sum \frac{f^n(x_0)}{n!}(x-x_0)^n$ is called ... of f about
page 36
Answer (Please select your correct option)
Power series
Maclaurin series
Taylor series
None of these
Mubashar Ali
Diplomaticsmile13@gamil.com
Whatsup num 0314-7654536
58:00


MTH631 Real Analysis II
Question No : 20 of 25
If f and g are integrable on S and $f(X) \leq g(X)$ for X in S, then ———
page 163
Answer (Please select your correct option)
$\int_a^b f(X)dX \geq \int_a^b g(X)dX$
$\int_a^b f(X)dX \leq \int_a^b g(X)dX$
$\int_a^b f(X)dX > \int_a^b g(X)dX$
None of these
Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536
Start Time: 11:05 AM
114:00

MTH631 Real Analysis II

Question No : 19 of 76

The lower integral $\underline{\int_a^b} f(x)d(x)$ of f over R is the supremum of all the ———.

page 155

Answer (Please select your correct option)

- lower sums
- upper sums
- average sums
- None of these

Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536

Start Time: 11:05 AM

114:00

19

MTH631 Real Analysis II

Question No : 18 of 26

The upper integral $\overline{\int_R} f(X)d(X)$ of f over R is the infimum of all the_____

page 135

Answer (Please select your correct option)

- lower sums
- upper sums
- average sums
- None of these

Mubashar Ali
[redacted]
Whatsup num 0314-7654536

MTH631 Real Analysis II

Question No : 17 of 26

A function f is said to be absolutely integrable on $[a,b)$ if f is locally integrable on $[a,b)$ and ———

page 145

Answer (Please select your correct option)

- $\int_a^b |f(x)|dx < \infty$
- $\int_a^b |f(x)|dx = \infty$
- $\int_a^b |f(x)|dx \geq \infty$
- None of these

Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536

Start Time: 11:05 AM

MTH631 Real Analysis II

Question No : 16 of 25

If f and g are locally integrable on $[a,b)$, $g(x) > 0$ and $f(x) \geq 0$ on some subintervals $[a_1,b)$ of $[a,b)$, and $\lim_{x \to b^-} \frac{f(x)}{g(x)} = M$. If $M = 0$ and $\int_a^b g(x)dx < \infty$, then

page 144

Answer (Please select your correct option)

- $\int_a^b f(x)dx$ and $\int_a^b g(x)dx$ converges together
- $\int_a^b f(x)dx = \infty$
- $\int_a^b f(x)dx < \infty$
- $\int_a^b f(x)dx \geq \infty$

Start Time: 11:05 AM

115:00

Time Left

16

Question Summary : (Attempted Question)


MTH631 Real Analysis II
Question No : 15 of 26
The function $f(x) = (1-x)^{-p}$ is locally integrable on _____
page 139
Answer (Please select your correct option)
[0,1]
[-1,1)
(0,1)
[0,1)
Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536
Start Time 11:05 AM
15

MTH631 Real Analysis II

Question No : 11 of 26

If $p(X)$ is a homogeneous polynomial of degree r in $X - X_0$ If $p(X) \leq 0$ for all $X \neq X_0$ Then p is called _____

page 134

Answer (Please select your correct option)

- positive semidefinite
- positive definite
- negative semidefinite
- negative definite

Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536

MTH631 Real Analysis II

Question No : 13 of 26

If value of a becomes zero in the Taylor series $\sum_{n=0}^{\infty} \frac{f^n(a)}{n!}(x-a)^n$, then the series is called ———

page 44

Answer (Please select your correct option)

- Maclaurin series
- Fourier series
- Power series
- None of these

Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536


MTH631 Real Analysis II
Question No : 32 of 36
If $h = f(r, \theta)$, then using the chain rule, $\frac{\partial h}{\partial r} =$ _____
Answer (Please select your correct option)
$\frac{\partial f}{\partial x}\frac{\partial x}{\partial r} + \frac{\partial f}{\partial y}\frac{\partial y}{\partial r}$
$\frac{\partial f}{\partial x} + \frac{\partial x}{\partial r} + \frac{\partial f}{\partial y} + \frac{\partial y}{\partial r}$
$\frac{\partial f}{\partial x}\frac{\partial x}{\partial r} + \frac{\partial f}{\partial y}\frac{\partial y}{\partial r}$
None of these
116:00
Time Left
Question Summary :
Attempted Question
1
2
3
4
5
6
7
8

MTH631 Real Analysis II

Question No : 11 of 26

If $f(x) = 3x + 5x^3$, then at $x = -1$, $df =$ _____

Answer (Please select your correct option)

- $18dy$
- $18dx$
- 18

Address http://server/vutms/client/Paper.aspx

MTH631 Real Analysis II

Question No : 19 of 26

A function of the form $L(X) = m_1 x_1 + m_2 x_2 + \ldots + m_n x_n$ for some constants m_i's is called ———

page 112

Answer (Please select your correct option)

- linear function
- quadratic function
- constant function
- None of these

Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536


http://server/vutes/client/Paper.aspx
MTH631 Real Analysis II
Question No : 9 of 26
For the function $f(x,y) = 3x^2y^3 + xy$ in two variables, $f_{xy}(0,0) =$ .....
Answer (Please select your correct option)
1
-1
0
None of these
Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536


http://server/vutes/client/Paper.aspx
MTH631 Real Analysis II
Question No : 8 of 26
A critical point at which a function attains its minimum value among all points where it is defined is called
Answer (Please select your correct option)
global maximum
global minimum
supremum
None of these
Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536

MTH631 Real Analysis II

Question No : 7 of 26

A $f_1, f_2, \ldots f_n$ exist on a neighborhood of X_0 and are continuous at X_0, then f is ——— at X_0.

page 115

Answer (Please select your correct option)

- uniformly converget
- piecewise convergent
- differentiable
- None of these

Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536

Start Time: 11:05 AM

MTH631 Real Analysis II

Question No : 6 of 26

A set A ⊆ R of real numbers is ———— if there exists a real number M ∈ R, such that $x \leq M$ for every $x \in A$.

Answer (Please select your correct option)

- bounded below
- uniformly continuous
- bounded above
- None of these

Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536

Start Time: 11:05 AM

MTH631 Real Analysis II

Question No : 5 of 26

In a vector valued function $G=(g_1,g_2,\ldots,g_n)$. Then $g_1,g_2,\ldots,g_n$ are called the ———— functions of G.

page 114

Answer (Please select your correct option)

- bijective
- surjective
- component
- None of these

Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536

Start Time: 11:05 AM


Address http://server/vutes/client/Paper.aspx
MTH631 Real Analysis II
Question No : 4 of 26
The quotient of two continuous functions is a ———— function wherever the denominator is non-zero.
Answer (Please select your correct option)
differentiable
continuous
composite
None of these
Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536

MTH631 Real Analysis II

Question No : 2 of 26

Any polygonally connected set either it is open or not, is always _____

page 82

Answer (Please select your correct option)

- connected
- disconnected
- a region

Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536

MTH631 Real Analysis II

Question No : 3 of 26

The set $S = \{(x,y), x^2 + y^2 \leq 1 \text{ or } x^2 + y^2 \geq 4\}$ is not a region in R^2, since it is _____

page 90

Answer (Please select your correct option)

- not connected
- connected
- polygonally connected
- None of these

Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536

Start Time: 11:05 AM

MTH631 Real Analysis II

Question No : 2 of 25

A set S is polygonally connected if every pair of points in S can be connected by a polygonal path lying ————————

page 89

Answer (Please select your correct option)

- outside the set S
- partially in S
- entirely in S
- None of these

Muhashar Ali
Diplomaticsmile2@ymail.com
Whatsup num 0314-7654536

Start Time: 11:05 AM

MTH631 Real Analysis II

Question No : 1 of 26

A set is closed if and only if it always contains all of its ——

page 86

Answer (Please select your correct option)

- limit points
- neighbourhood points
- interior points
- None of these

Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536

Address http://server/vutes/client/Paper.aspx

MTH631 Real Analysis II

Ouestion No : 26 of 26

Suppose that $f(x,y)=\begin{cases} 0 & \text{if x and y are rational,} \\ 1 & \text{if x is rational and y is irrational,} \\ 2 & \text{if x is irrational and y is rational,} \\ 3 & \text{if x and y are irrational.} \end{cases}$

Find

$$\overline{\int_R} f(x,y)\,d(x,y) \quad \text{and} \quad \underline{\int_R} f(x,y)\,d(x,y) \quad \text{if} \quad R=[a,b]\times[c,d].$$

Let $\mathbf{P}=\{R_1,R_2,\ldots,R_k\}$. Let

$$M_j=\sup\{f(x,y)\,|\,\mathbf{X}\in R_j\} \quad \text{and} \quad m_j=\inf\{f(x,y)\,|\,\mathbf{X}\in R_j\}.$$

Let j be arbitrary in $\{1,2,\ldots,k\}$. Since R_j contains a point $(\hat{x}_j,\hat{y}_j)$ with $\hat{x}_j$ and $\hat{y}_j$ irrational, $M_j=3$. Hence, $\overline{\int_R} f(x,y)\,d(x,y)=3(b-a)(d-c)$. Since R_j contains a point $(\tilde{x}_j,\tilde{y}_j)$ with $\tilde{x}_j$ and $\tilde{y}_j$ rational, $m_j=0$. Hence, $\underline{\int_R} f(x,y)\,d(x,y)=0$.

File Edit View Tools Help

Address http://server/vutes/client/Paper.aspx

MTH631 Real Analysis II

Question No : 25 of 26

Prove: If a nonempty subset of R^n is both open and closed, then $S=R^n$.

Suppose that $\mathbf{X}_0$ is in S. If $S \neq \mathbb{R}^n$, there is an $\mathbf{X}_1 \notin S$. Let

$$H = \{\tau \mid (1-t)\mathbf{X}_0 + t\mathbf{X}_1 \in S \text{ for } 0 \leq t < \tau\}.$$

Since S is open it contains a neighborhood of $\mathbf{X}_0$, so $H \neq \emptyset$. Since $\mathbf{X}_1$ is not in S, $\tau \leq 1$ for all τ in H. Let $\rho = \sup H$ and $\overline{\mathbf{X}} = (1-\rho)\mathbf{X}_0 + \rho\mathbf{X}_1$; then $\overline{\mathbf{X}}$ is a limit point of S and so in S, because S is closed. Since S is open, it contains some ϵ−neighborhood of $\overline{\mathbf{X}}$, so $\rho + \epsilon/2$ is in H. This contradicts the definition of ρ. Hence, $S = R^n$.

VU Examination System (CLIENT) VUTES 7 ... Fall 2019 (FinalTerm)

File Edit View Tools Help

Address http://server/vutes/client/login.aspx

MTH631 Real Analysis II

Question No : 24 of 26

If R is degenerate, then prove by using the definition of Riemann integral that $\int_R f(X)dA = 0$ if f is bounded on R.

Suppose, for example, that $R = [a_1, b_1] \times [a_2, b_2] \times \cdots \times [a_n, b_n]$ with $a_1 = b_1$. Then every partition $\mathbf{P}$ of R is of the form $\mathbf{P} = \{R_1, R_2, \ldots, R_k\}$, where $v(R_j) = 0$, $1 \leq j \leq k$. Therefore, every Riemann sum of f over R equals zero, so $\int_R f(\mathbf{X})d\mathbf{X} = 0$.

Answer (Please click here to Add Answer)

100%


Address
http://server/vutes/client/Paper.aspx
MTH631 Real Analysis II
Question No : 23 of 26
Determine whether the integral $\int_0^\infty \frac{1+\cos^2 x}{\sqrt{1+x^2}}dx$ converges or diverges.
$\frac{1+\cos^2 x}{\sqrt{1+x^2}} > \frac{1}{\sqrt{1+x^2}} = \frac{1}{x\sqrt{1+1/x^2}} > \frac{1}{x\sqrt{2}}$ if $x > 1$. Since $\int_1^\infty \frac{dx}{x} = \infty$,
$\int_0^\infty \frac{1+\cos^2 x}{\sqrt{1+x^2}}dx = \infty$


Address
http://server/vutes/client/Paper.aspx
MTH631 Real Analysis II
Question No : 22 of 26
Calculate $\frac{\partial f(X)}{\partial \Phi}$ where $f(x,y,z) = \log(1+x+y+z)$, $\Phi = (0,1,0)$.
$h(t) = f(x+\phi_1 t, y+\phi_2 t, z+\phi_3 t) = \log(1+x+y+z+(\phi_1+\phi_2+\phi_3)t)$; $h'(t) = \frac{\phi_1+\phi_2+\phi_3}{1+x+y+z+(\phi_1+\phi_2+\phi_3)t}$; $\frac{\partial f(\mathbf{X})}{\partial \acute{\mathbf{L}}} = h'(0) = \frac{\phi_1+\phi_2+\phi_3}{1+x+y+z}$; if $\acute{\mathbf{L}} = (0,1,0)$ then $\frac{\partial f(\mathbf{X})}{\partial \acute{\mathbf{L}}} = \frac{1}{1+x+y+z}$.

Question No : 20 of 26

A ———— surface in $\mathbb{R}^n$ has zero content.

PAGE 161

Answer (Please select your correct option)

- parabolic
- elliptical
- differentiable
- None of these

Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536

MTH631 Real Analysis II

Question No : 19 of 26

If f is integrable on a rectangle R, then ———

page 158

Answer (Please select your correct option)

- $\overline{\int_R} f(x)d(x) > \underline{\int_R} f(x)d(x) = \int_R f(x)d(x)$
- $\overline{\int_R} f(x)d(x) = \underline{\int_R} f(x)d(x) = \int_R f(x)d(x)$ ⇦
- $\overline{\int_R} f(x)d(x) > \underline{\int_R} f(x)d(x) > \int_R f(x)d(x)$
- None of these

Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536

Start Time: 11:01 AM


MTH631 Real Analysis II
Question No : 18 of 26
Marks: 1
The upper integral $\overline{\int_a^b} f(X)d(X)$ of f over R is the infimum of all the———
page 155
Answer (Please select your correct option)
lower sums
upper sums
average sums
None of these
Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536


H631 Real Analysis II
stion No : 17 of 26
Marks: 1 (Budgeted Time
f and g are locally integrable on [a,b), g(x) > 0 and f(x) ≥ 0 on some subintervals [a1,b) of [a,b), and lim f(x)/g(x) = M. If M = ∞ and ∫ g(x)dx = ∞,
page 144
swer (Please select your correct option)
∫ f(x)dx and ∫ g(x)dx converges together
Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536
∫ f(x)dx ≥ ∞
∫ f(x)dx = ∞
∫ f(x)dx < ∞



Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536


MTH631 Real Analysis II
Question No : 15 of 25
If $p(X)$ is a homogeneous polynomial of degree r in $X - X_0$ if $p(X) < 0$ for all $X \neq X_0$. Then p is called
page 134
Answer (Please select your correct option)
positive semidefinite
positive definite
negative semidefinite
negative definite
Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536

Question No : 14 of 26

If $p(X)$ is a homogeneous polynomial of degree r in $X - X_0$. If $p(X) \geq 0$ for all X, then p is called ———

page 134

Answer (Please select your correct option)

- positive semidefinite
- positive definite
- negative semidefinite
- negative definite

Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536

Start Time: 11:01 AM

Question No : 13 of 26

If value of a becomes zero in the Taylor series $\sum_{n=0}^{\infty} \frac{f^n(a)}{n!}(x-a)^n$, then the series is called ———

page 44

Answer (Please select your correct option)

- Maclaurin series
- Fourier series
- Power series
- None of these

Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536

MTH631 Real Analysis II

Question No : 12 of 26

If $h = f(x,y), x = f(r,\theta)$ and $y = f(r,\theta)$, then using the chain rule, $\frac{\partial h}{\partial \theta} =$ _____

page 127

Answer (Please select your correct option)

- $\frac{\partial f}{\partial r}\frac{\partial r}{\partial \theta} + \frac{\partial f}{\partial r}\frac{\partial r}{\partial \theta}$
- $\frac{\partial f}{\partial x}\frac{\partial x}{\partial \theta} + \frac{\partial f}{\partial y}\frac{\partial y}{\partial \theta}$
- $\frac{\partial f}{\partial x} + \frac{\partial x}{\partial \theta} + \frac{\partial f}{\partial y} + \frac{\partial y}{\partial \theta}$

Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536

Question No : 11 of 26

If $f(x)=3x+5x^3$, then at $x=-1$, $df=$ _____

Answer (Please select your correct option)

- $18dy$
- $18dx$
- 18

Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536

MTH631 Real Analysis II

Question No : 10 of 26

A function of the form $L(X) = m_1 x_1 + m_2 x_2 + \ldots + m_n x_n$ for some constants m_i's is called ———

page 112

Answer (Please select your correct option)

- linear function
- quadratic function
- constant function
- None of these

Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536

MTH631 Real Analysis II

Question No : 9 of 26

The rate of change of $f(x,y)$ in the direction of a unit vector is called the ———— of $f(x,y)$

Answer (Please select your correct option)

- differential
- directional derivative
- mixed derivative
- None of these

Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536


MTH631 Real Analysis II
Question No : 8 of 26
A critical point at which a function attains its minimum value among all points where it is defined is called
Answer (Please select your correct option)
global maximum
global minimum
supremum
None of these
Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536


Address http://server/vutms/client/login.aspx
MTH631 Real Analysis II
Question No : 7 of 26
Marks: 1
A function f is ———— on a subset S of R^n if S is contained in an open set on which $f_x, f_y, \ldots, f_z$ are continuous
page 116
Answer (Please select your correct option)
continuously differentiable
uniformly convergent
piecewise convergent
None of these
Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536


MTH631 Real Analysis II
Question No : 6 of 26
A set A ⊆ R of real numbers is ---------- if there exists a real number M ∈ R, such that x ≤ M for every x ∈ A
Answer (Please select your correct option)
bounded below
uniformly continuous
bounded above
None of these
Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536


MTH631 Real Analysis II
Question No : 5 of 26
Marks: 1 (Budgeted Time
Suppose $U_0 \in T$ is a limit point of T. ϕ is continuous at U_0 and f is continuous at $X_0 = G(U_0)$. Then ——— is continuous at U_0
page 102
Answer (Please select your correct option)
$f \circ g$
$\frac{f}{g}$
$\frac{g}{f}$
Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536
None of these


MTH631 Real Analysis II
Question No : 4 of 26
Marks: 1
If f and g are continuous functions at the point X_0 in R^n, then $\frac{f}{g}$ is continuous at X_0 provided that _____
page 100
Answer (Please select your correct option)
$g(X_0) = 0$
$g(X_0) \neq 0$
$f(X_0) = g(X_0) = 0$
None of these
Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536

MTH631 Real Analysis II

Question No : 3 of 26 Marks

For the function $f(x,y)=\frac{xy}{x^2+y^2}$, the limit of $f(x,y)$ as $(x,y)\to(0,0)$ along the line $y=x$ is ——

page 87

Answer (Please select your correct option)

- $\frac{1}{2}$
- $-\frac{1}{2}$
- 0
- 1

Mubashar Ali
Diplomaticsmile13@gmail.com
Whatsup num 0314-7654536

MTH631 Real Analysis II

Question No : 1 of 26

The singleton set is always —

page 81

Answer (Please select your correct option)

- disconnected
- polygonally disconnected
- connected
- None of these

MTH631 Real Analysis II

Question No : 2 of 26

Any polygonally connected set either it is open or not, is always _____

page 82

Answer (Please select your correct option)

- connected
- disconnected
- a region
- None of these

MTH631 Real Analysis II

Question No : 3 of 26 **Marks:** 1 (Budgeted Ti

For the function $f(x,y)=\frac{xy}{x^2+y^2}$, the limit of $f(x,y)$ as $(x,y)\to(0,0)$ along the line $y=x$ is ——

page 87

Answer (Please select your correct option)

- $\frac{1}{2}$
- $-\frac{1}{2}$
- 0
- 1

MTH631 Real Analysis II

Question No : 1 of 26 **Marks: 1 (Budgeted**

If f and g are continuous functions at the point X_0 in R^*, then $\frac{f}{g}$ is continuous at X_0 provided that _____

page 100

Answer (Please select your correct option)

- $g(X_0) = 0$
- $g(X_0) \neq 0$
- $f(X_0) = g(X_0) = 0$
- None of these

MTH631 Real Analysis II

Question No : 5 of 26 Marks: 1 (Budgeted Ti

Suppose $U_0 \in T$ is a limit point of T. g is continuous at U_0 and f is continuous at $X_0 = G(U_0)$. Then ——— is continuous at U_0.

page 102

Answer (Please select your correct option)

- $f \circ g$
- $\frac{f}{g}$
- $\frac{g}{f}$
- None of these

MTH631 Real Analysis II

Question No : 6 of 26

Marks: 1 (Budgeted T

A set A ⊆ R of real numbers is ———— if there exists a real number M ⊆ R, such that $x \leq M$ for every $x \in A$.

Answer (Please select your correct option)

- bounded below
- uniformly continuous
- bounded above
- None of these

MTH631 Real Analysis II

Question No : 7 of 26 **Marks:** 1 (Bud

A function f is ———— on a subset S of R^n if S is contained in an open set on which $f_{x_1}, f_{x_2}, \ldots f_{x_n}$ are continuous

page 116

Answer (Please select your correct option)

- continuously differentiable
- uniformly convergent
- piecewise convergent
- None of these

MTH631 Real Analysis II

Question No : 8 of 26 **Marks:** 1 (Budgeted

A critical point at which a function attains its minimum value among all points where it is defined is called _____

Answer (Please select your correct option)

- global maximum
- global minimum
- supremum
- None of these

MTH631 Real Analysis II

Question No : 9 of 26 Marks: 1 (Budgete

The rate of change of $f(x,y)$ in the direction of a unit vector is called the ———— of $f(x,y)$

Answer (Please select your correct option)

- differential
- directional derivative
- mixed derivative
- None of these

MTH631 Real Analysis II

Question No : 18 of 26

A function of the form $L(X) = m_1 x_1 + m_2 x_2 + \ldots + m_n x_n$ for some constants m_i's is called ———

page 112

Answer (Please select your correct option)

- linear function
- quadratic function
- constant function
- None of these

Rukhsar Kanwal
Mth631
Best Of luck

MTH631 Real Analysis II

Question No : 18 of 25

A function of the form $L(X) = m_1 x_1 + m_2 x_2 + \ldots + m_n x_n$ for some constants m_i's is called ——

page 112

Answer (Please select your correct option)

- linear function
- quadratic function
- constant function
- None of these

Education Information Expert
Prepared by Mubashar Ali
EDUCATION INFORMATION EXPERT

2.17.1 Linear Function

A *linear function* is a function of the form

$$L(\mathbf{X}) = m_1x_1 + m_2x_2 + \cdots + m_nx_n,$$

where m_1, m_2, $\ldots$, m_n are constants. From definition of differentiable at $\mathbf{X}_0$ if and only if there is a linear function L s can be approximated so well near $\mathbf{X}_0$ by

$$L(\mathbf{X}) - L(\mathbf{X}_0) = L(\mathbf{X} - \mathbf{X}_0)$$

MTH631 Real Analysis II

Question No : 9 of 25

For the function $f(x,y) = 3x^3y^3 + xy$ in two variables, $f_{xy}(0,0) =$.....

Answer (Please select your correct option)

- 1
- -1 (correct)
- 0
- None of these

Education Expert Prepared by Anbashar Ali
EDUCATION EXPERT

it is an *rth-order partial derivative of f*. The function

$$f(x,y) = 3x^2y^3 + xy$$

has partial derivatives ever[illegible]. It[illegible] first-order partial derivatives are

$$f_x(x,y) = 6xy^3 + y, \quad f_y(x,y) = 9x^2y^2 + x.$$

Its second-order partial derivatives are

$$\begin{aligned} f_{xx}(x,y) &= 6y^3, & f_{yy}(x,y) &= 18x^2y, \\ f_{xy}(x,y) &= 18xy^2 + 1, & f_{yx}(x,y) &= 18xy^2 + 1. \end{aligned}$$

There are eight third-order partial derivatives. Some examples are

$$f_{xxy}(x,y) = 18y^2, \quad f_{xyx}(x,y) = 18y^2, \quad f_{yxx}(x,y) = 18y^2.$$

Compute $f_{xx}(0,0)$, $f_{yy}(0,0)$, $f_{xy}(0,0)$, and $f_{yx}(0,0)$ if

$$f(x,y) = \begin{cases} \frac{(x^2y+xy^2)\sin(x-y)}{x^2+y^2}, & (x,y) \neq (0,0), \\ 0, & (x,y) = (0,0). \end{cases}$$

If $(x,y) \neq (0,0)$, the ordinary rules for differentiation, applied sepa and y, yield

$$\begin{aligned} f_x(x,y) &= \frac{(2xy+y^2)\sin(x-y)+(x^2y+xy^2)\cos(x-y)}{x^2+y^2} \\ &\quad - \frac{2x(x^2y+xy^2)\sin(x-y)}{(x^2+y^2)^2}, \quad (x,y) \neq (0,0), \end{aligned}$$

and

$$\begin{aligned} f_y(x,y) &= \frac{(x^2+2xy)\sin(x-y)-(x^2y+xy^2)\cos(x-y)}{x^2+y^2} \\ &\quad - \frac{2y(x^2y+xy^2)\sin(x-y)}{(x^2+y^2)^2}, \quad (x,y) \neq (0,0). \end{aligned}$$

These formulas do not apply if $(x,y) = (0,0)$, so we find $f_x(0,0)$ and $f_$ their definitions as difference quotients:

$$\begin{aligned} f_x(0,0) &= \lim_{x\to 0} \frac{f(x,0) - f(0,0)}{x} = \lim_{x\to 0} \frac{0-0}{x} = 0, \\ f_y(0,0) &= \lim_{y\to 0} \frac{f(0,y) - f(0,0)}{y} = \lim_{y\to 0} \frac{0-0}{y} = 0. \end{aligned}$$

Setting $y = 0$ in (2.32) and (2.33) yields

$$f_x(x,0) = 0, \quad f_y(x,0) = \sin x, \quad x \neq 0,$$

$$f_y(0,0) = \lim_{y\to 0} \frac{f(0,y)-f(0,0)}{y} = \lim_{y\to 0} \frac{0-0}{y} = 0.$$

Setting $y = 0$ in (2.32) and (2.33) yields

$$f_x(x,0) = 0, \quad f_y(x,0) = \sin x, \quad x \neq 0,$$

so

$$f_{xx}(0,0) = \lim_{x\to 0} \frac{f_x(x,0)-f_x(0,0)}{x} = \lim_{x\to 0} \frac{0-0}{x} = 0.$$

$$f_{yx}(0,0) = \lim_{x\to 0} \frac{f_y(x,0)-f_y(0,0)}{x} = \lim_{x\to 0} \frac{\sin x - 0}{x} = 1.$$

Setting $x = 0$ in (2.32) and (2.33) yields

$$f_x(0,y) = -\sin y, \quad f_y(0,y) = 0, \quad y \neq 0,$$

so

$$f_{xy}(0,0) = \lim_{y\to 0} \frac{f_x(0,y)-f_x(0,0)}{y} = \lim_{y\to 0} \frac{-\sin y - 0}{y} = -1,$$

$$f_{yy}(0,0) = \lim_{y\to 0} \frac{f_y(0,y)-f_y(0,0)}{y} = \lim_{y\to 0} \frac{0-0}{y} = 0.$$

2.16.2 Equality of Mixed Partial Derivatives

Theorem: Suppose that f, f_x, f_y, and f_{xy} exist on a neighborhood N o
and f_{xy} is continuous at (x_0, y_0).

Then $f_{yx}(x_0, y_0)$ exists, and

$$f_{yx}(x_0,y_0) = f_{xy}(x_0,y_0).$$

Proof: Suppose that $\varepsilon > 0$. Choose $\delta > 0$ so that the open

$$S_\delta = \{(x,y) : |x-x_0| < \delta, |y-y_0| < \delta\}$$

is in N.


MTH631 Real Analysis II
A critical point at which a function attains its minimum value among all points where it is defined is called
Answer (Please select your correct option)
global maximum
global minimum
supremum
None of these
In handouts , There is no Concept of globe max,min „ ther local max. and local min. & I think D is the right Answer
correct

2.18 Maxima and Minima

We say that $\mathbf{X}_0$ is a *local extreme point* of f if there is a $\delta > 0$ such that

$$f(\mathbf{X}) - f(\mathbf{X}_0)$$

does not change sign in $S_\delta(\mathbf{X}_0) \cap D_f$.

More specifically, $\mathbf{X}_0$ is a *local maximum point* if

$$f(\mathbf{X}) \leq f(\mathbf{X}_0)$$

or a *local minimum point* if

$$f(\mathbf{X}) \geq f(\mathbf{X}_0)$$

for all $\mathbf{X}$ in $S_\delta(\mathbf{X}_0) \cap D_f$.

MTH631 Real Analysis II

Question No : 7 of 26

A $f_1, f_2, \ldots f_n$ exist on a neighborhood of X_0 and are continuous at X_0, then f is ——— at X_0

page 115

Answer (Please select your correct option)

- uniformly converget
- piecewise convergent
- differentiable
- None of these

Education Information Expert
prepared by Anwashar Ali
EDUCATION INFORMATION EXPERT

2.17.3 A sufficient Condition for Differentiability

Theorem: If $f_{x_1}, f_{x_2}, \ldots, f_{x_n}$ exist on a neighborhood of $\mathbf{X}_0$ and are continuous at $\mathbf{X}_0$, then f is differentiable at $\mathbf{X}_0$.

Proof: Let $\mathbf{X}_0 = (x_{10}, x_{20}, \ldots, x_{n0})$ and suppose that $\varepsilon > 0$. Our assumptions imply that there is a $\delta > 0$ such that $f_{x_1}, f_{x_2}, \ldots, f_{x_n}$ are defined in the n-ball

$$S_\delta(\mathbf{X}_0) = \{\mathbf{X} : |\mathbf{X} - \mathbf{X}_0| < \delta\}$$

and

MTH631 Real Analysis II

Question No : 6 of 26

A set A ⊂ R of real numbers is ———————if there exists a real number M ⊂ R, such that x ≤ M for every x ⊂ A.

Answer (Please select your correct option)

- bounded below
- uniformly continuous
- bounded above
- None of these

May be Bounded above

or

may be the D option.. plz check

1.12 Pointwise and Uniform Bounded Functions

A sequence of functions $\{F_n\}$ on the set S is said to be pointwise bounded on S if the sequence of functions is bounded for every $x \in S$, that is, if there exists a finite valued function $\phi(x)$ defined on S such that

$$|F_n(x)| < \phi(x), \quad x \in S, n = 1, 2, 3, ...$$

We say that $\{F_n\}$ is uniformly bounded on S if there exist a number M such that

$$|F_n(x)| < M, \quad x \in S, n = 1, 2, 3, ...$$

conclude that $f(\mathbf{C}) = u$ for some $\mathbf{C}$ in S.

Theorem: A function f is uniformly continuous on a subset S of its domain in $\mathbb{R}^n$ if for every $\varepsilon > 0$ there is a $\delta > 0$ such that

$$|f(\mathbf{X}) - f(\mathbf{X}')| < \varepsilon$$

whenever

$$|\mathbf{X} - \mathbf{X}'| < \delta$$

and $\mathbf{X}, \mathbf{X}' \in S$.

Remark: We emphasize that δ must depend only on ε and S, and not on the particular points $\mathbf{X}$ and $\mathbf{X}'$.

Theorem: If f is continuous on a compact set S in $\mathbb{R}^n$, then f is uniformly continuous on S.

$|x - p_1| < \delta$.

Since $\{F_n\}$ is pointwise bounded, there exists $M_i < \infty$ such that

$$|F_n(p_i)| < M_i, n \in \mathbb{N}.$$

If we take

$$M = \max\{M_1, ..., M_r\},$$

then $|F_n(x)| < M + \varepsilon$ for every $x \in S$. This proves the first part of the theorem.

We say that $\{F_n\}$ is uniformly bounded on S if there exist a number M such that

$$|F_n(x)| < M, \quad x \in S, n = 1, 2, 3, \ldots.$$

MTH631 Real Analysis II

Question No : 5 of 26

In a vector valued function $G=(g_1,g_2,\ldots,g_n)$. Then $g_1,g_2,\ldots,g_n$ are called the ———— functions of G.

page 114

Answer (Please select your correct option)

- bijective
- surjective
- component
- None of these

Education Information Expert
Prepared by Anashar Ali
EDUCATION INFORMATION EXPERT

2.19 Differentiable Vector Valued Function

A vector-valued function $\mathbf{G} = (g_1, g_2, \ldots, g_n)$ is *differentiable* at

$$\mathbf{U}_0 = (u_{10}, u_{20}, \ldots, u_{m0})$$

if its component functions g_1, g_2, $\ldots$, g_n are differentiable at $\mathbf{U}_0$.

Lemma: Suppose that $\mathbf{G} = (g_1, g_2, \ldots, g_n)$ is differentiable at

$$\mathbf{U}_0 = (u_{10}, u_{20}, \ldots, u_{m0}),$$

and define

$$M = \left(\sum_{i=1}^{n} \sum_{j=1}^{m} \left(\frac{\partial g_i(\mathbf{U}_0}{\partial u_j} \right)^2 \right)^{1/2}.$$

Then, if $\epsilon > 0$, there is a $\delta > 0$ such that

MTH631 Real Analysis II

Question No : 4 of 25

The quotient of two continuous functions is a ———— function wherever the denominator is non-zero.

Answer (Please select your correct option)

- differentiable
- continuous
- composite
- None of these

Information Expert
Subashar Ali
Education for Prepared by
EDUCATION INFORMATION EXPERT

MTH631 Real Analysis II

Question No : 3 of 26

The set $S = ((x,y), x^2 + y^2 \leq 1 \text{ or } x^2 + y^2 \geq 4)$ is not a region in R^2, since it is _____

page 90

Answer (Please select your correct option)

- not connected
- connected
- polygonally connected
- None of these

Education Information Expert
Prepared by Shahbashar Ali
EDUCATION INFORMATION EXPERT

Example: Intervals are the only regions in $\mathbb{R}$. The n-ball $B_r(\mathbf{X}_0)$ is a region in $\mathbb{R}^n$, as is its closure $\overline{S}_r(\mathbf{X}_0)$. The set $S = \{(x, y) : x^2 + y^2 \leq 1 \quad \text{or} \quad x^2 + y^2 \geq 4\}$ is not a region in $\mathbb{R}^2$, since it is not connected.

The set S_1 obtained by adding the line segment

$$L_1: \quad \mathbf{X} = t(0, 2) + (1 - t)(0, 1), \quad 0 < t < 1,$$

MTH631 Real Analysis II

Question No : 2 of 25

A set S is polygonally connected if every pair of points in S can be connected by a polygonal path lying ————

page 89

Answer (Please select your correct option)

- outside the set S
- partially in S
- entirely in S
- None of these

Education Information Expert
Prepared by Shahzad Ali
EDUCATION INFORMATION EXPERT

2.8 Polygonally Connected Set

A set S is *polygonally connected* if every pair of points in S can be connected by a polygonal path lying entirely in S.

Theorem: An open set S in $\mathbb{R}^n$ is connected if and only if it is polygonally connected.

MTH631 Real Analysis II

Question No : 1 of 26

A set is closed if and only if it always contains all of its

page 86

Answer (Please select your correct option)

- limit points
- neighbourhood points
- interior points
- None of these

Education Information Expert Mubashar Ali
EDUCATION INFORMATION EXPERT

limit $\mathbf{X}$. Since $\mathbf{X}$ is a limit point of every S_k and every S_k is closed, $\mathbf{X}$ is in e
S_k (**A set is closed if and only if it contains all its limit points**). There
$\overline{\mathbf{X}} \in I$, so $I \neq \emptyset$. Moreover, $\overline{\mathbf{X}}$ is the only point in I, since if $\mathbf{Y} \in I$, then

MTH631 Real Analysis II

Question No : 19 of 26

The lower integral $\underline{\int_R} f(X)d(X)$ of f over R is the supremum of all the ——

page 135

Answer (Please select your correct option)

- lower sums
- upper sums
- average sums
- None of these

Education Information Expert Prepared by Subhashar Ali
EDUCATION INFORMATION EXPERT

$$s(\mathbf{P}) = \sum_{j=1}^{k} m_j V(R_j).$$

The *lower integral of* f *over* R, denoted by

$$\underline{\int_R} f(\mathbf{X})\, d\mathbf{X},$$

is the supremum of all lower sums.

Theorem: Let f be bounded on a rectangle R and le

Then

1. The upper sum $S(\mathbf{P})$ of f over $\mathbf{P}$ is the supremu
   sums of f over $\mathbf{P}$.

MTH631 Real Analysis II

Question No : 20 of 26

If f and g are integrable on S and $f(x) \leq g(x)$ for x in S, then ———

page163

Answer (Please select your correct option)

- $\int_S f(x)dx \geq \int_S g(x)dx$
- $\int_S f(x)dx \leq \int_S g(x)dx$
- $\int_S f(x)dx > \int_S g(x)dx$
- None of these

Information Expert
Mashar Ali
Education Information Expert Shared By Mashar Ali
EDUCATION INFORMATION EXPERT

Theorem: If f and g are integrable on S and $f(\mathbf{X}) \leq g(\mathbf{X})$ for $\mathbf{X}$ in S, then

$$\int_S f(\mathbf{X})\, d\mathbf{X} \leq \int_S g(\mathbf{X})\, d\mathbf{X}.$$

Theorem: If f is integrable on S, then so is $|f|$, and

$$\left|\int_S f(\mathbf{X})\, d\mathbf{X}\right| \leq \int_S |f(\mathbf{X})|\, d\mathbf{X}.$$

Theorem: If f and g are integrable on S, then so is the product fg.

MTH631 Real Analysis II

Question No : 19 of 26

The upper integral $\overline{\int_R} f(X)d(X)$ of f over R is the infimum of all the———

page 135

Answer (Please select your correct option)

- lower sums
- upper sums
- average sums
- None of these

Education Information Expert Prepared by Mubashar Ali
EDUCATION INFORMATION EXPERT

3.9 Upper and Lower Integrals

If f is bounded on a rectangle R in $\mathbb{R}^n$ and $\mathbf{P} = \{R_1, R_2, \ldots, R_k\}$ is a partition of R.

Let

$$M_j = \sup_{\mathbf{X} \in R_j} f(\mathbf{X}), \quad m_j = \inf_{\mathbf{X} \in R_j} f(\mathbf{X}).$$

The *upper sum* of f over $\mathbf{P}$ is

$$S(\mathbf{P}) = \sum_{j=1}^{k} M_j V(R_j).$$

MTH631 Real Analysis II

Question No : 17 of 26

A function f is said to be absolutely integrable on $[a,b)$ if f is locally integrable on $[a,b)$ and ———

page 145

Answer (Please select your correct option)

- $\int_a^b |f(x)|dx < \infty$ ⇦
- $\int_a^b |f(x)|dx = \infty$
- $\int_a^b |f(x)|dx \geq \infty$
- None of these

Education Information Expert Prepared by Mubashar Ali
EDUCATION INFORMATION EXPERT

3.2 Absolute integrability

We say that f is *absolutely integrable on* $[a,b)$ if f is locally integrable on $[a,b)$ and $\int_a^b |f(x)|\,dx < \infty$. In this case we also say that $\int_a^b f(x)\,dx$ *converges absolutely* or *is absolutely convergent*.

Remark: If f is nonnegative and integrable on $[a,b)$, then f is absolutely integrable on $[a,b)$, since $|f| = f$.

Example: The empty set and singleton sets are connected, because they cannot be represented as the union of two disjoint nonempty sets.

2.6 Heine-Borel Theorem

We are going to state and prove the Heine-Borel theorem for $\mathbb{R}^n$.

This theorem concerns *compact* sets. As in $\mathbb{R}$, a compact set in $\mathbb{R}^n$ is a closed and bounded set.

of *supremum* and *infimum* of a function on a set S.

Theorem: If f is continuous on a compact set S in $\mathbb{R}^n$, then f is bounded on S.

Rukhsar Kanwal
Mth631

MTH631 Real Analysis II

Question No : 1 of 26

The singleton set is always ---

page 81

Answer (Please select your correct option)

- disconnected
- polygonally disconnected
- connected — correct
- None of these

f S can be expressed in this way, then S is *disconnected*.

ample: The empty set and singleton sets are connected, because
esented as the union of two disjoint nonempty sets.


MTH631 Real Analysis II
Question No : 2 of 26
Any polygonally connected set either it is open or not, is always _____
page 82
Answer (Please select your correct option)
connected
disconnected
a region
None of these

, $B = \emptyset$, since if $B \neq \emptyset$, S would be
e, $A = S$, which completes the proof of necessity.

: **Any polygonally connected set, open or not, is connected.** T
A set (not open) may be connected but not polygonally connec

s in $\mathbb{R}^n$. A *region* S in $\mathbb{R}^n$ is the

MTH631 Real Analysis II

Question No : 3 of 26 Marks: 1 (Budgeted Tir

For the function $f(x,y)=\frac{xy}{x^2+y^2}$, the limit of $f(x,y)$ as $(x,y)\to(0,0)$ along the line $y=x$ is ——

page 87

Answer (Please select your correct option)

- $\frac{1}{2}$
- $-\frac{1}{2}$
- 0
- 1

Example: The function

$$f(x,y) = \frac{xy}{x^2+y^2}$$

is defined everywhere in $\mathbb{R}^2$ except at $(0,0)$. Does $\lim_{(x,y)\to(0,0)} f(x,y)$ exist?

If we try to answer this question by letting (x,y) approach $(0,0)$ along the line $y = x$, we see the functional values

$$f(x,x) = \frac{x^2}{2x^2} = \frac{1}{2}$$

and conclude that the limit is $1/2$.

MTH631 Real Analysis II

Question No : 1 of 26 **Marks:** 1 (Budgeted

If f and g are continuous functions at the point X_0 in R^n, then $\frac{f}{g}$ is continuous at X_0 provided that _____

page 100

Answer (Please select your correct option)

- $g(X_0) = 0$
- $g(X_0) \neq 0$
- $f(X_0) = g(X_0) = 0$
- None of these

valid for all p and q. The quotient

$$f(x) = \frac{h(x)}{g(x)} \tag{1.51}$$

of two power series

$$\begin{aligned} h(x) &= \sum_{n=0}^{\infty} c_n (x - x_0)^n, \quad |x - x_0| < R_1, \\ g(x) &= \sum_{n=0}^{\infty} b_n (x - x_0)^n, \quad |x - x_0| < R_2, \end{aligned}$$

can be represented as a power series

$$f(x) = \sum_{n=0}^{\infty} a_n (x - x_0)^n \tag{1.52}$$

with a positive radius of convergence, provided that

$$b_0 = g(x_0) \neq 0.$$

MTH631 Real Analysis II

Question No : 5 of 26 Marks: 1 (Budgeted Ti

Suppose $U_0 \in T$ is a limit point of T. d is continuous at U_0 and f is continuous at $X_0 = G(U_0)$. Then ——— is continuous at U_0.

page 102

Answer (Please select your correct option)

- $f \circ g$
- $\frac{f}{g}$
- $\frac{g}{f}$
- None of these

Theorem: Suppose that $\mathbf{U}_0$ is in T and is a limit point of T, $\mathbf{G}$ is continuous at $\mathbf{U}_0$, and f is continuous at $\mathbf{X}_0 = \mathbf{G}(\mathbf{U}_0)$. Then $h = f \circ \mathbf{G}$ is continuous at $\mathbf{U}_0$.

MTH631 Real Analysis II

Question No : 6 of 26

Marks: 1 (Budgeted 1

A set A ⊆ R of real numbers is _____ if there exists a real number M ⊆ R, such that x ≤ M for every x ⊆ A.

Answers (Please select your correct option)

- bounded below
- uniformly continuous
- bounded above
- None of these

uniformly continous

MTH631 Real Analysis II

Question No : 7 of 26 Marks: 1 (Bud

A function f is ———— on a subset S of R^n if S is contained in an open set on which $f_1, f_2, \ldots f_k$ are continuous

page 116

Answer (Please select your correct option)

- continuously differentiable
- uniformly convergent
- piecewise convergent
- None of these

2.17.4 Continuously Differentiable Function

We say that f is *continuously differentiable* on a subset S of $\mathbb{R}^n$ if S is contained in an open set on which $f_{x_1}, f_{x_2}, \ldots, f_{x_n}$ are continuous.

The above theorem implies that such a function is differentiable at each $\mathbf{X}_0$ in S.


MTH631 Real Analysis II
Question No : 8 of 26
A critical point at which a function attains its minimum value among all points where it is defined is called
global maximum
global minimum
supremum
None of these
I think D is right answer.
in hangouts there is not any
discussion about globle
max. &min.

MTH631 Real Analysis II

Question No : 9 of 26 Marks: 1 (Budgete

The rate of change of $f(x,y)$ in the direction of a unit vector is called the ———— of $f(x,y)$

Answer (Please select your correct option)

- differential
- directional derivative
- mixed derivative
- None of these

Proof: See lecture.

2.16 Directional Derivative

Let $\mathbf{\Phi}$ be a unit vector and $\mathbf{X}$ a point in $\mathbb{R}^n$.

The directional derivative of f at $\mathbf{X}$ in the direction of $\mathbf{\Phi}$ is defined by

$$\frac{\partial f(\mathbf{X})}{\partial \mathbf{\Phi}} = \lim_{t \to 0} \frac{f(\mathbf{X} + t\mathbf{\Phi}) - f(\mathbf{X})}{t}$$

if the limit exists.


MTH631 Real Analysis II
Question No : 18 of 26
A function of the form $L(X) = m_1 x_1 + m_2 x_2 + \ldots + m_n x_n$ for some constants m_i's is called ———
page 112
Answer (Please select your correct option)
linear function
quadratic function
constant function
None of these

2.17.1 Linear Function

A *linear function* is a function of the form

$$L(\mathbf{X}) = m_1 x_1 + m_2 x_2 + \cdots + m_n x_n,$$

where m_1, m_2, ..., m_n are constants. From definition of differentia

Question 1

Prove that $d(S) = d(\overline{S})$ for any set S in $\mathbb{R}^n$.

Proof:

Since $S \subset \overline{S}$, (A) $d(S) \leq d(\overline{S})$. If $\mathbf{X}$ and $\mathbf{Y}$ are in $\overline{S}$ and $\epsilon > 0$, there are points $\mathbf{X}_1, \mathbf{Y}_1$ in S such that $|\mathbf{X} - \mathbf{X}_1| < \epsilon$ and $|\mathbf{Y} - \mathbf{Y}_1| < \epsilon$. Then

$$|\mathbf{X} - \mathbf{Y}| \leq |\mathbf{X} - \mathbf{X}_1| + |\mathbf{X}_1 - \mathbf{Y}_1| + |\mathbf{Y}_1 - \mathbf{Y}| < d(S) + 2\epsilon.$$

Hence, $d(\overline{S}) < d(S) + \epsilon$. Let $\epsilon \to 0+$ to conclude that $d(\overline{S}) \leq d(S)$, which, with (A), implies that $d(S) = d(\overline{S})$.

Icecream Video Editor

Question 2

If a nonempty subset S of $\mathbb{R}^n$ is both open and closed, then $S = \mathbb{R}^n$.

Prove:

Suppose that $\mathbf{X}_0$ is in S. If $S \neq \mathbb{R}^n$, there is an $\mathbf{X}_1 \notin S$. Let

$$H = \{\tau \mid (1-t)\mathbf{X}_0 + t\mathbf{X}_1 \in S \text{ for } 0 \leq t < \tau\}.$$

Since S is open it contains a neighborhood of $\mathbf{X}_0$, so $H \neq \emptyset$. Since $\mathbf{X}_1$ is not in S, $\tau \leq 1$ for all τ in H. Let $\rho = \sup\ H$ and $\overline{\mathbf{X}} = (1-\rho)\mathbf{X}_0 + \rho\mathbf{X}_1$; then $\overline{\mathbf{X}}$ is a limit point of S and so in S, because S is closed. Since S is open, it contains some $\epsilon-$neighborhood of $\overline{\mathbf{X}}$, so $\rho + \epsilon/2$ is in H. This contradicts the definition of ρ. Hence, $S = R^n$.

Icecream Video Editor

Question 3

$$S = \{(x, y, x) \mid |x| \le 2, |y| \le 1, |z-2| \le 2\}$$

Ans:

$d(S)$ is the supremum of $\sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2 + (z_1 - z_2)^2}$

where $-2 \le x_1,\ x_2 \le 2, -1 \le y_1, y_2 \le 1, 0 \le z_1, z_2 \le 4$;

to maximize this function let, for example,

$x_1 = -2, x_2 = 2, y_1 = -1, y_2 = 1, z_1 = 0, z_2 = 4$; thus, $d(S) = \sqrt{4^2 + 2^2 + 4^2} = 6$.

Icecream Video Editor

Question 4

$$f(x, y, z) = \log(1 + x + y + z), \quad \mathbf{\Phi} = (0, 1, 0)$$

Ans:

$$\begin{aligned} h(t) &= f(x + \phi_1 t, y + \phi_2 t, z + \phi_3 t) \\ &= \log(1 + x + y + z + (\phi_1 + \phi_2 + \phi_3)t); \\ h'(t) &= \frac{\phi_1 + \phi_2 + \phi_3}{1 + x + y + z + (\phi_1 + \phi_2 + \phi_3)t}; \end{aligned}$$

$$\frac{\partial f(\mathbf{X})}{\partial \acute{\mathbf{L}}} = h'(0) = \frac{\phi_1 + \phi_2 + \phi_3}{1 + x + y + z}; \text{ if } \acute{\mathbf{L}} = (0, 1, 0)$$

then

$$\frac{\partial f(\mathbf{X})}{\partial \acute{\mathbf{L}}} = \frac{1}{1 + x + y + z}.$$

Icecream Video Editor

Question 5

If $h(u, v) = f(u^2 + v^2)$, then $vh_u - uh_v = 0$.

ANS:

$$h_u = 2uf'$$

$$h_v = 2vf'$$

$$vh_u - uh_v = 2(vu - uv)f'$$

$$vh_u - uh_v = 0.$$

Question 6

If $h(u, v) = f(\sin u + \cos v)$, then $h_u \sin v + h_v \cos u = 0$.

Ans:

$$h_u = f' \cos u$$
$$h_v = -f' \sin v$$
$$h_u \sin v + h_v \cos u = (\sin v \cos u - \cos u \sin v) f'$$
$$h_u \sin v + h_v \cos u = 0.$$

Icecream Video Editor

Question 7

If $h(u, v) = f(u/v)$, then $uh_u + vh_v = 0$.

Ans:

$$h_u = \frac{1}{v} f'$$

$$h_v = -\frac{u}{v^2} f'$$

$$uh_u + vh_v = \left(\frac{u}{v} - \frac{u}{v}\right) f'$$

$$uh_u + vh_v = 0.$$

Icecream Video Editor

Question 8

If $h(u, v) = f(g(u, v), -g(u, v))$, then $dh = (f_x - f_y)\, dg$.

Ans:

$$h_u = (f_x - f_y)g_u$$

$$h_v = (f_x - f_y)g_v$$

$$dh = h_u\, du + h_v\, dv$$

$$dh = (f_x - f_y)(g_u\, du + g_v\, dv)$$

$$dh = (f_x - f_y)\, dg.$$

Icecream Video Editor

Question 9

Let $h(r, \theta, z) = f(x, y, z)$, where $x = r\cos\theta$ and $y = r\sin\theta$.

Find h_r, h_θ, and h_z in terms of f_x, f_y, and f_z.

Ans:

$$h_r = f_x x_r + f_y y_r + f_z z_r$$

$$h_r = f_x \cos\theta + f_y \sin\theta$$

$$h_\theta = f_x x_\theta + f_y y_\theta + f_z z_\theta$$

$$h_\theta = f_x(-r\sin\theta) + f_y(r\cos\theta)$$

$$h_\theta = r(-f_x \sin\theta + f_y \cos\theta)$$

$$h_z = f_x x_z + f_y y_z + f_z = f_z.$$

Question 10

Let $h(r, \theta, \phi) = f(x, y, z)$, where $x = r \sin\phi \cos\theta$, $y = r \sin\phi \sin\theta$, and $z = r \cos\phi$. Find h_r, h_θ, and h_ϕ in terms of f_x, f_y, and f_z.

ANS:

$$h_r = f_x x_r + f_y y_r + f_z z_r$$

$$h_r = f_x \sin\phi \cos\theta + f_y \sin\phi \sin\theta + f_z \cos\phi$$

$$h_\theta = f_x x_\theta + f_y y_\theta + f_z z_\theta$$

$$h_\theta = r \sin\phi(-f_x \sin\theta + f_y \cos\theta)$$

$$h_\phi = f_x x_\phi + f_y y_\phi + f_z z_\phi$$

$$h_\phi = r(f_x \cos\phi \cos\theta + f_y \cos\phi \sin\theta - f_z \sin\phi.$$

Question 11

Let $h(\mathbf{U}) = f(\mathbf{G}(\mathbf{U}))$ and find $d_{\mathbf{U}_0}h$ by writing h explicitly as a function of $\mathbf{U}$.

$$\begin{aligned} f(x, y) &= 3x^2 + 4xy^2 + 3x, \\ g_1(u, v) &= ve^{u+v-1}, \\ g_2(u, v) &= e^{-u+v-1}, \end{aligned} \qquad (u_0, v_0) = (0, 1)$$

Ans:

$$h(\mathbf{U}) = 3v^2e^{2u+2v-2} + 4ve^{u+v-1}e^{-2u+2v-2} + 3ve^{u+v-1}$$

$$= 3v^2e^{2u+2v-2} + 4ve^{-u+3v-3} + 3ve^{u+v-1}$$

$$h_u(\mathbf{U}) = 6v^2e^{2u+2v-2} - 4ve^{-u+3v-3} + 3ve^{u+v-1}$$

$$h_u(\mathbf{U}_0) = 5$$

$$h_v(\mathbf{U}) = (6v + 6v^2)e^{2u+2v-2} + (4 + 12v)e^{-u+3v-3} + (3 + 3v)e^{u+v-1}$$

$$h_v(\mathbf{U}_0) = 34$$

$$d_{\mathbf{U}_0}h = h_u(\mathbf{U}_0)\,du + h_v(\mathbf{U}_0)\,dv = 5\,du + 34\,dv$$

Icecream Video Editor

Question 12

Let $h(\mathbf{U}) = f(\mathbf{G}(\mathbf{U}))$ and find $d_{\mathbf{U}_0}h$ by writing h explicitly as a function of $\mathbf{U}$.

$$
\begin{aligned}
f(x, y, z) &= e^{-(x+y+z)}, \\
g_1(u, v, w) &= \log u - \log v + \log w, \\
g_2(u, v, w) &= -2\log u - 3\log w, \\
g_3(u, v, w) &= \log u + \log v + 2\log w,
\end{aligned}
\qquad (u_0, v_0, w_0) = (1, 1, 1)
$$

Ans:

$$x(\mathbf{U}) + y(\mathbf{U}) + z(\mathbf{U}) = (\log u - \log v + \log w) + (-2\log u - 3\log w) + (\log u + \log v + 2\log w)$$

$$x(\mathbf{U}) + y(\mathbf{U}) + z(\mathbf{U}) = 0$$

hence

$$h(\mathbf{U}) \equiv 1$$

$$d_{v_0}h = 0.$$

Icecream Video Editor

Question 13

Let $h(\mathbf{U}) = f(\mathbf{G}(\mathbf{U}))$ and find $d_{\mathbf{U}_0}h$ by writing h explicitly as a function of $\mathbf{U}$.

$$f(x, y) = (x + y)^2,$$
$$g_1(u, v) = u \cos v, \qquad (u_0, v_0) = (3, \pi/2)$$
$$g_2(u, v) = u \sin v,$$

Ans:

$$h(\mathbf{U}) = u^2(\cos v + \sin v)^2$$
$$= u^2(1 + 2 \sin v \cos v)$$
$$= u^2(1 + \sin 2v)$$
$$h_u(\mathbf{U}) = 2u(1 + \sin 2v)$$
$$h_u(\mathbf{U}_0) = 6$$
$$h_v(\mathbf{U}) = 2u^2 \cos 2v;$$
$$h_v(\mathbf{U}_0) = -18$$
$$d_{\mathbf{U}_0}h = h_u(\mathbf{U}_0)\, du + h_v(\mathbf{U}_0)\, dv$$
$$d_{\mathbf{U}_0}h = 6\, du - 18\, dv.$$

Icecream Video Editor

Question 14

Let $h(\mathbf{U}) = f(\mathbf{G}(\mathbf{U}))$ and find $d_{\mathbf{U}_0} h$ by writing h explicitly as a function of $\mathbf{U}$.

$$
\begin{aligned}
f(x, y, z) &= x^2 + y^2 + z^2, \\
g_1(u, v, w) &= u \cos v \sin w, \\
g_2(u, v, w) &= u \cos v \cos w, \\
g_3(u, v, w) &= u \sin v;
\end{aligned}
\qquad (u_0, v_0, w_0) = (4, \pi/3, \pi/6)
$$

Ans:

$$h(\mathbf{U}) = u^2 \cos^2 v \sin^2 w + u^2 \cos^2 v \cos^2 w + u^2 \sin^2 v$$

$$h(\mathbf{U}) = u^2$$

$$h_u(\mathbf{U}) = 2u$$

$$h_v(\mathbf{U}) = 0$$

$$h_w(\mathbf{U}) = 0$$

$$h_u(\mathbf{U}_0) = 8;$$

$$h_v(\mathbf{U}_0) = 0;$$

$$h_w(\mathbf{U}_0) = 0$$

$$dh = 8\,du.$$

Question 15

Show that $(0, 0)$ is a critical point of each of the following functions, and that they have positive semidefinite second differentials at $(0, 0)$.

$$p(x, y) = x^2 - 2xy + y^2 + x^4 + y^4;$$

Ans:

$$p(x, y) = x^2 - 2xy + y^2 + x^4 + y^4$$

$$p_x(x, y) = 2x - 2y + 4x^3$$

$$p_y(x, y) = -2x + 2y + 4y^3.$$

Since $p_x(0, 0) = p_y(0, 0) = 0$, $(0, 0)$ is a critical point of p.

$$p_{xx}(x, y) = 2 + 12x^2$$

$$p_{xy}(x, y) = -2;$$

$$p_{yy}(x, y) = 2 + 12y^2$$

$$(d^{(2)}_{(0,0)}p)(x, y) = 2x^2 - 4xy + 2y^2$$

$(d^{(2)}_{(0,0)}p)(x, y) = 2(x - y)^2$ is positive semidefinite.

Question 16

Show that $(0, 0)$ is a critical point of each of the following functions, and that they have positive semidefinite second differentials at $(0, 0)$.

$$q(x, y) = x^2 - 2xy + y^2 - x^4 - y^4.$$

Ans:

$$q(x, y) = x^2 - 2xy + y^2 - x^4 - y^4$$

$$q_x(x, y) = 2x - 2y - 4x^3$$

$$q_y(x, y) = -2x + 2y - 4y^3$$

Since $q_x(0,0) = q_y(0,0) = 0$, $(0, 0)$ is a critical point of q.

$$q_{xx}(x, y) = 2 - 12x^2$$

$$q_{xy}(x, y) = -2$$

$$q_{yy}(x, y) = 2 - 12y^2$$

$$(d^{(2)}_{(0,0)}q)(x, y) = 2x^2 - 4xy + 2y^2$$

$(d^{(2)}_{(0,0)}q)(x, y) = 2(x - y)^2$ is positive semidefinite.

Video Editor

Question 17

If R is degenerate, then implies that $\int_R f(\mathbf{X})\, d\mathbf{X} = 0$ if f is bounded on R.

Proof:

Suppose, for example, that $R = [a_1, b_1] \times [a_2, b_2] \times \cdots \times [a_n, b_n]$ with $a_1 = b_1$.

Then every partition $\mathbf{P}$ of R is of the form $\mathbf{P} = \{R_1, R_2, \ldots, R_k\}$, where $v(R_j) = 0$, $1 \leq j \leq k$. Therefore, every Riemann sum of f over R equals zero, so $\int_R f(\mathbf{X}) d\mathbf{X} = 0$.

Icecream Video Editor

Question 18

Suppose that

$$f(x,y)=\begin{cases}0 & \text{if } x \text{ and } y \text{ are rational,}\\ 1 & \text{if } x \text{ is rational and } y \text{ is irrational,}\\ 2 & \text{if } x \text{ is irrational and } y \text{ is rational,}\\ 3 & \text{if } x \text{ and } y \text{ are irrational.}\end{cases}$$

Find $\overline{\int_R} f(x,y)\,d(x,y)$ and $\underline{\int_R} f(x,y)\,d(x,y)$ if $R=[a,b]\times[c,d]$.

Ans:

Let $\mathbf{P}=\{R_1,R_2,\ldots,R_k\}$. Let

$$M_j=\sup\{f(x,y)\,|\,\mathbf{X}\in R_j\} \quad \text{and} \quad m_j=\inf\{f(x,y)\,|\,\mathbf{X}\in R_j\}.$$

Let j be arbitrary in $\{1,2,\ldots,k\}$. Since R_j contains a point $(\hat{x}_j,\hat{y}_j)$ with $\hat{x}_j$ and $\hat{y}_j$ irrational, $M_j=3$. Hence, $\overline{\int_R} f(x,y)\,d(x,y)=3(b-a)(d-c)$. Since R_j contains a point $(\tilde{x}_j,\tilde{y}_j)$ with $\tilde{x}_j$ and $\tilde{y}_j$ rational, $m_j=0$. Hence, $\underline{\int_R} f(x,y)\,d(x,y)=0$.

Icecream Video Editor

Question 19

If S_1 and S_2 have zero content, then $S_1 \cup S_2$ has zero content.

Prove

If $\epsilon > 0$, there are rectangles $T_1, T_2, \ldots, T_r$ and $T_1', T_2', \ldots, T_s'$ such that $S_1 \subset \cup_{i=1}^{r} T_i$, $S_2 \subset \cup_{j=1}^{s} T_j'$, $\sum_{i=1}^{r} V(T_r) < \epsilon/2$, and $\sum_{j=1}^{s} V(T_j') < \epsilon/2$. Then $\{T_1, \ldots, T_m, T_1', \ldots, T_r'\}$ covers $S_1 \cup S_2$ with total content $< \epsilon$.

Icecream Video Editor

Question 20

If S_1 has zero content and $S_2 \subset S_1$, then S_2 has zero content.

prove

If $\epsilon > 0$, there are rectangles $T_1, T_2, \ldots, T_r$ and $T_1', T_2', \ldots, T_s'$ such that If $S_1 \subset \cup_{i=1}^{r} T_r$ and $S_2 \subset S_1$, then $S_2 \subset \cup_{i=1}^{r} T_r$.

Icecream Video Editor