


That all is collected by removing the mistakes of the one paper file, hope it helpful…!




In that the answer is "unique point"




In that the correct answer is the "Set of Real Numbers".




In this the exact answer is "an ordered triple".




It's right




This one is also wrong as the answer is "4th one".because the central point of spherical is pie/3 that should be same in cylindrical so, that's only occur in last one...



That's right



Here's answer is the "2nd" one...!



That can't say exactly guys, its formula is following...!

a.b = |a||b| cosØ
Ø=angle between the vectors

MTH301 Calculus II

Question No : 9 of 26 — Marks: 1 (Budgeted Time 1 Min)

Is the function $f(x, y)$ continuous at origin? If not, why?

$$f(x, y) = \begin{cases} 0 & \text{if } x \geq 0 \text{ and } y \geq 0 \\ 1 & \text{Otherwise} \end{cases}$$

Answer (Please select your correct option)

- $f(x, y)$ is continuous at origin
- $f(0, 0)$ is not defined
- $f(0, 0)$ is defined but $\lim_{(x,y)\to(0,0)} f(x, y)$ does not exist
- $f(0, 0)$ is defined and $\lim_{(x,y)\to(0,0)} f(x, y)$ exists but these two numbers are not equal.

That's correct answer…!

MTH301 Calculus II

Question No : 10 of 26 — Marks: 1 (Budgeted Time 1 Min)


Is the function $f(x, y)$ continuous at origin? If not, why?

$$f(x, y) = \begin{cases} \frac{3x^2 y}{x^2 + y^2} & \text{if } (x, y) \neq 0 \\ 0 & \text{if } (x, y) = 0 \end{cases}$$

Answer (Please select your correct option)

- $f(x, y)$ is continuous at origin
- $f(0, 0)$ is not defined
- $f(0, 0)$ is defined but $\lim_{(x,y)\to(0,0)} f(x, y)$ does not exist
- $f(0, 0)$ is defined and $\lim_{(x,y)\to(0,0)} f(x, y)$ exists but these two numbers are not equal.

But that's wrong the correc one is the 1st..




That's the first answer, as that was been define in the lecture 22…!




Here's the answer is the last one, you can check that on the start of lecture 11th, handout…!




Its answer is the "perpendicular"...!



That's right answer...!



This is also right one..!



"No conclusion can be drawn" is the answere...!

MTH301 Calculus II

Question No : 17 of 26 Marks: 1 (Budgeted Time 1 Min)

If $R = R_1 \cup R_2$, where R_1 and R_2 are no overlapping regions then

$$\iint_{R_1} f(x,y)dA + \iint_{R_2} f(x,y)dA =$$

Answer (Please select your correct option)

- $\iint_R f(x,y)dA$
- $\iint_{R_1} f(x,y)dA \cup \iint_{R_2} f(x,y)dA$
- $\iint_R f(x,y)dV$
- $\iint_{R_1} f(x,y)dA \cap \iint_{R_2} f(x,y)dA$

Its answer is the first one option…!

MTH301 Calculus II

Question No : 18 of 26 Marks: 1 (Budgeted Time 1 Min)

If $R = \{(x,y) / 0 \le x \le 2 and 1 \le y \le 4\}$, then

$$\iint_R (6x^2 + 4xy^3)dA =$$

Answer (Please select your correct option)

- $\int_1^4 \int_0^2 (6x^2 + 4xy^3)dydx$
- $\int_0^2 \int_1^4 (6x^2 + 4xy^3)dxdy$
- $\int_1^4 \int_0^2 (6x^2 + 4xy^3)dxdy$
- $\int_2^4 \int_0^1 (6x^2 + 4xy^3)dxdy$

Its answer is the 3rd option guys..!

MTH301 Calculus II

Question No : 19 of 26 Marks: 1 (Budgeted Time 1 Min)

If $R = \{(x,y) / 2 \le x \le 4 \text{ and } 0 \le y \le 1\}$, then

$\iint_R (4xe^{2y})dA =$

Answer (Please select your correct option)

- $\int_0^1 \int_2^4 (4xe^{2y})dydx$
- $\int_0^1 \int_2^4 (4xe^{2y})dxdy$
- $\int_1^4 \int_0^2 (4xe^{2y})dxdy$
- $\int_1^4 \int_0^2 (4xe^{2y})dydx$

It's right one...!

MTH301 Calculus II

Question No : 20 of 26 Marks: 1 (Budgeted Time 1 Min)


If $R = \{(x,y) / 0 \le x \le 4 \text{ and } 0 \le y \le 9\}$, then

$\iint_R (3x - 4x\sqrt{xy})dA =$

Answer (Please select your correct option)


- $\int_0^9 \int_0^4 (3x - 4x\sqrt{xy})dydx$
- $\int_0^4 \int_4^9 (3x - 4x\sqrt{xy})dxdy$
- $\int_4^9 \int_0^0 (3x - 4x\sqrt{xy})dxdy$
- $\int_0^4 \int_0^9 (3x - 4x\sqrt{xy})dydx$


This one's also right...



Its equation is

$$f_x(a)(x - x_0) + f_y(b)(y - y_0) + f_z(c)(z - z_0) = 0$$





These both can be solve by the following example the me gonna paste...!

Evaluate the following double integral.


$$\iint \left(3 + 2x - 3y^2\right) dx\, dy$$

Solution:


$$\iint\left(3+2x-3y^2\right)\,dx\,dy$$

$$\int(3x+2x^2-3xy^2)dy$$


$$3xy+2x^2y-xy^3$$



I didn't got this question dear's...! :(



You can use the above tangent equation to sove this...!



Its answer is the under mentioned...!

as our require is area of the following region

$V = \iint_R 1\,dA = \iint_R dA$ we can compute area by keeping f(x,y) const

$$= \int_0^1 \int_{x^2}^{x} dy\,dx$$

$$= \int_0^1 [x]_{x^2}^{x}$$

$$= \int_0^1 (x - x^2)$$

$$= \left[\frac{x^2}{2} - \frac{x^3}{3}\right]_0^1$$

$$= \left[\frac{1}{2} - \frac{1}{3}\right]$$

$$= \left[\frac{3-2}{6}\right]$$

$= \frac{1}{6}$ that's exactly what we require...!

Jis bahi nay yeh create ki file us ko bhi aur muj(Zaheer Abbas) ko bhi jis nay uplode ki, dono ko apni doiaon main zaror yad rakhna. Allah Tiala hum sub ki her tarah say madad farmay. (Aameen)