Quiz - UC Browser
quiz.vu.edu.pk/QuizQuestion.aspx?ver=91fd1d29-db8c-4bf4-863a-17bf18fabbb0

All Subjects Past Papers
www.badartoor.com

MC170201745: Shakeel Ahmad

MTH631:Quiz-2

Time Left 49 sec(s)

Quiz Start Time: 04:51 PM, 30 July 2018

Question # 4 of 10 (Start time: 04:53:47 PM, 30 July 2018) Total Marks: 1

. a point

$$\mathbf{X}_0$$

is local extreme point if

Select the correct option

Reload Math Equations

- for all $\mathbf{X} \in D_f$.
- partial derivatives exists and equal to zero.
- (●) does not change sign in $S_\delta(\mathbf{X}_0) \cap D_f$.
- None of these

Click to Save Answer & Move to Next Question

Type here to search
4:54 PM
7/30/2018

Quiz - UC Browser
quiz.vu.edu.pk/QuizQuestion.aspx?ver=c12a9001-5efa-4210-b725-b21789842693

MC170201745: Shakeel Ahmad

Time Left 73 sec(s)

MTH631:Quiz-2

Quiz Start Time: 04:51 PM, 30 July 2018

Question # 8 of 10 (Start time: 04:56:40 PM, 30 July 2018) Total Marks: 1

$f(x, y)$

as

$(x, y) \to (0, 0)$

Select the correct option

Reload Math Equations

- ○ $\frac{1}{2}$
- ● $-\frac{1}{2}$
- ○ 0
- ○ 1

Click to Save Answer & Move to Next Question

All Subjects Past Papers
www.badartoor.com

Type here to search
4:56 PM
7/30/2018

Quiz - UC Browser

quiz.vu.edu.pk/QuizQuestion.aspx?ver=fd9cba51-b9a9-4220-a661-c95ea1805961

MC170201745: Shakeel Ahmad

Time Left 40 sec(s)

MTH631:Quiz-2

Quiz Start Time: 04:51 PM, 30 July 2018

Question # 3 of 10 (Start time: 04:52:52 PM, 30 July 2018) **Total Marks: 1**

Identify the true statement(s)

Select the correct option

Reload Math Equations

- the function $f(x,y)=x^3+y^3$ has a local extreme point at origin.
- If $f_{x_i}(\mathbf{X}_0)=0$ then
- At every point where $f_{x_i}(\mathbf{X}_0)=0$ function has local extreme point.
- None of these

Click to Save Answer & Move to Next Question

All Subjects Past Papers
www.badartoor.com

Type here to search

4:53 PM
7/30/2018

Question # 9 of 10 (Start time: 06:48:56 PM, 31 July 2018)

Which statement(s) is(are) true about the function

$$f(x,y)=\frac{x^2+y^2}{x-y}$$

Select the correct option

○	$f_x(x,y)=\frac{2x}{x-y}-\frac{x^2+y^2}{(x+y)^2}$
○	$f_y(x,y)=\frac{y}{x-y}+\frac{x^2+y^2}{(x-y)^2}$.
○	$f_y(x,y),\ f_x(x,y)$ are continuous everywhere.
○	f is differentiable every where except at the points where

For the function

$$f(x,y)=\frac{xy}{x^2+y^2},$$

Select the correct option

○	$\frac{1}{2}$
○	$-\frac{1}{2}$
○	0
○	1

Quiz - UC Browser
quiz.vu.edu.pk/QuizQuestion.aspx?ver=7ad831c0-d1af-498e-9473-d867b99b8312
All Subjects Past Papers
www.badartoor.com

MC170201745: Shakeel Ahmad

MTH631:Quiz-2

Time Left 74 sec(s)

Quiz Start Time: 04:51 PM, 30 July 2018

Question # 5 of 10 (Start time: 04:54:32 PM, 30 July 2018) Total Marks: 1

$$df =$$

Select the correct option

Reload Math Equations

- 9
- 9dx
- -1
- None of these

Click to Save Answer & Move to Next Question

Type here to search
4:54 PM
7/30/2018

M. 31 July 2018)

A critical point at which a function attains its maximum value among all points where it is defined is called _____

Select the correct option

○	global minimum
○	infimum
○	global maximum
○	None of these

All Subjects Past Papers
www.badartoor.com

Question # 5 of 10 (Start time: 06:44:50 PM, 31 July 2018)

For the functions $f(x) = 3x^2 + 5x^3$, the differential at x = - 1 exists because

Select the correct option

○	$\lim_{x \to -1} \frac{f(x)}{x+1} = \infty.$
○	$\lim_{x \to -1} \frac{9(x+1)}{x+1} = 0.$
○	$\lim_{x \to -1} \frac{f(x) + 2 - 9(x+1)}{x+1} = 0.$
○	$\lim_{x \to -1} \frac{f(x) + 2 - 9(x+1)}{x+1} = 1.$

Question # 8 of 10 (Start time: 06:47:23 PM, 31 July 2018)

Identify the false statement(s)

Select the correct option

○	The partial derivative is a special case of directional derivative.
○	For a functions of n variables $f(\mathbf{X})$ the partial derivative with respect to variable
○	The partial derivative of the function $f(x, y, z) = 3xyz + 2x^2 + z^2$ with respect to third variable is
○	None of these

Start your online journey from | Calculus/Extrema and Points | LMS-Virtual University of Paki | Quiz
quiz.vu.edu.pk/QuizQuestion.aspx?ver=86cb911e-6972-4b10-87a0-55af82933c83

MC170201745: Shakeel Ahmad

Time Left 66 sec(s)

MTH631:Quiz-2

Quiz Start Time: 04:51 PM, 30 July 2018

Question # 2 of 10 (Start time: 04:51:15 PM, 30 July 2018) Total Marks: 1

A set A ⊂ R of real numbers is ----------if there exists a real number m ∈ R, such that x ≥ m for every x ∈ A.

All Subjects Past Papers
www.badartoor.com

Select the correct option

- bounded below
- uniformly continuous
- bounded above
- None of these

Click to Save Answer & Move to Next Question

All Subjects Past Papers
www.badartoor.com

80%
Type here to search
4:51 PM
7/30/2018

LMS-Virtual University of Paki
Quiz
quiz.vu.edu.pk/QuizQuestion.aspx?ver=d42f00bb-04a3-4aaa-a72f-2aacd013d18a
Google

MC170202886: Aqsa Zunaira

Time Left 41 sec(s)

MTH631:Quiz-2

Quiz Start Time: 03:53 PM, 30 July 2018

Question # 9 of 10 (Start time: 04:01:11 PM, 30 July 2018) Total Marks: 1

along the line

$$y = -x$$

is

Select the correct option

Reload Math Equations

- $\frac{1}{2}$
- $-\frac{1}{2}$
- 0
- 1

Click to Save Answer & Move to Next Question

All Subjects Past Papers
www.badartoor.com

80%
Type here to search
4:02 PM
7/30/2018

LMS-Virtual University of Paki
Quiz
quiz.vu.edu.pk/QuizQuestion.aspx?ver=e89e3140-59d7-457c-addc-b5f6a85e05c1
Google

MC170202886: Aqsa Zunaira

Time Left 75 sec(s)

MTH631:Quiz-2

Quiz Start Time: 03:53 PM, 30 July 2018

Question # 8 of 10 (Start time: 04:00:51 PM, 30 July 2018) Total Marks: 1

A sufficient condition for a function of several variables to be differentiable at point is

All Subjects Past Papers
www.badartoor.com

Select the correct option

Reload Math Equations

- only the partial derivative should exists at that point.
- All partial derivatives exists and are continuous at that point
- only limit at that point should exists.
- None of these

Click to Save Answer & Move to Next Question

80%
Type here to search
4:01 PM
7/30/2018

LMS-Virtual University of Paki
Quiz
quiz.vu.edu.pk/QuizQuestion.aspx?ver=b1853d56-ba52-4c4c-9dde-f47f415767d3
Google

MC170202886: Aqsa Zunaira

Time Left 43 sec(s)

MTH631:Quiz-2

Quiz Start Time: 03:53 PM, 30 July 2018

Question # 7 of 10 (Start time: 03:59:57 PM, 30 July 2018) Total Marks: 1

Identify the false statement(s)

All Subjects Past Papers
www.badartoor.com

Select the correct option

Reload Math Equations

- The partial derivative is a special case of directional derivative.
- $\frac{\partial f(\mathbf{X})}{\partial \mathbf{E}_2}$.
- with respect to third variable is $f_z = 2z$
- None of these

Click to Save Answer & Move to Next Question

80%
Type here to search
4:00 PM
7/30/2018

LMS-Virtual University of Paki
Quiz
quiz.vu.edu.pk/QuizQuestion.aspx?ver=58c92713-3a77-4009-ac92-2838017dcf17
Google

MC170202886: Aqsa Zunaira

Time Left 31 sec(s)

MTH631:Quiz-2

Quiz Start Time: 03:53 PM, 30 July 2018

Question # 1 of 10 (Start time: 03:53:17 PM, 30 July 2018) Total Marks: 1

and

$f_{yx}(x_0, y_0)$

exists. Then,

All Subjects Past Papers
www.badartoor.com

Select the correct option

Reload Math Equations

- (●) $f_{yx}(x_0, y_0) = f_{xy}(x_0, y_0)$
- ( ) $f_{xx}(x_0, y_0) = f_{yy}(x_0, y_0)$
- ( ) $f_x(x_0, y_0) = f_y(x_0, y_0)$
- ( ) None of these

Click to Save Answer & Move to Next Question

80%
Type here to search
3:54 PM
7/30/2018

LMS-Virtual University of Paki
Quiz
quiz.vu.edu.pk/QuizQuestion.aspx?ver=17ad3fa3-3141-4ce8-a300-2435194323f1
Google

MC170202886: Aqsa Zunaira

Time Left 58 sec(s)

MTH631:Quiz-2

Quiz Start Time: 03:53 PM, 30 July 2018

Question # 3 of 10 (Start time: 03:55:28 PM, 30 July 2018) Total Marks: 1

A critical point at which a function attains its maximum value among all points where it is defined is called .....

All Subjects Past Papers
www.badartoor.com

Select the correct option

- global minimum
- infimum
- global maximum
- None of these

Click to Save Answer & Move to Next Question

80%
Type here to search
3:56 PM
7/30/2018

LMS-Virtual University of Paki
Quiz
quiz.vu.edu.pk/QuizQuestion.aspx?ver=88d81915-2062-4901-be11-9ffd9c15a3f0
Google

MC170202886: Aqsa Zunaira

Time Left 31 sec(s)

MTH631:Quiz-2

Quiz Start Time: 03:53 PM, 30 July 2018

Question # 4 of 10 (Start time: 03:56:48 PM, 30 July 2018) Total Marks: 1

Which expression represents the differential of a function of n variables at the point X0 ?

All Subjects Past Papers
www.badartoor.com

Select the correct option

Reload Math Equations

- $L(X) = f_{x_1}(X_0) + f_{x_2}(X_0) + \ldots + f_{x_n}(X_0)$
- $L(X) = f(X_0)x_1 + f(X_0)x_2 + \ldots + f(X_0)x_n$
- $L(X) = f_{x_1}(X_0)x_1 + f_{x_2}(X_0)x_2 + \ldots + f_{x_n}(X_0)x_n$
- None of these

Click to Save Answer & Move to Next Question

80%
Type here to search
3:57 PM
7/30/2018

Start your online journey from | Calculus/Extrema and Points | LMS-Virtual University of Paki | Quiz
quiz.vu.edu.pk/QuizQuestion.aspx?ver=1c5fffa8-ed8f-4dc2-b807-abcfa81a6c44

MC170201745: Shakeel Ahmad

Time Left 82 sec(s)

MTH631:Quiz-2

Quiz Start Time: 04:51 PM, 30 July 2018

Question # 1 of 10 (Start time: 04:51:01 PM, 30 July 2018) Total Marks: 1

A point on the graph of a function at which the tangent line is parallel to x-axis is called the .....

All Subjects Past Papers
www.badartoor.com

Select the correct option

- (●) critical point
- ( ) point of inflection
- ( ) point of relative extrema
- ( ) None of these

Click to Save Answer & Move to Next Question

80%
Type here to search
4:51 PM
7/30/2018

LMS-Virtual University of Paki
Quiz
quiz.vu.edu.pk/QuizQuestion.aspx?ver=2337c52c-f727-45b9-aae8-8850b430de30
Google

MC170202886: Aqsa Zunaira

Time Left 19 sec(s)

MTH631:Quiz-2

Quiz Start Time: 03:53 PM, 30 July 2018

Question # 5 of 10 (Start time: 03:57:57 PM, 30 July 2018)

Total Marks: 1

Which statement(s) is(are) true about the function

$$f(x,y)=\frac{x^2+y^2}{x-y}$$

All Subjects Past Papers
www.badartoor.com

Select the correct option

Reload Math Equations

- $f_x(x,y)=\frac{2x}{x-y}-\frac{x^2+y^2}{(x+y)^2}$
- $f_y(x,y)=\frac{y}{x-y}+\frac{x^2+y^2}{(x-y)^2}.$
- $f_y(x,y),\ f_x(x,y)$ are continuous everywhere.
- is differentiable every where except at the points where $y=x$

Click to Save Answer & Move to Next Question

80%
Type here to search
3:59 PM
7/30/2018

Select the correct option

○	the function $f(x,y)=x^3+y^3$ has a local extreme point at origin.
○	may or may not has a local extreme point at $\mathbf{X}_0$
○	At every point where $f_{x_i}(\mathbf{X}_0)=0$ function has local extreme point.
○	None of these

If

is continuous at $f(x, y)$

Select the correct option

○	$f_{yx}(x_0, y_0) = f_{xy}(x_0, y_0)$
○	$f_{xx}(x_0, y_0) = f_{yy}(x_0, y_0)$
○	$f_x(x_0, y_0) = f_y(x_0, y_0)$
○	None of these

Quiz - UC Browser

quiz.vu.edu.pk/QuizQuestion.aspx?ver=2f2c4fbb-8e6d-4b94-8603-5d62bc6ca4a1

MC170201745: Shakeel Ahmad

Time Left 76 sec(s)

MTH631:Quiz-2

Quiz Start Time: 04:51 PM, 30 July 2018

Question # 10 of 10 (Start time: 04:57:08 PM, 30 July 2018)

Total Marks: 1

A tangent plane to the surface

$$z = f(x, y)$$

is always the limit of the ----------

All Subjects Past Papers
www.badartoor.com

Select the correct option

Reload Math Equations

- (●) secant plane
- ( ) tangent line
- ( ) normal plane
- ( ) None of these

Click to Save Answer & Move to Next Question

Type here to search

4:57 PM
7/30/2018

For a function of several variables ,

, a point $f(\mathbf{X})$

Select the correct option

○	for all $f(\mathbf{X}) - f(\mathbf{X}_0)$ $\mathbf{X} \in D$
○	partial derivatives exists and equal to zero.
○	If $f(\mathbf{X}) - f(\mathbf{X}_0)$ does not change sign in
○	None of these

Quiz - UC Browser
quiz.vu.edu.pk/QuizQuestion.aspx?ver=8ff06fa7-b0a5-491c-ac6b-6ef352ea51c0

MC170201745: Shakeel Ahmad

Time Left 74 sec(s)

MTH631:Quiz-2

Quiz Start Time: 04:51 PM, 30 July 2018

Question # 7 of 10 (Start time: 04:56:19 PM, 30 July 2018) Total Marks: 1

and

$f_{yx}(x_0, y_0)$

exists. Then,

All Subjects Past Papers
www.badartoor.com

Select the correct option

Reload Math Equations

- (●) $f_{yx}(x_0, y_0) = f_{xy}(x_0, y_0)$
- ( ) $f_{xx}(x_0, y_0) = f_{yy}(x_0, y_0)$
- ( ) $f_x(x_0, y_0) = f_y(x_0, y_0)$
- ( ) None of these

Click to Save Answer & Move to Next Question

Type here to search
4:56 PM
7/30/2018

Question # 3 of 10 (Start time: 06:43:31 PM, 31 July 2018)

A tangent plane to the surface

$$z = f(x, y)$$

is always the limit of the ----------

Select the correct option

○	secant plane
○	tangent line
○	normal plane
○	None of these

A sufficient condition for a function of several variables to be differentiable at point is

Select the correct option

○	only the partial derivative should exists at that point
○	All partial derivatives exists and are continuous at that point
○	only limit at that point should exists
○	None of these

ct the correct option

the function $f(x,y) = x^2 + y^2$ has a local extreme point at origin.
then $f_x(X_0) = 0$ f
At every point where $f_x(X_0) = 0$ function has local extreme point.
None of these

Question # 10 of 10 (Start time: 06:49:35 PM, 31 July 2018)

Identify the true statement(s)

Select the correct option

○	the function $f(x,y)=x^3+y^3$ has a local extreme point at origin.
○	If $f_x(\mathbf{X}_0)=0$ then
○	At every point where $f_x(\mathbf{X}_0)=0$ function has local extreme point
○	None of these

Quiz - UC Browser
quiz.vu.edu.pk/QuizQuestion.aspx?ver=d81f8970-01ed-4c13-85fc-0af31a99c5ea

MC170201745: Shakeel Ahmad

Time Left 27 sec(s)

MTH631:Quiz-2

Quiz Start Time: 04:51 PM, 30 July 2018

Question # 6 of 10 (Start time: 04:54:54 PM, 30 July 2018)

Total Marks: 1

For the function

$$f(x,y) = 3x^2y^3 + xy$$

in two variables.

All Subjects Past Papers
www.badartoor.com

Select the correct option

Reload Math Equations

- ○ 0
- ○ -1
- ● 1
- ○ None of these

Click to Save Answer & Move to Next Question

Type here to search
4:55 PM
7/30/2018

limit of uniformly covergent s | LMS-Virtual University of Pak | Quiz
quiz.vu.edu.pk/QuizQuestion.aspx?ver=a9545596-6079-4fb0-a724-4e5c2631d83b
Google

MC170203385: Maffia Bibi

Time Left 83 sec(s)

MTH631:Quiz-1

Quiz Start Time: 03:04 PM, 31 May 2018

Question # 10 of 10 (Start time: 03:07:17 PM, 31 May 2018) Total Marks: 1

If we have

$$f(x)=\sum_{n=0}^{\infty}a_n(x-x_0)^n,\quad |x-x_0|<R,$$

Select the correct option

Reload Math Equations

- $a_n=\frac{f^{(n)}(0)}{n!}.$
- $a_n=\frac{f^{(2n)}(x_0)}{n!}.$
- $a_n=\frac{f(nx_0)}{n!}.$
- $a_n=\frac{f^{(n)}(x_0)}{n!}.$

All Subjects Past Papers
www.badartoor.com

Click to Save Answer & Move to Next Question

90%
Type here to search
3:07 PM
5/31/2018

The values of x in a specific interval for which a given power series converges is called

Select the correct option

○	Interval of continuity
○	ineterval of convergence
○	radius of convergence
○	None of these

Question # 8 of 10 (Start time: 08:07:35 PM, 01 June 2018)

If we have

$$f(x)=\sum_{n=0}^{\infty} a_n(x-x_0)^n, \quad |x-x_0|<R,$$

Select the correct option

○	$a_n=\frac{f^{(n)}(0)}{n!}.$
○	$a_n=\frac{f^{(2n)}(x_0)}{n!}.$
○	$a_n=\frac{f(nx_0)}{n!}.$
○	$a_n=\frac{f^{(n)}(x_0)}{n!}.$

Question # 10 of 10 (Start time: 08:08:38 PM, 01 June 2018)

If $f(x) = \sin x$, then

Select the correct option

○	$f^{2m+1}(0) = 1, m \geq 0$
○	$f^{2m}(0) = 1$
○	$f''(x) = \sin x$
○	$\|f^{(k)}\|_{(-\infty,\infty)} = 1,\ k > 0$

limit of uniformly covergent s | LMS-Virtual University of Pak | Quiz
quiz.vu.edu.pk/QuizQuestion.aspx?ver=d456606b-0286-42ea-b662-280b88a4be28
Google

MC170203385: Maffia Bibi

Time Left 84 sec(s)

MTH631:Quiz-1

Quiz Start Time: 03:04 PM, 31 May 2018

Question # 1 of 10 (Start time: 03:04:50 PM, 31 May 2018) Total Marks: 1

The limit of a uniformly convergent series of integrable functions is always

All Subjects Past Papers
www.badartoor.com

Select the correct option

Reload Math Equations

- not integrable
- differentiable
- integrable
- None of these

Click to Save Answer & Move to Next Question

90%
Type here to search
3:05 PM
5/31/2018

limit of uniformly covergent s | LMS-Virtual University of Pak | Quiz
quiz.vu.edu.pk/QuizQuestion.aspx?ver=4d010cba-ffb0-492e-a3df-08547d405e36
Google

MC170201823: Abid Farooq

Time Left 83 sec(s)

MTH631:Quiz-1

Quiz Start Time: 03:19 PM, 31 May 2018

Question # 3 of 10 (Start time: 03:20:46 PM, 31 May 2018) Total Marks: 1

If we have

$$f(x)=\sum_{n=0}^{\infty} a_n(x-x_0)^n, \quad |x-x_0|<R,$$

Select the correct option

Reload Math Equations

- $a_n=\frac{f^{(n)}(0)}{n!}.$
- $a_n=\frac{f^{(2n)}(x_0)}{n!}.$
- $a_n=\frac{f(nx_0)}{n!}.$
- $a_n=\frac{f^{(n)}(x_0)}{n!}.$

All Subjects Past Papers
www.badartoor.com

Click to Save Answer & Move to Next Question

90%
Type here to search
3:21 PM
5/31/2018

limit of uniformly covergent s | LMS-Virtual University of Pak | Quiz
quiz.vu.edu.pk/QuizQuestion.aspx?ver=e257da42-f5d0-48de-9ca8-cea023b73326
Google

MC170203385: Maffia Bibi

Time Left 82 sec(s)

MTH631:Quiz-1

Quiz Start Time: 03:04 PM, 31 May 2018

Question # 6 of 10 (Start time: 03:06:15 PM, 31 May 2018) Total Marks: 1

If a series converges absolutely uniformly on S, then

All Subjects Past Papers
www.badartoor.com

Select the correct option

- (●) it converges uniformly on S
- ( ) it diverges
- ( ) it is differentiable on S
- ( ) None of these

Click to Save Answer & Move to Next Question

90%
Type here to search
3:06 PM
5/31/2018

If a series converges absolutely uniformly on S, then

Select the correct option

- it converges uniformly on S
- it diverges
- it is differentiable on S
- None of these

Question # 3 of 10 (Start time: 08:04:27 PM, 01 June 2018)

The term by term differentiation of the series is $S(x) = \sum_{n=0}^{\infty}(-1)^n \frac{x^{2n+1}}{(2n+1)!}$

Select the correct option

○	$\sum_{n=0}^{\infty}(-1)\frac{x^{2n+1}}{(2n+1)!}$
○	$\sum_{n=0}^{\infty}(-1)^n\frac{x^{2n}}{(2n)!}$
○	$\sum_{n=0}^{\infty}\frac{x^{2n+1}}{(2n+1)!}$
○	None of these.

limit of uniformly covergent s | LMS-Virtual University of Pak | Quiz
quiz.vu.edu.pk/QuizQuestion.aspx?ver=5603618e-d6a7-4846-bef5-f396d7c21563
Google

MC170203385: Maffia Bibi

All Subjects Past Papers
www.badartoor.com

Time Left 85 sec(s)

MTH631:Quiz-1

Quiz Start Time: 03:04 PM, 31 May 2018

Question # 2 of 10 (Start time: 03:05:01 PM, 31 May 2018) Total Marks: 1

If $f(x) = \sin x$, then

Select the correct option

Reload Math Equations

- $f^{2m+1}(0) = 1, m \geq 0$
- $f^{2m}(0) = 1$
- $f''(x) = \sin x$
- $\|f^{(k)}\|_{(-\infty,\infty)} = 1,\ k \geq 0$

Click to Save Answer & Move to Next Question

90%
Type here to search
3:05 PM
5/31/2018

limit of uniformly covergent s | LMS-Virtual University of Pak | Quiz
quiz.vu.edu.pk/QuizQuestion.aspx?ver=369b048b-a086-4908-984c-91cca7b47039
Google

MC170201823: Abid Farooq

Time Left 82 sec(s)

MTH631:Quiz-1

Quiz Start Time: 03:19 PM, 31 May 2018

Question # 2 of 10 (Start time: 03:20:33 PM, 31 May 2018) Total Marks: 1

The infinite series expansion of $\cos x$ for $-\infty < x < \infty$ is given by

All Subjects Past Papers
www.badartoor.com

Select the correct option

Reload Math Equations

- ● $\sum_{n=0}^{\infty}(-1)^n \frac{x^{2n}}{2n!}.$
- ○ $\sum_{n=0}^{\infty}(-1)^n \frac{x^n}{n!}.$
- ○ $\sum_{n=0}^{\infty}(-1)^n \frac{x^{2n}}{(n+1)!}.$
- ○ None of these

Click to Save Answer & Move to Next Question

90%
Type here to search
3:21 PM
5/31/2018

limit of uniformly covergent s | LMS-Virtual University of Pak | Quiz
quiz.vu.edu.pk/QuizQuestion.aspx?ver=deb4d9c8-65a0-4dd9-b0ec-be8cabe21b5f
Google

MC170203385: Maffia Bibi

Time Left 83 sec(s)

MTH631:Quiz-1

Quiz Start Time: 03:04 PM, 31 May 2018

Question # 8 of 10 (Start time: 03:06:56 PM, 31 May 2018) Total Marks: 1

Fourier series expansion of a function is a representation of the function as a linear combination of

All Subjects Past Papers
www.badartoor.com

Select the correct option

- ○ sines and tangents.
- ○ cosines and tangents.
- ○ tangents and cotangents.
- ● sines and cosines.

Click to Save Answer & Move to Next Question

90%
Type here to search
3:07 PM
5/31/2018

limit of uniformly covergent s | LMS-Virtual University of Pak | Quiz
quiz.vu.edu.pk/QuizQuestion.aspx?ver=3bcf5bdd-b62d-4e12-a975-f4b8feb2d2f0
Google

MC170203385: Maffia Bibi

Time Left 79 sec(s)

MTH631:Quiz-1

Quiz Start Time: 03:04 PM, 31 May 2018

Question # 4 of 10 (Start time: 03:05:24 PM, 31 May 2018) Total Marks: 1

The infinite series expansion of $\cos x$ for $-\infty < x < \infty$ is given by

All Subjects Past Papers
www.badartoor.com

Select the correct option

Reload Math Equations

- ● $\sum_{n=0}^{\infty}(-1)^n \frac{x^{2n}}{2n!}.$
- ○ $\sum_{n=0}^{\infty}(-1)^n \frac{x^n}{n!}.$
- ○ $\sum_{n=0}^{\infty}(-1)^n \frac{x^{2n}}{(n+1)!}.$
- ○ None of these

Click to Save Answer & Move to Next Question

90%
Type here to search
3:05 PM
5/31/2018

limit of uniformly covergent s | LMS-Virtual University of Pak | Quiz
quiz.vu.edu.pk/QuizQuestion.aspx?ver=ac5d9297-6af9-4ef8-81d9-c525f40bf057
Google

MC170201823: Abid Farooq

Time Left 58 sec(s)

MTH631:Quiz-1

Quiz Start Time: 03:19 PM, 31 May 2018

Question # 7 of 10 (Start time: 03:21:28 PM, 31 May 2018) **Total Marks: 1**

Let f be defined by a power series with finite radius of convergence R. Then $\lim_{x\to(x_0+R)+} f(x) = \sum_{n=0}^{\infty} a_n R^n$ only if

Select the correct option

Reload Math Equations

All Subjects Past Papers
www.badartoor.com

- ○ $\sum_{n=0}^{\infty} b_n R^n$ is convergent.
- ● $\sum_{n=0}^{\infty} (-1)^n a_n R^n$ converges.
- ○ $\sum_{n=0}^{\infty} a_n R^n$ converges.
- ○ None of the above

Click to Save Answer & Move to Next Question

90%
Type here to search
3:22 PM
5/31/2018

limit of uniformly covergent s | LMS-Virtual University of Pak | Quiz
quiz.vu.edu.pk/QuizQuestion.aspx?ver=4da02c42-b110-4d55-9499-5be01403b36a
Google

MC170201823: Abid Farooq

Time Left 78 sec(s)

MTH631:Quiz-1

Quiz Start Time: 03:19 PM, 31 May 2018

Question # 10 of 10 (Start time: 03:22:52 PM, 31 May 2018) **Total Marks: 1**

The power series expansion of $f(x) = \sin x$ about $x_0 = 0$ is

All Subjects Past Papers
www.badartoor.com

Select the correct option

Reload Math Equations

- $\sum_{n=0}^{\infty} \frac{x^{2n}}{(2n)!}, \quad -\infty < x < \infty,$
- $\sin x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}, \quad -\infty < x < \infty.$
- $\sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}, \quad -\infty < x < \infty,$
- $\sum_{n=0}^{\infty} \frac{x^{2n+1}}{(2n+1)!}, \quad -\infty < x < \infty.$

Click to Save Answer & Move to Next Question

90%
Type here to search
3:23 PM
5/31/2018

limit of uniformly covergent s | LMS-Virtual University of Pak | Quiz
quiz.vu.edu.pk/QuizQuestion.aspx?ver=c7886443-637e-4ecb-8add-b3959eaf6b0b
Google

MC170201823: Abid Farooq

Time Left 82 sec(s)

MTH631:Quiz-1

Quiz Start Time: 03:19 PM, 31 May 2018

Question # 4 of 10 (Start time: 03:20:57 PM, 31 May 2018) Total Marks: 1

Fourier series expansion of a function is a representation of the function as a linear combination of

All Subjects Past Papers
www.badartoor.com

Select the correct option

- ○ sines and tangents.
- ○ cosines and tangents.
- ○ tangents and cotangents.
- ● sines and cosines.

Click to Save Answer & Move to Next Question

90%
Type here to search
3:21 PM
5/31/2018

Question # 1 of 10 (Start time: 08:02:38 PM, 01 June 2018)

The limit of a uniformly convergent series of integrable functi... always

Select the correct option

○	not integrable
○	differentiable
○	integrable
○	None of these

limit of uniformly covergent s | LMS-Virtual University of Pak | Quiz
quiz.vu.edu.pk/QuizQuestion.aspx?ver=0b270788-5683-4ba2-8a62-c647b953f749
Google

MC170201823: Abid Farooq

Time Left 42 sec(s)

MTH631:Quiz-1

Quiz Start Time: 03:19 PM, 31 May 2018

Question # 1 of 10 (Start time: 03:19:30 PM, 31 May 2018) **Total Marks: 1**

The term by term differentiation of the series

$$\sum_{n=0}^{\infty}(-1)^n\frac{x^{2n}}{(2n)!}, \quad -\infty < x < \infty$$

Select the correct option

Reload Math Equations

- $\sum_{n=0}^{\infty}\frac{x^{2n}}{(2n)!}, \quad -\infty < x < \infty.$
- $\sum_{n=0}^{\infty}(-1)^n\frac{x^{2n}}{(2n)!}, \quad -\infty < x < \infty.$
- $\sum_{n=0}^{\infty}(-1)^n\frac{x^{2n+1}}{(2n+1)!}, \quad -\infty < x < \infty$
- $\sum_{n=20}^{\infty}(-1)^n\frac{x^{2n+1}}{(2n+1)!}, \quad -\infty < x < \infty.$

All Subjects Past Papers
www.badartoor.com

Click to Save Answer & Move to Next Question

90%
Type here to search
3:20 PM
5/31/2018

limit of uniformly covergent s | LMS-Virtual University of Pak | Quiz
quiz.vu.edu.pk/QuizQuestion.aspx?ver=d1fc62b2-f57a-4533-902c-982407d31e33
Google

MC170203385: Maffia Bibi

Time Left 81 sec(s)

MTH631:Quiz-1

Quiz Start Time: 03:04 PM, 31 May 2018

Question # 3 of 10 (Start time: 03:05:10 PM, 31 May 2018) **Total Marks: 1**

Let f be defined by a power series with finite radius of convergence R. Then $\lim_{x\to(x_0+R)^-} f(x) = \sum_{n=0}^{\infty} a_n R^n$ only if

Select the correct option

Reload Math Equations

All Subjects Past Papers
www.badartoor.com

- ○ $\sum_{n=0}^{\infty} b_n R^n$ is uniformly convergent.
- ● $\sum_{n=0}^{\infty} a_n R^n$ converges
- ○ $\sum_{n=0}^{\infty} (-1)^n a_n R^n$ converges.
- ○ None of these.

Click to Save Answer & Move to Next Question

90%
Type here to search
3:05 PM
5/31/2018

Question # 5 of 10 (Start time: 08:05:49 PM, 01 June 2018)

Which of these functions is not uniformly continuous on (0, 1)?

Select the correct option

- x^2
- $1/x^2$
- $\sin x$
- $\frac{\sin x}{x}$

limit of uniformly covergent s | LMS-Virtual University of Pak | Quiz
quiz.vu.edu.pk/QuizQuestion.aspx?ver=87f53ba0-f7c8-4e1d-b71a-0877238d909e
Google

MC170203385: Maffia Bibi

Time Left 65 sec(s)

MTH631:Quiz-1

Quiz Start Time: 03:04 PM, 31 May 2018

Question # 7 of 10 (Start time: 03:06:27 PM, 31 May 2018) Total Marks: 1

The radius of convergence of the power series $\sum_{n=1}^{\infty} a_n(x-x_0)^n$ is

All Subjects Past Papers
www.badartoor.com

Select the correct option

Reload Math Equations

- ○ $R = \lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right|$
- ● $\frac{1}{R} = \lim_{n\to\infty}\left|\frac{a_{n+1}}{a_n}\right|$
- ○ $\frac{1}{R} = \lim_{n\to\infty}\left|\frac{a_n}{a_{n+1}}\right|$
- ○ None of these

Click to Save Answer & Move to Next Question

90%
Type here to search
3:07 PM
5/31/2018

limit of uniformly covergent s | LMS-Virtual University of Pak | Quiz
quiz.vu.edu.pk/QuizQuestion.aspx?ver=6974f85b-0250-47b2-923c-652fd95428c5
Google

MC170201823: Abid Farooq

Time Left 82 sec(s)

MTH631:Quiz-1

Quiz Start Time: 03:19 PM, 31 May 2018

Question # 8 of 10 (Start time: 03:22:21 PM, 31 May 2018) Total Marks: 1

Which of these functions is not uniformly continuous on (0, 1)?

All Subjects Past Papers
www.badartoor.com

Select the correct option

Reload Math Equations

- ○ x^2
- ● $1/x^2$
- ○ $\sin x$
- ○ $\frac{\sin x}{x}$

Click to Save Answer & Move to Next Question

90%
Type here to search
3:22 PM
5/31/2018

limit of uniformly covergent s | LMS-Virtual University of Pak | Quiz
quiz.vu.edu.pk/QuizQuestion.aspx?ver=2007668e-cf09-4db0-ab00-e04b740dcbc8
Google

MC170203385: Maffia Bibi

Time Left 83 sec(s)

MTH631:Quiz-1

Quiz Start Time: 03:04 PM, 31 May 2018

Question # 9 of 10 (Start time: 03:07:07 PM, 31 May 2018) **Total Marks: 1**

The values of x in a specific interval for which a given power series converges is called

All Subjects Past Papers
www.badartoor.com

Select the correct option

Reload Math Equations

- ○ Interval of continuity
- ● ineterval of convergence
- ○ radius of convergence
- ○ None of these

Click to Save Answer & Move to Next Question

90%
Type here to search
3:07 PM
5/31/2018

The term by term differentiation of the series

$$\sum_{n=0}^{\infty}(-1)^n\frac{x^{2n}}{(2n)!}, \quad -\infty < x < \infty$$

Select the correct option

○	$\sum_{n=0}^{\infty}\frac{x^{2n}}{(2n)!}, \quad -\infty < x < \infty.$
○	$\sum_{n=0}^{\infty}(-1)^n\frac{x^{2n}}{(2n)!}, \quad -\infty < x < \infty.$
○	$\sum_{n=0}^{\infty}(-1)^n\frac{x^{2n+1}}{(2n+1)!}, \quad -\infty < x < \infty$
○	$\sum_{n=20}^{\infty}(-1)^n\frac{x^{2n+1}}{(2n+1)!}, \quad -\infty < x < \infty.$

limit of uniformly covergent s | LMS-Virtual University of Pak | Quiz

quiz.vu.edu.pk/QuizQuestion.aspx?ver=6dfb3d9a-3dc9-4d56-a236-465707bdfbf0

All Subjects Past Papers
www.badartoor.com

MC170201823: Abid Farooq

Time Left 84 sec(s)

MTH631:Quiz-1

Quiz Start Time: 03:19 PM, 31 May 2018

Question # 6 of 10 (Start time: 03:21:18 PM, 31 May 2018) Total Marks: 1

In the definition of uniform convergence the natural number N always depends on

Select the correct option

Reload Math Equations

- ● ε (epsilon)
- ○ η
- ○ β
- ○ None of these

Click to Save Answer & Move to Next Question

90%

Type here to search 3:21 PM 5/31/2018

Question # 2 of 10 (Start time: 08:03:43 PM, 01 June 2018)

The power series expansion of $f(x) = \sin x$ about $x_0 = 0$ is

Select the correct option

○	$\sum_{n=0}^{\infty} \frac{x^{2n}}{(2n)!}, \quad -\infty < x < \infty,$
○	$\sin x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(-n)!}, \quad -\infty < x < \infty$
○	$\sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}, \quad -\infty < x < \infty,$
○	$\sum_{n=0}^{\infty} \frac{x^{2n+1}}{(2n+1)!}, \quad -\infty < x < \infty$

Question # 4 of 10 (Start time: 08:05:11 PM. 01 June 2018)

In the definition of uniform convergence the natural number N always depends on

Select the correct option

○	ϵ (epsilon)
○	η
○	β
○	None of these

limit of uniformly covergent s | LMS-Virtual University of Pak | Quiz
quiz.vu.edu.pk/QuizQuestion.aspx?ver=1ded4355-9861-446f-a53d-ff94fc55c18b
Google

MC170201823: Abid Farooq

Time Left 83 sec(s)

MTH631:Quiz-1

Quiz Start Time: 03:19 PM, 31 May 2018

Question # 5 of 10 (Start time: 03:21:08 PM, 31 May 2018) Total Marks: 1

If $f(x) = \sin x$, then

All Subjects Past Papers
www.badartoor.com

Select the correct option

Reload Math Equations

○	$f^{2m+1}(0) = 1, m \geq 0$
○	$f^{2m}(0) = 1$
○	$f''(x) = \sin x$
●	$\|f^{(k)}\|_{(-\infty,\infty)} = 1,\ k \geq 0$

Click to Save Answer & Move to Next Question

90%
Type here to search
3:21 PM
5/31/2018