
ZOO-506 Addition 2

CURRENT PAPER FILE (2021)

Lion group of zoologist

Special instruction from sir Arslan

Zoo 506 ka paper files arha he compelet papers solve krny k liy ap ko lion ki meri old file yaad krni hogi us file video b mery channel py jaoof he ko b dekh skty hen us k baad ye file is file me curre nt paper me jo question hen jo purani lion ki file me nehi thy type krdiy gy hen so apko 2no file krni hongi…



yee old file he is file k baad ye current question ki file yaad kry mcq b isi file me majood hen …

1. damages of red cotton bug

Red cotton bug: *Dysdercus cingulatus*

Family : Pyrrhocoridae

Order : Hemiptera

Symptom of damage

Red stained lint and rotting bolls.

Inner boll wall with warty growth or water soaked spots

Young bolls abort and turn dark brown

The bacterium *Nematospora gossypii enters the site of injury and stains the* fiber.c

1. Physical control method name

The extremes of both temperature and humidity, either high or low, are not only effective barriers in the natural control of insects

But they are also useful in artificial control

Light and electricity are other factors that come under this heading

Physical Factor Controlling Insects

1. Heat

2. Cold

3. Moisture

4. Light 5. Electricity

2. Symptoms of mealy bugs

Sugarcane Mealy Bug

SYMPTOMS OF DAMAGE

1. Nymph and adult suck juice from cane in group & reduces vigour of the plant
2. It also attack roots
3. Honey dew secretion leads to development of Sooty mould which gives blackish appearance to canes.

3. how new plant species cause Insects outbreak

Factors of insect outbreak: In the course of the development of agriculture, man has increased the number and magnitude of his **entomological problems**

Introduction of new plant species: Before the settlement of this country there were, of course, native insects attacking the various plants growing here.

4. Which device are use in electro cuting of insect

Devices are available for electrocuting insects

a. I-Some in the form of **electrical window screens** for flies

b. II-Others as **electric light traps** for moths and beetles

5. How agg of sugarcane top borer look like .

- Dull white elongate eggs
- Covered by buff coloured hairs from female anal truft

Dead heart in grown up canes reddish brown in color which cannot be easily pulled

6. Methods of plowing and cultivation in cultural

Time and method of plowing and cultivating form a fourth factor in the cultural control of insects.

1. Many insect pests overwinter in larval or pupal cells in the ground
2. These cells can often be broken up by fall plowing so the insects will not survive the winter

3. Cultivation in the summer frequently breaks up pupation cells
4. But plowing or cultivation to be effective must be timed according to the life history of the insect that is being controlled, otherwise it is useless.
5. For the control of some insects it is necessary to **refrain from cultivation** at certain times
6. The **depth of plowing** varies sometimes according to the habits of the insect.

7. Characteristics of order Orthoptera

Characteristics 0f order Orthoptera

1. This order includes: Indian House Crickets, Field Cricket, Short-horned, Grasshoppers, Katydid (Long-horned Grasshoppers)
2. Mouthparts: Chewing
3. Metamorphosis: Incomplete Metamorphosis
4. Damage: Chewing leaves, Crop grains
5. Over 20,000 species worldwide

8. Name of the sugarcane early shoot borer.

121 Pests of Sugarcane

Insect pest	Scientific name	Family	Order
EARLY SHOOT BORER	*Chilo infuscatellus*	Pyralidae	Lepidoptera
INTERNODE BORER	*Chilo sacchariphagus indicus*	Pyralidae	Lepidoptera
TOP BORER	*Scirpophaga nivella*	Pyralidae	Lepidoptera

9. What is ETL???

The economic threshold Level (**ETL**) is the population density at which control measures should be determined to prevent an increasing pest population from reaching the Economic injury level.

10. Symptoms of mealy bug

Symptoms of mealy bug feeding include stunting, chlorosis, defoliation, and wilting. Because mealybugs feed on sugary plant juices (**photosynthates**), their waste contains a high sugar content and is referred to as honeydew.

11. Some features of neuroptera

1. 9-Neuroptera

- **Includes:** Green Lacewing, Antlion
- **Mouthparts**: adults have hypognathous (directed downward) mouthparts and unique piercing– sucking larval jaws composed of the mandibles (directed downward) plus maxillae
- *Metamorphosis:* Complete Metamorphosis
- Both are beneficial insects
- Antlions creat a sand pit to capture unsuspecting prey



.

12. features of isopetera

11- Isoptera

- **Includes:** Dry-wood termite, Subterranean termite
- **Mouthparts:** Chewing Mouthparts
- **Metamorphosis:** Incomplete Metamorphosis
- Live for 15 years.
- Lay 1 egg every 15 seconds
- Eat wood. Can destroy a house in 2-3 years
- Found in every US state except Alaska.
- Recycle wood in the soil





13. Characteristics of Arthropod (5) mrks

- No one feature will serve to separate the arthropods from' all other animals
  But the possession by an animal of several of those here described will enable the observer to determine in each case whether he is examining one of this group.
  In arthropods the body is composed of a series of more or less similar pieces or segments Placed one behind another, the line of attachment of these to each other being usually somewhat evident on parts of the body at least
  This character is also shown, and indeed more clearly, in some worms, such as the common earthworm.

- Another character of the arthropods is the presence of jointed legs (or appendages of some kind), as is indicated by the name of the group, and these are not possessed by worms
  The surface of the body is covered by a secretion which hardens on exposure to the air, forming an outside shell or external skeleton called exoskeleton
  There being practically no internal structures supporting the soft organs within this shell except as ingrowths from the outside
  Distinctive Characters of Arthropods Groups

Factors of insect outbreak

Factors of insect outbreak

1. New plants were introduced by the settlers and grown in greater abundance than the wild and scattered ones
2. An insect finding in any of these an acceptable food had at once a more abundant supply and rapid multiplication thus became possible and resulted in increase of the pest to injurious numbers
3. **II- Accidental introduction insect**: A second factor has been the accidental introduction of many insect pests from foreign countries
4. In the United States such forms have often failed entirely to maintain themselves
5. Unfortunately on the other hand some others have frequently found conditions favorable to a rapid increase
6. These multiply unchecked by their natural enemies which in most cases have not accompanied them to this country
7. **III- Destruction of insectivorous birds**: A third factor is that with the increasing occupation of the country, insectivorous birds have been destroyed or frightened away
8. Although some birds have adjusted themselves to the new conditions it is not likely that their increase makes up for the loss of other species

What is the physical appearance of white grub adult?

They are a creamy-white color with a rusty orange head and six legs at the front of their body. Grubs are **C-shaped** and their bodies appear slick and shiny. A caterpillar is totally different in appearance from the butterfly it finally becomes the white grub.

Why classification of insects are important?

It is important to be able to identify different species of insects. With the large number of insect species in the world, it is critical to have a system for organising them. ... Classification involves grouping organisms into a series of hierarchical categories: kingdom, phylum, class, order, family, genus and species.

Tillage, Intelligent cultivation AND destruction of vegetable ?

- ✓ Tillage: Tillage is the agricultural preparation of soil by mechanical agitation of various types, such as digging, stirring, and overturning
- ✓ The destruction of vegetable trash and weeds, frequently called clean culture, is also an important factor, especially in the control of hibernating insects.
- ✓ Intelligent cultivation and use of fertilizers is well as pruning to remove injured or diseased parts will aid in insect control

What are the names of biological control of parasites

- ✓ Trichogramma chilonis
- ✓ T.japonicum
- ✓ Isotima javensis
- ✓ Bacillus popillae(Bacteria)
- ✓ Beauveria bassiana(fungus)

Sir Arslan

Addition 3 continues……..

Sir Arslan