Which one of the following assumptions is TRUE for Two-Way ANOVA?

Select the correct option

Reload Mat

- Groups may or may not have equal sample sizes.
- Samples must be independent.
- Population variances must be different.
- The population must be Away from a normal distribution.

Click to Save Answer & Move to Nex

ZOO103 - Principles of Systematics (Quiz-3) Quiz Start Time: 08:27 PM, 07

Question # 2 of 10 (Start time: 08:28:31 PM, 07 July 2023) Tot

_____ is used to compare means of more than two samples.

Select the correct option

Reload Math Eq

- Range
- ANOVA
- T-Test
- Z-Test

Click to Save Answer & Move to Next Qu

Question # 3 of 10 (Start time: 08:29:21 PM, 07 July 2023) **Tot**

Which one of the following is True?

Select the correct option

Reload Math Eq

○	A one-way ANOVA uses one independent variable, while a Two-way ANOVA uses two dependent variables.
○	A one-way ANOVA uses one dependent variable, while a Two-way ANOVA uses two independent variables.
○	A one-way ANOVA uses one independent variable, while a Two-way ANOVA uses one dependent and one independent variable.
○	A one-way ANOVA uses one independent variable, while a Two-way ANOVA uses two independent variables.

Click to Save Answer & Move to Next Qu

Which one of the following is TRUE for T-Test?

Select the correct option

Reload M

- Sample size should be 50 or less than 50.
- Samples should be taken randomly from the population
- Standard deviation must be known.
- All randomly collected samples should have the different variances

Click to Save Answer & Move to N

Which one of the following is WRONG for ANOVA?

Select the correct option

Reload Mat

○	Heterogeneity of variances.
○	Number of samples must be more than two.
○	Data should be normally distributed.
○	Samples must be independent.

Click to Save Answer & Move to Nex

Question # 6 of 10 (Start time: 06:05:25 PM, 07 July 2020)

In two-way ANOVA, there are

Select the correct option

Reload Math

- Dependent variables
- Independent variables
- Depends upon situation
- One is dependent and other two variables are independent

Click to Save Answer & Move to Next

ZOO103 - Principles of Systematics (Quiz-3) **Quiz Start Time: 08:39 PM, 07 July 2023**

Question # 3 of 10 (Start time: 08:41:04 PM, 07 July 2023) **Total Marks:** 1

Which one of the following is TRUE for T-Test?

Select the correct option

Reload Math Equations

○	Sample size should be 50 or less than 50.
○	Standard deviation must be known.
○	All randomly collected samples should have the different variances
○	Samples should be taken randomly from the population

Click to Save Answer & Move to Next Question

sec(s)

ZOO103 - Principles of Systematics (Quiz-3)

Quiz Start Time: 08:39 PM, 07 July 2023

Question # 1 of 10 (Start time: 08:39:28 PM, 07 July 2023)

Total Marks: 1

_____ is positive square root of variance.

Select the correct option

Reload Math Equations

- Standard deviation
- Confidence interval
- Statistical inference
- Statistical dispersion

Click to Save Answer & Move to Next Question

Question # 10 of 10 (Start time: 08:34:59 PM, 07 July 2023) **Total**

For two sample t-test, the two populations:

Select the correct option

Reload Math Equa

- ○ Vary according to situation
- ○ Must be dependent
- ○ Must be independent
- ○ May be dependent or independent

Click to Save Answer & Move to Next Quest

Question # 2 of 10 (Start time: 08:40:14 PM, 07 July 2023) **Total Marks:** 1

Which one of the following assumptions is TRUE for Two-Way ANOVA?

Select the correct option

Reload Math Equations

○	Samples must be independent.
○	Population variances must be different.
○	The population must be Away from a normal distribution.
○	Groups may or may not have equal sample sizes.

Click to Save Answer & Move to Next Question

DF (within group) = _____

Select the correct option

Reload

- Dividing the SS (between group) with DF (between group)
- (Group mean-Grand mean)2 × Number of observation in each group
- Total number of observation – Total number of groups
- Total number of groups – 1

Click to Save Answer & Move to

Question # 7 of 10 (Start time: 08:32:35 PM, 07 July 2023) **Total M**

According to Maloney et al., Most abundant natural predator is:

Select the correct option

Reload Math Equat

- Spider
- Brown Bear
- Leopard
- Lion

Click to Save Answer & Move to Next Questi

Time Left sec(s)

ZOO103 - Principles of Systematics (Quiz-3)

Quiz Start Time: 08:58 PM, 07 July 2023

Question # 2 of 10 (Start time: 08:59:08 PM, 07 July 2023)

Total Marks: 1

_____ is positive square root of variance.

Select the correct option

Reload Math Equations

- Statistical dispersion
- Confidence interval
- Statistical inference
- Standard deviation

Click to Save Answer & Move to Next Question

ZOO103 - Principles of Systematics (Quiz-3)

Quiz Start Time: 09:10 PM, 07 July 2023

Question # 1 of 10 (Start time: 09:10:21 PM, 07 July 2023)

Total Marks: 1

Which one of the following is TRUE for T-Test?

Select the correct option

Reload Math Equations

- Samples should be taken randomly from the population
- Standard deviation must be known.
- All randomly collected samples should have the different variances
- Sample size should be 50 or less than 50.

Click to Save Answer & Move to Next Question

inal text
this translation
feedback will be used to help improve Google Translate

ZOO103 - Principles of Systematics (Quiz-3) Quiz Start Time: 09:10 PM, 07 July 2023

Question # 2 of 10 (Start time: 09:11:39 PM, 07 July 2023) Total Marks: 1

Which one of the following assumptions is TRUE for Two-Way ANOVA?

Select the correct option

Reload Math Equations

- The population must be Away from a normal distribution.
- Population variances must be different.
- Samples must be independent.
- Groups may or may not have equal sample sizes.

Click to Save Answer & Move to Next Question

inal text
this translation
feedback will be used to help improve Google Translate

sec(s)

ZOO103 - Principles of Systematics (Quiz-3) Quiz Start Time: 09:10 PM, 07 July 2023

Question # 3 of 10 (start time: 09:12:38 PM, 07 July 2023) Total Marks: 1

DF (between) =

Select the correct option

Reload Math Equations

- (Each observation − Grand mean) 2
- Total number of Observations − Total number of Groups
- Total number of Groups − 1
- (Group mean-Grand mean)2 * Number of observation in each group

Click to Save Answer & Move to Next Question

inal text
this translation
feedback will be used to help improve Google Translate

Question # 5 of 10 (Start time: 09:14:28 PM, 07 July 2023) **Total Marks:** 1

According to Maloney et al., Most abundant natural predator is:

Select the correct option

Reload Math Equations

- Leopard
- Brown Bear
- Lion
- Spider

Click to Save Answer & Move to Next Question

inal text
this translation
feedback will be used to help improve Google Translate

Question # 6 of 10 (Start time: 09:15:45 PM, 07 July 2023) Total Marks: 1

In two-way ANOVA, there are

Select the correct option

Reload Math Equations

- Independent variables
- Depends upon situation
- One is dependent and other two variables are independent
- Dependent variables

Click to Save Answer & Move to Next Question

inal text
this translation
feedback will be used to help improve Google Translate

Question # 7 of 10 (start time: 09:16:37 PM, 07 July 2023) Total Marks: 1

Which one of the following is TRUE for ANOVA?

Select the correct option

Reload Math Equations

- Used only for numerical data.
- It is a Non-parametric test.
- Used to compare means of Less than two samples.
- Used only for Alpha-numeric data.

Click to Save Answer & Move to Next Question

inal text
this translation
feedback will be used to help improve Google Translate

Question # 8 of 10 (Start time: 09:17:34 PM, 07 July 2023) **Total Marks:** 1

Which one of the following is True?

Select the correct option

Reload Math Equations

- A one-way ANOVA uses one independent variable, while a Two-way ANOVA uses two dependent variables.
- A one-way ANOVA uses one dependent variable, while a Two-way ANOVA uses two independent variables.
- A one-way ANOVA uses one independent variable, while a Two-way ANOVA uses one dependent and one independent variable.
- A one-way ANOVA uses one independent variable, while a Two-way ANOVA uses two independent variables.

Click to Save Answer & Move to Next Question

inal text
this translation
feedback will be used to help improve Google Translate

Question # 9 of 10 (Start time: 09:18:27 PM, 07 July 2023) **Total Marks:** 1

Which one of the following is WRONG for ANOVA?

Select the correct option

Reload Math Equations

- Samples must be independent.
- Number of samples must be more than two.
- Heterogeneity of variances.
- Data should be normally distributed.

Click to Save Answer & Move to Next Question

inal text
this translation
feedback will be used to help improve Google Translate

Question # 10 of 10 (Start time: 09:19:19 PM, 07 July 2023) **Total Marks:** 1

For two sample t-test, the two populations:

Select the correct option

Reload Math Equations

- May be dependent or independent
- Must be dependent
- Vary according to situation
- Must be independent

Click to Save Answer & Move to Next Question

inal text
this translation
feedback will be used to help improve Google Translate

Question # 1 of 10 (Start time: 09:20:45 PM, 07 July 2023)

Which one of the following assumptions is TRUE for Two-Way ANOVA?

Select the correct option

- ○ Groups may or may not have equal sample sizes.
- ○ Samples must be independent.
- ○ The population must be Away from a normal distribution.
- ○ Population variances must be different.

Question # 2 of 10 (Start time: 09:21:33 PM, 07 July 2023)

Which one of the following is True?

Select the correct option

- A one-way ANOVA uses one independent variable, while a Two-way ANOVA uses one dependent and one independent variable.
- A one-way ANOVA uses one independent variable, while a Two-way ANOVA uses two dependent variables.
- A one-way ANOVA uses one dependent variable, while a Two-way ANOVA uses two independent variables.
- A one-way ANOVA uses one independent variable, while a Two-way ANOVA uses two independent variables.

Which one of the following is TRUE for ANOVA?

Select the correct option

- Used only for Alpha-numeric data.
- Used to compare means of Less than two samples.
- Used only for numerical data.
- It is a Non-parametric test.

Question # 4 of 10 (Start time: 09:23:09 PM, 07 July 2023)

For two sample t-test, the two populations:

Select the correct option

- ○ Must be independent
- ○ Vary according to situation
- ○ Must be dependent
- ○ May be dependent or independent

T-test is used to find out whether the means of two samples are statistically different from each other or not.

Select the correct option

- True
- False

Question # 6 of 10 (Start time: 09:24:11 PM, 07 July 2023)

---------- is positive square root of variance.

Select the correct option

- ○ Standard deviation
- ○ Statistical inference
- ○ Statistical dispersion
- ○ Confidence interval

Which one of the following is TRUE for T-Test?

Select the correct option

- ○ All randomly collected samples should have the different variances
- ○ Standard deviation must be known.
- ○ Sample size should be 50 or less than 50.
- ○ Samples should be taken randomly from the population

Question # 8 of 10 (Start time: 09:25:07 PM, 07 July 2023)

---------- is used to compare means of more than two samples.

Select the correct option

- T-Test
- Range
- ANOVA
- Z-Test

Question # 9 of 10 (Start time: 09:25:33 PM, 07 July 2023)

Which one of the following is WRONG for ANOVA?

Select the correct option

- Samples must be independent.
- Heterogeneity of variances.
- Data should be normally distributed.
- Number of samples must be more than two.

In two-way ANOVA, there are

Select the correct option

- One is dependent and other two variables are independent
- Depends upon situation
- Dependent variables
- Independent variables