Download More Files from VUAnswer.com

ZOO501 CURRENT FINAL TERM MCQS

1. Due to **Loss** in hydrogen ions, pH rises.
2. As a result of gastrulation, ectoderm, mesoderm and **Endoderm** are formed.
3. Male leopard **Frogs** make their sperm in summer.
4. **Differential gene transcription** is step for gene expression regulation.
5. Coding regions of DNA **exon.**
6. How many domains are present for transcription **3**
7. pdx1 is present in **Pancreatic** And Duodenal1 gene.
8. Bacteria transmit genes from one cell to other by **sex pili.**
9. Cells rearrange their **DNA** to create new immunoglobin and antigen receptor gene.
10. Sea urchins contain **25,000 to 50,000** mRNA.
11. In many species, sperm are attracted towards eggs by **chemotaxis**.
12. Acrosomal protein mediating recognition by protein **bindin.**
13. Sea water has **High** sodium ion conc.
14. Eggs have sparse **equally** spaced yolk are icolecithel.
15. Cytoplasm of blastodisc become embryo this meroblastic division called **discoidal.**
16. **The micromere** constitute major signaling center.
17. Osteoblasts secrete collagen proteoglycan **matrix** that is able to bind calcium salts
18. Tgf beta family secreted from the surface **ectoderm** and roof of neural tube.
19. Dendrites are numerous and do not extend far from the neuronal cell body or **soma,** axons may extend for several feet.
20. One agent of ventral specification is **sonic** hedgegog protein.
21. In frogs and chicks, **secondary** nueralation is seen in neural tube.
22. The cells at dorsal most portion of neural tube become **neural crest cells.**
23. About **33%** of identical twins have two complete and separate chorions
24. Mammals undergo holoblastic **rotational** cleavage.
25. In birds, **gravity** is critical in determining anterior posterior axis
26. Decidua becomes rich in blood vessels that will provide **oxygen** and nutrients to embryo.
27. Mammalian oocyte is released from ovary and swept by fimbriae into **the oviduct.**
28. Fertilization in chick egg occurs in **oviduct.**
29. Developmental decisions are usually **irreversible.**
30. All multicellular organisms arise by a slow process of progressive change called **development**.
31. Major class of cadherin includes **N- cadherin.**
32. Only **small %** of genome is expressed in partial cell.
33. Cortical are **granules** are membrane bound structure containing proteolytic enzymes.
34. Small subunits of MPF **cyclin-dependent kinase.**
35. It secretes a **cellulose** tunic (which gave the name "tunicates" to these creatures).
36. Inner surface cells thin 30×5um process **filopodia.**
37. β-catenin is Important for Inducing capacity of **micromeres.**
38. In fishes , First **12** divisions occur synchronously form a mouln of cells that sits an animal pole of large yolk cell.
39. Fertilization occurs in **ampulla** of oviduct a close, a region close to ovary.
40. In amphibians most of the xenopus most of tadpole is formed by **primary neurulation** and tail of neural tube formed by 2ndry neuralation.
41. Neural fold migrates to midline of embryo and fusing to form **neural tube** beneath the overlying ectoderm.

Join VU BIO ANIMALS

Download More Files from VUAnswer.com

42. **Tissues** form nervous system in the fly or in the frog.
43. The marginal zones to form clusters of neurons called **nuclei.**
44. Paracrine factors instruct **Myotome cells** to become muscles by inducing them to synthesize MyoD protein.
45. Sclerix is expressed in mesenchyme **from** the scleretome.
46. Acrosomal reaction in **sea urchins** is often highly specific to each species.
47. Cells are arrange their.. to creat new immunoglobin
48. In frogs and chicks, **secondary neurulation** is usually seen in the neural tube of the lumbar (abdominal) and tail vertebrae.
49. Regulation of gene expression by **several level.**
50. Which type of neuralation in fish **exclusively secondary**.
51. sonic hedgehocy is a type of **protein.**
52. **Blastomeres** have two parts of cell cycle.
53. Number of cortical granules Sea urchin **15000.**
54. How many strategies involve in cell commitment **2.**
55. Resting membrane potential of the egg is **70mv.**
56. One chemotactic molecule, a 14-amino acid peptide called **resact**. Download More Files from VUAnswer.com
57. Force for sperm propulsion is provided by **dynin.**
58. Regulative development arises from **Conditional Specification.**
59. Human brain contains **10^{12} glial cells.**
60. Fish have discoidal **telolicithal** pattern.
61. Process of osteogensis **2**
62. The mitotic spindle and contractile ring are **perpendicular** to each other.
63. Gastrulation in Snails is accomplished primarily by **epiboly.**
64. The first cell movement of fish gastrulation is the **epiboly** of the blastoderm.
65. This fetal organ the **Chorion** is derived primarily from embryonic trophoblast cells.
66. Secondary neurulation involves the making of a **medullary cord** and its subsequent hollowing into neural tube
67. The mantle zone cells differentiate into both **neurons** and **glia.**
68. The cerebral cortex in humans has **six** layers.
69. Paracrine factors such as **TGF** and FGF7 are important in normal skin development.
70. The **fifth** phase involves the invasion of the cartilage model by blood vessels.

71. Mode of commitment_____3___
72. strageries of commitment_____2___
73. Rana pipiens usually lays around __2500 eggs_____while the bullfrog, Rana catesbiana, can lay as many as 20,000
74. Multinucleated type of cell form._____syncytiotrophoblast
75. slow block, initiated by the intracellular release of_____ calcium ions
76. There are_____ two____ major modes of bone formation, or **osteogenesis**
77. Mutations of the ___*SOX9* gene____cause camptomelic dysplasia,
78. Human brain contains_____**10^{12}** glial cells_____
79. Telolecithal is cleavage is found in eggs of _____Birds and Fishes
80. ____Pax6 _____ plays a major role in eye formation
81. ___Pax1_____and Scleraxis are the transcription factors are thought to activate cartilage-specific genes
82. The eggs of insects have their yolk in the ____center_____ (i.e., they are centrolecithal),
83. Egg activates due to which ion more _____Ca_____
84. Which dermis give rise to brain _____Ectoderm_____
85. Which dermis give rise to eye _____Ectoderm_____
86. Mod of commitment in insects _____syncytial specification_____
87. Regulative development arises from_____ conditional specification
88. Nucleosome is composed of an octamer of histone proteins wrapped with two loops containing approximately ___140 base pairs___of DNA
89. the membrane potential of the sperm becomes ___more negative___ as potassium ions leave the sperm
90. Paracrine factors such as TGF- and FGF7 are important in normal _____skin development
91. In mouse ZP3 is attached to_____zona pellucid

Join VU BIO

Download More Files from VUAnswer.com

92. It is estimate ____egg is 25000 to 50000 mRNA.
Sea Urchin
93. Fertilin protein is important for sperm-egg fusion in Mammals
94. Fast block to polyspermy is initiated by Sodium
95. Transition from fertilization to cleavage is caused by factor MPF
96. Eggs having sparse and equally yolk space are called isolethial
97. Archenteron is used for primitive gut
98. Spiral cleavage found in Snails
99. Sonic hedgehog is a type of protein

1. Which type of neurulation in fish secondary
2. Nural crest divides functionally in domains 4
3. Myogenic bHLH proteins are specific for development of muscle cell
4. Zygote which divide mitotically to produce all the cells of the body.
5. Tatratology the study of birth defect.
6. Identical twins occur in roughly 0.25 of human births.
7. Requlatively development is seen in most vertebrate embroys , and it is obviously critical in development of identical twins.
8. Insect such as drosophila use all three mods of specification to commit their cells to particular fates.
9. The speed of metamorphosis is directly linked with environment pressures
10. In acetabularia unicellular organisms, development us controlled at both the transciptional and translational level.
11. The individuals of each species, however is divided into two mating types Plus and minus calamadomonas.
12. 140 base pairs of DNA nucleosome.
13. Viteline envelop contains at least eight different glycoproteins.
14. The first evidence suggestion the importance of sperm in reproduction came from a series experiment performed by Lazzaro Spallanzani
15. The volume of the sea urchin is about 200 Pico liters
16. Hindbrain Rhombencephalon
17. The most crucial, difference between mammalian cleavage and all other types involves the phenomenon of compaction
18. The embryo contains three germ layers. The ectoderm, the endoderm and the mesoderm.
19. The neural form keel fish
20. Spiral holoblastic clevage is characteristic of several animals groups, including **annelid** worms, some flatworms, and most mollusks (snails).
21. Diploblastic animals are those who have ectoderm and endoderm but no true mesoderm (hydra)
22. Even distribution of yolk in ovum isolecithal
23. Size of bindin protein 30,500-Da
24. which has two parts in cell cycle options Blastomere
25. After the blastoderm cells have covered about half the zebra fish yolk cell, a thickening occurs throughout the margin of the epibolizing blastoderm. This thickening is called the **germ ring**
26. Which forms spinal cord, hind brain and mid brain chordamesoderm.
27. This streak is first visible in birds as a thickening of the Epiblast at the posterior region of the embryo, just anterior to koller sickle.
28. It will induce the uterine cells of the mammals to from the matrenal portion of the placenta, the Decidua.
29. Between the area pellucida and the area opaca is a thin layer of cells called the **marginal zone** (or marginal belt).
30. The mitotic division of the nucleus is called karyokinesis
31. Fishes, reptiles , and birds undergo discoidal meroblastic cleavage, wherein the early cell divisions do not cut through the yolk of the egg.
32. The fast block, initiated by sodium influx into the cell in sea urchin.
33. The cells adjacent to the chordamesoderm, the paraxial mesoderm cells, are the precursors of the mesodermal somites in the fish.
34. The rudiment of the central nervous system, is called neurulation.
35. Blastula produced by radial cleavage has no blastocoel and are called stereoblastulae.
36. Fraternal twins are the result of two separate fertilization events.
37. The prechordal plate induces the formation of the forebrain; the chordamesoderm induces the formation of the midbrain, hindbrain, and spinal cord.

Download More Files from VUAnswer.com

38. The resulting larvae of snail is called _____glochidia_____which resemble tiny bear traps.
39. D
40. Fertilization in the absence of sex occur through the __Binary Fission__
41. The macromeres divide meridionally, forming a tier of eight (8) cells
42. cells, derived from the micromeres, are called the primary mesenchyme.
43. Molluscus follow which type of cleavage _____spiral holoblastic cleavage
44. A mammalian blastula is called a blastocyst.
45. Which of the following undergo discoidal cleavage __fishes, reptiles and birds
46. The egg jelly factors that initiate the Acrosomal reaction in sea urchins are often highly specific to each species
47. The Metencephalon gives rise to the cerebellum
48. In primary neurulation, the original ectoderm is divided into three sets of cells
49. The growing region at the tip of the lip is called the chordoneural hinge
50. Neural Tube will form ____Brain and spinal cord
51. Each ICM (Inner Cell Mass) cell has the same potency
52. Variable cleavages produce no invariant fate assignments to cells
53. In early embryos of these insects, cell division is not complete
54. The neural crest cells originate at the dorsal most region of the neural tube
55. The human β-globin gene has an enhancer in its 3′ UTR, roughly 700 base pairs
56. the SOX9 gene, which encodes a DNAbinding protein
57. the entire region of calcified spicules becomes surrounded by compact mesenchymal cells that form the Periosteum
58. The second step is the alignment of the myoblasts together into chains
59. Cleavages sphere is known as __Blastula
60. Osteoclasts are probably derived from the same precursors as macrophage blood cells
61. charby desired to know the cause of birth defects
62. The direct conversion of mesenchymal tissue into bone is called intramembranous ossification
63. Those cells that remain integral components of the neural tube lining become ependymal cells
64. The original type of trophoblast cells constitute a layer called the cytotrophoblast
65. The major structural characteristic of avian, reptilian, & mammalian gastrulation is the primitive streak
66. The Blastomere cell cycle is governed by the synthesis and degradation of_____cyclin
67. Both have Anurans __and _. __Toad and frog
68. They have a notochord as larvae (chordates), but they lack vertebrae.
69. Acrosomal vesicles is derived from Golgi apparatus and contains __Enzymes__
70. The large subunit of MPF is _cyclin B
71. ZP3 is the specific glycoprotein in the mouse
72. MITF necessary for the production of __Pigment cells
73. Mutation of sox9 gene cause ____camptomelic dysplasia
74. Between the blastoderm and the yolk is a space called the sub-germinal cavity
75. The second step of gastrulation involves the involution of the mesoderm
76. The first cleavage furrow appears centrally in the blastodisc
77. Fertilization occurs in the ampulla of the oviduct
78. Mother provides fetus with __nutrients and oxygen
79. The first segregation of cells within the inner cell mass results in the formation of the hypoblast (sometimes called the primitive endoderm)
80. The fetal portion of placenta formed by __Chorion__
81. Paracrine factor instruct myotome cells to become__ Muscles
82. When myoblasts stop dividing, secrete fibronectin

Join VU BIO ANIMALS

Download More Files from VUAnswer.com

83. Retinoic Acid comes from ___Somites___
84. Hox genes pattern the anterior-posterior axis and help to specify positions along that ____axis
85. The telencephalon will eventually form the cerebral hemispheres, and the diencephalon will form the thalamic and hypothalamic brain regions
86. Endochondral ossification involves the formation of cartilage
87. J.Hammerling exchanged nuclei b/w two morphologically distinct species
88. When plus individuals initiate fusion by extending a fertilization tube. Which contacts & fuses with a specific site on the minus individual.
89. Promoter is enhancer which is thought to regulate transcription.
90. Identical twins share a common __Chorion
91. In Birds,___pH____differences appear crucial for distinguishing dorsal from ventral
92. Which regulate function of Promoter ____Enhancer
93. The space between two neurons, where transmitter are released ___synapse
94. integral components of the neural tube lining become ependymal
95. Mantle cells differentiate into____ Gray matter
96. The peripheral ring of blastoderm cells that have not shed their deep cells constitutes the area __opaca.
97. During fertilization, the actin molecules fuse to form the long microfilaments.
98. A condition in which cell lack the instructions for development is ____naive
1. metamorphosis is initiated by hormones from the tadpole's thyroid gland
2. Brain has 100 billion neurons
3. Release of Ca from intracellular storage controlled by __dye ___Aequorin_
4. Nervous system develop by ____Neural Tube
5. Allows the mRNA to exit the nucleus____poly a tail
6. Contact and ____between sperm and egg ___Recognition
7. The first evidence suggesting the importance of sperm in reproduction came from a series of experiments performed by ____in the late 1700s. __Lazzaro Spallanzani
8. The early embryonic cells need a supply of energy and amino acids which is provided by yolk proteins in the egg.
9. The depression is called the primitive groove

ZOO501 CURRENT MCQS MIDTERM (2018) UPDATED

1. Red blood cells make_____Globins
2. how many strategies for developmental commitment _____2
3. Ribosomes produce in sea urchin _____10^{12}
4. Resact is isolated from egg jelly of _____sea urchin
5. ___is a discipline that studies embryonic & other developmental processes____Developmental Biology
6. 0nly___small___ %age of genome expressed in particular cell
7. Digestive & respiratory system formed by_____ Endoderm
8. Progesterone work for signals the egg to resume its _____meiotic division
9. Insect characteristics are ..... specification_____ syncytial
10. In Chlamydomonas haploid chromosome has.....copy _____one
11. Modes of commitments are..... _____ 3
12. After fertilization developmental process proceed by _____Cleavage
13. Ca ion responsible for cortical granules stores in _____Endoplasmic Reticulum of Eggs
14. Seawater has a particularly high ___ ion concentration _____Na (Sodium)
15. There are ___approaches of anatomy_____4(four)
16. The major cell adhesion molecules appear to be the Cadherins.
17. It is estimate ____egg is 25000 to 50000 mRNA. _____ *Sea Urchin*
18. Fusion of ____and sperm cell plasma membrane. _____ Egg
19. Resact diffuse readily in _____sea water.
20. The requirement of capacitance varies from species to _____Species.

Download More Files from VUAnswer.com

21. The division of embryo into 3 _____Germ layers
22. Cell potency___≥___fate
23. Zona pellucida_____ also initiate the acrosomal reaction after sperm bound to it.
24. Mammalian egg is surrounded by layer called. _____Cumulus
25. The volume of the sea urchin is about ___200_____ Picoliters.
26. The sperm travel by the amoeboid motion of ____ lamellipodial extensions___of the cell membrane.
27. The sex and __Reproduction_____ process included into fertilization.
28. The term spermatozoa means.. _____Sperm Animals
29. In 1824____J. L. Prevost and J. B. Dumasclaimed__ sperm is not parasite.
30. 4 cadherin classes are found in _____Vertebrates____ embryos. .
31. The second major group of protostomes are the _____Lophotrochozoa__
32. Sperm discovered in_____1678__
33. Transcription factors have ___three___ major domains
34. How many Amphibian oocytes produces ribosomes as_____10^{12}
35. Enhancers can also be used to inhibit _____transcription___
36. In deutrosomes the mouth opening would formed after... opening ____Anal__
37. Germs cell formed in...._____specialized portion of egg cytoplasm
38. Syncytial specification found in ____insects___
39. Third anatomical approach to developmental biology is ____Teratology___
40. The mature sea urchin is metabolically sluggish and activated by. _____Sperm__
41. The resting membrane potential is.....mv. _____70mV
42. Fusion is a __Active___process that is activated by fusogenic proteins.
43. Axoneme s formed by. _____Microtubules__
44. Eukaryotic genes are contained within a complex of DNA and protein called__Chromatin___
45. The negative enhancers are called____Silencers___
46. The ability of embryonic cells to change their fates is called.....___Regulation___
47. ____Pax6___transcription factor found in mammalian eye, nervous system and pancreas.
48. E- Cadherin is expressed on all early mammalian embryonic cells.
49. Annelids, molluscs and flatworms belong to..... ___ Lophotrochozoa
50. Potency___ of cell decrease if cell is committed.
51. Fertilization in Rana Pipiens usually lasts for about __3 (three)_ years.
52. An uncommitted cell can be described as ___Naive___
53. Lying immediately beneath the plasma membrane of the egg is a thin ___Shell__
54. Acroome can be considered as modified secretory_____ **vesicles**
55. Most marine organisms release their ___gametes___ into the environment
56. The fast block method is used to prevent_____Polyspermy__
57. Autonomous specification was first demonstrated in_____1887
58. In mouse ZP3 is attached to_____zona pellucid
59. TAFs used in_____transcription factor activity
60. in slow block method which is cortical granule protein_____ hyalin and peroxidase
61. Egg shell formed _____weight of egg._____9 – 12%
62. potency is ___intrinsic___ property
63. cytoplasm that contains many nuclei called _____ syncytium
64. Ectoderm forms..... __ epidermis, nervous system, and pigment cells.
65. according to dutch sperm are ...___ parasitic animals (spermatozoa)
66. Acrosome & nucleus constitute the head of the sperm.
67. cortical granules are _____membrane bound structures
68. Acrosomal reaction has 2 (two) component
69. Acrosome-reacted sperm transfer their binding from ZP3 to the adjacent ZP2 molecules
70. Ca & bicarbonate ions maybe critical in activating cAMP
71. Peroxidase enzyme hardens fertilization envelope
72. The Fast block to polyspermy is achieved by changing the electric potential
73. P-cadherin stands for _____Placental Cadherin
74. Most transcription factor can bind to specific DNA sequences
75. MITF protein necessary for production of pigments cell and their pigments.
76. Outside of plasma membrane is the vitelline envelope.
77. Mouse __**Fertilin**___ is localized to posterior plasma membrane of the sperm **head**
78. Finally the fourth cortical granules protein ____hyaline___ forms a coating around egg.

Join VU BIO ANIMALS

Download More Files from VUAnswer.com

79. The cleavage stages are called.____Blastomere
80. Enhancers are.....DNA Sequences
81. The mitotic spindle and contractile ring are perpendicular to each other
82. J.Whittaker stained blastomeres for the presence of the enzyme acetyl cholinesterase
83. Protocadherins are involved in separating notochord from mesoderm in __Xenopus gastrulation
84. Pax6 factor needed for mammalian eye
85. Which stands comparative embryology is ____First
86. Cell type specification precedes any large-scale embryonic cell migration
87. Series of extremely rapid divisions in zygote is ___Cleavage
88. In protosome, mouth is formed first than Deuterostomes: forms.
89. Northern blots, in situ hybridization, & the PCR can show which cells are transcribing particular genes.
90. Coding regions of DNA are called __Exon
91. Na and __K__ Are important in fast block polyspermy.
92. The force for sperm propulsion is provided by __dynein
93. Spallenazani demonstrated that filtered __toad__ semen devoid of sperm would not fertilize egg.
94. Mesoderm generates the____KIDNEY.
95. Sperms are attracted towards egg of their species by CHEMOTAXIS.
96. Mathematical modeling seeks to desire developmental phenomenon in terms of EQUATIONS.
97. A region that is not translated in proteins is called UNTRANSLATED region.
98. Invariant cleavage produce the ___SAME___ lineages in embryo
99. Germ layers are formed as a result of ___GASTRULATION
100. RNA polymerase binds with PROMOTOR in start of transcription
101. The region of globular actin molecule lies between nuclear and acrosomal vesicle in ___SEA URCHIN
102. Which technique is used in situ hybridization ___ Northern blots
103. Conditional specification involves interaction with RELATIVE POSITIONS cells
104. Particular proteins or carbohydrate on the sperm surface __LOST__ during capacitation
105. __LOSS___ of hydrogen ion causes pH to rise.
106. Peroxidase enzyme hardens the fertilization envelope by crosslinking TYROSINE residues
107. FERTILIZATION transmits genes from parents to offspring.
108. Regulation of gene expression accomplished by SELECTIVE MESSENGER RNA TRANSLATION
109. Development contact genotype and PHENOTYPE
110. Nucleosome is composed of OCTAMER of histone protein.

1. Knowledge of gene activity in humans can be obtained by candidate GENE MAPPING & POSITIONAL CLONING
2. Anatomical abnormalities are caused by MUTANT GENES
3. All multicellular organisms arise by a slow process of progressive change called DEVELOPMNET
4. Animals development is the division of embryo into three layers Ectoderm, Mesoderm and___ENDODERM
5. The ions which are responsible for increasing granule reactions____Ca (Calcium)
6. The zona pellucida is a fglycoprotein matrix, which is synthesized and secreted by the growing oocyte
7. In several other species, certain regions of the membrane are specialized for sperm recognition & fusion
8. An intrinsic program has been activated within the cell that causes it to follow a particular pathway of development
9. Activation of egg metabolism to start development
10. Insects such as Drosophila use all three modes of specification to commit their cells to particular fates
11. Nuclear control of cell morphogenesis and the interaction of nucleus and cytoplasm are well studied in Acetabularia.
12. Upon fertilization, the intracellular calcium ion concentration of the egg increases greatly
13. Bacteria are able to transmit genes from one individual to another by means of sex pili
10. +ve and – ve mating take place *Chlamydomonas*

Join VU BIO ANIMALS

Download More Files from VUAnswer.com