MUHAMMAD IMRAN

ZOO501-Developmental-Biology

Quiz No.2

Topics 85 to 119

1. This streak is first visible as a thickening of the …..at the posterior region of the embryo, just anterior to Koller's sickle_____ **epiblast**
2. The prechordal ……. induces the formation of the forebrain; the chordamesoderm induces the formation of the midbrain, hindbrain, and spinal cord_____ **plate**
3. discoidal meroblastic cleavage occur in _____ **birds and reptiles**
4. Between the area pellucida and the area opaca is a thin layer of cells called the …..or marginal belt _____ **marginal zone**
5. It will induce the uterine cells to form the maternal portion of the placenta, the ….. _____ **decidua**
6. Chorion is derived from ……. Cell_____ **trophoblast**
7. only the cytoplasm of the blastodisc becomes the embryo, this type of meroblastic cleavage is called discoidal present in _____ **fishes**
8. Neurulation in ……is exclusively secondary. _____**fishes**
9. The mammalian oocyte is released from the ovary and swept by the ……into the oviduct ._____ **fimbriae**
10. The cells adjacent to the chordamesoderm, the paraxial mesoderm cells, are the precursors of the in…… fishe_____ **mesodermal somites.**
11. The rudiment of the central nervous system, is called _____ **neurulation**
12. Fraternal twins are the result of ….separate fertilization events. _____**two**
13. Fishes, ……., and birds undergo discoidal meroblastic cleavage, wherein the early cell divisions do not cut through the yolk of the egg._____ **reptiles**

Best of luck