LEARNING COEFFICIENTS AND ACTIVE-BLOCK ARRANGEMENTS IN SCALAR RESIDUAL NETWORKS

BOBBY FABER

Abstract. We study squared-error population losses of scalar polynomial residual networks in their raw weights. At the identity target, odd activation degree permits cancellation and produces a depth-dependent transition, including a logarithmic factor at the critical depth. At an arbitrary realization, nonzero active blocks separate consecutive inactive groups: the local learning coefficient is the sum of a regular active contribution and the identity contributions of those groups. This gives realizations of the same function with different local exponents or multiplicities. An ordered-chain classification gives exact fiber sizes and a global formula for every realizable target. The fiber classification, counts, and local and global power/logarithmic Laplace orders are certified in Lean under the stated sparse Gram and prior assumptions.

1. Introduction

A depth-five cubic residual network represents $f(x) = x + x^3$ with its sole nonzero weight at any of its five positions, yet these realizations have different local loss geometry. We describe this geometry by a pair (λ, m): the local learning coefficient λ and multiplicity m give the order $N^{-\lambda}(\log N)^{m-1}$ of the local integral of e^{-NK_f}, where K_f is the squared-error population loss and $N \to \infty$ is a positive concentration parameter. At positions 1 or 5, the pair is $(13/12, 1)$; at positions 2 or 4, it is $(7/6, 2)$; at the center, it is $(7/6, 1)$. Thus the same function has local integral orders $N^{-13/12}$, $N^{-7/6}\log N$, and $N^{-7/6}$. Figure 1 and Table 1 display the three cases.

Learning coefficients capture singular geometry beyond parameter counting or Jacobian rank (Wei et al., 2023; Lau et al., 2025). Here every realization has five parameters and raw coefficient-Jacobian rank one, but the local pairs differ. A local analysis near one realization therefore need not describe the geometry near another realization of the same function. Equivalently, the local low-loss volume has order $\epsilon^{\lambda}(\log(1/\epsilon))^{m-1}$ as the tolerance $\epsilon \downarrow 0$: smaller λ gives more volume, and m distinguishes logarithmic scales when the exponents agree.

An inactive residual block is the identity, so moving the sole active block preserves the output. Perturbing inactive weights, however, introduces terms at different stages of composition. The active block separates the resulting cancellation opportunities.

For odd degree r, signed rth powers can cancel within a consecutive group of inactive blocks. Active blocks separate these groups into independent leading loss directions, so their positions determine the available cancellation regimes. A group's contribution changes regime when its length reaches r,

Date: 7 November 2026.

2020 *Mathematics Subject Classification.* 62F15; 14E15, 68T07.

Key words and phrases. singular learning theory, residual networks, real log canonical threshold, Newton polyhedra, Lean.

where a logarithm appears. Theorem 2 adds these group contributions to the regular contribution of the active weights. Theorem 3 classifies every target fiber and selects its global pair: local pairs depend on the inactive-group lengths, while the global pair depends only on degree, depth, and active count.

The architecture is the scalar monomial specialization of the polynomial ResNets studied by Dahlgren et al. (2026). Related work connects sparse subnetworks to singularities of polynomial function spaces (Shahverdi et al., 2026). Here we compute local loss-volume exponents, giving exact examples relevant to local learning coefficient estimation (Lau et al., 2025). Exact coefficients are also known for reduced-rank regression and multilayer linear networks (Aoyagi and Watanabe, 2005; Aoyagi, 2024). Our calculations use the singular learning framework and monomial rules of Watanabe (2009); Lin (2017); they concern deterministic population-loss integrals.

2. Model and assumptions

Fix integers $r \geqslant 2$ and $L \geqslant 1$. The parameters are independent raw weights $w \in \mathbb{R}^L$, with

$$h_0(x) = x, \qquad h_\ell(x) = h_{\ell-1}(x) + w_\ell^r h_{\ell-1}(x)^r, \quad 1 \leqslant \ell \leqslant L. \tag{1}$$

This is the bias-free scalar specialization of $h \mapsto h + \sigma(wh)$ with $\sigma(z) = z^r$ (Dahlgren et al., 2026, Definition 2.1), hence the coefficient w^r. Replacing it by an independent coefficient c changes the geometry at zero: $w \mapsto c = w^r$ is not a local diffeomorphism there, and raw Lebesgue measure induces a density proportional to $|c|^{1/r-1}$ on each branch. A smooth positive prior in c therefore defines a different local integral. Write $F_c(x) = x + cx^r$, so $h_L = F_{w_L^r} \circ \cdots \circ F_{w_1^r}$. The target f is realizable in this architecture. For an input probability measure μ, let

$$K_f(w) = \int_{\mathbb{R}} (h_L(w, x) - f(x))^2 \, d\mu(x). \tag{2}$$

Put $\nu = (r^L - 1)/(r - 1)$ and $d_j = 1 + j(r - 1)$ for $1 \leqslant j \leqslant \nu$. Throughout, assume the sparse feature Gram matrix

$$G_{ij} = \int_{\mathbb{R}} x^{d_i + d_j} \, d\mu(x) \quad (1 \leqslant i, j \leqslant \nu) \quad \text{is finite and positive definite.} \tag{3}$$

Then K_f is comparable to the squared coefficient-error norm. A finite absolute moment of order $2r^L$ and infinite support of μ suffice: a nonzero univariate polynomial has finitely many zeros.

At a realization w^*, let π be a smooth positive density on a full neighborhood. Take a nonnegative smooth compact cutoff χ, supported in a sufficiently small neighborhood and equal to one near w^*. Define the local pair by

$$Z_{N,w^*} = \int_{\mathbb{R}^L} \chi(w) e^{-N K_f(w)} \pi(w) \, dw \asymp N^{-\lambda} (\log N)^{m-1}, \qquad N \longrightarrow \infty. \tag{4}$$

The symbol $\asymp$ denotes bounds by positive constant multiples. For every loss treated below, λ is also the local integrability threshold: K_f^{-s} is locally integrable for $s < \lambda$ and diverges at $s = \lambda$. This needs no resolution. For $s > 0$ it follows from (4) through the Mellin identity

$\int K_f^{-s}\chi\pi\,dw = \Gamma(s)^{-1}\int_0^\infty t^{s-1}Z_{t,w^*}\,dt$, since the zero set of K_f meets a small $\operatorname{supp}\chi$ only in the null set $\{w^*\}$. For $s \leqslant 0$, integrability is trivial. The threshold statement is certified in Lean on small balls with bounded measurable densities (Section 7). The integer m records the logarithmic power, which our calculations determine separately. For this analytic loss, m is also the order of the first positive pole of the local zeta integral, by the usual resolution framework (Lin, 2017, Theorem 2.10 and Proposition 3.2). We use this pole reading only as an interpretation. Our normalization uses a *squared* loss; Lin's coefficient-ideal threshold is 2λ (his zeta convention is $(\sum_i f_i^2)^{-z/2}$). A measurable density bounded above and away from zero already suffices for the order in (4); smoothness supplies the customary pole interpretation.

All local constants may depend on r, L, w^* and μ; the estimates are pointwise in the nonzero active weights. Global statements use a nonnegative measurable density on $\mathbb{R}^L$ with finite mass, continuous and strictly positive at every point of the target fiber. Probability normalization and independence are unnecessary.

3. The identity-target phase transition

Write $K_0 = K_\chi$. Effective coefficients $u_i = w_i^r$ will be used only as algebraic notation: the integration variables remain the raw weights. Set

$$S(w) = \sum_{i=1}^{L} w_i^r, \qquad Q(w) = \sum_{i=1}^{L} w_i^{2r}.$$

LEAN ✓
SparseIdentityCertificate

Theorem 1 (Identity target). *Under (3), near $w = 0$,*

$$K_0(w) \asymp S(w)^2 + Q(w)^2. \tag{5}$$

The local pair is

$$(\lambda, m) = \begin{cases} (L/(2r), 1), & r \text{ even or } L < r, \\ (1/2, 2), & r \text{ odd},\ L = r, \\ ((L+r)/(4r), 1), & r \text{ odd},\ L > r. \end{cases} \tag{6}$$

Moreover, K_0 is coercive and has zero set $\{0\}$. Thus the same pair governs the global integral under a finite-mass prior continuous and positive at zero.

The first two output error coefficients are exactly S and

$$B = r\sum_{i<j} u_i u_j = \frac{r}{2}(S^2 - Q).$$

All higher coefficients have higher homogeneous degree in u. The Gram bound and this identity give (5); Appendix A supplies the uniform estimate and coercivity argument.

For even r, the summands of S are nonnegative, so $S^2 \asymp \|w\|^{2r}$ and the radial exponent is $L/(2r)$. For odd r and $L \geqslant 2$, write $w = \rho\theta$ with $\theta \in \mathbb{S}^{L-1}$. The angular function $\tau(\theta) = \sum_i \theta_i^r$ has a nonempty regular zero set. Near it, transverse coordinates $t = \tau(\theta)$ reduce the loss and volume to

$$\rho^{2r}(t^2 + \rho^{2r}), \qquad \rho^{L-1}\,d\rho\,dt\,d\eta \tag{7}$$

up to bounded positive factors. A finite collection of local charts with bounded positive densities suffices. In the thin region $|t| \leqslant \rho^r$, cancellation changes the radial scale to ρ^{4r} while the angular width contributes ρ^r. This gives $(L+r)/(4r)$. The competing exponent is $L/(2r)$; their crossing at $L = r$ produces one logarithm. Appendix B computes both exponents and the logarithm using two explicit sectors.

Remark (Newton polyhedra). The transition is where the Newton prediction stops agreeing with the pair. We use the Newton polyhedron, Newton distance and nondegeneracy of Varchenko (1976) and Arnold et al. (1988). The degree-$2r$ part of K_0 is $G_{11}S^2$, and the other terms have degree at least $3r$. The Newton polyhedron of K_0 at zero is therefore $\{\alpha \in \mathbb{R}^L_{\geqslant 0} : \sum_i \alpha_i \geqslant 2r\}$. Its Newton distance is $2r/L$. The diagonal meets the compact face $\sum_i \alpha_i = 2r$ in its relative interior, so the Newton prediction is $(L/(2r), 1)$. This prediction is always an upper bound for λ: $S^2 \leqslant LQ$ gives $K_0 \leqslant C\sum_i w_i^{2r}$ near zero, and that sum has the stated exponent.

- *Even* r, *or* $L = 1$. The face polynomial $G_{11}S^2$ has no zeros off the coordinate hyperplanes. K_0 is comparable to $\sum_i w_i^{2r}$, and the prediction is exact. The even comparison (10) below is likewise comparable to a sum of monomials. Its Newton distance, together with the face the diagonal meets, gives (11) directly (Lin, 2017).
- *Odd* r *and* $L \geqslant 2$. The face polynomial $G_{11}S^2$ vanishes to second order on $\{S = 0\} \cap (\mathbb{R}^*)^L$. That set is nonempty: it contains $(1, \dots, 1, -(L-1)^{1/r})$. So K_0 is Newton-degenerate for every $L \geqslant 2$. For $L < r$ the prediction still agrees with the pair, because the cancellation sector is dominated. At $L = r$ it misses the logarithm. For $L > r$ it overestimates λ, because the cancellation region $|t| \leqslant \rho^r$ lowers it to $(L+r)/(4r) < L/(2r)$.

4. The local arrangement theorem

Each inactive group inherits the identity-target geometry of Theorem 1; the active weights contribute regular directions. For an inactive group of size n, define $A_r(0) = 0$, $M_r(0) = 1$, and, for $n \geqslant 1$,

$$A_r(n) = \begin{cases} n/(2r), & r \text{ even}, \\ n/(2r) - (n-r)_+/(4r), & r \text{ odd}, \end{cases} \qquad M_r(n) = \begin{cases} 2, & r \text{ odd and } n = r, \\ 1, & \text{otherwise}. \end{cases} \tag{8}$$

Here $(a)_+ = \max(a, 0)$. These are the identity pairs of Theorem 1, with the empty group contributing an integral equal to one.

Fix w^* with exactly k nonzero weights, at positions $i_1 < \cdots < i_k$. Put $c_j = (w^*_{i_j})^r \neq 0$. There are $k+1$ consecutive inactive groups, of lengths $n_0, \dots, n_k$, before, between, and after the active blocks; $k + \sum_g n_g = L$. Write their raw weights as $a_g = (a_{g,1}, \dots, a_{g,n_g})$ and put

$$S_g = \sum_{i=1}^{n_g} a_{g,i}^r, \qquad Q_g = \sum_{i=1}^{n_g} a_{g,i}^{2r}.$$

The active deviations $\delta_j = w_{i_j}^r - c_j$ are valid local coordinates because $r(w_{i_j}^*)^{r-1} \neq 0$. Only these active coordinates are changed.

LEAN ✓
ArrangementCertificate

Theorem 2 (Local arrangement). *Let $f = h_L(w^*, \cdot)$ and assume (3), with any nonzero active values and any inactive group sizes $n_g \geqslant 0$. For odd $r \geqslant 3$, on a sufficiently small full neighborhood of w^*,*

$$K_f \asymp \sum_{j=1}^{k} \delta_j^2 + \sum_{g=0}^{k} (S_g^2 + Q_g^2). \tag{9}$$

For even $r \geqslant 2$ the comparison is

$$K_f \asymp \sum_{j=1}^{k} \delta_j^2 + \sum_{\text{inactive } i} |w_i|^{2r}. \tag{10}$$

Under the local prior conditions of Section 2, for every $k \geqslant 0$ with total depth $L \geqslant 1$,

$$\boxed{\lambda = \frac{k}{2} + \sum_{g=0}^{k} A_r(n_g), \qquad m = 1 + \sum_{g=0}^{k} \big(M_r(n_g) - 1\big).} \tag{11}$$

Independence of the retained directions follows by induction on active count: each new active block multiplies old directions by a polynomial coprime to the new active direction, while the new group directions have strictly higher degree.

Independence controls the retained coefficient directions, but cancellations can make those coefficients much smaller than the raw perturbation. A remainder estimate solely in the raw distance would then be insufficient: the omitted terms must be small relative to the retained coefficients themselves. Appendix C proves $\|E\| \leqslant C\varepsilon \|z\|$, where z collects the retained active and group coefficients and $\varepsilon \to 0$ throughout the raw neighborhood. Absorbing this error proves the loss comparisons. Their model integral factors on product boxes: active directions contribute $k/2$, and group logarithmic powers add. Bounded positive prior and Jacobian factors preserve the result, without requiring a product prior.

5. Same function, different local geometry

Take $r = 3$ and $f(x) = x + x^3$. Its complete fiber consists of the unit coordinate vectors e_j, $1 \leqslant j \leqslant L$: a composition with q nonzero blocks has degree 3^q, so there is exactly one active coefficient, which must equal one. Figure 1 shows how placement partitions the inactive weights. Table 1 applies (11) to all those points at depths four and five.




FIGURE 1. The three distinct local pairs for the depth-five cubic target $f(x) = x + x^3$. Boxes denote full residual blocks, labeled by raw weight; black boxes are active, and brackets mark consecutive inactive groups. The endpoint, one-position-inward, and center placements have group lengths $(0, 4)$, $(1, 3)$, and $(2, 2)$, respectively. Only the second has a group at the critical length $r = 3$, giving $m = 2$. By (11), positions 5 and 4 have the same pairs as 1 and 2; see Table 1.

Depth L	Active position j	Inactive lengths (n_0, n_1)	Local (λ, m)
4	1, 4	$(0, 3), (3, 0)$	$(1, 2)$
4	2, 3	$(1, 2), (2, 1)$	$(1, 1)$
5	1, 5	$(0, 4), (4, 0)$	$(13/12, 1)$
5	2, 4	$(1, 3), (3, 1)$	$(7/6, 2)$
5	3	$(2, 2)$	$(7/6, 1)$

TABLE 1. The same cubic target at every listed realization. Positions are counted in composition order.

At depth four, moving the active block changes the logarithmic factor. At depth five, it can change the exponent itself. All these points have raw coefficient-Jacobian rank one; that rank does not determine either pair. The theorem also permits more than one logarithm: two groups of length r give $m = 3$. Such a local statement need not describe the globally dominant realization.

REMARK (WHAT A LOCAL LEARNING COEFFICIENT SEES). Let $\varpi = \chi\pi\, dw$ be the local measure of (4). Write $Z(t) = \int e^{-tK_f}\, d\varpi$ and $E_t[K_f] = Z(t)^{-1} \int K_f e^{-tK_f}\, d\varpi$; here t plays the role of $n\beta$ for a tempered posterior. K_f is bounded on the compact support, so differentiating under the integral gives $(\log Z)' = -E_t[K_f]$. Integrating this and applying (4), for each fixed $t_0 > 0$,

$$\int_{t_0}^{T} E_t[K_f]\, dt = \lambda \log T - (m-1) \log\log T + O(1), \qquad T \to \infty. \tag{12}$$

Dividing by $\log T$ shows that the average of $tE_t[K_f]$ against dt/t tends to λ. The multiplicity enters only at the $\log\log T$ scale.

- At depth five, the endpoint and center placements differ by $1/12$ in λ.
- At depth five, positions 2 and 3 share $\lambda = 7/6$. The integral at position 2 minus the integral at position 3 is $-\log\log T + O(1)$.
- At depth four, all four placements share $\lambda = 1$.

Asymptotically, as $\log T \to \infty$ and with no rate, the averaged quantity therefore separates placements whose exponents differ. The value of λ alone does not separate placements that differ only in multiplicity.

At finite T, the correction $-(m-1)\log\log T/\log T$ and the uncontrolled $O(1)/\log T$ can exceed gaps of this size. Take position 2, with $(7/6, 2)$, against position 1, with $(13/12, 1)$. Even ignoring the $O(1)$ term, the averaged quantity orders them correctly only once $\log T > 46$.

This is a statement about the deterministic population integral. It does not cover the estimator of Lau et al. (2025), which uses the empirical loss and a Gaussian localization. Nor does it assert the pointwise limit $tE_t[K_f] \to \lambda$. That limit follows from the full resolution asymptotic expansion for analytic losses, but not from the two-sided bounds (4) used here.

6. Full fibers and global selection

The arrangement theorem describes one realization. The global integral combines the local contributions over the complete target fiber.

LEAN ✓
UnifiedNetworkCertificate

Theorem 3 (Full fibers and global selection)**.** *Assume* (3) *and* $0 \leqslant k \leqslant L$. *Let* $f = F_{c_k} \circ \cdots \circ F_{c_1}$, *with* $c_j \neq 0$ *and, for even* r, $c_j > 0$. *For* $k = 0$, *set* $f = x$. *The complete effective fiber consists of the ordered coefficients* $c_1, \dots, c_k$ *placed in any* k *positions, with all other coefficients zero. Its raw fiber* $\mathcal{F}_f = \{w : K_f(w) = 0\}$ *has size*

$$\#\mathcal{F}_f = \begin{cases} \binom{L}{k}, & r \text{ odd}, \\ 2^k\binom{L}{k}, & r \text{ even}. \end{cases}$$

For every prior satisfying the global conditions of Section 2,

$$\int_{\mathbb{R}^L} e^{-nK_f(w)}\pi(w)\,dw \asymp n^{-[k/2+A_r(L-k)]}(\log n)^{M_r(L-k)-1}. \tag{13}$$

Repeated coefficients and opposite signs are included whenever realizable.

The classification follows by removing outer active blocks successively. For normalized polynomials H, G with constant coefficient zero and linear coefficient one, equality $H + cH^r = G + dG^r$ with $c, d \neq 0$ forces $H = G$ and $c = d$. A leading-coefficient and degree argument proves this recovery step (Appendix D).

The identity loss is coercive, and the $L^2(\mu)$ triangle inequality gives

$$\sqrt{K_f(w)} \geqslant \sqrt{K_0(w)} - \|f - x\|_{L^2(\mu)}.$$

Thus K_f has a positive gap outside small neighborhoods of its finite fiber, making the complementary integral exponentially small. Among local contributions, minimize λ, then maximize m among ties. For $n = L - k$, the inequality $\sum_g (n_g - r)_+ \leqslant (n - r)_+$ gives (13) (Appendix D). The cubic examples have global pairs $(1, 2)$ and $(13/12, 1)$; a prior positive near another realization and supported in a sufficiently small neighborhood of it instead retains its local pair.

Scope. The results cover every active count in scalar, bias-free polynomial residual networks and concern deterministic population-loss integrals. Priors are in raw weights: a smooth positive prior in effective coefficients has different geometry at zero. Higher-dimensional, biased, and more general architectures are outside the proved scope.

7. Formal verification

The pairs, fibers and counts of all three theorems are formalized in Lean 4 for localizations on small balls with bounded measurable densities. The transfer to the smooth cutoffs of (4) is on paper. The repository records the certified statements and their exact assumptions (Appendix E).

The unified certificate covers all $r \geqslant 2$, positive depths, and active counts: complete raw fibers and exact counts, local arrangement pairs, and the global pair. It uses (3) and Section 2's global prior conditions; standalone local certificates allow measurable priors bounded above and away from zero. Inverse-power criteria cover all real exponents locally and positive exponents globally. Lean also certifies endpoint divergence and two-sided Laplace bounds with the stated logarithmic powers. The zeta-pole interpretation uses the cited resolution theory; meromorphic continuation and leading asymptotic constants are not part of the Lean certificate. Neither are the two remarks: the Newton-polyhedron comparison in Section 3 and the averaged identity (12). Their comparisons with Theorems 1 and 2 are proved here, and the Newton-polyhedron framework is cited.

Appendix A. Coefficient reduction, zeros, and coercivity

Write $h_\ell = xT_\ell(x^{r-1})$. The exact recursion

$$T_0(y) = 1, \qquad T_\ell(y) = T_{\ell-1}(y) + u_\ell y T_{\ell-1}(y)^r$$

shows that $h_L - x = \sum_{j=1}^{\nu} P_j(u)x^{d_j}$, where P_j is homogeneous of degree j. Comparing the first two coefficients gives $P_1 = S$ and $P_2 = B = r(S^2 - Q)/2$. At depth one the absent P_2 is set to zero, consistently with $Q = S^2$.

In a fixed small ball in effective variables, $|P_j(u)| \leqslant C_j \|u\|^j \leqslant C_j Q$ for $j \geqslant 2$. Hence the Gram upper bound gives $K_0 \leqslant C(S^2 + Q^2)$. Conversely, with $\gamma > 0$ the smallest eigenvalue of G,

$$K_0 \geqslant \gamma(S^2 + B^2), \qquad Q = S^2 - 2B/r \leqslant S^2 + 2|B|/r. \tag{14}$$

Locally $S^4 \leqslant CS^2$, so $Q^2 \leqslant C(S^2 + B^2)$. This proves (5), including depth one. Globally (14) gives

$$Q \leqslant K_0/\gamma + (2/r)\sqrt{K_0/\gamma}.$$

Since $Q = \sum_i w_i^{2r} \to \infty$ as $\|w\| \to \infty$, K_0 is coercive. Also $K_0 = 0$ implies $S = B = Q = 0$, hence $w = 0$. For a finite-mass prior the integral outside any sufficiently small fixed neighborhood is therefore $O(e^{-bN})$ for some $b > 0$.

Appendix B. Angular sectors and the logarithmic factor

For even r, $S \asymp \|w\|^r$ and (5) reduces the integral to an ordinary radial integral with density ρ^{L-1} and loss ρ^{2r}. The pair is $(L/(2r), 1)$. At depth one the loss is exactly a positive constant times w^{2r} for either parity.

Suppose r is odd and $L \geqslant 2$, and use the Euclidean sphere. Let $\tau(\theta) = \sum_i \theta_i^r$, $q(\theta) = \sum_i \theta_i^{2r}$. Compactness gives $0 < c \leqslant q \leqslant C$. The set $\Sigma = \{\tau = 0\}$ is nonempty, since it contains $(2^{-1/2}, -2^{-1/2}, 0, \dots, 0)$. At a point of Σ, Euler's identity gives $\theta \cdot \nabla\tau = r\tau = 0$, whereas $\nabla\tau \neq 0$. Thus the tangential derivative is nonzero. The inverse function theorem supplies smooth local coordinates (t, η) with $t = \tau$, and surface density bounded above and below on smaller coordinate boxes. A finite collection of these boxes covers a neighborhood of the compact set Σ. On the remaining compact angular set, $|\tau|$ has a positive lower bound.

Restrict each chart to the inverse image of a product box $|t| < b$, $\eta \in V$ contained in its image; finitely many smaller boxes cover Σ, and bounded larger boxes give upper bounds. The smaller boxes give lower bounds, while the unused coordinates contribute finite positive volume (one in dimension zero). For radial cutoff $R > 0$, the product domain lies between $(0, \rho_0) \times (-\rho_0^r, \rho_0^r)$ and $(0, \rho_1) \times (-\rho_1^r, \rho_1^r)$, where $\rho_0 = \min(R, b^{1/r})$ and $\rho_1 = \max(R, b^{1/r})$. These inclusions justify the following model integrals.

The reduced loss equals $\rho^{2r}(t^2 + \rho^{2r}q^2)$ and is comparable to (7). In a product box $0 < \rho < \rho_0$, $|t| < \rho_0^r$, split into two sectors for each sign $\sigma \in \{-1, 1\}$:

$$\begin{aligned}
&\mathcal{C}: \quad t = \sigma\rho^r\nu, \quad 0 < \nu < 1, && \rho^{L-1}\,d\rho\,dt = \rho^{L+r-1}\,d\rho\,d\nu,\\
&\mathcal{O}: \quad t = \sigma z^r, \quad \rho = zy, && \rho^{L-1}|d\rho\,dt| = rz^{L+r-1}y^{L-1}\,dz\,dy,\\
&\qquad 0 < z < \rho_0,\ 0 < y < 1.
\end{aligned}$$

Apart from boundaries of measure zero these are exactly $|t| < \rho^r$ and $\rho^r < |t|$. The loss is comparable to ρ^{4r} in $\mathcal{C}$ and to $z^{4r}y^{2r}$ in $\mathcal{O}$. Sector $\mathcal{C}$ has pair

$((L+r)/(4r),1)$. For $\mathcal{O}$, substitute $a = z^{4r}$ and $b = y^{2r}$. Up to constants its integral is

$$\int_0^{\rho_0^{4r}} \int_0^1 e^{-Nab} a^{\alpha-1} b^{\beta-1}\, db\, da, \qquad \alpha = \frac{L+r}{4r}, \quad \beta = \frac{L}{2r}.$$

The inner integral is comparable to $\min\{1, (Na)^{-\beta}\}$. Splitting at $a = N^{-1}$ gives order $N^{-\min(\alpha,\beta)}$, with a factor $\log N$ exactly when $\alpha = \beta$. The same pair follows directly from the integer monomial $z^{4r}y^{2r}$ with density $z^{L+r-1}y^{L-1}$ by Lin (2017, Proposition 2.1). The noncancellation region contributes $O(N^{-\beta})$, so it does not change the resulting pair. The same integrals with the exponential replaced by the negative power diverge when either exponent reaches its critical value. Positive contributions from the chart boxes cannot cancel.

Finally, if $cF \leqslant K \leqslant CF$, then the corresponding Laplace integrals satisfy $Z_F(CN) \leqslant Z_K(N) \leqslant Z_F(cN)$ for the same positive amplitude. Thus both the exponent and logarithmic power transfer to the actual loss, not just the power-integrability threshold. For the pole interpretation, replace the cutoff by a small closed box around the isolated zero, on which the density is smooth and positive. The difference is supported away from the zero and contributes an exponentially small Laplace term and a zeta term holomorphic near the positive real axis. Resolution on that box supplies the stated interpretation.

Appendix C. Proof of the local arrangement theorem

We first identify independent leading coefficient directions. We then control the higher-order terms uniformly, including along cancellation paths. Finally, we transfer the coefficient comparison to the population loss and factor the local integral.

Fix $r, k, (n_g)$ and nonzero active coefficients $c_1, \dots, c_k$. Constants in the coefficient estimates depend only on these fixed data. Use coefficient ℓ^1 norms on polynomial spaces and sum norms on their direct sums; use the Euclidean norm on each effective group vector $u_g = (a^r_{g,i})_i$. Norm equivalence will be used only in fixed finite dimension.

The auxiliary composition map. Set $d_h = 1 + h(r-1)$, $\nu_g = (r^{n_g} - 1)/(r-1)$, and

$$\mathcal{P}_g = \operatorname{span}\{x^{d_h} : 1 \leqslant h \leqslant \nu_g\}, \qquad \mathcal{P} = \mathbb{R}^k \oplus \bigoplus_{g=0}^{k} \mathcal{P}_g.$$

Thus $\mathcal{P}_g = \{0\}$ for an empty group and $\mathcal{P}_g = \operatorname{span}\{x^r\}$ for a singleton. For $\nu = (\delta, p_0, \dots, p_k) \in \mathcal{P}$, define

$$\Psi(\nu) = (\mathrm{id} + p_k) \circ F_{c_k+\delta_k} \circ \cdots \circ (\mathrm{id} + p_1) \circ F_{c_1+\delta_1} \circ (\mathrm{id} + p_0). \tag{15}$$

For $k = 0$ this means $\Psi(p_0) = \mathrm{id} + p_0$. This is a polynomial map into the finite-dimensional space of polynomials of degree at most r^L: degrees multiply, and $k + \sum_g n_g = L$. Every output has linear term x, and its other degrees are 1 modulo $r-1$, so all output differences lie in the sparse feature space of (3). Let $\mathcal{H}_g$ denote the composition of inactive group g. For its error $p_g = \mathcal{H}_g - x$, $\Psi(\nu) = h_L$ and $\Psi(0) = f$.

The differential can be identified exactly. Put $\widehat{H}_0 = x$, $\widehat{H}_j = F_{c_j} \circ \widehat{H}_{j-1}$, and $T_j = F_{c_k} \circ \cdots \circ F_{c_{j+1}}$, with $T_k = \mathrm{id}$. Define the fixed polynomials

$$U_j = (T_j' \circ \widehat{H}_j)\widehat{H}_{j-1}^r, \quad V_g = (T_g' \circ \widehat{H}_g)\widehat{H}_g^r, \quad W_g = (T_g' \circ \widehat{H}_g)\widehat{H}_g^{2r-1}.$$

The chain rule gives, for arbitrary auxiliary group perturbations $q_g \in \mathcal{P}_g$,

$$D\Psi(0)[\alpha, q_0, \dots, q_k] = \sum_{j=1}^{k} \alpha_j U_j + \sum_{g=0}^{k} (T_g' \circ \widehat{H}_g)(q_g \circ \widehat{H}_g). \tag{16}$$

In particular V_g and W_g are the columns for the first two group coefficients when those coefficients exist.

Independence for $r \geqslant 3$. First consider the virtual family U_j, V_g, W_g in $\mathbb{R}[x]$, including both directions at every group. At $k = 0$ the family is x^r, x^{2r-1}. Suppose the family at j active blocks is independent and has degrees at most $(2r-1)d$, where $d = r^j$ is the degree of the partial output H. Appending F_e, $e \neq 0$, multiplies all old columns by $P = 1 + reH^{r-1}$ and adds the active column H^r and the last-group columns $\widetilde{H}^r, \widetilde{H}^{2r-1}$, where $\widetilde{H} = H + eH^r$.

The multiplied old degrees are at most $(3r-2)d$; the last two degrees are $r^2 d$ and $r(2r-1)d$, strictly greater and distinct for $r \geqslant 3$. Their coefficients in any relation therefore vanish. The remaining relation is $PR + tH^r = 0$. Since $P - reH^{r-1} = 1$, P and H^r are coprime. The polynomial P is nonconstant, so $P \mid tH^r$ forces $t = 0$, then $R = 0$. This proves independence and preserves the degree bound. Delete both group columns for an empty group and W_g for a singleton. The remaining columns are independent and, by (16), belong to the actual sparse output space. The virtual columns need not belong to that space and are used only to prove independence before restriction.

Uniform control of omitted coefficients. Each actual inactive group has

$$\mathcal{H}_g = x + S_g x^r + B_g x^{2r-1} + R_g, \qquad B_g = \frac{r}{2}(S_g^2 - Q_g), \qquad \|R_g\| \leqslant C_g \|u_g\|_2 Q_g. \tag{17}$$

Indeed its coefficient of x^{d_h} is homogeneous of degree h in u_g. On $\|u_g\|_2 \leqslant 1$, the sum of the finitely many terms with $h \geqslant 3$ is bounded by $C_g \|u_g\|_2^3$, and $Q_g = \|u_g\|_2^2$. Empty and singleton groups have $R_g = 0$; also $B_g = 0$ for a singleton.

For odd r, let z consist of the active deviations, S_g for $n_g \geqslant 1$, and B_g for $n_g \geqslant 2$. Let J insert these scalars as the corresponding auxiliary coordinates; its norm is one. With

$$\varepsilon = \sum_j |\delta_j| + \sum_g \|u_g\|_2, \qquad R = (0, R_0, \dots, R_k),$$

we have the *exact identity* $v = Jz + R$. Shrink the neighborhood so $\varepsilon \leqslant 1$ and $|S_g| \leqslant 1$ for every group. Since

$$Q_g \leqslant S_g^2 + \frac{2}{r}|B_g| \leqslant |S_g| + \frac{2}{r}|B_g|,$$

(17) gives fixed constants $C_0, C_1 > 0$ such that

$$\|z\| \leqslant C_0 \varepsilon, \qquad \|R\| \leqslant C_1 \varepsilon \|z\|. \tag{18}$$

The first bound follows also directly from $|S_g| \leqslant \sqrt{n_g}\,\|u_g\|_2$ and $|B_g| \leqslant C\,\|u_g\|_2^2$. No division by S_g, B_g, or $\|z\|$ is involved, so these bounds include simultaneous cancellations and $z = 0$.

Lemma C.1 (Taylor estimate with a controlled remainder)**.** *Let Ψ be a C^2 map between finite-dimensional normed spaces on a neighborhood of the closed unit ball. Let J be linear and suppose $v = Jz + R$, $0 \leqslant \varepsilon \leqslant 1$, and (18) holds. Set $M = \|J\| + C_1$, and assume $MC_0\varepsilon \leqslant 1$. If H_* bounds $\left\|D^2\Psi\right\|$ on that ball, then, with $A = D\Psi(0)J$,*

$$\Psi(v) - \Psi(0) = Az + E, \qquad \|E\| \leqslant \left(\|D\Psi(0)\|\,C_1 + \frac{H_*}{2}M^2C_0\right)\varepsilon\,\|z\|\,. \tag{19}$$

Proof. We have $\|v\| \leqslant M\,\|z\| \leqslant MC_0\varepsilon \leqslant 1$. Taylor's formula with its integral remainder gives the exact decomposition

$$E = D\Psi(0)R + \int_0^1 (1-t)D^2\Psi(tv)[v,v]\,dt.$$

The first term is bounded by $\|D\Psi(0)\|\,C_1\varepsilon\,\|z\|$. The second is bounded by

$$(H_*/2)\,\|v\|^2 \leqslant (H_*/2)M^2\,\|z\|^2 \leqslant (H_*/2)M^2C_0\varepsilon\,\|z\|\,.$$

The integral includes all mixed terms of the composition map, involving active deviations, group coefficients, or both. □

Apply the lemma to (15); a finite H_* exists by compactness because Ψ is polynomial. The retained-column independence gives $a > 0$ with $\|Az\| \geqslant a\,\|z\|$. Let C_* be the parenthesized constant in (19). Choose ε small enough that $MC_0\varepsilon \leqslant 1$ and $C_*\varepsilon \leqslant a/2$, in addition to the preceding group bounds. Then

$$\frac{a}{2}\,\|z\| \leqslant \|h_L - f\| \leqslant (\|A\| + a/2)\,\|z\|\,.$$

All these restrictions hold on a full neighborhood of the fixed raw realization because $\varepsilon \to 0$ there. Norm equivalence, (17), and the sparse Gram bounds give

$$K_f \asymp \sum_j \delta_j^2 + \sum_g (S_g^2 + B_g^2) \asymp \sum_j \delta_j^2 + \sum_g (S_g^2 + Q_g^2).$$

For the last comparison use $Q_g \leqslant S_g^2 + 2|B_g|/r$ and $B_g = r(S_g^2 - Q_g)/2$ with bounded S_g.

Even degree, including $r = 2$. Now $u_g \geqslant 0$, so $Q_g \leqslant S_g^2$. Retain $z = (\delta, S_g : n_g \geqslant 1)$ and set $R_g' = \mathcal{H}_g - x - S_g x^r$. Homogeneity of all omitted coefficients, now of degree at least two, gives

$$\left\|R_g'\right\| \leqslant C_g Q_g \leqslant C_g S_g^2 \leqslant C_g\sqrt{n_g}\,\varepsilon S_g.$$

Thus $v = Jz + (0, R_0', \ldots, R_k')$ obeys the same lemma. The retained family U_j, V_g is independent by the previous induction with old degree bound rd, multiplied bound $(2r-1)d$, and new last-group degree $r^2 d > (2r-1)d$ for every $r \geqslant 2$; coprimality separates the new active column. No two-direction independence is used at $r = 2$. Absorption yields $K_f \asymp \sum_j \delta_j^2 + \sum_g S_g^2$. Finally

$$\sum_{i\text{ in }g} |w_i|^{2r} \leqslant S_g^2 \leqslant n_g \sum_{i\text{ in }g} |w_i|^{2r}$$

for each nonempty group, proving (10).

Integration. The Jacobian determinant of the active coordinate map is $\prod_j r(w_{i_j}^*)^{r-1} \neq 0$. Its inverse absolute Jacobian and the prior are bounded above and away from zero locally. Inner and outer product boxes in $(\delta, a_0, \dots, a_k)$ bound the cutoff integral. On each box the comparison integral factors into Gaussian active integrals and identity-group integrals. Theorem 1, or the usual addition rule (Lin, 2017, Proposition 3.7), gives (11): exponents add and $m - 1 = \sum_g (M_r(n_g) - 1)$. Empty groups contribute one. All comparison losses vanish only at the base point, so shrinking cutoffs does not change the pair. Integration uses $(\delta, a_0, \dots, a_k)$; z records coefficients rather than raw coordinates.

Appendix D. Full fibers and global selection

Ordered recovery. Every active block has degree r, so a network with q active blocks has degree r^q. Equal outputs therefore have the same active count. To recover their coefficients, suppose

$$H + cH^r = G + dG^r, \qquad c, d \neq 0,$$

where $H(0) = G(0) = 0$ and $H'(0) = G'(0) = 1$. Equality of output degrees gives $\deg H = \deg G = D \geqslant 1$. Choose $a \neq 0$ so that $U = aG$ and H have the same leading coefficient. Comparing leading output coefficients gives $d = ca^r$, hence

$$c(H - U) \sum_{i=0}^{r-1} H^i U^{r-1-i} = G - H.$$

The sum has degree $(r-1)D$: its leading coefficient is r times the $(r-1)$st power of the common leading coefficient. If $H - U \neq 0$, its zero constant coefficient forces degree at least one. The left side would then have degree greater than D, while the right side has degree at most D. Thus $H = U$; the linear coefficients give $a = 1$, and consequently $H = G$ and $c = d$. Iterating this step recovers the ordered active chain, including repeated or opposite coefficients. The empty chain gives the identity.

Fiber size. The positive-definite feature Gram makes zero population loss equivalent to polynomial equality. The recovered chain can occupy any k of the L positions, in its fixed order. Each placement has one raw realization for odd r and 2^k independent sign choices for even r. Different supports or signs give different vectors, even when active coefficients repeat. This proves the stated count, including the singleton identity fiber at $k = 0$.

Global selection. Coercivity from Appendix A and the L^2 triangle inequality make every sublevel set of K_f compact. Delete disjoint small neighborhoods of the finite fiber. Choose a large ball outside which $K_f \geqslant 1$. Within the ball, the complement of the open fiber neighborhoods is compact and zero-free, so continuity gives a positive minimum whenever that complement is nonempty. Set $b = 1$ if that complement is empty; otherwise take the smaller of its minimum and one. Then $K_f \geqslant b > 0$ on the entire remaining set, so the prior-weighted tail is at most $e^{-bN} \int \pi$. Continuity and positivity of the prior at each fiber point give local upper and positive lower bounds. Each fiber neighborhood therefore has the pair of Theorem 2 with a positive contribution. A finite sum of terms $N^{-\lambda_i}(\log N)^{m_i - 1}$ has the smallest λ_i and then the largest m_i among ties. Several realizations with that same pair affect the multiplicative constants, not the logarithmic power; multiplicities do not add across disjoint neighborhoods.

For even r, every placement has pair $(k/2 + (L-k)/(2r), 1)$. For odd r, write $n = L - k = \sum_g n_g$. Then

$$\sum_g A_r(n_g) = \frac{n}{2r} - \frac{1}{4r} \sum_g (n_g - r)_+, \qquad \sum_g (n_g - r)_+ \leqslant (n - r)_+.$$

If $n < r$, all placements have the same exponent and $m = 1$. If $n = r$, all exponents agree and concentrating the inactive blocks in one gap gives $m = 2$,

the largest multiplicity. If $n > r$, the inequality is strict whenever at least two gaps are nonempty: if any gap exceeds r, the other nonempty gaps strictly reduce the sum of positive parts from $n - r$; if none does, that sum is zero. Hence concentration in one gap uniquely describes the minimizing patterns, and these have $m = 1$. Such a placement exists before, between, or after the ordered active blocks. When $n = 0$ these descriptions coincide. This proves (13), without requiring adjacent active blocks or comparing leading constants.

Appendix E. Lean source and audit record

The proof source is available at https://github.com/bubskee/residual-slt-in-a-haystack. The reproducibility snapshot, containing the proof source and audit, is

`2865e70a9ca7f559dabfce6aa32e13cf4fc88259`

It uses Lean `v4.35.0` and mathlib commit

`56a005855ea17e0b0958f73ec801ae140b0a1679`

The principal declarations, all in namespace `ResidualSLT`, are

- `unified_residual_network_certificate`: full fibers, exact counts, and local/global learning pairs;
- `sparse_identity_network_rlct_laplace_certificate`: the identity pair;
- `arrangement_raw_network_rlct_laplace_certificate_of_depth_pos`: the local arrangement pair;
- `network_full_raw_fiber`: ordered active-chain classification;
- `raw_network_fiber_cardinality`: exact raw-fiber size.

Exact hypotheses are recorded in `paper/formalization_status.md`. From the repository root, run:

```
git checkout 2865e70a9ca7f559dabfce6aa32e13cf4fc88259
cd lean
lake exe cache get
lake build
lake env lean -DwarningAsError=true -DautoImplicit=false \
  ../paper/checks/NetworkAudit.lean
```

The audit prints the certificate definitions and the types and axiom dependencies of these five declarations. Every dependency set is exactly `propext`, `Classical.choice`, and `Quot.sound`; none contains `sorryAx`. The logs are in `paper/checks/`: `lean-build.txt` and `NetworkAudit.txt`.

Tool and computational resource disclosure. Following the disclosure recommendation of the Leiden Declaration on Artificial Intelligence and Mathematics (Alper et al., 2026), we report substantial AI assistance. Astra and Opus 5.5 assisted with writing, proof development, and Lean formalization. Fable 5.1 provided adversarial review and attempted to find errors in the arguments. The author retains responsibility for the results, arguments, and citations.

Acknowledgments. The author thanks BlueDot Impact for financial support.

References

Jarod Alper, Michael J. Barany, Alain Chavarri Villarello, Sander Dahmen, Walter Dean, Karthik Ganapathy, Michael Harris, David Holmes, Mateja Jamnik, Steven Kelk, Bryna Kra, Ursula Martin, Bartosz Naskręcki, Rodrigo Ochigame, Jim Portegies, and Johannes Schmitt. Leiden Declaration on Artificial Intelligence and Mathematics. Zenodo, 2026. URL https://doi.org/10.5281/zenodo.20302944. 2 June 2026.

Miki Aoyagi. Consideration on the learning efficiency of multiple-layered neural networks with linear units. *Neural Networks*, 172:106132, 2024. doi: 10.1016/j.neunet.2024.106132.

Miki Aoyagi and Sumio Watanabe. Stochastic complexities of reduced rank regression in Bayesian estimation. *Neural Networks*, 18(7):924–933, 2005. doi: 10.1016/j.neunet.2005.03.014.

V. I. Arnold, S. M. Gusein-Zade, and A. N. Varchenko. *Singularities of Differentiable Maps, Volume II: Monodromy and Asymptotics of Integrals*. Birkhäuser, Boston, 1988.

Isabel Dahlgren, Giovanni Luca Marchetti, and Kathlén Kohn. Polynomial residual networks are generically identifiable. arXiv:2610.33738v1 [cs.LG], 2026. URL https://arxiv.org/abs/2610.33738v1. 24 October 2026.

Edmund Lau, Zach Furman, George Wang, Daniel Murfet, and Susan Wei. The local learning coefficient: A singularity-aware complexity measure. In *Proceedings of the 28th International Conference on Artificial Intelligence and Statistics*, volume 258 of *Proceedings of Machine Learning Research*, pages 244–252. PMLR, 2025. URL https://proceedings.mlr.press/v258/lau25a.html.

Shaowei Lin. Ideal-theoretic strategies for asymptotic approximation of marginal likelihood integrals. arXiv:1003.5338v3 [stat.CO], 2017. URL https://arxiv.org/abs/1003.5338v3. Version 3, 13 February 2017; first posted 2010.

Vahid Shahverdi, Giovanni Luca Marchetti, and Kathlén Kohn. Learning on a razor's edge: Identifiability and singularity of polynomial neural networks. arXiv:2505.11846v3 [cs.LG], 2026. URL https://arxiv.org/abs/2505.11846v3. Version 3, 11 June 2026; published at ICLR 2026.

A. N. Varchenko. Newton polyhedra and estimation of oscillating integrals. *Functional Analysis and Its Applications*, 10(3):175–196, 1976.

Sumio Watanabe. *Algebraic Geometry and Statistical Learning Theory*, volume 25 of *Cambridge Monographs on Applied and Computational Mathematics*. Cambridge University Press, 2009. ISBN 9780521864671. doi: 10.1017/CBO9780511800474.

Susan Wei, Daniel Murfet, Mingming Gong, Hui Li, Jesse Gell-Redman, and Thomas Quella. Deep learning is singular, and that's good. *IEEE Transactions on Neural Networks and Learning Systems*, 34(12):10473–10486, 2023. doi: 10.1109/TNNLS.2022.3167409. First published online 30 June 2022.

Independent researcher