

Separating Variables in Bivariate Polynomial Ideals

Manfred Buchacher

joint work with
Manuel Kauers and Gleb Pogudin

What is the problem?

What is the problem?

Given an ideal $I \subseteq \mathbb{K}[x, y]$, compute $I \cap (\mathbb{K}[x] + \mathbb{K}[y])$.

What is the problem?

Given an ideal $I \subseteq \mathbb{K}[x, y]$, compute $I \cap (\mathbb{K}[x] + \mathbb{K}[y])$.

Why is the problem interesting?

What is the problem?

Given an ideal $I \subseteq \mathbb{K}[x, y]$, compute $I \cap (\mathbb{K}[x] + \mathbb{K}[y])$.

Why is the problem interesting?

Intersection of \mathbb{K} -algebras.

Let $u, v \in \mathbb{K}[t_1, \dots, t_n]$. The intersection $\mathbb{K}[u] \cap \mathbb{K}[v]$ can be computed by determining pairs $(f, g) \in \mathbb{K}[x] \times \mathbb{K}[y]$ such that $f(u) = g(v)$, i.e. such that $f(x) - g(y) \in \langle x - u, y - v \rangle \cap \mathbb{K}[x, y]$.

What is the problem?

Given an ideal $I \subseteq \mathbb{K}[x, y]$, compute $I \cap (\mathbb{K}[x] + \mathbb{K}[y])$.

Why is the problem interesting?

Intersection of \mathbb{K} -algebras.

Let $u, v \in \mathbb{K}[t_1, \dots, t_n]$. The intersection $\mathbb{K}[u] \cap \mathbb{K}[v]$ can be computed by determining pairs $(f, g) \in \mathbb{K}[x] \times \mathbb{K}[y]$ such that $f(u) = g(v)$, i.e. such that $f(x) - g(y) \in \langle x - u, y - v \rangle \cap \mathbb{K}[x, y]$.

An elimination procedure for Laurent series as in Mireille Bousquet-Mélou's proof of the algebraicity of the generating function of Gessel's walks.

Definition

Definition

Let $p \in \mathbb{K}[x, y]$. It is **seperated**, if there is a $(f, g) \in \mathbb{K}[x] \times \mathbb{K}[y]$ such that

$$p = f - g.$$

Definition

Let $p \in \mathbb{K}[x, y]$. It is **seperated**, if there is a $(f, g) \in \mathbb{K}[x] \times \mathbb{K}[y]$ such that

$$p = f - g.$$

It is **separable**, if there is a $q \in \mathbb{K}[x, y] \setminus \{0\}$ such that qp is separated.

Definition

Let $p \in \mathbb{K}[x, y]$. It is **separated**, if there is a $(f, g) \in \mathbb{K}[x] \times \mathbb{K}[y]$ such that

$$p = f - g.$$

It is **separable**, if there is a $q \in \mathbb{K}[x, y] \setminus \{0\}$ such that qp is separated.

Let $I \subseteq \mathbb{K}[x, y]$ be an ideal. Then

$$A(I) := \{(f, g) \in \mathbb{K}[x] \times \mathbb{K}[y] \mid f - g \in I\}$$

is the **algebra of separated polynomials** of I .

Definition

Let $p \in \mathbb{K}[x, y]$. It is **separated**, if there is a $(f, g) \in \mathbb{K}[x] \times \mathbb{K}[y]$ such that

$$p = f - g.$$

It is **separable**, if there is a $q \in \mathbb{K}[x, y] \setminus \{0\}$ such that qp is separated.

Let $I \subseteq \mathbb{K}[x, y]$ be an ideal. Then

$$A(I) := \{(f, g) \in \mathbb{K}[x] \times \mathbb{K}[y] \mid f - g \in I\}$$

is the **algebra of separated polynomials** of I .

Problem

Given generators of an ideal $I \subseteq \mathbb{K}[x, y]$, determine a set of generators for the algebra $A(I)$ of separated polynomials.

Examples

Is the polynomial $x^3 - y^3$ separable?

Is the polynomial $x^3 - y^3$ separable?

Yes. It is even separated.

Is the polynomial $x^3 - y^3$ separable?

Yes. It is even separated.

The associated algebra of separated polynomials is

$$A(\langle x^3 - y^3 \rangle) = \mathbb{K}[(x^3, y^3)].$$

Is $x^2 + xy + y^2$ separable?

Is $x^2 + xy + y^2$ separable?

Yes, because $(x - y)(x^2 + xy + y^2) = x^3 - y^3$.

Is $x^2 + xy + y^2$ separable?

Yes, because $(x - y)(x^2 + xy + y^2) = x^3 - y^3$.

The associated algebra of separated polynomials is

$$A(\langle x^2 + xy + y^2 \rangle) = \mathbb{K}[(x^3, y^3)].$$

Is $x(x^2 + xy + y^2)$ still separable?

Is $x(x^2 + xy + y^2)$ still separable?

No. It is a multiple of x and involves y .

Is $x(x^2 + xy + y^2)$ still separable?

No. It is a multiple of x and involves y .

The algebra of separated polynomials is

$$A(\langle x(x^2 + xy + y^2) \rangle) = \mathbb{K}[(1, 1)].$$

What is $A(I)$ for the ideal I generated by

$$(x^2 - xy + y^2)(x^3 - 2xy^2 - 1) \quad \text{and} \quad (x^2 - xy + y^2)(y^3 - 2x^2y - 1)?$$

What is $A(I)$ for the ideal I generated by

$$(x^2 - xy + y^2)(x^3 - 2xy^2 - 1) \quad \text{and} \quad (x^2 - xy + y^2)(y^3 - 2x^2y - 1)?$$

A list of generators for $A(I)$ is \times

$$(x^{12} - 2x^6, y^{12} - 2y^6),$$

$$(9x^{15} - 26x^9 + 17x^3, 9y^{15} - 26y^9 + 17y^3),$$

$$(81x^{18} - 323x^6, 81y^{18} - 323y^6),$$

$$(81x^{21} - 539x^9 + 458x^3, 81y^{21} - 539y^9 + 458y^3).$$

General structure of the Algorithm

General structure of the Algorithm

- 1 Write $I = I_0 \cap I_1$ with I_0 zero-dimensional and I_1 principal.

General structure of the Algorithm

- 1 Write $I = I_0 \cap I_1$ with I_0 zero-dimensional and I_1 principal.
- 2 Compute generators of $A(I_0)$ via Gröbner basis computations and linear algebra.

General structure of the Algorithm

- 1 Write $I = I_0 \cap I_1$ with I_0 zero-dimensional and I_1 principal.
- 2 Compute generators of $A(I_0)$ via Gröbner basis computations and linear algebra.
- 3 Compute generators of $A(I_1)$ by making a suitable ansatz and linear algebra.

General structure of the Algorithm

- 1 Write $I = I_0 \cap I_1$ with I_0 zero-dimensional and I_1 principal.
- 2 Compute generators of $A(I_0)$ via Gröbner basis computations and linear algebra.
- 3 Compute generators of $A(I_1)$ by making a suitable ansatz and linear algebra.
- 4 Compute the intersection $A(I) = A(I_0) \cap A(I_1)$.

Zero-Dimensional Ideals

When I is zero-dimensional, there are

$$p, q \in \mathbb{K}[x, y] \setminus \{0\} \quad \text{such that}$$

$$I \cap \mathbb{K}[x] = \langle p \rangle \quad \text{and} \quad I \cap \mathbb{K}[y] = \langle q \rangle.$$

When I is zero-dimensional, there are

$$p, q \in \mathbb{K}[x, y] \setminus \{0\} \quad \text{such that}$$

$$I \cap \mathbb{K}[x] = \langle p \rangle \quad \text{and} \quad I \cap \mathbb{K}[y] = \langle q \rangle.$$

Clearly,

$$\mathbb{K}[x] \cdot (p, 0) + \mathbb{K}[y] \cdot (0, q) \subseteq A(I).$$

When I is zero-dimensional, there are

$$p, q \in \mathbb{K}[x, y] \setminus \{0\} \quad \text{such that}$$

$$I \cap \mathbb{K}[x] = \langle p \rangle \quad \text{and} \quad I \cap \mathbb{K}[y] = \langle q \rangle.$$

Clearly,

$$\mathbb{K}[x] \cdot (p, 0) + \mathbb{K}[y] \cdot (0, q) \subseteq A(I).$$

Consequently,

$$(f, g) \in A(I) \iff (\text{rem}(f, p), \text{rem}(g, q)) \in A(I).$$

When I is zero-dimensional, there are

$$p, q \in \mathbb{K}[x, y] \setminus \{0\} \quad \text{such that}$$

$$I \cap \mathbb{K}[x] = \langle p \rangle \quad \text{and} \quad I \cap \mathbb{K}[y] = \langle q \rangle.$$

Clearly,

$$\mathbb{K}[x] \cdot (p, 0) + \mathbb{K}[y] \cdot (0, q) \subseteq A(I).$$

Consequently,

$$(f, g) \in A(I) \iff (\text{rem}(f, p), \text{rem}(g, q)) \in A(I).$$

It is therefore sufficient to find all pairs $(f, g) \in A(I)$ with

$$\deg_x f < \deg_x p \quad \text{and} \quad \deg_y g < \deg_y q.$$

Algorithm

Algorithm

Input: $I \subseteq \mathbb{K}[x, y]$ of dimension zero.

Output: generators of $A(I)$.

Algorithm

Input: $I \subseteq \mathbb{K}[x, y]$ of dimension zero.

Output: generators of $A(I)$.

1 If $I = \langle 1 \rangle$, return $\{(1, 0), (x, 0), (0, 1), (0, y)\}$.

Algorithm

Input: $I \subseteq \mathbb{K}[x, y]$ of dimension zero.

Output: generators of $A(I)$.

- 1 If $I = \langle 1 \rangle$, return $\{(1, 0), (x, 0), (0, 1), (0, y)\}$.
- 2 Compute $p \in \mathbb{K}[x]$ and $q \in \mathbb{K}[y]$ such that

$$I \cap \mathbb{K}[x] = \langle p \rangle \text{ and } I \cap \mathbb{K}[y] = \langle q \rangle.$$

Algorithm

Input: $I \subseteq \mathbb{K}[x, y]$ of dimension zero.

Output: generators of $A(I)$.

- 1 If $I = \langle 1 \rangle$, return $\{(1, 0), (x, 0), (0, 1), (0, y)\}$.
- 2 Compute $p \in \mathbb{K}[x]$ and $q \in \mathbb{K}[y]$ such that

$$I \cap \mathbb{K}[x] = \langle p \rangle \quad \text{and} \quad I \cap \mathbb{K}[y] = \langle q \rangle.$$

- 3 Make an ansatz $h = \sum_{i=0}^{\deg_x p-1} a_i x^i - \sum_{j=0}^{\deg_y q-1} b_j y^j$ with undetermined coefficients a_i, b_j .

Algorithm

Input: $I \subseteq \mathbb{K}[x, y]$ of dimension zero.

Output: generators of $A(I)$.

- 1 If $I = \langle 1 \rangle$, return $\{(1, 0), (x, 0), (0, 1), (0, y)\}$.
- 2 Compute $p \in \mathbb{K}[x]$ and $q \in \mathbb{K}[y]$ such that

$$I \cap \mathbb{K}[x] = \langle p \rangle \quad \text{and} \quad I \cap \mathbb{K}[y] = \langle q \rangle.$$

- 3 Make an ansatz $h = \sum_{i=0}^{\deg_x p-1} a_i x^i - \sum_{j=0}^{\deg_y q-1} b_j y^j$ with undetermined coefficients a_i, b_j .
- 4 Compute the normal form of h with respect to a Gröbner basis of I and equate its coefficients to zero.

Algorithm

Input: $I \subseteq \mathbb{K}[x, y]$ of dimension zero.

Output: generators of $A(I)$.

- 1 If $I = \langle 1 \rangle$, return $\{(1, 0), (x, 0), (0, 1), (0, y)\}$.
- 2 Compute $p \in \mathbb{K}[x]$ and $q \in \mathbb{K}[y]$ such that

$$I \cap \mathbb{K}[x] = \langle p \rangle \quad \text{and} \quad I \cap \mathbb{K}[y] = \langle q \rangle.$$

- 3 Make an ansatz $h = \sum_{i=0}^{\deg_x p-1} a_i x^i - \sum_{j=0}^{\deg_y q-1} b_j y^j$ with undetermined coefficients a_i, b_j .
- 4 Compute the normal form of h with respect to a Gröbner basis of I and equate its coefficients to zero.
- 5 Solve the resulting linear system over \mathbb{K} for the unknowns a_i, b_j and let $(f_1, g_1), \dots, (f_d, g_d)$ be the pairs of polynomials corresponding to a basis of the solution space.

Algorithm

Input: $I \subseteq \mathbb{K}[x, y]$ of dimension zero.

Output: generators of $A(I)$.

- 1 If $I = \langle 1 \rangle$, return $\{(1, 0), (x, 0), (0, 1), (0, y)\}$.
- 2 Compute $p \in \mathbb{K}[x]$ and $q \in \mathbb{K}[y]$ such that

$$I \cap \mathbb{K}[x] = \langle p \rangle \quad \text{and} \quad I \cap \mathbb{K}[y] = \langle q \rangle.$$

- 3 Make an ansatz $h = \sum_{i=0}^{\deg_x p-1} a_i x^i - \sum_{j=0}^{\deg_y q-1} b_j y^j$ with undetermined coefficients a_i, b_j .
- 4 Compute the normal form of h with respect to a Gröbner basis of I and equate its coefficients to zero.
- 5 Solve the resulting linear system over \mathbb{K} for the unknowns a_i, b_j and let $(f_1, g_1), \dots, (f_d, g_d)$ be the pairs of polynomials corresponding to a basis of the solution space.
- 6 Return $(f_1, g_1), \dots, (f_d, g_d), (p, 0), \dots, (x^{\deg_x p-1} p, 0), (0, q), \dots, (0, y^{\deg_y q-1} q)$.

Principal Ideals

Assume that

$$p \in \mathbb{K}[x, y] \setminus (\mathbb{K}[x] \cup \mathbb{K}[y]).$$

Assume that

$$p \in \mathbb{K}[x, y] \setminus (\mathbb{K}[x] \cup \mathbb{K}[y]).$$

Proposition [Fried and MacRae, 1969]

If p is separable, then there is a separated multiple which divides any other separated multiple of it.

Assume that

$$p \in \mathbb{K}[x, y] \setminus (\mathbb{K}[x] \cup \mathbb{K}[y]).$$

Proposition [Fried and MacRae, 1969]

If p is separable, then there is a separated multiple which divides any other separated multiple of it.

Proposition [Fried and MacRae, 1969]

Let f, g, F, G be nonconstant polynomials. Then $f(x) - g(y)$ divides $F(x) - G(y)$ if and only if there is a polynomial r such that

$$F = r \circ f \quad \text{and} \quad G = r \circ g.$$

Assume that

$$p \in \mathbb{K}[x, y] \setminus (\mathbb{K}[x] \cup \mathbb{K}[y]).$$

Proposition [Fried and MacRae, 1969]

If p is separable, then there is a separated multiple which divides any other separated multiple of it.

Proposition [Fried and MacRae, 1969]

Let f, g, F, G be nonconstant polynomials. Then $f(x) - g(y)$ divides $F(x) - G(y)$ if and only if there is a polynomial r such that

$$F = r \circ f \quad \text{and} \quad G = r \circ g.$$

Theorem

If I is principal, then $A(I)$ is simple.

Definition

A function ω from the set of monomials in x and y to \mathbb{R} is called a **weight function** if there are $\omega_x, \omega_y \in \mathbb{Z}_{>0}$ such that $\omega(x^i y^j) = \omega_x i + \omega_y j$ for all $i, j \in \mathbb{Z}_{\geq 0}$.

Definition

A function ω from the set of monomials in x and y to \mathbb{R} is called a **weight function** if there are $\omega_x, \omega_y \in \mathbb{Z}_{>0}$ such that $\omega(x^i y^j) = \omega_x i + \omega_y j$ for all $i, j \in \mathbb{Z}_{\geq 0}$.

The sum of terms of p of maximal weight is denoted by $lp_\omega(p)$.

Definition

A function ω from the set of monomials in x and y to \mathbb{R} is called a **weight function** if there are $\omega_x, \omega_y \in \mathbb{Z}_{>0}$ such that $\omega(x^i y^j) = \omega_x i + \omega_y j$ for all $i, j \in \mathbb{Z}_{\geq 0}$.

The sum of terms of p of maximal weight is denoted by $lp_\omega(p)$.

Theorem

If p is separable and P is its minimal separated multiple, then there is a unique weight function ω such that

Definition

A function ω from the set of monomials in x and y to \mathbb{R} is called a **weight function** if there are $\omega_x, \omega_y \in \mathbb{Z}_{>0}$ such that $\omega(x^i y^j) = \omega_x i + \omega_y j$ for all $i, j \in \mathbb{Z}_{\geq 0}$.

The sum of terms of p of maximal weight is denoted by $lp_\omega(p)$.

Theorem

If p is separable and P is its minimal separated multiple, then there is a unique weight function ω such that

(a) $lp_\omega(p)$ involves at least two monomials, and

Definition

A function ω from the set of monomials in x and y to \mathbb{R} is called a **weight function** if there are $\omega_x, \omega_y \in \mathbb{Z}_{>0}$ such that $\omega(x^i y^j) = \omega_x i + \omega_y j$ for all $i, j \in \mathbb{Z}_{\geq 0}$.

The sum of terms of p of maximal weight is denoted by $lp_\omega(p)$.

Theorem

If p is separable and P is its minimal separated multiple, then there is a unique weight function ω such that

- (a) $lp_\omega(p)$ involves at least two monomials, and
- (b) the minimal separated multiple of $lp_\omega(p)$ is $lp_\omega(P)$.

An Example

Is the polynomial

$$p(x, y) = x^3 + x^2y + xy^2 + y^3 + x^2 + xy + y^2$$

separable?

Is the polynomial

$$p(x, y) = x^3 + x^2y + xy^2 + y^3 + x^2 + xy + y^2$$

separable?

Yes, because

$$(x - y)p(x, y) = x^4 + x^3 - y^4 - y^3.$$

Is $p(x, y) = x^3 + x^2y + xy^2 + y^3 + x^2 - xy + y^2$ separable?

Is $p(x, y) = x^3 + x^2y + xy^2 + y^3 + x^2 - xy + y^2$ separable?

Its leading part $lp(p)$ is $x^3 + x^2y + xy^2 + y^3$, and the minimal separated multiple of $lp(p)$ is $x^4 - y^4$.

Is $p(x, y) = x^3 + x^2y + xy^2 + y^3 + x^2 - xy + y^2$ separable?

Its leading part $lp(p)$ is $x^3 + x^2y + xy^2 + y^3$, and the minimal separated multiple of $lp(p)$ is $x^4 - y^4$.

Make an ansatz

$$P(x, y) = x^4 - y^4 + \sum_{i+j < 4} p_{ij} x^i y^j$$

for the minimal separated multiple P of p , divide it by p , and set the coefficients of the remainder equal to zero.

Is $p(x, y) = x^3 + x^2y + xy^2 + y^3 + x^2 - xy + y^2$ separable?

Its leading part $lp(p)$ is $x^3 + x^2y + xy^2 + y^3$, and the minimal separated multiple of $lp(p)$ is $x^4 - y^4$.

Make an ansatz

$$P(x, y) = x^4 - y^4 + \sum_{i+j < 4} p_{ij} x^i y^j$$

for the minimal separated multiple P of p , divide it by p , and set the coefficients of the remainder equal to zero.

The resulting linear system does not have a solution, and therefore, p is not separable.

The Homogeneous Case

Proposition

Let ω be a weight function, and let p satisfy

$$lp_{\omega}(p) = p.$$

Proposition

Let ω be a weight function, and let p satisfy

$$lp_{\omega}(p) = p.$$

Then p is separable if and only if

Proposition

Let ω be a weight function, and let p satisfy

$$lp_{\omega}(p) = p.$$

Then p is separable if and only if

(a) p involves a monomial only in x , and

Proposition

Let ω be a weight function, and let p satisfy

$$lp_{\omega}(p) = p.$$

Then p is separable if and only if

- (a) p involves a monomial only in x , and
- (b) all the roots of $p(x, 1)$ in $\overline{\mathbb{K}}$ are distinct and the ratio of every two of them is a root of unity.

Proposition

Let ω be a weight function, and let p satisfy

$$lp_{\omega}(p) = p.$$

Then p is separable if and only if

- (a) p involves a monomial only in x , and
- (b) all the roots of $p(x, 1)$ in $\overline{\mathbb{K}}$ are distinct and the ratio of every two of them is a root of unity.

Moreover, if p is separable and N is the minimal number such that the ratio of every pair of roots of $p(x, 1)$ is an N -th root of unity, then the weight of the minimal separated multiple is $N\omega_x$.

Reduction to the Homogeneous Case

Reduction to the Homogeneous Case

J. W. S. Cassels, *Factorization of polynomials in several variables*,
Proceedings of the 15th Scandinavian Congress Oslo 1968, 1969

For

$$P(x, y) = f(x) - g(y)$$

consider the auxiliary equations

$$f(x) = t \quad \text{and} \quad g(y) = t,$$

For

$$P(x, y) = f(x) - g(y)$$

consider the auxiliary equations

$$f(x) = t \quad \text{and} \quad g(y) = t,$$

and their solutions

$$\alpha_0, \dots, \alpha_{m-1} \quad \text{and} \quad \beta_0, \dots, \beta_{n-1} \quad \text{over} \quad \overline{\mathbb{K}(t)}.$$

For

$$P(x, y) = f(x) - g(y)$$

consider the auxiliary equations

$$f(x) = t \quad \text{and} \quad g(y) = t,$$

and their solutions

$$\alpha_0, \dots, \alpha_{m-1} \quad \text{and} \quad \beta_0, \dots, \beta_{n-1} \quad \text{over} \quad \overline{\mathbb{K}(t)}.$$

The Galois group G of $\overline{\mathbb{K}(t)}/\mathbb{K}(t)$ acts on $\mathbb{Z}_m \times \mathbb{Z}_n$ by

$$\pi(i, j) = (i', j') \quad :\Longleftrightarrow \quad (\pi(\alpha_i), \pi(\beta_j)) = (\alpha_{i'}, \beta_{j'}).$$

Consider the map

$$p(x, y) \mapsto T = \{(i, j) \mid p(\alpha_i, \beta_j) = 0\}.$$

Consider the map

$$p(x, y) \mapsto T = \{(i, j) \mid p(\alpha_i, \beta_j) = 0\}.$$

It is a bijection between factors of $f(x) - g(y)$ and (invariant) subsets

$$T \subseteq \mathbb{Z}_m \times \mathbb{Z}_n \quad \text{with} \quad G \cdot T = T.$$

Consider the map

$$p(x, y) \mapsto T = \{(i, j) \mid p(\alpha_i, \beta_j) = 0\}.$$

It is a bijection between factors of $f(x) - g(y)$ and (invariant) subsets

$$T \subseteq \mathbb{Z}_m \times \mathbb{Z}_n \quad \text{with} \quad G \cdot T = T.$$

Furthermore,

$$\text{if } T \subseteq \bar{T} \text{ are invariant, then } p_T(x, y) \mid p_{\bar{T}}(x, y).$$

Consider the map

$$p(x, y) \mapsto T = \{(i, j) \mid p(\alpha_i, \beta_j) = 0\}.$$

It is a bijection between factors of $f(x) - g(y)$ and (invariant) subsets

$$T \subseteq \mathbb{Z}_m \times \mathbb{Z}_n \quad \text{with} \quad G \cdot T = T.$$

Furthermore,

$$\text{if } T \subseteq \bar{T} \text{ are invariant, then } p_T(x, y) \mid p_{\bar{T}}(x, y).$$

It restricts to a bijection between separated factors and (separated) invariant subsets $T \subseteq \mathbb{Z}_m \times \mathbb{Z}_n$ such that

$$\chi_T(i, -) = \chi_T(i', -) \quad \text{or} \quad \chi_T(i, -) \cdot \chi_T(i', -) = 0 \quad \text{for all } i, i' \in \mathbb{Z}_m.$$

Consider the map

$$p(x, y) \mapsto T = \{(i, j) \mid p(\alpha_i, \beta_j) = 0\}.$$

It is a bijection between factors of $f(x) - g(y)$ and (invariant) subsets

$$T \subseteq \mathbb{Z}_m \times \mathbb{Z}_n \quad \text{with} \quad G \cdot T = T.$$

Furthermore,

$$\text{if } T \subseteq \bar{T} \text{ are invariant, then } p_T(x, y) \mid p_{\bar{T}}(x, y).$$

It restricts to a bijection between separated factors and (separated) invariant subsets $T \subseteq \mathbb{Z}_m \times \mathbb{Z}_n$ such that


$$\chi_T(i, -) = \chi_T(i', -) \quad \text{or} \quad \chi_T(i, -) \cdot \chi_T(i', -) = 0 \quad \text{for all } i, i' \in \mathbb{Z}_m.$$

In particular, $\mathbb{Z}_m \times \mathbb{Z}_n$ is invariant and separated, and corresponds to the separated factor $f(x) - g(y)$.


An Example

An Example


The factors of $x^6 - y^6$ in $\mathbb{Q}[x, y]$.




1




1




$x - y$




1




$x - y$




$x + y$




1




$x - y$




$x + y$




$x^2 - y^2$




1




$x - y$




$x + y$




$x^2 - y^2$




$x^2 - xy + y^2$




1




$x - y$




$x + y$




$x^2 - y^2$




$x^2 - xy + y^2$




$x^2 + xy + y^2$




$$1$$




$$x - y$$




$$x + y$$




$$x^2 - y^2$$




$$x^2 - xy + y^2$$




$$x^2 + xy + y^2$$




$$x^3 - y^3$$




1




$x - y$




$x + y$




$x^2 - y^2$




$x^2 - xy + y^2$




$x^2 + xy + y^2$




$x^3 - y^3$




$x^3 - 2x^2y + 2xy^2 - y^3$




1




$x - y$




$x + y$




$x^2 - y^2$




$x^2 - xy + y^2$




$x^2 + xy + y^2$




$x^3 - y^3$




$x^3 - 2x^2y + 2xy^2 - y^3$




$x^3 + 2x^2y + 2xy^2 + y^3$




1




$x - y$




$x + y$




$x^2 - y^2$




$x^2 - xy + y^2$




$x^2 + xy + y^2$




$x^3 - y^3$




$x^3 - 2x^2y + 2xy^2 - y^3$




$x^3 + 2x^2y + 2xy^2 + y^3$




$x^3 + y^3$




$$1$$




$$x - y$$




$$x + y$$




$$x^2 - y^2$$




$$x^2 - xy + y^2$$




$$x^2 + xy + y^2$$




$$x^3 - y^3$$




$$x^3 - 2x^2y + 2xy^2 - y^3$$




$$x^3 + 2x^2y + 2xy^2 + y^3$$




$$x^3 + y^3$$




$$x^4 + x^2y^2 + y^4$$




1




$x - y$




$x + y$




$x^2 - y^2$




$x^2 - xy + y^2$




$x^2 + xy + y^2$




$x^3 - y^3$




$x^3 - 2x^2y + 2xy^2 - y^3$




$x^3 + 2x^2y + 2xy^2 + y^3$




$x^3 + y^3$




$x^4 + x^2y^2 + y^4$




$x^4 - x^3y + xy^3 - y^4$




1




$x - y$




$x + y$




$x^2 - y^2$




$x^2 - xy + y^2$




$x^2 + xy + y^2$




$x^3 - y^3$




$x^3 - 2x^2y + 2xy^2 - y^3$




$x^3 + 2x^2y + 2xy^2 + y^3$




$x^3 + y^3$




$x^4 + x^2y^2 + y^4$




$x^4 - x^3y + xy^3 - y^4$




$x^4 + x^3y - xy^3 - y^4$




1




$x - y$




$x + y$




$x^2 - y^2$




$x^2 - xy + y^2$




$x^2 + xy + y^2$




$x^3 - y^3$




$x^3 - 2x^2y + 2xy^2 - y^3$




$x^3 + 2x^2y + 2xy^2 + y^3$




$x^3 + y^3$




$x^4 + x^2y^2 + y^4$




$x^4 - x^3y + xy^3 - y^4$




$x^4 + x^3y - xy^3 - y^4$




$x^5 - x^4y + \dots - y^5$




1




$x - y$




$x + y$




$x^2 - y^2$




$x^2 - xy + y^2$




$x^2 + xy + y^2$




$x^3 - y^3$




$x^3 - 2x^2y + 2xy^2 - y^3$




$x^3 + 2x^2y + 2xy^2 + y^3$




$x^3 + y^3$




$x^4 + x^2y^2 + y^4$




$x^4 - x^3y + xy^3 - y^4$




$x^4 + x^3y - xy^3 - y^4$




$x^5 - x^4y + \dots - y^5$




$x^5 + x^4y + \dots + y^5$




1




$x - y$




$x + y$




$x^2 - y^2$




$x^2 - xy + y^2$




$x^2 + xy + y^2$




$x^3 - y^3$




$x^3 - 2x^2y + 2xy^2 - y^3$




$x^3 + 2x^2y + 2xy^2 + y^3$




$x^3 + y^3$




$x^4 + x^2y^2 + y^4$




$x^4 - x^3y + xy^3 - y^4$




$x^4 + x^3y - xy^3 - y^4$



$x^5 - x^4y + \dots - y^5$



$x^5 + x^4y + \dots + y^5$



$x^6 - y^6$

Definition

Definition

Let T be an invariant subset of $\mathbb{Z}_n \times \mathbb{Z}_m$. The **separable closure** T^{sep} of T is defined by

$$T^{\text{sep}} := \bigcap_{\substack{S \supseteq T \\ S \text{ inv, sep}}} S.$$

Theorem

If p is separable and P is its minimal separated multiple, then there is a unique weight function ω such that

- (a) $lp_{\omega}(p)$ involves at least two monomials, and
- (b) the minimal separated multiple of $lp_{\omega}(p)$ is $lp_{\omega}(P)$.

Sketch of Proof

Sketch of Proof

Assume $\alpha_i, \beta_j \in \mathbb{K}^{\text{Puisseux}}(t^{-1})$, and define

$$\bar{\alpha}_i := \text{lt}(\alpha_i) \quad \text{and} \quad \bar{\beta}_j := \text{lt}(\beta_j), \text{ and}$$

$$T := \{(i, j) \mid p(\alpha_i, \beta_j) = 0\} \quad \text{and} \quad \bar{T} := \{(i, j) \mid \text{lp}_\omega(p)(\bar{\alpha}_i, \bar{\beta}_j) = 0\}.$$

Sketch of Proof

Assume $\alpha_i, \beta_j \in \mathbb{K}^{\text{Puisseux}}(t^{-1})$, and define

$$\bar{\alpha}_i := \text{lt}(\alpha_i) \quad \text{and} \quad \bar{\beta}_j := \text{lt}(\beta_j), \text{ and}$$

$$T := \{(i, j) \mid p(\alpha_i, \beta_j) = 0\} \quad \text{and} \quad \bar{T} := \{(i, j) \mid \text{lp}_\omega(p)(\bar{\alpha}_i, \bar{\beta}_j) = 0\}.$$

Since

$$p(\alpha_i, \beta_j) = 0 \quad \implies \quad \text{lp}_\omega(p)(\bar{\alpha}_i, \bar{\beta}_j) = 0,$$

we have

$$T \subseteq \bar{T}, \quad \text{and hence} \quad T^{\text{sep}} \subseteq \bar{T}^{\text{sep}}.$$

Sketch of Proof

Assume $\alpha_i, \beta_j \in \mathbb{K}^{\text{Puisseux}}(t^{-1})$, and define

$$\bar{\alpha}_i := \text{lt}(\alpha_i) \quad \text{and} \quad \bar{\beta}_j := \text{lt}(\beta_j), \text{ and}$$

$$T := \{(i, j) \mid p(\alpha_i, \beta_j) = 0\} \quad \text{and} \quad \bar{T} := \{(i, j) \mid \text{lp}_\omega(p)(\bar{\alpha}_i, \bar{\beta}_j) = 0\}.$$

Since

$$p(\alpha_i, \beta_j) = 0 \implies \text{lp}_\omega(p)(\bar{\alpha}_i, \bar{\beta}_j) = 0,$$

we have

$$T \subseteq \bar{T}, \quad \text{and hence} \quad T^{\text{sep}} \subseteq \bar{T}^{\text{sep}}.$$

If P is the minimal separated multiple of p , then

$$T^{\text{sep}} = \mathbb{Z}_m \times \mathbb{Z}_n, \quad \text{and hence} \quad \bar{T}^{\text{sep}} = \mathbb{Z}_m \times \mathbb{Z}_n,$$

and $\text{lp}_\omega(P)$ is the minimal separated multiple of $\text{lp}_\omega(p)$.

Arbitrary Bivariate Ideals

Let $I = I_0 \cap I_1$ be such that I_0 is zero-dimensional and I_1 principal. Given a set of generators of $A(I_0)$ and the generator of $A(I_1)$, how can we determine a set of generators of

$$A(I) = A(I_0) \cap A(I_1)?$$

Lemma

Lemma

Let $I_0 \subseteq \mathbb{K}[x, y]$ be a zero-dimensional ideal. There is a finite-dimensional \mathbb{K} -subspace V of $\mathbb{K}[x] \times \mathbb{K}[y]$ such that

$$V \oplus A(I_0) = \mathbb{K}[x] \times \mathbb{K}[y].$$

Lemma

Let $I_0 \subseteq \mathbb{K}[x, y]$ be a zero-dimensional ideal. There is a finite-dimensional \mathbb{K} -subspace V of $\mathbb{K}[x] \times \mathbb{K}[y]$ such that

$$V \oplus A(I_0) = \mathbb{K}[x] \times \mathbb{K}[y].$$

Moreover, given $(f, g) \in \mathbb{K}[x] \times \mathbb{K}[y]$, we can compute $(\tilde{f}, \tilde{g}) \in V$ such that

$$(f, g) - (\tilde{f}, \tilde{g}) \in A(I_0).$$

Algorithm

Algorithm

Input: $\alpha \in \mathbb{K}[x] \times \mathbb{K}[y]$, and $A(I_0)$ and V as before, and a finite set $S = \{s_1, \dots, s_m\}$ of elements of \mathbb{N} .

Output: a basis of the vector space of polynomials p such that $\text{supp}(p) \subseteq S$ and $p(\alpha) \in A(I_0)$.

Algorithm

Input: $\alpha \in \mathbb{K}[x] \times \mathbb{K}[y]$, and $A(I_0)$ and V as before, and a finite set $S = \{s_1, \dots, s_m\}$ of elements of \mathbb{N} .

Output: a basis of the vector space of polynomials p such that $\text{supp}(p) \subseteq S$ and $p(\alpha) \in A(I_0)$.

1 For $i = 1, \dots, m$, compute $r_i \in V$ such that $\alpha^{s_i} - r_i \in A(I_0)$.

Algorithm

Input: $\alpha \in \mathbb{K}[x] \times \mathbb{K}[y]$, and $A(I_0)$ and V as before, and a finite set $S = \{s_1, \dots, s_m\}$ of elements of \mathbb{N} .

Output: a basis of the vector space of polynomials p such that $\text{supp}(p) \subseteq S$ and $p(\alpha) \in A(I_0)$.

- 1 For $i = 1, \dots, m$, compute $r_i \in V$ such that $\alpha^{s_i} - r_i \in A(I_0)$.
- 2 Compute a basis B of the space of all $(c_1, \dots, c_m) \in \mathbb{K}^m$ with $c_1 r_1 + \dots + c_m r_m = 0$.

Algorithm

Input: $\alpha \in \mathbb{K}[x] \times \mathbb{K}[y]$, and $A(I_0)$ and V as before, and a finite set $S = \{s_1, \dots, s_m\}$ of elements of \mathbb{N} .

Output: a basis of the vector space of polynomials p such that $\text{supp}(p) \subseteq S$ and $p(\alpha) \in A(I_0)$.

- 1 For $i = 1, \dots, m$, compute $r_i \in V$ such that $\alpha^{s_i} - r_i \in A(I_0)$.
- 2 Compute a basis B of the space of all $(c_1, \dots, c_m) \in \mathbb{K}^m$ with $c_1 r_1 + \dots + c_m r_m = 0$.
- 3 For every element $(c_1, \dots, c_m) \in B$, return $c_1 t^{s_1} + \dots + c_m t^{s_m}$.

A set of generators of $\mathcal{A}(I_0) \cap \mathcal{A}(I_1)$ can then be computed by iteratively applying the algorithm as follows:

A set of generators of $\mathcal{A}(I_0) \cap \mathcal{A}(I_1)$ can then be computed by iteratively applying the algorithm as follows:

Algorithm

A set of generators of $\mathcal{A}(I_0) \cap \mathcal{A}(I_1)$ can then be computed by iteratively applying the algorithm as follows:

Algorithm

Input: a zero-dimensional ideal I_0 and a generator α of $\mathcal{A}(I_1)$.

Output: a set of generators for $\mathcal{A}(I_0) \cap \mathcal{A}(I_1)$.

A set of generators of $A(I_0) \cap A(I_1)$ can then be computed by iteratively applying the algorithm as follows:

Algorithm

Input: a zero-dimensional ideal I_0 and a generator α of $A(I_1)$.

Output: a set of generators for $A(I_0) \cap A(I_1)$.

- 1 Compute a basis of a vector space V for which $V \oplus A(I_0) = \mathbb{K}[x] \times \mathbb{K}[y]$.

A set of generators of $A(I_0) \cap A(I_1)$ can then be computed by iteratively applying the algorithm as follows:

Algorithm

Input: a zero-dimensional ideal I_0 and a generator α of $A(I_1)$.

Output: a set of generators for $A(I_0) \cap A(I_1)$.

- 1 Compute a basis of a vector space V for which $V \oplus A(I_0) = \mathbb{K}[x] \times \mathbb{K}[y]$.
- 2 Set $G = \emptyset$, $\Delta = \emptyset$.

A set of generators of $A(I_0) \cap A(I_1)$ can then be computed by iteratively applying the algorithm as follows:

Algorithm

Input: a zero-dimensional ideal I_0 and a generator α of $A(I_1)$.

Output: a set of generators for $A(I_0) \cap A(I_1)$.

- 1 Compute a basis of a vector space V for which $V \oplus A(I_0) = \mathbb{K}[x] \times \mathbb{K}[y]$.
- 2 Set $G = \emptyset$, $\Delta = \emptyset$.
- 3 While $\gcd(\Delta) \neq 1$, do:

A set of generators of $A(I_0) \cap A(I_1)$ can then be computed by iteratively applying the algorithm as follows:

Algorithm

Input: a zero-dimensional ideal I_0 and a generator α of $A(I_1)$.

Output: a set of generators for $A(I_0) \cap A(I_1)$.

- 1 Compute a basis of a vector space V for which $V \oplus A(I_0) = \mathbb{K}[x] \times \mathbb{K}[y]$.
- 2 Set $G = \emptyset$, $\Delta = \emptyset$.
- 3 While $\gcd(\Delta) \neq 1$, do:
- 4 Select a set $S \subseteq \mathbb{N} \setminus \langle \Delta \rangle$ with $|S| > \dim V$ and find a polynomial p with $p(\alpha) \in A(I_0)$ and $\text{supp}(p) \subseteq S$.

A set of generators of $A(I_0) \cap A(I_1)$ can then be computed by iteratively applying the algorithm as follows:

Algorithm

Input: a zero-dimensional ideal I_0 and a generator α of $A(I_1)$.

Output: a set of generators for $A(I_0) \cap A(I_1)$.

- 1 Compute a basis of a vector space V for which $V \oplus A(I_0) = \mathbb{K}[x] \times \mathbb{K}[y]$.
- 2 Set $G = \emptyset$, $\Delta = \emptyset$.
- 3 While $\gcd(\Delta) \neq 1$, do:
 - 4 Select a set $S \subseteq \mathbb{N} \setminus \langle \Delta \rangle$ with $|S| > \dim V$ and find a polynomial p with $p(\alpha) \in A(I_0)$ and $\text{supp}(p) \subseteq S$.
 - 5 $G = G \cup \{p\}$, $\Delta = \Delta \cup \{\deg p\}$

A set of generators of $A(I_0) \cap A(I_1)$ can then be computed by iteratively applying the algorithm as follows:

Algorithm

Input: a zero-dimensional ideal I_0 and a generator α of $A(I_1)$.

Output: a set of generators for $A(I_0) \cap A(I_1)$.

- 1 Compute a basis of a vector space V for which $V \oplus A(I_0) = \mathbb{K}[x] \times \mathbb{K}[y]$.
- 2 Set $G = \emptyset$, $\Delta = \emptyset$.
- 3 While $\gcd(\Delta) \neq 1$, do:
 - 4 Select a set $S \subseteq \mathbb{N} \setminus \langle \Delta \rangle$ with $|S| > \dim V$ and find a polynomial p with $p(\alpha) \in A(I_0)$ and $\text{supp}(p) \subseteq S$.
 - 5 $G = G \cup \{p\}$, $\Delta = \Delta \cup \{\deg p\}$
- 6 Find a basis of the vector space of polynomials p with $p(\alpha) \in A(I_0)$ and $\text{supp}(p) \subseteq S = \mathbb{N} \setminus \langle \Delta \rangle$ and add the resulting polynomials to G .

A set of generators of $A(I_0) \cap A(I_1)$ can then be computed by iteratively applying the algorithm as follows:

Algorithm

Input: a zero-dimensional ideal I_0 and a generator α of $A(I_1)$.

Output: a set of generators for $A(I_0) \cap A(I_1)$.

- 1 Compute a basis of a vector space V for which $V \oplus A(I_0) = \mathbb{K}[x] \times \mathbb{K}[y]$.
- 2 Set $G = \emptyset$, $\Delta = \emptyset$.
- 3 While $\gcd(\Delta) \neq 1$, do:
 - 4 Select a set $S \subseteq \mathbb{N} \setminus \langle \Delta \rangle$ with $|S| > \dim V$ and find a polynomial p with $p(\alpha) \in A(I_0)$ and $\text{supp}(p) \subseteq S$.
 - 5 $G = G \cup \{p\}$, $\Delta = \Delta \cup \{\deg p\}$
- 6 Find a basis of the vector space of polynomials p with $p(\alpha) \in A(I_0)$ and $\text{supp}(p) \subseteq S = \mathbb{N} \setminus \langle \Delta \rangle$ and add the resulting polynomials to G .
- 7 Return G

An Example

To compute $A(I_0) \cap A(I_1)$ for

$$I_0 = \langle x^3 - 2xy + y^2, y^3 - 2x^2y - 1 \rangle \quad \text{and} \quad I_1 = \langle x^2 - xy + y^2 \rangle,$$

we find

To compute $A(I_0) \cap A(I_1)$ for

$$I_0 = \langle x^3 - 2xy + y^2, y^3 - 2x^2y - 1 \rangle \quad \text{and} \quad I_1 = \langle x^2 - xy + y^2 \rangle,$$

we find $A(I_1) = \mathbb{K}(x^3, -y^3)$,

To compute $A(I_0) \cap A(I_1)$ for

$$I_0 = \langle x^3 - 2xy + y^2, y^3 - 2x^2y - 1 \rangle \quad \text{and} \quad I_1 = \langle x^2 - xy + y^2 \rangle,$$

we find $A(I_1) = \mathbb{K}(x^3, -y^3)$, and $V = \bigoplus_{i=0}^8 \mathbb{K} \cdot (0, y^i)$ such that

$$\bigoplus_{i=0}^8 \mathbb{K} \cdot (0, y^i) \oplus A(I_0) = \mathbb{K}[x] \times \mathbb{K}[y].$$

To compute $A(I_0) \cap A(I_1)$ for

$$I_0 = \langle x^3 - 2xy + y^2, y^3 - 2x^2y - 1 \rangle \quad \text{and} \quad I_1 = \langle x^2 - xy + y^2 \rangle,$$

we find $A(I_1) = \mathbb{K}(x^3, -y^3)$, and $V = \bigoplus_{i=0}^8 \mathbb{K} \cdot (0, y^i)$ such that

$$\bigoplus_{i=0}^8 \mathbb{K} \cdot (0, y^i) \oplus A(I_0) = \mathbb{K}[x] \times \mathbb{K}[y].$$

By making an ansatz for a polynomial p with $\deg(p) \leq 10$ such that $p((x^3, -y^3)) \in A(I_0)$, we find $p = t^4 - 2t^2$,

To compute $A(I_0) \cap A(I_1)$ for

$$I_0 = \langle x^3 - 2xy + y^2, y^3 - 2x^2y - 1 \rangle \quad \text{and} \quad I_1 = \langle x^2 - xy + y^2 \rangle,$$

we find $A(I_1) = \mathbb{K}(x^3, -y^3)$, and $V = \bigoplus_{i=0}^8 \mathbb{K} \cdot (0, y^i)$ such that

$$\bigoplus_{i=0}^8 \mathbb{K} \cdot (0, y^i) \oplus A(I_0) = \mathbb{K}[x] \times \mathbb{K}[y].$$

By making an ansatz for a polynomial p with $\deg(p) \leq 10$ such that $p((x^3, -y^3)) \in A(I_0)$, we find $p = t^4 - 2t^2$, and, in the next step, a polynomial $q = 9t^5 - 26t^3 + 17$ with support in $S = \{1, 2, 3, 5, 6, 7, 9, 10, 11, 13\}$.

To compute $A(I_0) \cap A(I_1)$ for

$$I_0 = \langle x^3 - 2xy + y^2, y^3 - 2x^2y - 1 \rangle \quad \text{and} \quad I_1 = \langle x^2 - xy + y^2 \rangle,$$

we find $A(I_1) = \mathbb{K}(x^3, -y^3)$, and $V = \bigoplus_{i=0}^8 \mathbb{K} \cdot (0, y^i)$ such that

$$\bigoplus_{i=0}^8 \mathbb{K} \cdot (0, y^i) \oplus A(I_0) = \mathbb{K}[x] \times \mathbb{K}[y].$$

By making an ansatz for a polynomial p with $\deg(p) \leq 10$ such that $p((x^3, -y^3)) \in A(I_0)$, we find $p = t^4 - 2t^2$, and, in the next step, a polynomial $q = 9t^5 - 26t^3 + 17$ with support in $S = \{1, 2, 3, 5, 6, 7, 9, 10, 11, 13\}$.

Since $\gcd(4, 5) = 1$, the set $S = \mathbb{N} \setminus \langle 4, 5 \rangle$ is finite,

To compute $A(I_0) \cap A(I_1)$ for

$$I_0 = \langle x^3 - 2xy + y^2, y^3 - 2x^2y - 1 \rangle \quad \text{and} \quad I_1 = \langle x^2 - xy + y^2 \rangle,$$

we find $A(I_1) = \mathbb{K}(x^3, -y^3)$, and $V = \bigoplus_{i=0}^8 \mathbb{K} \cdot (0, y^i)$ such that

$$\bigoplus_{i=0}^8 \mathbb{K} \cdot (0, y^i) \oplus A(I_0) = \mathbb{K}[x] \times \mathbb{K}[y].$$

By making an ansatz for a polynomial p with $\deg(p) \leq 10$ such that $p((x^3, -y^3)) \in A(I_0)$, we find $p = t^4 - 2t^2$, and, in the next step, a polynomial $q = 9t^5 - 26t^3 + 17$ with support in $S = \{1, 2, 3, 5, 6, 7, 9, 10, 11, 13\}$.

Since $\gcd(4, 5) = 1$, the set $S = \mathbb{N} \setminus \langle 4, 5 \rangle$ is finite, and the space of polynomials whose support is contained in S is generated by $81t^6 - 323t^3$, $81t^7 - 539t^3 + 458$, and $6561t^{11} - 191125t^3 + 184564$.

The implementation of the algorithm can be found on
<http://kauers.de/software/separate.m>

The implementation of the algorithm can be found on
<http://kauers.de/software/separate.m>

Thank you for your attention.