

Separated Variables on Plane Algebraic Curves

Manfred Buchacher

Institut für Algebra
Johannes Kepler Universität Linz

Supported by the LIT with the grant LIT-2022-11-YOU-214

Let $r \in \mathbb{C}(x, y)$, and $p \in \mathbb{C}[x, y]$.

Let $r \in \mathbb{C}(x, y)$, and $p \in \mathbb{C}[x, y]$.

Problem

Solve

$$r + q(x, y)p = f(x) - g(y)$$

for $q, f, g \in \mathbb{C}(x, y)$.

Some **field theoretic** interpretations

Some **field theoretic** interpretations

Intersections of fields

Some **field theoretic** interpretations

Intersections of fields

$$q(x, y)p = f(x) - g(y)$$

Some **field theoretic** interpretations

Intersections of fields

$$f(x) \equiv g(y) \pmod{p}$$

Some **field theoretic** interpretations

Intersections of fields

$$\mathbb{C}(x) \cap \mathbb{C}(y) \pmod{p}$$

Some **field theoretic** interpretations

Intersections of fields

$$\mathbb{C}(x) \cap \mathbb{C}(y) \pmod{p}$$

Field membership

Some **field theoretic** interpretations

Intersections of fields

$$\mathbb{C}(x) \cap \mathbb{C}(y) \pmod{p}$$

Field membership

$$r + q(x, y)p = f(x)$$

Some **field theoretic** interpretations

Intersections of fields

$$\mathbb{C}(x) \cap \mathbb{C}(y) \pmod{p}$$

Field membership

$$r \equiv f(x) \pmod{p}$$

Some **field theoretic** interpretations

Intersections of fields

$$\mathbb{C}(x) \cap \mathbb{C}(y) \pmod{p}$$

Field membership

$$r \in \mathbb{C}(x) \pmod{p}$$

Some **field theoretic** interpretations

Intersections of fields

$$\mathbb{C}(x) \cap \mathbb{C}(y) \pmod{p}$$

Field membership

$$r \in \mathbb{C}(x) \pmod{p}$$

Sum-decomposition

Some **field theoretic** interpretations

Intersections of fields

$$\mathbb{C}(x) \cap \mathbb{C}(y) \pmod{p}$$

Field membership

$$r \in \mathbb{C}(x) \pmod{p}$$

Sum-decomposition

$$r + q(x, y)p = f(x) - g(y)$$

Some **field theoretic** interpretations

Intersections of fields

$$\mathbb{C}(x) \cap \mathbb{C}(y) \pmod{p}$$

Field membership

$$r \in \mathbb{C}(x) \pmod{p}$$

Sum-decomposition

$$r \equiv f(x) - g(y) \pmod{p}$$

Some **field theoretic** interpretations

Intersections of fields

$$\mathbb{C}(\mathbf{x}) \cap \mathbb{C}(\mathbf{y}) \pmod{p}$$

Field membership

$$r \in \mathbb{C}(\mathbf{x}) \pmod{p}$$

Sum-decomposition

$$r \in \mathbb{C}(\mathbf{x}) + \mathbb{C}(\mathbf{y}) \pmod{p}$$

These problems arise in

- enumerative combinatorics
- computer vision
- parameter identification in ODE models
- algebraic independence of solutions of ODEs
- designing diffractive optical systems

Define

$$F(\mathfrak{r}, \mathfrak{p}) := \{(f, g) \in \mathbb{C}(\mathfrak{x}) \times \mathbb{C}(\mathfrak{y}) : f - g \in \mathfrak{r} + \langle \mathfrak{p} \rangle\}.$$

Define

$$F(r, p) := \{(f, g) \in \mathbb{C}(x) \times \mathbb{C}(y) : f - g \in r + \langle p \rangle\}.$$

Prop

Let (f, g) be any element of $F(r, p)$.

Define

$$F(r, p) := \{(f, g) \in \mathbb{C}(x) \times \mathbb{C}(y) : f - g \in r + \langle p \rangle\}.$$

Prop

Let (f, g) be any element of $F(r, p)$. Then

$$F(r, p) = (f, g) + F(0, p).$$

Define

$$F(r, p) := \{(f, g) \in \mathbb{C}(x) \times \mathbb{C}(y) : f - g \in r + \langle p \rangle\}.$$

Prop

Let (f, g) be any element of $F(r, p)$. Then

$$F(r, p) = (f, g) + F(0, p).$$

Furthermore,

$$F(p) \equiv F(0, p)$$

is a simple field.

The problem of computing $F(r, p)$ splits into

- 1) finding a **generator** of $F(p)$, and
- 2) determining **any** element of $F(r, p)$.

The problem of computing $F(r, p)$ splits into

- 1) finding a **generator** of $F(p)$, and
- 2) determining **any** element of $F(r, p)$.

Strategy

The non-linear problem of solving

$$r + qp = f - g$$

is reduced to a **linear** problem. The reduction is based on the computation of the **poles** of f and g and their **multiplicities**.

1) Finding a generator of $F(p)$

1) Finding a generator of $F(\mathfrak{p})$

Poles

There is a generator $(f, g) \in F(p)$ such that

$$f(\infty) = \infty.$$

There is a generator $(f, g) \in F(p)$ such that

$$f(\infty) = \infty.$$

If (s_1, s_2) is a root of p , then

$$f(s_1) = \infty \quad \text{iff} \quad g(s_2) = \infty.$$

There is a generator $(f, g) \in F(p)$ such that

$$f(\infty) = \infty.$$

If (s_1, s_2) is a root of p , then

$$f(s_1) = \infty \quad \text{iff} \quad g(s_2) = \infty.$$

Def

Let \sim be the smallest equivalence relation on $\{p = 0\}$ such that

$$(x_0, y_0) \sim (x_1, y_1) \quad \text{whenever} \quad x_0 = x_1 \text{ or } y_0 = y_1.$$

There is a generator $(f, g) \in F(p)$ such that

$$f(\infty) = \infty.$$

If (s_1, s_2) is a root of p , then

$$f(s_1) = \infty \quad \text{iff} \quad g(s_2) = \infty.$$

Def

Let \sim be the smallest equivalence relation on $\{p = 0\}$ such that

$$(x_0, y_0) \sim (x_1, y_1) \quad \text{whenever} \quad x_0 = x_1 \text{ or } y_0 = y_1.$$

The equivalence class of (x_0, y_0) is called the **orbit** of (x_0, y_0) .

Thm

The coordinates of the orbit of ∞ are **poles** of f and g , respectively. The orbit is **finite**, and it is **exhaustive**.

Thm

The coordinates of the orbit of ∞ are **poles** of f and g , respectively. The orbit is **finite**, and it is **exhaustive**. If

$$F(p) \cong \mathbb{C},$$

it might however be infinite.

1) Finding a generator of $F(p)$

Multiplicities

The problem is reduced to one for **homogeneous** polynomials by studying p **locally** at the elements of the **orbit** of ∞ .

Any $\omega \in \mathbb{Z}^2$ induces a **grading** on $\mathbb{C}[x, y]$ by

$$\omega(ax^i y^j) = \omega_x i + \omega_y j.$$

The **leading part** of $p(x, y)$ is the sum of terms of maximal (weighted) degree $\omega(p)$. It is denoted by $lp_\omega(p)$.

Any $\omega \in \mathbb{Z}^2$ induces a **grading** on $\mathbb{C}[x, y]$ by

$$\omega(ax^i y^j) = \omega_x i + \omega_y j.$$

The **leading part** of $p(x, y)$ is the sum of terms of maximal (weighted) degree $\omega(p)$. It is denoted by $lp_\omega(p)$.

The weighted degrees of the terms of

$$p = xy - 1 - y - x^2y - x^2y^2$$

with respect to $\omega = (1, -2)$ are

$$-1, 0, -2, 0, -2.$$

Any $\omega \in \mathbb{Z}^2$ induces a **grading** on $\mathbb{C}[x, y]$ by

$$\omega(ax^i y^j) = \omega_x i + \omega_y j.$$

The **leading part** of $p(x, y)$ is the sum of terms of maximal (weighted) degree $\omega(p)$. It is denoted by $lp_\omega(p)$.

The weighted degrees of the terms of

$$p = xy - 1 - y - x^2y - x^2y^2$$

with respect to $\omega = (1, -2)$ are

$$-1, 0, -2, 0, -2.$$

Its leading part is

$$lp_\omega(p) = -1 - x^2y.$$

Let us consider the **example** with

$$p = xy - 1 - y - x^2y - x^2y^2.$$

Let us consider the **example** with

$$p = xy - 1 - y - x^2y - x^2y^2.$$

The elements of the orbit of ∞ are


$$(\infty, 0), (\infty, -1), (0, -1) \quad \text{and} \quad (0, \infty).$$

Let us consider the **example** with

$$p = xy - 1 - y - x^2y - x^2y^2.$$

The elements of the orbit of ∞ are

$$(\infty, 0), (\infty, -1), (0, -1) \quad \text{and} \quad (0, \infty).$$



The leading part of p associated with $(\infty, 0)$ is

$$lp_{\omega}(p) = -1 - x^2y.$$

The leading part of p associated with $(\infty, 0)$ is

$$\text{lp}_\omega(p) = -1 - x^2y.$$

Since

$$F(\text{lp}_\omega(p)) = \mathbb{C}((x^2, -y^{-1})),$$

there is a $k \in \mathbb{N}$ such that

$$(m(\infty, f), m(0, g)) = k \cdot (2, 1).$$

The analysis of the other poles is done analogously. It results in the following 1-parameter family for their multiplicities.

| f | | g | |
|----------|------|----------|------|
| ∞ | $2k$ | ∞ | k |
| 0 | $2k$ | 0 | k |
| | | -1 | $2k$ |

Making the ansatz

$$f = \frac{f_0 + f_1x + \cdots + f_4x^4}{x^2} \quad \text{and} \quad g = \frac{g_0 + g_1y + \cdots + g_4y^4}{y(1+y)^2}$$

and

$$q = \frac{q_{0,0} + q_{1,0}x + \cdots + q_{2,1}x^2y}{x^2y(1+y)^2}$$

we find that

$$F(p) = \mathbb{C} \left(\left(\frac{(1-x)^2(1+x+x^2)}{x^2}, -\frac{(1+y+y^2)^2}{y(1+y)^2} \right) \right).$$

Thm

If $F(p)$ is non-trivial, then choosing $k = 1$ results in the multiplicities of the poles of a generator.

If $F(p)$ is non-trivial, then every orbit is finite.

The question of how the procedure can be turned into an algorithm raises the following

If $F(p)$ is non-trivial, then every orbit is finite.

The question of how the procedure can be turned into an algorithm raises the following

Problem

Compute an upper bound on the size of a finite orbit.

Before we explain how to compute $F(r, p)$,
let us consider the following simpler problem.

Before we explain how to compute $F(r, p)$,
let us consider the following simpler problem.

Problem

Solve

$$r + q(x, y)p = f(x)$$

for $q, f \in \mathbb{C}(x, y)$.

2) Determining an element of $F(r, p)$

2) Determining an element of $F(r, p)$

Poles

Let (s_1, s_2) be a root of p such that

$$r(s_1, s_2) = f(s_1) - g(s_2).$$

Let (s_1, s_2) be a root of p such that

$$r(s_1, s_2) = f(s_1) - g(s_2).$$

Then

$$f(s_1) = \infty \quad \text{or} \quad g(s_2) = \infty \quad \text{if} \quad r(s_1, s_2) = \infty.$$

Let (s_1, s_2) be a root of p such that

$$r(s_1, s_2) = f(s_1) - g(s_2).$$

Then

$$f(s_1) = \infty \quad \text{or} \quad g(s_2) = \infty \quad \text{if} \quad r(s_1, s_2) = \infty.$$

If

$$r(s_1, s_2) < \infty, \quad \text{then} \quad f(s_1) = \infty \quad \text{iff} \quad g(s_2) = \infty.$$

Do the orbits of the poles of r provide all poles of f and g ?

Do the orbits of the poles of r provide all poles of f and g ?

No!

Do the orbits of the poles of r provide all poles of f and g ?

No!

It is also necessary to consider the orbits of its **roots** if $F(p)$ is non-trivial, and the **finite orbits** if $F(p)$ is trivial.

Do the orbits of the poles of r provide all poles of f and g ?

No!

It is also necessary to consider the orbits of its **roots** if $F(p)$ is non-trivial, and the **finite orbits** if $F(p)$ is trivial.

Prop [Bell, Moosa, Topaz, Bellaïche]

If $F(p)$ is trivial, then the number of finite orbits is finite.

2) Determining an element of $F(r, p)$

Multiplicities

(Upper) bounds on the multiplicities are derived from

- the local behavior of r ,
- its (weighted) degrees $\omega(r)$,
- and the generators of $F(\mathfrak{lp}_\omega(\mathfrak{p}))$.

Let (x_1, y_1) and (x_2, y_2) be roots of p .

Let (x_1, y_1) and (x_2, y_2) be roots of p .

Problem

Decide whether (x_2, y_2) is in (some part of) the orbit of (x_1, y_1) .

Let (x_1, y_1) and (x_2, y_2) be roots of p .

Problem

Decide whether (x_2, y_2) is in (some part of) the orbit of (x_1, y_1) .

Problem

How do the semi-algorithms generalize?

Some references

- *Separating Variables in Bivariate Polynomial Ideals*
- *Separating Variables in Bivariate Polynomial Ideals: the Local Case*
- *Separated Variables on Plane Algebraic Curves*

Some references

- *Separating Variables in Bivariate Polynomial Ideals*
- *Separating Variables in Bivariate Polynomial Ideals: the Local Case*
- *Separated Variables on Plane Algebraic Curves*
- *Galoisian structure of large steps walks confined in the first quadrant*, Pierre Bonnet and Charlotte Hardouin

Some references

- *Separating Variables in Bivariate Polynomial Ideals*
- *Separating Variables in Bivariate Polynomial Ideals: the Local Case*
- *Separated Variables on Plane Algebraic Curves*
- *Galoisian structure of large steps walks confined in the first quadrant*, Pierre Bonnet and Charlotte Hardouin

Thank you!