

$$x(1 - t(x^{-1} + x))F(x; t) = x - tF(0; t)$$

Manfred Buchacher

Austrian Academy of Sciences

Johann Radon Institute for Computational and Applied Mathematics

I did my PhD studies from 2017 to 2021 at the Institute for Algebra at JKU Linz under the supervision of Manuel Kauers, and I stayed there for another few months as a post-doc.

I did my PhD studies from 2017 to 2021 at the Institute for Algebra at JKU Linz under the supervision of Manuel Kauers, and I stayed there for another few months as a post-doc.

Research interests:

Algorithmic enumerative combinatorics, computer and constructive algebra, discrete differential equations and the elementary power series algebra to solve them.

$$x(1 - t(x^{-1} + x))F(x; t) = x - tF(0; t)$$

$$x(1 - t(x^{-1} + x))F(x; t) = x - tF(0; t)$$

Goal: Solve the equation for $F(x; t)$ in $\mathbb{Q}[x][[t]]$.

$$F(x;t) = 1 + txF(x;t) - t\Delta F(x;t),$$

$$F(x;t) = 1 + tx F(x;t) - t \Delta F(x;t),$$

where Δ acts on $\mathbb{Q}[x][[t]]$ by

$$\Delta F(x;t) := (F(x;t) - F(0;t))/x.$$

Combinatorial meaning

Definition

A lattice walk is a sequence P_0, P_1, \dots, P_n of points of \mathbb{Z}^d . We call P_0 and P_n its starting and end point, respectively, the consecutive differences $P_{i+1} - P_i$ are its steps, and n is its length.

Definition

A lattice walk is a sequence P_0, P_1, \dots, P_n of points of \mathbb{Z}^d . We call P_0 and P_n its starting and end point, respectively, the consecutive differences $P_{i+1} - P_i$ are its steps, and n is its length.

Example


The sequence $(0, 1, 0, 1, 2, 1, 2, 1, 2, 3)$ is a lattice walk on \mathbb{N} that starts at 0, ends at 3, has length 9, and whose steps are $1, -1, 1, 1, -1, 1, -1, 1, 1$.

Definition

A lattice walk is a sequence P_0, P_1, \dots, P_n of points of \mathbb{Z}^d . We call P_0 and P_n its starting and end point, respectively, the consecutive differences $P_{i+1} - P_i$ are its steps, and n is its length.

Example

The sequence $(0, 1, 0, 1, 2, 1, 2, 1, 2, 3)$ is a lattice walk on \mathbb{N} that starts at 0, ends at 3, has length 9, and whose steps are $1, -1, 1, 1, -1, 1, -1, 1, 1$.



Let $f(i; n)$ be the number of walks on \mathbb{N} that start at 0, end at i and consist of (precisely) n steps taken from $\{-1, 1\}$, and let

$$F(x; t) := \sum_{n \geq 0} \left(\sum_{i \geq 0} f(i; n) x^i \right) t^n.$$

Let $f(i; n)$ be the number of walks on \mathbb{N} that start at 0, end at i and consist of (precisely) n steps taken from $\{-1, 1\}$, and let

$$F(x; t) := \sum_{n \geq 0} \left(\sum_{i \geq 0} f(i; n) x^i \right) t^n.$$

Then $F(x; t)$ solves the functional equation, since

$$\begin{aligned} f(i; n+1) &= f(i-1; n) + f(i+1, n) \\ f(i; 0) &= \delta_{i,0} \quad \text{and} \quad f(i; n) = 0, \quad i < 0. \end{aligned}$$

Existence and uniqueness

By extracting the coefficient of t^n of

$$F(x; t) = 1 + t(x^{-1} + x)F(x; t) - tx^{-1}F(0; t),$$

we find that

$$[t^n]F(x; t) = \begin{cases} 1, & \text{if } n = 0, \\ (x^{-1} + x)[t^{n-1}]F(x; t) - x^{-1}[t^{n-1}]F(0; t) & \text{else.} \end{cases}$$

By extracting the coefficient of t^n of

$$F(x; t) = 1 + t(x^{-1} + x)F(x; t) - tx^{-1}F(0; t),$$

we find that

$$[t^n]F(x; t) = \begin{cases} 1, & \text{if } n = 0, \\ (x^{-1} + x)[t^{n-1}]F(x; t) - x^{-1}[t^{n-1}]F(0; t) & \text{else.} \end{cases}$$

This recurrence relation has a unique solution, and therefore this is also true for the functional equation.

The map

$$F(x; t) \mapsto 1 + t(x^{-1} + x)F(x; t) - tx^{-1}F(0; t)$$

represents a contraction on $\mathbb{Q}[x][[t]]$ with respect to

$$d(F, G) := 2^{-\text{val}(F-G)},$$

where

$$\text{val}(F) := \min\{n : [t^n]F \neq 0\}.$$

The map

$$F(x; t) \mapsto 1 + t(x^{-1} + x)F(x; t) - tx^{-1}F(0; t)$$

represents a contraction on $\mathbb{Q}[x][[t]]$ with respect to

$$d(F, G) := 2^{-\text{val}(F-G)},$$

where

$$\text{val}(F) := \min\{n : [t^n]F \neq 0\}.$$

Banach's fixed point theorem implies the existence and uniqueness of a solution of the functional equation.

What does it mean to solve a discrete differential equation?

What does it mean to solve a discrete differential equation?

It means to decide whether its solution is **rational**, **algebraic**, **D-finite** or **D-algebraic**, and in case it is, to find an algebraic or differential equation for it.

How can a discrete differential equation be solved?

How can a discrete differential equation be solved?

There are essentially 2 different approaches:

How can a discrete differential equation be solved?

There are essentially 2 different approaches:

- the **analytic** approach: involving complex analysis, boundary value problems and Galois theory of difference equations, and

How can a discrete differential equation be solved?

There are essentially 2 different approaches:

- the **analytic** approach: involving complex analysis, boundary value problems and Galois theory of difference equations, and
- the formal or **algebraic** one: involving elementary power series algebra, constructive and computer algebra.

We focus on formal and algebraic methods, and show how discrete differential equations can be solved using only operations such as

$$+, \cdot, \circ \quad \text{and} \quad [x^>].$$

We focus on formal and algebraic methods, and show how discrete differential equations can be solved using only operations such as

$$+, \cdot, \circ \quad \text{and} \quad [x^>].$$

The latter operator acts on series by discarding all terms involving non-positive powers in x , e.g.

$$[x^>] \left(x^{-2} + x^{-1} + 1 + x + x^2 \right) = x + x^2.$$

The classical kernel method

Rewrite the functional equation,

$$x(1 - t(\bar{x} + x))F(x; t) = x - tF(0; t),$$

where $\bar{x} := x^{-1}$,

Rewrite the functional equation,

$$x(1 - t(\bar{x} + x))F(x; t) = x - tF(0; t),$$

where $\bar{x} := x^{-1}$, replace x by the power series root

$$x_0(t) := \frac{1 - \sqrt{1 - 4t^2}}{2t} \quad \text{of} \quad K(x; t) := x(1 - t(\bar{x} + x)),$$

Rewrite the functional equation,

$$x(1 - t(\bar{x} + x))F(x; t) = x - tF(0; t),$$

where $\bar{x} := x^{-1}$, replace x by the power series root

$$x_0(t) := \frac{1 - \sqrt{1 - 4t^2}}{2t} \quad \text{of} \quad K(x; t) := x(1 - t(\bar{x} + x)),$$

and solve the resulting equation

$$0 = x_0(t) - tF(0; t)$$

Rewrite the functional equation,

$$x(1 - t(\bar{x} + x))F(x; t) = x - tF(0; t),$$

where $\bar{x} := x^{-1}$, replace x by the power series root

$$x_0(t) := \frac{1 - \sqrt{1 - 4t^2}}{2t} \quad \text{of} \quad K(x; t) := x(1 - t(\bar{x} + x)),$$

and solve the resulting equation

$$0 = x_0(t) - tF(0; t)$$

to find that

$$F(0; t) = x_0(t)/t, \quad \text{and} \quad F(x; t) = \frac{x - x_0(t)}{x(1 - t(\bar{x} + x))}.$$

Theorem [Bousquet-Mélou, Jehanne]

Solutions of ordinary discrete differential equations are **algebraic** (over a field of rational functions).

Theorem [Bousquet-Mélou, Jehanne]

Solutions of ordinary discrete differential equations are **algebraic** (over a field of rational functions).

Theorem [Buchacher, Kauers]

(The components of) solutions of systems of ordinary linear discrete differential equations are **algebraic** (over a field of rational functions).

The orbit-sum method

The equation

$$x(1 - t(\bar{x} + x))F(x; t) = x - tF(0; t)$$

The equation

$$x(1 - t(\bar{x} + x))F(x; t) = x - tF(0; t)$$

can be solved by replacing x by \bar{x} ,

$$\bar{x}(1 - t(\bar{x} + x))F(\bar{x}; t) = \bar{x} - tF(0; t),$$

The equation

$$x(1 - t(\bar{x} + x))F(x; t) = x - tF(0; t)$$

can be solved by **replacing** x by \bar{x} ,

$$\bar{x}(1 - t(\bar{x} + x))F(\bar{x}; t) = \bar{x} - tF(0; t),$$

subtracting the latter equation from the former,

$$xF(x; t) - \bar{x}F(\bar{x}; t) = \frac{x - \bar{x}}{1 - t(\bar{x} + x)},$$

The equation

$$x(1 - t(\bar{x} + x))F(x; t) = x - tF(0; t)$$

can be solved by **replacing** x by \bar{x} ,

$$\bar{x}(1 - t(\bar{x} + x))F(\bar{x}; t) = \bar{x} - tF(0; t),$$

subtracting the latter equation from the former,

$$xF(x; t) - \bar{x}F(\bar{x}; t) = \frac{x - \bar{x}}{1 - t(\bar{x} + x)},$$

and **discarding** non-positive powers in x ,

$$xF(x; t) = [x^>] \frac{x - \bar{x}}{1 - t(\bar{x} + x)}.$$

The generalization of the orbit-sum method to partial discrete differential equations of higher order raises the following questions:

The generalization of the orbit-sum method to partial discrete differential equations of higher order raises the following questions:

- How can we compute in a finite field extension of $\mathbb{Q}(x_1, \dots, x_n)$ in an effective and efficient way?

The generalization of the orbit-sum method to partial discrete differential equations of higher order raises the following questions:

- How can we compute in a finite field extension of $\mathbb{Q}(x_1, \dots, x_n)$ in an effective and efficient way?
- Given $p \in \mathbb{Q}[x_1, \dots, x_n][y]$, how can we solve

$$p(x_1, \dots, x_n, y) = 0$$

for y in terms of series in x_1, \dots, x_n ?

The generalization of the orbit-sum method to partial discrete differential equations of higher order raises the following questions:

- How can we compute in a finite field extension of $\mathbb{Q}(x_1, \dots, x_n)$ in an effective and efficient way?
- Given $p \in \mathbb{Q}[x_1, \dots, x_n][y]$, how can we solve

$$p(x_1, \dots, x_n, y) = 0$$

for y in terms of series in x_1, \dots, x_n ?

- How can we do effective arithmetic for multivariate algebraic series?

The generalization of the orbit-sum method to partial discrete differential equations of higher order raises the following questions:

- How can we compute in a finite field extension of $\mathbb{Q}(x_1, \dots, x_n)$ in an effective and efficient way?
- Given $p \in \mathbb{Q}[x_1, \dots, x_n][y]$, how can we solve

$$p(x_1, \dots, x_n, y) = 0$$

for y in terms of series in x_1, \dots, x_n ?

- How can we do effective arithmetic for multivariate algebraic series?
- How can we compute the convex hull of the support of a multivariate algebraic series?

Wiener-Hopf factorization

The coefficient polynomial of $F(x; t)$ in

$$x(1 - t(\bar{x} + x))F(x; t) = x - tF(0; t)$$

factors into

$$-t(x - x_0)(x - x_1) = -t(1 - \bar{x}x_0)x(x - x_1),$$

The coefficient polynomial of $F(x; t)$ in

$$x(1 - t(\bar{x} + x))F(x; t) = x - tF(0; t)$$

factors into

$$-t(x - x_0)(x - x_1) = -t(1 - \bar{x}x_0)x(x - x_1),$$

where

$$(1 - \bar{x}x_0)^{\pm 1} \in \mathbb{Q}[\bar{x}][[t]] \quad \text{and} \quad (x - x_1)^{\pm 1} \in \mathbb{Q}[x][[t]].$$

The coefficient polynomial of $F(x; t)$ in

$$x(1 - t(\bar{x} + x))F(x; t) = x - tF(0; t)$$

factors into

$$-t(x - x_0)(x - x_1) = -t(1 - \bar{x}x_0)x(x - x_1),$$

where

$$(1 - \bar{x}x_0)^{\pm 1} \in \mathbb{Q}[\bar{x}][[t]] \quad \text{and} \quad (x - x_1)^{\pm 1} \in \mathbb{Q}[x][[t]].$$

So

$$-t(x - x_1)F(x; t) = \frac{x - tF(0; t)}{(1 - \bar{x}x_0)x}$$

with the lhs in $\mathbb{Q}[x][[t]]$ and the rhs in $\mathbb{Q}[\bar{x}][[t]]$.

The coefficient polynomial of $F(x; t)$ in

$$x(1 - t(\bar{x} + x))F(x; t) = x - tF(0; t)$$

factors into

$$-t(x - x_0)(x - x_1) = -t(1 - \bar{x}x_0)x(x - x_1),$$

where

$$(1 - \bar{x}x_0)^{\pm 1} \in \mathbb{Q}[\bar{x}][[t]] \quad \text{and} \quad (x - x_1)^{\pm 1} \in \mathbb{Q}[x][[t]].$$

So

$$-t(x - x_1)F(x; t) = \frac{x - tF(0; t)}{(1 - \bar{x}x_0)x}$$

with the lhs in $\mathbb{Q}[x][[t]]$ and the rhs in $\mathbb{Q}[\bar{x}][[t]]$. Therefore

$$-t(x - x_1)F(x; t) = [x^0] \frac{x - tF(0; t)}{(1 - \bar{x}x_0)x} = 1,$$

The coefficient polynomial of $F(x; t)$ in

$$x(1 - t(\bar{x} + x))F(x; t) = x - tF(0; t)$$

factors into

$$-t(x - x_0)(x - x_1) = -t(1 - \bar{x}x_0)x(x - x_1),$$

where

$$(1 - \bar{x}x_0)^{\pm 1} \in \mathbb{Q}[\bar{x}][[t]] \quad \text{and} \quad (x - x_1)^{\pm 1} \in \mathbb{Q}[x][[t]].$$

So

$$-t(x - x_1)F(x; t) = \frac{x - tF(0; t)}{(1 - \bar{x}x_0)x}$$

with the lhs in $\mathbb{Q}[x][[t]]$ and the rhs in $\mathbb{Q}[\bar{x}][[t]]$. Therefore

$$-t(x - x_1)F(x; t) = [x^0] \frac{x - tF(0; t)}{(1 - \bar{x}x_0)x} = 1,$$

and so

$$F(x; t) = -\frac{1}{t(x - x_1)}.$$

A generalization to systems of ordinary linear discrete differential equations raises the questions:

A generalization to systems of ordinary linear discrete differential equations raises the questions:

- Given $A \in \mathbb{Q}(x, t)^{n \times n}$, does it factor into

$$A = A_- A_+,$$

where

$$A_-^{\pm 1} \in \mathbb{Q}[\bar{x}]((t))^{n \times n} \quad \text{and} \quad A_+^{\pm 1} \in \mathbb{Q}[x]((t))^{\pm 1}?$$

A generalization to systems of ordinary linear discrete differential equations raises the questions:

- Given $A \in \mathbb{Q}(x, t)^{n \times n}$, does it factor into

$$A = A_- A_+,$$

where

$$A_{\pm}^{\pm 1} \in \mathbb{Q}[\bar{x}]((t))^{n \times n} \quad \text{and} \quad A_{\pm}^{\pm 1} \in \mathbb{Q}[x]((t))^{\pm 1}?$$

- If there is such a factorization, how can it be determined?

Compositional inverses and Lagrange inversion

Replacing t in

$$x(1 - t(\bar{x} + x))F(x; t) = x - tF(0; t),$$

by the power series root

$$t_0(x) := \frac{x}{1 + x^2} \quad \text{of} \quad x(1 - t(\bar{x} + x)) \in \mathbb{Q}(x)[t],$$

Replacing t in

$$x(1 - t(\bar{x} + x))F(x; t) = x - tF(0; t),$$

by the power series root

$$t_0(x) := \frac{x}{1 + x^2} \quad \text{of} \quad x(1 - t(\bar{x} + x)) \in \mathbb{Q}(x)[t],$$

results in

$$\frac{x}{1 + x^2} F\left(0; \frac{x}{1 + x^2}\right) = x,$$

Replacing t in

$$x(1 - t(\bar{x} + x))F(x; t) = x - tF(0; t),$$

by the power series root

$$t_0(x) := \frac{x}{1 + x^2} \quad \text{of} \quad x(1 - t(\bar{x} + x)) \in \mathbb{Q}(x)[t],$$

results in

$$\frac{x}{1 + x^2} F\left(0; \frac{x}{1 + x^2}\right) = x,$$

i.e. $\frac{x}{1+x^2}$ and $xF(0; x)$ are compositional inverse to each other.

Theorem [Lagrange Inversion]

Let $H(t) \in \mathbb{Q}[[t]]$ be of valuation 1, and let $G(t)$ be its compositional inverse, i.e. such that $G(H(t)) = t$. Then

$$[t^n]G(t) = \frac{1}{n}[t^{-1}]H^{-n}(t), \quad n \neq 0.$$

Theorem [Lagrange Inversion]

Let $H(t) \in \mathbb{Q}[[t]]$ be of valuation 1, and let $G(t)$ be its compositional inverse, i.e. such that $G(H(t)) = t$. Then

$$[t^n]G(t) = \frac{1}{n}[t^{-1}]H^{-n}(t), \quad n \neq 0.$$

Applying this to

$$H(t) = (t^{-1} + t)^{-1}$$

we find that

$$[t^{n+1}]tF(0;t) = \frac{1}{n+1}[t^{-1}](t^{-1} + t)^{n+1},$$

Theorem [Lagrange Inversion]

Let $H(t) \in \mathbb{Q}[[t]]$ be of valuation 1, and let $G(t)$ be its compositional inverse, i.e. such that $G(H(t)) = t$. Then

$$[t^n]G(t) = \frac{1}{n}[t^{-1}]H^{-n}(t), \quad n \neq 0.$$

Applying this to

$$H(t) = (t^{-1} + t)^{-1}$$

we find that

$$[t^{n+1}]tF(0;t) = \frac{1}{n+1}[t^{-1}](t^{-1} + t)^{n+1},$$

which means that

$$f(0;n) = \begin{cases} 0, & \text{if } n \not\equiv 0 \pmod{2}, \\ \frac{1}{n/2+1} \binom{n}{n/2} & \text{else.} \end{cases}$$

The series $F(0; t)$ is algebraic (and consequently also $F(x; t)$) because

$$G(H(t)) = t \quad \text{implies} \quad H(G(t)) = t.$$

The series $F(0; t)$ is algebraic (and consequently also $F(x; t)$) because

$$G(H(t)) = t \quad \text{implies} \quad H(G(t)) = t.$$

Therefore

$$P(t, F(0; t)) = 0 \quad \text{for} \quad P(t, s) = t^2 s^2 - s + 1.$$

Guess and prove

We guess an annihilating polynomial $P(x, t, y)$ for $F(x; t)$ by

We guess an annihilating polynomial $P(x, t, y)$ for $F(x; t)$ by

1. making an ansatz

$$P(x, t, y) = \sum_{i,j,k=0}^d p_{ijk} x^i t^j y^k$$

for some $d \in \mathbb{N}$ and undetermined p_{ijk} 's,

We guess an annihilating polynomial $P(x, t, y)$ for $F(x; t)$ by

1. making an ansatz

$$P(x, t, y) = \sum_{i,j,k=0}^d p_{ijk} x^i t^j y^k$$

for some $d \in \mathbb{N}$ and undetermined p_{ijk} 's,

2. computing a truncation

$$F_0(x; t) := \sum_{n=0}^{n_0} \left(\sum_{i \geq 0} f(i; n) x^i \right) t^n$$

of $F(x; t)$ up to some order n_0 , and

We guess an annihilating polynomial $P(x, t, y)$ for $F(x; t)$ by

1. making an ansatz

$$P(x, t, y) = \sum_{i,j,k=0}^d p_{ijk} x^i t^j y^k$$

for some $d \in \mathbb{N}$ and undetermined p_{ijk} 's,

2. computing a truncation

$$F_0(x; t) := \sum_{n=0}^{n_0} \left(\sum_{i \geq 0} f(i; n) x^i \right) t^n$$

of $F(x; t)$ up to some order n_0 , and

3. setting coefficients of $x^i t^j$, $j \leq n_0$, in $P(x, t, F_0(x; t))$ equal to zero, and solving the resulting linear system.

For $d = 2$ and $n_0 = 8$ we find

$$P(x, t, y) = 1 - (1 - 2xt)y - xt(1 - t(x^{-1} + x))y^2.$$

For $d = 2$ and $n_0 = 8$ we find

$$P(x, t, y) = 1 - (1 - 2xt)y - xt(1 - t(x^{-1} + x))y^2.$$

We can prove that the guess is correct by

For $d = 2$ and $n_0 = 8$ we find

$$P(x, t, y) = 1 - (1 - 2xt)y - xt(1 - t(x^{-1} + x))y^2.$$

We can prove that the guess is correct by

1. showing that $P(x, t, y)$ has a (unique) root

For $d = 2$ and $n_0 = 8$ we find

$$P(x, t, y) = 1 - (1 - 2xt)y - xt(1 - t(x^{-1} + x))y^2.$$

We can prove that the guess is correct by

1. showing that $P(x, t, y)$ has a (unique) root

$$F_{\text{cand}}(x; t) \in \mathbb{Q}[x][[t]], \text{ and}$$

2. verifying that $F_{\text{cand}}(x, t)$ satisfies the functional equation.

Theorem [Implicit Function Theorem]

Let \mathbb{K} be a field of characteristic 0, and let $A(x, y) \in \mathbb{K}[[t, y]]$ s.t.

$$A(0, 0) = 0 \quad \text{and} \quad \frac{\partial A}{\partial y}(0, 0) \neq 0.$$

Then there is a unique $f(t) \in \mathbb{K}[[t]]$ with

$$f(0) = 0 \quad \text{and} \quad A(t, f(t)) = 0.$$

Theorem [Implicit Function Theorem]

Let \mathbb{K} be a field of characteristic 0, and let $A(x, y) \in \mathbb{K}[[t, y]]$ s.t.

$$A(0, 0) = 0 \quad \text{and} \quad \frac{\partial A}{\partial y}(0, 0) \neq 0.$$

Then there is a unique $f(t) \in \mathbb{K}[[t]]$ with

$$f(0) = 0 \quad \text{and} \quad A(t, f(t)) = 0.$$

This guarantees the existence of a root $F_{\text{cand}}(x; t)$ of P in $\mathbb{Q}(x)[[t]]$, which actually lies in $\mathbb{Q}[x][[t]]$.

By performing closure properties of algebraic functions we observe that

$$A(y) := y(1 - 4t^2 - t^2y^2)$$

is an annihilating polynomial for

$$x(1 - t(\bar{x} + x))F_{\text{cand}}(x; t) - x - tF_{\text{cand}}(0; t).$$

By performing closure properties of algebraic functions we observe that

$$A(y) := y(1 - 4t^2 - t^2y^2)$$

is an annihilating polynomial for

$$x(1 - t(\bar{x} + x))F_{\text{cand}}(x; t) - x - tF_{\text{cand}}(0; t).$$

By plugging some truncation of this series into the second factor of A we find that y is its minimal polynomial.

Looking for an additional equation

Instead of solving the seemingly underdetermined equation

$$x(1 - t(\bar{x} + x))F(x; t) = x - tF(0; t)$$

one can derive an additional equation and solve the resulting system.

Instead of solving the seemingly underdetermined equation

$$x(1 - t(\bar{x} + x))F(x; t) = x - tF(0; t)$$

one can derive an additional equation and solve the resulting system.

The "first passage" decomposition leads to

$$F(0; t) = 1 + t^2 F(0; t)^2.$$

Instead of solving the seemingly underdetermined equation

$$x(1 - t(\bar{x} + x))F(x; t) = x - tF(0; t)$$

one can derive an additional equation and solve the resulting system.

The "first passage" decomposition leads to

$$F(0; t) = 1 + t^2 F(0; t)^2.$$

The "arch" decomposition implies the equation

$$F(0; t) = \frac{1}{1 - t^2 F(0; t)}$$

Instead of solving the seemingly underdetermined equation

$$x(1 - t(\bar{x} + x))F(x; t) = x - tF(0; t)$$

one can derive an additional equation and solve the resulting system.

The "first passage" decomposition leads to

$$F(0; t) = 1 + t^2 F(0; t)^2.$$

The "arch" decomposition implies the equation

$$F(0; t) = \frac{1}{1 - t^2 F(0; t)}$$

The additional equation allows to determine $F(x; t)$ by solving a system of polynomial equations.

Note: Although the second equation is equivalent to the first, the latter suggests to write $F(0;t)$ as a continued fraction

$$F(0;t) = \frac{1}{1 - \frac{t^2}{1 - \frac{t^2}{1 - \dots}}}.$$

Note: Although the second equation is equivalent to the first, the latter suggests to write $F(0;t)$ as a continued fraction

$$F(0;t) = \frac{1}{1 - \frac{t^2}{1 - \frac{t^2}{1 - \dots}}}.$$

This indicates that there is a connection between the enumeration of restricted lattice walks and continued fractions and orthogonal polynomials.

The reflection principle

The orbit-sum method showed that

$$F(x; t) = [x^{\geq}] \frac{1 - \bar{x}^2}{1 - t(\bar{x} + x)}.$$

The orbit-sum method showed that

$$F(x; t) = [x^{\geq}] \frac{1 - \bar{x}^2}{1 - t(\bar{x} + x)}.$$

The combinatorial interpretation of this equation is that


walks in \mathbb{Z} from 0 to i not lying in \mathbb{N}

=


walks in \mathbb{Z} from -2 to i

A bijection is given by the reflection principle:


A bijection is given by the reflection principle:



A bijection is given by the reflection principle:



A bijection is given by the reflection principle:



This implies that

$$\begin{aligned} f(i; n) &= \# \text{ walks in } \mathbb{Z} \text{ from } 0 \text{ to } i \text{ of length } n - \\ &\quad \# \text{ walks in } \mathbb{Z} \text{ from } -2 \text{ to } i \\ &= \binom{n}{\frac{n+i}{2}} - \binom{n}{\frac{n+i+2}{2}} = \frac{1}{\frac{n+i}{2} + 1} \binom{n}{\frac{n+i}{2}}. \end{aligned}$$

The cycle lemma / Spitzer's lemma

Langrange inversion implies that

$$f(0; n) = \frac{1}{n+1} [t^{-1}] (t^{-1} + t)^{n+1}$$

Langrange inversion implies that

$$f(0; n) = \frac{1}{n+1} [t^{-1}] (t^{-1} + t)^{n+1}$$


Is there a combinatorial explanation for this identity?

Lemma [Spitzer's lemma]

Let $a_1, a_2, \dots, a_N \in \mathbb{R}$ be s.t. $a_1 + a_2 + \dots + a_N = 0$ and no other partial sum of consecutive a_i 's read cyclically vanishes. Then there is a unique cyclic permutation $a_i, a_{i+1}, \dots, a_N, a_1, \dots, a_{i-1}$ s.t. the sum of the first j numbers is non-negative for $j = 1, 2, \dots, N$.



Lemma [Spitzer's lemma]

Let $a_1, a_2, \dots, a_N \in \mathbb{R}$ be s.t. $a_1 + a_2 + \dots + a_N = 0$ and no other partial sum of consecutive a_i 's read cyclically vanishes. Then there is a unique cyclic permutation $a_i, a_{i+1}, \dots, a_N, a_1, \dots, a_{i-1}$ s.t. the sum of the first j numbers is non-negative for $j = 1, 2, \dots, N$.



Lemma [Spitzer's lemma]

Let $a_1, a_2, \dots, a_N \in \mathbb{R}$ be s.t. $a_1 + a_2 + \dots + a_N = 0$ and no other partial sum of consecutive a_i 's read cyclically vanishes. Then there is a unique cyclic permutation $a_i, a_{i+1}, \dots, a_N, a_1, \dots, a_{i-1}$ s.t. the sum of the first j numbers is non-negative for $j = 1, 2, \dots, N$.



Theorem

Let $r, s \in \mathbb{N}_+$ be relative prime. Then the number of walks in \mathbb{Z}^2 from $(0, 0)$ to (r, s) with steps taken from $\{(1, 0), (0, 1)\}$ staying weakly below $ry = sx$ equals $\frac{1}{r+s} \binom{r+s}{r}$.