Years From Posed To Resolved: The Solvability Atlas

MAT 124 Statistics Project

Gianangelo Dichio

2026-10-13

Abstract

DRAFT for Gianangelo to rewrite in his own words. Target 100 to 150 words.

Objective: estimate the mean number of years a research problem stays open, from the year it was posed to the year it was resolved, for the solved problems in the Solvability Atlas. The data are 213 cleaned rows from the atlas problem table built on 2026-10-13, one row per solved problem with both years recorded. Descriptive work used all 213 rows; inference used a sample of 50 rows drawn with replacement from the cleaned records under seed 124. The full-data mean is 34.30 years and the median is 20.00 years. Year posed and years to resolve have $r = -0.68$ and $R^2 = 0.46$. The 95% t interval from the sample is 18.69 to 34.51 years. The two-sided test of $\mu = 50$ gives $t = -5.94$ and $p < 0.0001$, so the sample provides evidence against the benchmark at $\alpha = 0.05$.

1 Data Form

Approval. The assignment lists approved public sources. This dataset is the author's own and sits outside that list, so the instructor's approval of the source is pending. Assignment 1 asks for three dataset choices; this form covers the first choice, and the second and third choices are still to be added.

Source line. Solvability Atlas problem table, Bucket Foundation; code MIT, each row under the licence in its `licence` column (CC BY-SA 4.0 for Wikipedia rows, Apache-2.0 for formal-conjectures rows, MIT for atlas rows, and four rows that cite an AAAS headline); download: https://github.com/bucket-foundation/bucket-foundation/blob/dev/tools/solvability-atlas/surface/table.csv.

Source. Title: Solvability Atlas problem table. Creator: Bucket Foundation, Gianangelo Dichio. Accessed 2026-10-13. Data period: problems posed from 1506 to 2023 and resolved from 1676 to 2026. Files attached: `data/original.csv` and `data/cleaned.csv`.

Population. Solved research problems recorded in the atlas with a posed year and a resolved year. One row is one problem. The file holds 5209 problems, open and solved, compiled from public problem lists.

Collection. Observational. The rows were compiled from public lists: Wikipedia problem lists, science timelines and prize competition pages, the formal-conjectures repository and the atlas's own 71 curated problems. The sampling method of those lists is not reported. Limits on representation:

a list favours well-known problems, and a problem enters this study only when both of its years were recorded, which leaves out most rows.

Variables. X: year posed, calendar year, quantitative, interval level. Y: years from posed to resolved, in years, quantitative, ratio level. Category 1: branch, with labels applied, biophysics, chemistry, cosmology, information, mathematics, mind, physics; categorical, nominal. Category 2: formal status, the state of a machine-checked proof, with labels none, partial, proved, statement; categorical, ordinal.

Feasibility. Total rows: 5209. Rows complete on the four variables after the filters: 213. Filters and missing-value rules are the seven cleaning rules in the Data Check. No category is derived.

Question. For solved research problems in the cleaned atlas table, is the mean time from posed to resolved equal to 50 years? Expected relationship: problems posed later have fewer years to resolve, because a recently posed problem that is already solved cannot have been open for long.

Hypotheses. $H_0 : \mu = 50$ years. $H_1 : \mu \neq 50$ years. The benchmark of 50 years is hypothetical. It was fixed in the code before the sample was drawn, and it stands for two research generations of 25 years each.

2 Descriptive Statistics

2.1 Data Check

Original rows: 5209. Removed: 4996. Retained: 213. Row IDs are the atlas `id` values and are kept in every file. Rules apply in order, and a row is counted under the first rule it meets. The excluded row IDs, one per line with the rule, are in `data/excluded_rows.csv`.

Cleaning rule	Rows removed
R1 posed year missing	4522
R2 resolved year missing	470
R3 status other than solved	2
R4 variant of another row	2
R5 resolved year before posed year	0
R6 branch or formal status missing	0
R7 duplicate id	0

Missing values: a row missing either year is removed under R1 or R2, and no value is imputed. Duplicate records: no ID repeats, and a variant row that restates part of another problem is removed under R4. Implausible values: a resolved year earlier than the posed year would be removed under R5, and no row met that rule.

2.2 Tables

Counts and relative frequencies of branch, denominator 213 problems.

Branch	Count	Relative frequency
physics	46	0.2160
mathematics	36	0.1690
information	36	0.1690
cosmology	28	0.1315
applied	26	0.1221
biophysics	20	0.0939
chemistry	14	0.0657
mind	7	0.0329
Total	213	1.0000

Counts and relative frequencies of formal status, denominator 213 problems.

Formal status	Count	Relative frequency
none	197	0.9249
proved	11	0.0516
partial	3	0.0141
statement	2	0.0094
Total	213	1.0000

Branch by formal status. Each cell holds the count and, in parentheses, the conditional row proportion; the denominator is the row total for that branch.

Branch	none	partial	proved	statement	Row total
applied	26 (1.0000)	0 (0.0000)	0 (0.0000)	0 (0.0000)	26
biophysics	20 (1.0000)	0 (0.0000)	0 (0.0000)	0 (0.0000)	20
chemistry	14 (1.0000)	0 (0.0000)	0 (0.0000)	0 (0.0000)	14
cosmology	28 (1.0000)	0 (0.0000)	0 (0.0000)	0 (0.0000)	28
information	32 (0.8889)	0 (0.0000)	4 (0.1111)	0 (0.0000)	36
mathematics	24 (0.6667)	3 (0.0833)	7 (0.1944)	2 (0.0556)	36
mind	7 (1.0000)	0 (0.0000)	0 (0.0000)	0 (0.0000)	7
physics	46 (1.0000)	0 (0.0000)	0 (0.0000)	0 (0.0000)	46
Column total	197	3	11	2	213

Grouped Y. Classes are $[\text{lower}, \text{upper})$ in years. Sturges' rule gives $1 + \log_2 213 = 8.73$, about nine classes; the range of 387 years over nine classes is 43.00 years, rounded to a width of 40 years so that boundaries fall on multiples of 40.

Class, years	Midpoint	Count	Rel. freq.	Cum. count	Cum. rel. freq.
0 to under 40	20	147	0.6901	147	0.6901
40 to under 80	60	48	0.2254	195	0.9155
80 to under 120	100	11	0.0516	206	0.9671
120 to under 160	140	2	0.0094	208	0.9765
160 to under 200	180	1	0.0047	209	0.9812
200 to under 240	220	0	0.0000	209	0.9812
240 to under 280	260	0	0.0000	209	0.9812
280 to under 320	300	0	0.0000	209	0.9812
320 to under 360	340	2	0.0094	211	0.9906
360 to under 400	380	2	0.0094	213	1.0000

2.3 Summaries

Y is years from posed to resolved. Variance uses the $n-1$ denominator. Quartiles and the percentile follow QUARTILE.INC and PERCENTILE.INC, which is linear interpolation in `numpy.percentile`.

Statistic	Value, years
n	213
Mean	34.30
Median	20.00
Mode	0, 1 (13 rows each)
Sample variance	2903.60
Sample standard deviation	53.89
Minimum	0.00
Q1	5.00
Median	20.00
Q3	45.00
Maximum	387.00
Range	387.00
IQR	40.00
90th percentile	70.00

Fences: lower $= Q1 - 1.5 \times IQR = 5.00 - 1.5(40.00) = -55.00$; upper $= Q3 + 1.5 \times IQR = 45.00 + 1.5(40.00) = 105.00$. Flagged outliers: 7. Row IDs with their values: `fermat` (357), `fourcolor` (124), `kepler` (387), `sd-anthrax-as-a-microbial-disease` (330), `sd-collapse-of-a-massive-star-into-a-bl` (143), `sd-rings-of-uranus` (188), `sd-colour-of-the-sky` (365).

With the flagged rows: mean 34.30, sample SD 53.89, n = 213. Without them: mean 26.27, sample SD 25.51, n = 206. Both results are kept, and every later calculation uses all 213 rows.

z-score for `fermat` (Fermat's Last Theorem): $z = (357 - 34.30)/53.89 = 5.99$. This problem stayed open 5.99 sample standard deviations above the mean.

Within 1, 2 and 3 sample SDs of the mean: 0.9390, 0.9718 and 0.9812 of rows as proportions. The histogram is right-skewed with skewness 4.39, and the mean minus one SD is negative while Y has a floor of zero, so a bell-shaped model fits poorly; the proportions sit well above 0.68 at 1 SD and below 0.997 at 3 SDs. Chebyshev's theorem guarantees at least 75% within 2 SDs for any distribution, and the data show 0.9718.

From class midpoints: mean 41.41 and sample SD 51.76, against 34.30 and 53.89 from the raw data. The differences are 7.11 and −2.12 years.

Branch	Count	Mean Y, years
applied	26	20.69
biophysics	20	33.20
chemistry	14	25.93
cosmology	28	32.39
information	36	18.64
mathematics	36	66.94
mind	7	44.29
physics	46	31.35

Weighted mean of the branch means: $26(20.69) + 20(33.20) + 14(25.93) + 28(32.39) + 36(18.64) + 36(66.94) + 7(44.29) + 46(31.35)$, divided by 213, equals 34.30. The overall mean is 34.30, so the two agree.

2.4 Relationships


Paired count: 213. $r = -0.68$. Regression equation: $\hat{y} = 848.99 + (-0.42)x$, with x the year posed and $\hat{y}$ the years to resolve. $R^2 = 0.46$. Sum of squared residuals: 331048.09.

Slope: each later posed year goes with 0.42 fewer years to resolve on average. Intercept: 848.99 years is the fitted value at year 0, far outside the observed years 1506 to 2023, so it has no practical reading. R^2: year posed accounts for 46.22% of the variation in years to resolve. The relationship is negative, roughly linear at the old end and weak to moderate in strength. The residuals fan out: they are wide for problems posed before 1900 and narrow for recent ones, and they are bounded below because Y cannot be negative.


Prediction at x = 1950, inside the observed range: $\hat{y} = 848.99 + (-0.42)(1950) = 27.96$ years. For `fermat` (Fermat's Last Theorem): x = 1637, observed y = 357, fitted 159.75, residual 197.25 years.

Extrapolation: the line gives negative years for posed years after 2016, so it cannot be used beyond the data. Possible confounder: branch, since older rows come from mathematics and physics. Causal limit: the data are observational and hold solved problems only. A problem posed in 2010 can appear here with at most 16 years, so part of the negative slope comes from which rows can be in the table, and no causal claim follows.


2.5 Graphs And Interpretation




Graph 1: physics has the most problems, 46, and mind the fewest, 7.



Graph 2: the two largest branches, physics and mathematics, hold 38.50% of the problems.



Graph 3: 147 of 213 problems fall in the first class, and counts drop off to a long right tail.




Graph 4: the dots pile up from 0 to about 45 years, thin out by 100, and leave gaps before the largest values.

Graph 5. Stem plot of years to resolve. Key: `12 | 3` means 123 years; the stem is tens of years and each leaf is one problem.


```
0 | 000000000000011111111111112222222223333333334444444445555555666666777888889999 9
1 | 000011222223555566777778999
2 | 00222223444445566778
3 | 111122334566667888899
4 | 00112233344555666789999
5 | 112233456779
6 | 01223347
7 | 00139
8 | 003589
```

```
 9 | 19
10 | 014
11 |
12 | 4
13 |
14 | 3
15 |
16 |
17 |
18 | 8
19 |
20 |
21 |
22 |
23 |
24 |
25 |
26 |
27 |
28 |
29 |
30 |
31 |
32 |
33 | 0
34 |
35 | 7
36 | 5
37 |
38 | 7
```


Graph 5: 203 of 213 leaves sit on stems 0 to 9, and stems 20 to 32 are empty.




Graph 6: the polygon peaks at the first midpoint, 20 years, and falls steeply.




Graph 7: a proportion of 0.6901 of problems resolved in under 40 years and 0.9155 in under 80.




Graph 8: the box runs from 5 to 45 years with the median at 20, and 7 points sit beyond the upper fence of 105.00.




Graph 9: years to resolve fall as the posed year rises, $r = -0.68$.



Graph 10: the residuals spread wider at large fitted values and have a hard lower edge, so the linear model's equal-spread assumption fails.



Graph 11: with the axis cut, the smallest branch looks close to empty and differences between branches look larger than they are.



Graph 12: from a zero baseline the bar heights are proportional to the counts.

Shape: one cluster near zero, strong right skew, gaps above 200 years and 7 flagged values. The median, 20.00 years, is the main measure of centre, because the long right tail pulls the mean up

to 34.30.

3 Inferential Statistics

3.1 Probability

Cutoff c = 50 years. A: a problem took at least 50 years. B: the problem is in physics.

	A	not A	Total
B	7	39	46
not B	36	131	167
Total	43	170	213

$P(A) = 43/213 = 0.2019$. $P(B) = 46/213 = 0.2160$. $P(A \text{ and } B) = 7/213 = 0.0329$. $P(A \text{ or } B) = 82/213 = 0.3850$. $P(A \mid B) = 7/46 = 0.1522$. $P(A)P(B) = 0.0436$, against $P(A \text{ and } B) = 0.0329$: the two differ by 0.0107, so A and B are dependent in these data. $C(10, 3) = 120$ counts the ways to choose which 3 of 10 draws are the successes.

3.2 Models

Binomial. K counts A-successes in 10 uniform draws with replacement, p = 0.2019. Conditions: a fixed number of 10 trials; two outcomes, A or not A; a constant p because draws are with replacement; independent draws. $P(K = 3) = C(10, 3)\, p^3(1-p)^7 = 0.2037$. $P(K \geq 1) = 1 - (1-p)^{10} = 0.8951$. Mean $= 10p = 2.02$. SD $= \sqrt{10p(1-p)} = 1.27$.


Normal. Full-data mean $\mu = 34.30$ and population SD $\sigma = 53.76$ years. $P(Y \geq 50) = 0.3851$ under the model, against the observed proportion 0.2019. Modelled 90th percentile: 103.19 years, against 70.00 observed. The histogram is right-skewed and the model puts probability 0.2617 below zero years, so the fit is weak and these two results are model exercises.

3.3 Sampling

Reference population: all N = 213 cleaned rows, $\mu = 34.30$ and $\sigma = 53.76$ years. Method: `numpy.random.default_rng(124).integers(1, N + 1, 50)` draws 50 independent uniform positions with replacement. Draws that repeat an earlier row: 1. Positions, row IDs and values are saved in `sample_50.csv`.

Simulation: 500 samples of size 5 under seed 125 and 500 of size 50 under seed 126, separate from the saved sample. Every draw ID is in `simulation_draws.csv` and all 1,000 means are in `simulation_means.csv`.

Sample size	Mean of means	SD of means	μ	$\sigma/\sqrt{n}$
n = 5	35.20	24.86	34.30	24.04
n = 50	35.26	7.84	34.30	7.60



Both histograms centre near μ, marked by the line. The spread shrinks from 24.86 at n = 5 to 7.84 at n = 50, close to $\sigma/\sqrt{n}$. At n = 5 the means keep the right skew of Y; at n = 50 the shape is closer to symmetric, as the central limit theorem predicts.

3.4 Confidence Interval

From the saved sample: n = 50, $\bar{x} = 26.60$, s = 27.84, SE $= 27.84/\sqrt{50} = 3.94$, df = 49, $t^* = 2.01$, margin $= 2.01 \times 3.94 = 7.91$. Interval: 26.60 ± 7.91, from 18.69 to 34.51 years.

Checks: the draws are independent by design. The sample is right-skewed, with median 13.50 and maximum 104 years, and n = 50 is large enough for the t procedure to be approximate; outliers widen the interval. Interpretation: we are 95% confident that the mean time from posed to resolved for the 213 cleaned atlas problems is between 18.69 and 34.51 years. Across repeated samples of 50 drawn this way, about 95% of the intervals built by this method would contain the reference mean. This interval contains the full-data mean of 34.30 years. The sample is kept as drawn.

3.5 Hypothesis Test

$H_0 : \mu = 50$, $H_1 : \mu \neq 50$, $\alpha = 0.05$. The benchmark of 50 years is hypothetical. $t = (26.60 - 50)/3.94 = -5.94$, df = 49, p-value < 0.0001. Decision: reject H_0. Conclusion: the sample provides evidence that the mean time to resolve differs from 50 years.

The decision agrees with the interval: 50 is outside 18.69 to 34.51. The p-value is the probability, when the mean is 50 years, of a sample mean at least this far from 50 in either direction. A Type I error would be concluding the mean differs from 50 years when it equals 50. A Type II error would be keeping 50 years when the mean differs. The sample mean minus the benchmark is −23.40 years, 46.80% of the benchmark.

Analysis

DRAFT for Gianangelo to rewrite in his own words. Target 150 to 200 words.

Years from posed to resolved are right-skewed. Most solved problems in the atlas closed within a few decades: the median is 20.00 years and the third quartile is 45.00, while the mean is 34.30 because 7 problems sit beyond the upper fence of 105.00 years. The median is the better centre. Year posed and years to resolve move in opposite directions, with $r = -0.68$ and $R^2 = 0.46$; part

of that slope comes from the table holding solved problems only, since a recent problem that is already solved has a short time by construction. The saved sample of 50 rows has mean 26.60 years, and its 95% interval, 18.69 to 34.51, contains the full-data mean of 34.30. The test of the 50-year benchmark gives $p < 0.0001$, so the sample leads us to reject H_0. The sample mean differs from the full-data mean by -7.70 years, or 1.95 standard errors of 3.94.

Conclusion

DRAFT for Gianangelo to rewrite in his own words. Target 100 to 150 words.

The research question asks whether solved atlas problems took 50 years on average. The sample provides evidence against H_0: $t = -5.94$, $p < 0.0001$, and the interval runs from 18.69 to 34.51 years. Two limits apply. The table holds solved problems with both years recorded, so long-open problems that are still unsolved are missing and the mean understates how long problems stay open. The rows come from public lists that favour famous problems. One improvement: record a posed year for every row in the atlas, open or solved, so that open problems can enter as still running. The claim covers the 213 cleaned rows and no wider population of research problems.

References

Bucket Foundation, Dichio, G. (2026). *Solvability Atlas problem table*, version of 2026-10-13. https://github.com/bucket-foundation/bucket-foundation/blob/dev/tools/solvability-atlas/surface/table.csv. Accessed 2026-10-13.

Wikipedia contributors. Problem lists, science timelines and prize competition pages. CC BY-SA 4.0. The URL for each row is in `problems-sourced.tsv` in the atlas folder.

Google DeepMind. *formal-conjectures*. Apache-2.0. https://github.com/google-deepmind/formal-conjectures.

Benchmark: $\mu_0 = 50$ years is hypothetical and has no outside source.

Software: Python with numpy, pandas, scipy and matplotlib; `stats_project.py` reproduces every number under seed 124.