

Study

travling at 105km/h for duration of 1h 50min

traffic slows you down to 70km/h. how long will it take

$v = \frac{d}{t}$

$d = v \cdot t$

$= (105\,km/h)(1.833h)$

$= 192.4965\,km$

$t = \frac{d}{v}$

$= \frac{192.4965km}{70km/h}$

$= 2.74995h$

$= 2h\ 45min$

$\therefore$ 55min Longer drive

read question carfully...

it ask how much Longer will it take to drive

A car stopped at traffic light

distance from the light $x(t) = bt^2 - ct^3$, where $b = 2.40 m/s^2$, $c = 0.120 m/s^3$

a) Calculate V_{avg} $t=0$ to $t=10$

$V(t) = 2bt - 3ct^2$

$V(10-0) = 2b(10) - 3c(10)^2$

$= 48 - 36$

$= 12 m/s$

b) find instant V at $t = 0, 5, 10$

$V(0) = 2bt - 3ct^2$

$= 0 m/s$

$V(5) = 2bt - 3ct^2$

$= 15 m/s$

$V(10) = 2bt - 3ct^2$

$= 12 m/s$

C) When is the car at rest again

$0 = 2bt - 3ct^2$

$0 = 4.8t - 0.36t^2$

$0 = t(4.8 - 0.36t)$

$t = 0$

$0.36t = 4.8$

$t = 13.33s$

textbook says answer is $13.3 m/s$?

Speed?

Cars velocity: $V_x(t) = \alpha + \beta t^2$, where $\alpha = 3.00 m/s$ and $\beta = 0.100 m/s^3$

a) calculate avg velocity 0-5s

$V(5) = 5.50 m/s$

$V(0) = 3.00 m/s$

$$a = \frac{5.50 - 3.00}{5.00 - 0}$$


$$= 0.500 m/s^2$$

b) instant a at 0 and 5

$a(0) = 2(0.100)(0) = 0 m/s^2$

$a(5) = 2(0.100)(5) = 1 m/s^2$

c) graphs of v and a from 0-5s



$a(t) = 2(0.100)(t)$

$=$

1 -> 0.2
2 -> 0.4
3 -> 0.6
4 -> 0.8
5 -> 1

Object has cards $(1.1, 3.4)$ at $t_1 = 0$

$(5.3, -0.5)$ at $t_2 = 3.0\text{s}$

a) find the components of avg velocity

$\Delta d_y = y_2 - y_1 = -3.9$


$\Delta d_x = x_2 - x_1 = 4.2$

$$V = \frac{\Delta d}{\Delta t}$$

$V_y = -1.3\text{m/s}$

$V_x = 1.4\text{m/s}$

b) magnitude and direction



$r = \sqrt{1.4^2 + 1.3^2}$

$= 1.9\text{m/s}$ ← magnitude

$\theta = \tan^{-1}\left(\frac{1.3}{1.4}\right)$

$= 43^\circ$ ← direction

$\alpha = 317^\circ$

Bird flying given by $x(t) = \alpha t$ and $y(t) = 3.0m - \beta t^2$

where $\alpha = 2.4 m/s$ and $\beta = 1.2 m/s^2$


a) Sketch path from 1 – 2s

$x(0) = 2.4(0) = 0$ $\quad y(0) = 3.0m - (1.2)(0)^2 = 3.0$

$x(1) = 2.4(1) = 2.4$ $\quad y(1) = 3.0m - (1.2)(1)^2 = 1.8$

$x(2) = 2.4(2) = 4.8$ $\quad y(2) = 3.0m - (1.2)(2)^2 = -1.8$

a)



b) find vector for $\vec{V}$ and $\vec{a}$ as function of time

$V(t) = V_x(t)\hat{i} + V_y(t)\hat{j}$ $\qquad a(t) = a_x(t)\hat{i} + a_y(t)\hat{j}$

$V_x(t) = \alpha$ $\qquad a_x(t) = 0$


$V_y(t) = -2\beta t$ $\qquad a_y(t) = -2\beta$

$\boxed{V(t) = (2.4\hat{i} - 2\beta t\hat{j})\, m/s}$ $\qquad \boxed{a(t) = -2\beta\hat{j}\, m/s^2}$

c) find mag and direction for $\vec{V}$ and $\vec{a}$ at $t = 2.0s$

$V_x(2) = \alpha = 2.4 m/s$ $\qquad a_x(2) = 0$

$V_y(2) = -2(1.2)(2) = -4.8 m/s$ $\qquad a_y(2) = -2(1.2) = -2.4 m/s^2$




$r = \sqrt{2.4^2 + 4.8^2}$

$= \boxed{5.4 m/s}$

$\theta = \tan^{-1}\left(\frac{4.8}{2.4}\right)$

$= 63° \Rightarrow \boxed{297°}$




$\boxed{\text{mag} = 2.4 m/s^2}$

$\boxed{\theta = 270°}$

d) Speeding up, turning right

book slides off table, speed of 1.10 m/s. it hits the floor at 0.480 s



horizontal velocity is constant throughout

a) hight above floor

$$h = \cancel{v_i \Delta t^0} + \frac{1}{2} g \Delta t^2$$

$$= \frac{1}{2} g \Delta t^2$$

$$= \frac{1}{2}(9.80 m/s^2)(0.480 s)^2$$

$$= 1.13 m$$

b) horizontal distance

$$d = v_{ox} t$$


$$= (1.10 m/s)(0.480 s)$$

$$= 0.528 m$$

c) horizontal and vertical components befor hits ground

$$v_x = 1.10 m/s$$

$$v_y = -gt = -4.70 m/s$$



$$r = \sqrt{1.1^2 + 4.7^2}$$

$$= 4.83 m/s$$

$$\theta = \tan^{-1}\left(\frac{4.7}{1.1}\right)$$

$$= -76.8^\circ$$

$$= 283^\circ$$


d)

two jump off a cliff

- Person 1 reaches the ground in 2.7s dropping vertically
- Person 2 jumps out with 95 cm/s

a) how far out will they jump

$95 \frac{cm}{s} \times \frac{1m}{100 cm} = 0.95 m/s$



$h = \frac{1}{2} g t^2$

$= \frac{1}{2}(9.8)(2.7)^2$

$= 35.721m$

Constant

$V_x = 0.95 m/s$

$V = \frac{d}{t}$


$d = v \cdot t$

$= (0.95 m/s)(2.70s)$


$= 2.57m$

we only look at horizontal numbers to find horizontal distance

ball drops off a table on earth



Same experament but on a different planet



a) whats the planets g?

V_x constant

$t = \sqrt{\frac{2h}{g}}$

$d = V_0 t$

$= V_0 \sqrt{\frac{2h}{g}}$

on earth $D = V_0 \sqrt{\frac{2h}{g_{earth}}}$

on planet $D = V_0 \sqrt{\frac{2h}{g_x}}$

$$\frac{1.93D}{D} = \frac{\sqrt{\frac{2h}{g_x}}}{\sqrt{\frac{2h}{g_{earth}}}} \Rightarrow 1.93 = \sqrt{\frac{g_e}{g_x}}$$

$$1.93^2 = \frac{g_e}{g_x}$$

$$\frac{g_e}{1.93^2} = g_x$$

$$2.63 m/s^2 = g_x$$

textbook says answer is $1.28 m/s^2$. GPT says wrong