

Subgroups of Hyperbolic Groups

A Default Beamer Presentation

David Buckley

Mathematics Department
University of Warwick

Postgraduate Group Theory Conference, 2009

Cayley Graphs

- Group G generated by set X .
- Cayley graph of G is $\text{Cay}(G, X)$

Cayley Graphs

- Group G generated by set X .
- Cayley graph of G is $\text{Cay}(G, X)$:
- Vertex-set G , edge set $\{(g, gx) : g \in G, x \in X\}$.

Cayley Graphs

- Group G generated by set X .
- Cayley graph of G is $\text{Cay}(G, X)$:
- Vertex-set G , edge set $\{(g, gx) : g \in G, x \in X\}$.
- Subgroup $H \leq G$.
- Coset Cayley graph of G relative to H

Cayley Graphs


- Group G generated by set X .
- Cayley graph of G is $\text{Cay}(G, X)$:
- Vertex-set G , edge set $\{(g, gx) : g \in G, x \in X\}$.
- Subgroup $H \leq G$.
- Coset Cayley graph of G relative to H :
- Vertex-set $\{Hg : g \in G\}$, edge set $\{(Hg, Hgx) : g \in G, x \in X\}$.

Hyperbolic Space

- Hyperbolic spaces curve "inwards."
- This makes geodesic triangles thin.


Hyperbolic Space

- Hyperbolic spaces curve "inwards."
- This makes geodesic triangles thin.




Hyperbolic Space

- Hyperbolic spaces curve "inwards."
- This makes geodesic triangles thin.



Hyperbolic Space

- Hyperbolic spaces curve "inwards."
- This makes geodesic triangles thin.



- Hyperbolic if all geodesic triangles are δ -thin.

Divergence

- Metric space X .
- A map $e : [0, \infty] \rightarrow [0, \infty]$ is a divergence function if:
 - When $r_1, r_2 \subset X$ two rays which start at the same point
 - and $d(r_1(a), r_2(a)) > e(0)$ for some $a > 0$
 - then $d_{a+b}(r_1(a+b), r_2(a+b)) > e(b)$ for all $b > 0$ (d_l means distance outside of $B_l(1)$).

Divergence

- Metric space X .
- A map $e : [0, \infty] \rightarrow [0, \infty]$ is a divergence function if:
 - When $r_1, r_2 \subset X$ two rays which start at the same point
 - and $d(r_1(a), r_2(a)) > e(a)$ for some $a > 0$
 - then $d_{a+b}(r_1(a+b), r_2(a+b)) > e(b)$ for all $b > 0$ (d_l means distance outside of $B_l(1)$).
- Hyperbolic spaces have exponential divergence functions.

Divergence

- Metric space X .
- A map $e : [0, \infty] \rightarrow [0, \infty]$ is a divergence function if:
 - When $r_1, r_2 \subset X$ two rays which start at the same point
 - and $d(r_1(a), r_2(a)) > e(0)$ for some $a > 0$
 - then $d_{a+b}(r_1(a+b), r_2(a+b)) > e(b)$ for all $b > 0$ (d_l means distance outside of $B_l(1)$).
- Hyperbolic spaces have exponential divergence functions.
- Will assume divergence is strict:
- Then $e(b) < d_{a+b}(r_1(a+b), r_2(a+b)) < ke(b)$ for all $b > 0$.

Hyperbolic Groups

- The Cayley graph of a finitely generated group is a geodesic metric space.
- A group is hyperbolic if its Cayley graph is hyperbolic.
- Examples:

Hyperbolic Groups

- The Cayley graph of a finitely generated group is a geodesic metric space.
- A group is hyperbolic if its Cayley graph is hyperbolic.
- Examples:
 - Finite groups.

Hyperbolic Groups

- The Cayley graph of a finitely generated group is a geodesic metric space.
- A group is hyperbolic if its Cayley graph is hyperbolic.
- Examples:
 - Finite groups.
 - Free groups of finite rank.

Hyperbolic Groups

- The Cayley graph of a finitely generated group is a geodesic metric space.
- A group is hyperbolic if its Cayley graph is hyperbolic.
- Examples:
 - Finite groups.
 - Free groups of finite rank.
 - Groups acting properly and cocompactly on hyperbolic spaces.

Hyperbolic Groups


- The Cayley graph of a finitely generated group is a geodesic metric space.
- A group is hyperbolic if its Cayley graph is hyperbolic.
- Examples:
 - Finite groups.
 - Free groups of finite rank.
 - Groups acting properly and cocompactly on hyperbolic spaces.
- Non-example: $\mathbb{Z} \times \mathbb{Z}$

Hyperbolic Groups

- The Cayley graph of a finitely generated group is a geodesic metric space.
- A group is hyperbolic if its Cayley graph is hyperbolic.
- Examples:
 - Finite groups.
 - Free groups of finite rank.
 - Groups acting properly and cocompactly on hyperbolic spaces.
- Non-example: $\mathbb{Z} \times \mathbb{Z}$
- Fix $G = \langle X | R \rangle$ a finitely generated group with δ -hyperbolic Cayley graph Γ .
- Let $A = X \cup X^{-1}$.


Quasiconvex Subgroups

- Let w be a word in A^* .
- Let α be the path starting at 1 and labelled by w .
- Let $H < G$ any subgroup.
- $d(w, H)$ is the max distance from a point on α to closest point in H .



Quasiconvex Subgroups


- Let w be a word in A^* .
- Let α be the path starting at 1 and labelled by w .
- Let $H < G$ any subgroup.
- $d(w, H)$ is the max distance from a point on α to closest point in H .



- H ε -quasiconvex if $d(w, H) \leq \varepsilon$ for any $w \in H$ which labels a geodesic.

Quasiconvex Subgroups

- Let w be a word in A^* .
- Let α be the path starting at 1 and labelled by w .
- Let $H < G$ any subgroup.
- $d(w, H)$ is the max distance from a point on α to closest point in H .



- H ϵ -quasiconvex if $d(w, H) \leq \epsilon$ for any $w \in H$ which labels a geodesic.
- For instance, $\langle a^2 \rangle$ is a 1-quasiconvex subgroup of $F(a, b)$.

Cut points

- Connected topological space X
- Some point $x \in X$
- x is a cut point if $X \setminus \{x\}$ is disconnected
- Example: Every point in \mathbb{R} is a cut point

Cut points

- Connected topological space X
- Some point $x \in X$
- x is a cut point if $X \setminus \{x\}$ is disconnected
- Example: Every point in \mathbb{R} is a cut point
- Subset $Y \subset X$
- Y is a cut set if $X \setminus Y$ is disconnected
- Example: Every closed 1-ball in $\mathbb{R} \times [0, 1]$ is a cut set.

Cut points

- Connected topological space X
- Some point $x \in X$
- x is a cut point if $X \setminus \{x\}$ is disconnected
- Example: Every point in \mathbb{R} is a cut point
- Subset $Y \subset X$
- Y is a cut set if $X \setminus Y$ is disconnected
- Example: Every closed 1-ball in $\mathbb{R} \times [0, 1]$ is a cut set.
- Y cuts set A from set B if $A \setminus Y$ and $B \setminus Y$ lie in different components of $X \setminus Y$.

Ends

- Metric space X
- Geodesic rays r_1, r_2 (path isometric to $[0, \infty]$)
- Say $r_1 \equiv r_2$ if there is no bounded set Y which cuts r_1 from r_2 .
- Space of ends is space of equivalence classes of rays r .

Ends

- Metric space X
- Geodesic rays r_1, r_2 (path isometric to $[0, \infty]$)
- Say $r_1 \equiv r_2$ if there is no bounded set Y which cuts r_1 from r_2 .
- Space of ends is space of equivalence classes of rays r .
- Compare to boundary - equivalence classes of geodesic rays under $r \equiv s$ if there exists M such that $d(x, s) \leq M$ for all $x \in r$ and vice versa.

Number of Ends

- Interested in *number* of ends $n(X)$.
- For a Cayley graph there are 0, 1, 2 or ∞ ends.

Number of Ends

- Interested in *number* of ends $n(X)$.
- For a Cayley graph there are 0, 1, 2 or ∞ ends.
- Example: Finite groups have 0-ended Cayley graphs (there are no geodesic rays).

Number of Ends

- Interested in *number* of ends $n(X)$.
- For a Cayley graph there are 0, 1, 2 or ∞ ends.
- Example: Finite groups have 0-ended Cayley graphs (there are no geodesic rays).
- Example: Hyperbolic triangle groups have 1-ended Cayley graphs (since they represent tilings of hyperbolic 2-space).

Number of Ends

- Interested in *number* of ends $n(X)$.
- For a Cayley graph there are 0, 1, 2 or ∞ ends.
- Example: Finite groups have 0-ended Cayley graphs (there are no geodesic rays).
- Example: Hyperbolic triangle groups have 1-ended Cayley graphs (since they represent tilings of hyperbolic 2-space).
- Example: \mathbb{Z} has a 2-ended Cayley graph (there's precisely 2 directions).

Number of Ends

- Interested in *number* of ends $n(X)$.
- For a Cayley graph there are 0, 1, 2 or ∞ ends.
- Example: Finite groups have 0-ended Cayley graphs (there are no geodesic rays).
- Example: Hyperbolic triangle groups have 1-ended Cayley graphs (since they represent tilings of hyperbolic 2-space).
- Example: \mathbb{Z} has a 2-ended Cayley graph (there's precisely 2 directions).
- Example: Non-cyclic free groups have ∞ -ended Cayley graphs (any string of positive powers like abb can be repeated indefinitely to give a unique end).

Problem


- G is a hyperbolic group with Cayley graph Γ .
- G has 1 end.
- H is a quasiconvex subgroup.
- Γ has a strict divergence function e .
- What can we say about ends of coset Cayley graph?

Example

- Let G be the hyperbolic triangle group $\langle x, y | x^2, y^3, (xy)^7 \rangle$.
- Let $H = \langle xyxy^{-1} \rangle$ (infinite, infinite index).

Example

- Let G be the hyperbolic triangle group $\langle x, y | x^2, y^3, (xy)^7 \rangle$.
- Let $H = \langle xyxy^{-1} \rangle$ (infinite, infinite index).



A Neat Trick

- Suppose w , a and b are words.
- Also, that w is long enough (about $\varepsilon + 2\delta$).

A Neat Trick

- Suppose w , a and b are words.
- Also, that w is long enough (about $\varepsilon + 2\delta$).
- Suppose wa does not have much cancellation
- (Roughly, $|wa| > |w| + |a| - k$).

A Neat Trick

- Suppose w , a and b are words.
- Also, that w is long enough (about $\varepsilon + 2\delta$).
- Suppose wa does not have much cancellation
- (Roughly, $|wa| > |w| + |a| - k$).
- Then Hwa does not have much cancellation.
- (Roughly, $d(H, Hwa) > d(H, Hw) + |a| - k - \varepsilon - 2\delta$).

Divergence Made Useful

- Divergence is a bit of a pain to use, but strict divergence is (obviously) easier!
- Suppose $d(r_1(a), r_2(a)) > e(0)$ with a minimal.
- If α is a path connecting $r_1(a+b)$ to $r_2(a+b)$, then there exists a word w of length nearly a and near-suffixes $r'_1 =_G w^{-1}r_1$ and $r'_2 =_G w^{-1}r_2$ such that:

Divergence Made Useful

- Divergence is a bit of a pain to use, but strict divergence is (obviously) easier!
- Suppose $d(r_1(a), r_2(a)) > e(0)$ with a minimal.
- If α is a path connecting $r_1(a+b)$ to $r_2(a+b)$, then there exists a word w of length nearly a and near-suffixes $r'_1 =_G w^{-1}r_1$ and $r'_2 =_G w^{-1}r_2$ such that:
- wr'_1 and wr'_2 don't have much cancellation
- (Roughly $e^{-1}(k)$ where k is the k you got along with e).

Divergence Made Useful

- Divergence is a bit of a pain to use, but strict divergence is (obviously) easier!
- Suppose $d(r_1(a), r_2(a)) > e(0)$ with a minimal.
- If α is a path connecting $r_1(a+b)$ to $r_2(a+b)$, then there exists a word w of length nearly a and near-suffixes $r'_1 =_G w^{-1}r_1$ and $r'_2 =_G w^{-1}r_2$ such that:
- wr'_1 and wr'_2 don't have much cancellation
- (Roughly $e^{-1}(k)$ where k is the k you got along with e).
- for all paths x to points in α , there is $x' =_G w^{-1}x$ such that wx' don't have much cancellation.

Wave Hands = Proof

- So there's only a finite number of ends of G relative to H .

Wave Hands = Proof

- So there's only a finite number of ends of G relative to H .
- Proof? You want proof?


Wave Hands = Proof

- So there's only a finite number of ends of G relative to H .
- Proof? You want proof?
- Take a huge ball $B_K(H)$.

H
●


Wave Hands = Proof

- So there's only a finite number of ends of G relative to H .
- Proof? You want proof?
- Let Hx, Hy be points in not in the ball, and in infinite components.




Wave Hands = Proof

- So there's only a finite number of ends of G relative to H .
- Proof? You want proof?
- Since they're in infinite components, may as well assume $d(Hx, H) \geq 2K$ and $d(Hy, H) \geq 2K$.




Wave Hands = Proof

- So there's only a finite number of ends of G relative to H .
- Proof? You want proof?
- There's a path α connecting x to y in the group Cayley graph



Wave Hands = Proof

- So there's only a finite number of ends of G relative to H .
- Proof? You want proof?
- There's a magical path w which doesn't cancel too much with x , y , or paths from 1 to points on α .



Wave Hands = Proof

- So there's only a finite number of ends of G relative to H .
- Proof? You want proof?
- If w is big enough, all points Hx on $H\alpha$ have $d(H, Hx)$ a bit lower than K . Thus $H\alpha$ connects Hx and Hy and lies outside the ball.
- Basically, this means the component a point Hx lies in is determined completely by its behaviour inside a small ball around H .
- There's only a finite number of short prefix words, so only a finite number of ends.