

# Conjugacy in Hyperbolic Groups

## A Short Recipe for Testing Conjugacy in Quasiconvex Subgroups

David Buckley

Mathematics Department  
University of Warwick


Postgraduate Group Theory Conference, 2008

# Hyperbolic Space

- Hyperbolic spaces curve "inwards."
- This makes geodesic triangles thin.


# Hyperbolic Space

- Hyperbolic spaces curve "inwards."
- This makes geodesic triangles thin.




# Hyperbolic Space

- Hyperbolic spaces curve "inwards."
- This makes geodesic triangles thin.



# Hyperbolic Space

- Hyperbolic spaces curve "inwards."
- This makes geodesic triangles thin.



- Hyperbolic if all geodesic triangles are  $\delta$ -thin.

# Hyperbolic Groups

- The Cayley graph of a finitely generated group is a geodesic metric space.
- A group is hyperbolic if its Cayley graph is hyperbolic.
- Examples:

# Hyperbolic Groups

- The Cayley graph of a finitely generated group is a geodesic metric space.
- A group is hyperbolic if its Cayley graph is hyperbolic.
- Examples:
  - Finite groups.

# Hyperbolic Groups

- The Cayley graph of a finitely generated group is a geodesic metric space.
- A group is hyperbolic if its Cayley graph is hyperbolic.
- Examples:
  - Finite groups.
  - Free groups of finite rank.



# Hyperbolic Groups

- The Cayley graph of a finitely generated group is a geodesic metric space.
- A group is hyperbolic if its Cayley graph is hyperbolic.
- Examples:
  - Finite groups.
  - Free groups of finite rank.
  - Groups acting properly and cocompactly on hyperbolic spaces.

# Hyperbolic Groups

- The Cayley graph of a finitely generated group is a geodesic metric space.
- A group is hyperbolic if its Cayley graph is hyperbolic.
- Examples:
  - Finite groups.
  - Free groups of finite rank.
  - Groups acting properly and cocompactly on hyperbolic spaces.
- Non-example:  $\mathbb{Z} \times \mathbb{Z}$

# Hyperbolic Groups


- The Cayley graph of a finitely generated group is a geodesic metric space.
- A group is hyperbolic if its Cayley graph is hyperbolic.
- Examples:
  - Finite groups.
  - Free groups of finite rank.
  - Groups acting properly and cocompactly on hyperbolic spaces.
- Non-example:  $\mathbb{Z} \times \mathbb{Z}$
- Fix  $G = \langle X | R \rangle$  a finitely generated group with  $\delta$ -hyperbolic Cayley graph  $\Gamma$ .
- Let  $A = X \cup X^{-1}$ .

# Quasiconvex Subgroups

- Let  $w$  be a word in  $A^*$ .
- Let  $\alpha$  be the path starting at 1 and labelled by  $w$ .
- Let  $H < G$  any subgroup.


# Quasiconvex Subgroups

- Let  $w$  be a word in  $A^*$ .
- Let  $\alpha$  be the path starting at 1 and labelled by  $w$ .
- Let  $H < G$  any subgroup.
- $d(w, H)$  is the max distance from a point on  $\alpha$  to closest point in  $H$ .



# Quasiconvex Subgroups


- Let  $w$  be a word in  $A^*$ .
- Let  $\alpha$  be the path starting at 1 and labelled by  $w$ .
- Let  $H < G$  any subgroup.
- $d(w, H)$  is the max distance from a point on  $\alpha$  to closest point in  $H$ .



- $H$   $\epsilon$ -quasiconvex if  $d(w, H) \leq \epsilon$  for any  $w \in H$  which labels a geodesic.

# Quasiconvex Subgroups

- Let  $w$  be a word in  $A^*$ .
- Let  $\alpha$  be the path starting at 1 and labelled by  $w$ .
- Let  $H < G$  any subgroup.
- $d(w, H)$  is the max distance from a point on  $\alpha$  to closest point in  $H$ .



- $H$   $\varepsilon$ -quasiconvex if  $d(w, H) \leq \varepsilon$  for any  $w \in H$  which labels a geodesic.
- For instance,  $\langle a^2 \rangle$  is a 1-quasiconvex subgroup of  $F(a, b)$ .

# The Problem

- We're given a word  $u \in A^*$ .
- In time  $O(|u|)$ , we must:



# The Problem

- We're given a word  $u \in A^*$ .
- In time  $O(|u|)$ , we must:
  - Check if there exists  $w \in A^*$  such that  $wuw^{-1} \in H$ .

# The Problem

- We're given a word  $u \in A^*$ .
- In time  $O(|u|)$ , we must:
  - Check if there exists  $w \in A^*$  such that  $wuw^{-1} \in H$ .
  - Return such a  $w$ , if one exists.

# The Problem

- We're given a word  $u \in A^*$ .
- In time  $O(|u|)$ , we must:
  - Check if there exists  $w \in A^*$  such that  $wuw^{-1} \in H$ .
  - Return such a  $w$ , if one exists.
- We assume  $G$  and  $H$  are fixed.
- Exponential running time with respect to  $\delta + \varepsilon$ .

# Shortest Words

- Given any word  $w \in A^*$ , find some  $w' \in A^*$  such that:
  - $w' =_G w$ , and
  - $|w'|$  is minimal

# Shortest Words

- Given any word  $w \in A^*$ , find some  $w' \in A^*$  such that:
  - $w' =_G w$ , and
  - $|w'|$  is minimal
- Shapiro's algorithm in "The linearity of the conjugacy problem in word hyperbolic groups" (Epstein+Holt, 2006).
- Runs in time linear in  $O(|w|)$ .

# Shortest Words

- Given any word  $w \in A^*$ , find some  $w' \in A^*$  such that:
  - $w' =_G w$ , and
  - $|w'|$  is minimal
- Shapiro's algorithm in “The linearity of the conjugacy problem in word hyperbolic groups” (Epstein+Holt, 2006).
- Runs in time linear in  $O(|w|)$ .
- Assume that **all** words are shortest.
- If not, just run Shapiro's algorithm.

# Minimal Conjugates

- Given any word  $u \in A^*$ , find some  $u', w \in A^*$  such that:
  - $u' =_G w^{-1}uw$ , and
  - $|u'|$  is minimal

# Minimal Conjugates

- Given any word  $u \in A^*$ , find some  $u', w \in A^*$  such that:
  - $u' =_G w^{-1}uw$ , and
  - $|u'|$  is minimal
- Strictly speaking, do not have this.



# Minimal Conjugates

- Given any word  $u \in A^*$ , find some  $u', w \in A^*$  such that:
  - $u' =_G w^{-1}uw$ , and
  - $|u'|$  is minimal
- Strictly speaking, do not have this.
- Do have an algorithm computing this for  $u^M$  where  $M$  is some constant (dependent on  $G$ ).
- Sufficient; compute with  $u^M$ , then use centraliser of  $u^M$  to check everything for  $u$ .

# Minimal Conjugates

- Given any word  $u \in A^*$ , find some  $u', w \in A^*$  such that:
  - $u' =_G w^{-1}uw$ , and
  - $|u'|$  is minimal
- Strictly speaking, do not have this.
- Do have an algorithm computing this for  $u^M$  where  $M$  is some constant (dependent on  $G$ ).
- Sufficient; compute with  $u^M$ , then use centraliser of  $u^M$  to check everything for  $u$ .
- We therefore assume  $u$  in the problem is its own minimal conjugate.

# The Conjugacy Problem

- Given any words  $u, v \in A^*$ :
  - Check if there is a  $g \in A^*$  with  $u =_G g^{-1}vg$ .
  - Return some such  $g$  if one exists.

# The Conjugacy Problem

- Given any words  $u, v \in A^*$ :
  - Check if there is a  $g \in A^*$  with  $u =_G g^{-1}vg$ .
  - Return some such  $g$  if one exists.
- Solved by Epstein and Holt (2006).
- Runs in time  $O(|u| + |v|)$ .

# Idea

- Suppose  $g \in G$  has  $gug^{-1} \in H$  as required.

# Idea

- Suppose  $g \in G$  has  $gug^{-1} \in H$  as required.
- Pick  $h \in H$  and  $w \in A^*$  such that  $w =_G hg$  and  $|w|$  is minimal.
- Then  $wuw^{-1} = v \in H$ .
- Aim is to either:

# Idea

- Suppose  $g \in G$  has  $gug^{-1} \in H$  as required.
- Pick  $h \in H$  and  $w \in A^*$  such that  $w =_G hg$  and  $|w|$  is minimal.
- Then  $wuw^{-1} = v \in H$ .
- Aim is to either:
  - find an  $O(1)$  bound on the length of  $w$ , or

# Idea

- Suppose  $g \in G$  has  $gug^{-1} \in H$  as required.
- Pick  $h \in H$  and  $w \in A^*$  such that  $w =_G hg$  and  $|w|$  is minimal.
- Then  $wuw^{-1} = v \in H$ .
- Aim is to either:
  - find an  $O(1)$  bound on the length of  $w$ , or
  - find an  $O(1)$  bound on the length of  $v$




# Idea

- Suppose  $g \in G$  has  $gug^{-1} \in H$  as required.
- Pick  $h \in H$  and  $w \in A^*$  such that  $w =_G hg$  and  $|w|$  is minimal.
- Then  $wuw^{-1} = v \in H$ .
- Aim is to either:
  - find an  $O(1)$  bound on the length of  $w$ , or
  - find an  $O(1)$  bound on the length of  $v$
- Bounded number of checks taking time  $O(|u|)$  means  $O(|u|)$  time.
- Upper bounds will all depend on constants  $\delta$  and  $\varepsilon$ .


# Bounding $|w|$

- Suppose a point on  $u$  corresponds to a point on  $v$ .
- $m$  and  $n$  distances to the meeting points on  $w$  from ends of  $u$ .



## Bounding $|w|$


- Suppose a point on  $u$  corresponds to a point on  $v$ .
- $m$  and  $n$  distances to the meeting points on  $w$  from ends of  $u$ .



- If  $\min\{m, n\} > \delta$  there is a suffix  $w'$  of  $w$  such that  $|w'uw'^{-1}| < |u|$  (contradiction!).


# Bounding $|w|$

- $p$  and  $q$  distances to meeting points on  $w$  from ends of  $v$ .



## Bounding $|w|$


- $p$  and  $q$  distances to meeting points on  $w$  from ends of  $v$ .



- If  $\max\{p, q\} > 2\delta + \varepsilon$  there is a word  $w' \in A^*$  such that  $|w'| < |w|$  and  $hw' =_G w$  for some  $h \in H$  (contradiction!).

## Bounding $|w|$


- $p$  and  $q$  distances to meeting points on  $w$  from ends of  $v$ .



- If  $\max\{p, q\} > 2\delta + \varepsilon$  there is a word  $w' \in A^*$  such that  $|w'| < |w|$  and  $hw' =_G w$  for some  $h \in H$  (contradiction!).

## Bounding $|w|$


- $p$  and  $q$  distances to meeting points on  $w$  from ends of  $v$ .



- If  $\max\{p, q\} > 2\delta + \varepsilon$  there is a word  $w' \in A^*$  such that  $|w'| < |w|$  and  $hw' =_G w$  for some  $h \in H$  (contradiction!).
- $|w| = m + p = n + q$ , so  $|w| \leq 3\delta + \varepsilon$ .

## Bounding $|w|$

- $p$  and  $q$  distances to meeting points on  $w$  from ends of  $v$ .




- If  $\max\{p, q\} > 2\delta + \varepsilon$  there is a word  $w' \in A^*$  such that  $|w'| < |w|$  and  $hw' =_G w$  for some  $h \in H$  (contradiction!).
- $|w| = m + p = n + q$ , so  $|w| \leq 3\delta + \varepsilon$ .
- $|w| \in O(1)$  and there  $O(1)$  possibilities for  $w$ .
- Check if  $wuw^{-1} \in H$  for all  $|w| \leq 3\delta + \varepsilon$ .




# Bounding $|v|$

- Suppose no point on  $u$  corresponds to a point on  $v$ .



# Bounding $|v|$


- Suppose no point on  $u$  corresponds to a point on  $v$ .



- By the same argument as before,  $\max\{p, q\} \leq 2\delta + \varepsilon$ .
- $|v| = p + q \leq 4\delta + 2\varepsilon$ .

# Bounding $|v|$

- Suppose no point on  $u$  corresponds to a point on  $v$ .



- By the same argument as before,  $\max\{p, q\} \leq 2\delta + \varepsilon$ .
- $|v| = p + q \leq 4\delta + 2\varepsilon$ .
- So  $|v| \in O(1)$  and there are  $O(1)$  possibilities for  $v$ .
- For each  $|v| \leq 4\delta + 2\varepsilon$ , check if there is  $w$  such that  $wuw^{-1} = v$ .

## Finished Product

- Can with some more work extend this to checking if, given a list  $a_1, \dots, a_n$ , there is one  $w$  such that  $wa_iw^{-1} \in H$  for all  $i$ .

## Finished Product

- Can with some more work extend this to checking if, given a list  $a_1, \dots, a_n$ , there is one  $w$  such that  $wa_iw^{-1} \in H$  for all  $i$ .
- Useful because if there is,  $\langle a_1, \dots, a_n \rangle$  conjugates into  $H$ .