

A Canonical Curvature Invariant of the Riemann Ξ -Function and Its Numerical Proximity to the Fine-Structure Constant

Author:

Weerasooriya Mahavidanage Buddhika
Independent Researcher - Toronto, Canada
buddhika.official@gmail.com

Abstract

We introduce a geometric framework for studying the completed Riemann Ξ -function through the logarithmic landscape

$$L(s) = \log |\Xi(s)|^2.$$

By examining transverse perturbations of this landscape along the critical line, we define a curvature function $K(\gamma)$ that measures the local rigidity of the Ξ -landscape. Evaluation at the central symmetry point $s = \frac{1}{2}$ yields a distinguished quantity

$$K_0 = \frac{\Xi''(1/2)}{\Xi(1/2)},$$

which we identify as a canonical curvature invariant of the completed Riemann function.

A central contribution of this work is the demonstration that, although raw curvature values depend on coordinate parametrization, a chain-rule-normalized curvature remains invariant under a broad class of linear and nonlinear reparameterizations. Extensive numerical tests show stability at approximately 10^{-12} relative precision, providing strong evidence that K_0 represents a genuine geometric property of the Ξ -landscape rather than a coordinate artifact.

Using this invariant, we define the associated dimensionless spectral quantity

$$N_{\Xi} = \frac{2\pi}{K_0},$$

and obtain the numerical value

$$N_{\Xi} = 135.970291691707664960149 \dots$$

This number emerges entirely from the geometry of the completed Riemann function and contains no fitted parameters or physical inputs. Remarkably, it lies within approximately 0.78% of the inverse fine-structure constant,

$$\alpha^{-1} = 137.035999084 \dots$$

While no mathematical theorem or physical mechanism connecting these quantities is established, the proximity is documented as an empirical observation worthy of further investigation.

We additionally formulate the **Curvature-Floor Conjecture**,

$$K(\gamma) \geq K(0),$$

which is supported by extensive numerical exploration over large intervals of the critical line. If proven, the conjecture would identify the central symmetry point as the location of minimal transverse rigidity of the Ξ -landscape and elevate K_0 from a local curvature quantity to a global geometric invariant of the completed Riemann function.

The results presented here establish a new geometric perspective on the Riemann Ξ -function, introducing a canonical curvature invariant, a corresponding spectral count, and a collection of open mathematical questions concerning spectral geometry, rigidity, and the structure of the Riemann landscape. The work does not claim a derivation of the fine-structure constant or a physical theory connecting number theory to fundamental physics. Rather, it identifies a previously unrecognized geometric invariant of the completed Riemann function whose mathematical properties and unexpected numerical relationships merit further study.

Contents

Abstract.....	2
1. Introduction	7
1.1 Motivation.....	7
1.2 The Importance of Canonical Geometric Invariants	8
1.3 Overview of the Present Result.....	9
2. The Riemann Ξ -Function	10
2.1 Definition of the Completed Ξ -Function	10
2.2 Symmetry and the Functional Equation	11
2.3 The Critical Line and Spectral Interpretation	12
3. The Logarithmic Ξ -Landscape	13
3.1 Definition of the Landscape Function	13
3.2 Longitudinal and Transverse Directions	14
3.3 Local Curvature Structure.....	16
4. Definition of the Curvature Function	18
4.1 Transverse Curvature Along the Critical Line	18
4.2 Derivation of $K(\gamma)$	20
4.3 Positivity Properties.....	21
4.4 Numerical Evaluation Procedure.....	22
5. The Canonical Curvature Invariant K_0	24
5.1 Definition of the Curvature Floor	24
5.2 The Value at $\gamma = \mathbf{0}$	25
5.3 Derivation of	26
5.4 Numerical Evaluation	28
5.5 High-Precision Stability Tests	29
6. Canonicity Under Reparameterization	31
6.1 The Coordinate-Dependence Problem	31
6.2 Chain-Rule Normalization	32
6.3 Linear Rescaling Tests	34

6.4 Nonlinear Reparameterization Tests	36
6.5 Numerical Evidence for Canonicity	37
7. The Spectral Count N_E	39
7.1 Motivation for a Dimensionless Spectral Measure	39
7.2 Definition	40
7.3 Numerical Value	41
7.4 Interpretation as a Geometric Spectral Quantity	42
8. Comparison with the Fine-Structure Constant	43
8.1 Numerical Comparison	43
8.2 Relative Difference	44
8.3 Significance of the 0.8% Proximity	45
8.4 Possible Interpretations	46
8.4.1 Numerical Coincidence	46
8.4.2 Indirect Structural Relationship	46
8.4.3 Deeper Physical Connection	47
8.5 Present Status	48
9. The Curvature-Floor Conjecture	49
9.1 Statement of the Conjecture	49
9.2 Numerical Evidence	51
9.3 Relation to Spectral Rigidity	52
9.4 Connections to the Riemann Zero Spectrum	53
9.5 Open Mathematical Questions	54
10. Discussion	56
10.1 What Has Been Established	56
10.2 What Has Not Been Established	58
10.3 Relation to Number Theory	59
10.4 Relation to Mathematical Physics	60
10.5 Future Directions	61
11. Conclusion	63

11.1 Summary of Results	63
11.2 The Canonical Invariant $K\mathbf{0}$	64
11.3 The Spectral Count $N\mathbf{E}$	65
11.4 Open Problems	66
11.5 Outlook	67
Appendix A. Computational Verification Package.....	68
References	75

1. Introduction

1.1 Motivation

The Riemann zeta function occupies a central position in modern mathematics through its deep connection to the distribution of prime numbers and the structure of analytic number theory. Its completed form,

$$\Xi(s) = \frac{1}{2} s(s-1) \pi^{-s/2} \Gamma\left(\frac{s}{2}\right) \zeta(s),$$

is particularly significant because it is an entire function and satisfies the exact functional symmetry

$$\Xi(s) = \Xi(1-s).$$

This symmetry places the critical line $\text{Re}(s) = 1/2$ at the geometric center of the theory and suggests that the completed function possesses a rich internal structure extending beyond its role in prime number theory.

Historically, research surrounding the zeta function has focused primarily on the location of its zeros and their relation to the Riemann Hypothesis. Comparatively less attention has been devoted to geometric quantities derived from the local shape of the Ξ -landscape itself. In particular, one may ask whether the completed function possesses canonical geometric invariants that characterize its stability, rigidity, or intrinsic curvature.

The present work originates from this question. Rather than studying individual zeros directly, we investigate the geometry of the logarithmic landscape generated by $\Xi(s)$ and define a transverse curvature function along the critical line. Numerical exploration reveals the existence of a distinguished curvature scale associated with the symmetry point $s = 1/2$. Remarkably, this quantity appears to be independent of coordinate choices after proper normalization and yields a dimensionless number that is unexpectedly close to the inverse fine-structure constant.

The purpose of this paper is not to claim a physical derivation of any fundamental constant. Rather, our objective is to establish the mathematical existence of a canonical geometric invariant of the Ξ -function and to document its numerical properties in a precise and reproducible manner.

1.2 The Importance of Canonical Geometric Invariants

A central principle of geometry is that physically or mathematically meaningful quantities should not depend on arbitrary coordinate choices. Quantities that vary under simple reparameterizations generally reflect properties of the chosen coordinates rather than properties of the underlying object itself. Consequently, the search for canonical invariants plays a fundamental role throughout mathematics and theoretical physics.

Examples abound. In differential geometry, scalar curvature provides an invariant characterization of local geometric structure. In spectral geometry, eigenvalues of self-adjoint operators encode intrinsic information about a manifold independent of coordinate descriptions. In dynamical systems, invariant measures and Lyapunov exponents characterize properties that persist under changes of variables.

The same principle applies to the Riemann Ξ -function. If one defines a curvature-like quantity associated with the Ξ -landscape, it is essential to determine whether that quantity represents genuine geometry or merely an artifact of parametrization. Any candidate invariant must therefore survive coordinate transformations and admit a normalization procedure that removes arbitrary scale dependence.

A major focus of the present work is precisely this issue. We identify a transverse curvature quantity and demonstrate numerically that, after chain-rule normalization, it remains stable under a broad class of coordinate transformations, including linear scalings and nonlinear reparameterizations. This evidence suggests that the resulting quantity is not merely a coordinate-dependent curvature measure but rather a genuine geometric invariant of the Ξ -landscape.

1.3 Overview of the Present Result

The main result of this paper is the identification of a distinguished curvature invariant associated with the completed Riemann function. Beginning from the logarithmic landscape generated by $\Xi(s)$, we define a transverse curvature function $K(\gamma)$ along the critical line. Evaluation at the symmetry point yields

$$K_0 = \frac{\Xi''(1/2)}{\Xi(1/2)} = 0.04620998623306458 \dots$$

We then investigate the behavior of this quantity under coordinate transformations. While the raw curvature changes as expected under rescaling, a chain-rule-normalized version remains invariant to numerical precision across a wide range of linear and nonlinear reparameterizations. This provides strong evidence that K_0 represents a canonical geometric property of the Ξ -landscape rather than a coordinate artifact.

From this invariant we define the associated dimensionless spectral quantity

$$N_\Xi = \frac{2\pi}{K_0} = 135.9702916917 \dots$$

A notable observation is that this number lies within approximately 0.8% of the inverse fine-structure constant,

$$\alpha^{-1} = 137.035999084 \dots$$

No physical identification is assumed in the present work, and no claim is made that these quantities are equal. Rather, the numerical proximity is recorded as an empirical fact whose significance remains unknown.

The paper proceeds by defining the curvature construction, establishing its canonicity properties, presenting high-precision numerical results, and formulating a curvature-floor conjecture that emerges naturally from the observed structure of the Ξ -landscape. The broader mathematical and physical implications of these findings are discussed, while carefully distinguishing established results from open questions.

2. The Riemann Ξ -Function

2.1 Definition of the Completed Ξ -Function

The Riemann zeta function is initially defined for complex numbers with $\text{Re}(s) > 1$ by the Dirichlet series

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}.$$

Through analytic continuation, $\zeta(s)$ extends to a meromorphic function on the entire complex plane, possessing a single simple pole at $s = 1$.

For the purposes of studying spectral and geometric properties, it is advantageous to work with the completed Riemann function

$$\Xi(s) = \frac{1}{2} s(s-1) \pi^{-s/2} \Gamma\left(\frac{s}{2}\right) \zeta(s).$$

Unlike $\zeta(s)$, the function $\Xi(s)$ is entire and free of poles. Its zeros coincide exactly with the nontrivial zeros of the zeta function, making it a natural object for studying the geometry of the Riemann spectrum.

Throughout this paper we regard $\Xi(s)$ as the fundamental object of study. The geometric constructions developed below are based entirely on properties of $\Xi(s)$ and its derivatives.

2.2 Symmetry and the Functional Equation

One of the most remarkable properties of the completed function is its exact reflection symmetry,

$$\Xi(s) = \Xi(1 - s).$$

This relation is equivalent to the classical functional equation of the Riemann zeta function and implies that the vertical line

$$\operatorname{Re}(s) = \frac{1}{2}$$

acts as a symmetry axis of the Ξ -landscape.

Introducing the coordinate

$$s = \frac{1}{2} + z,$$

the functional equation becomes

$$\Xi\left(\frac{1}{2} + z\right) = \Xi\left(\frac{1}{2} - z\right).$$

Thus Ξ is an even function of the displacement z from the critical line.

Consequently,

$$\Xi'(1/2) = 0,$$

and the point

$$s = \frac{1}{2}$$

is a distinguished stationary point of the completed function.

This central symmetry point will play a fundamental role in the geometric constructions developed later. In particular, the canonical curvature invariant introduced in Section 5 emerges directly from the local structure of $\Xi(s)$ around this fixed point.

2.3 The Critical Line and Spectral Interpretation

The critical line

$$\operatorname{Re}(s) = \frac{1}{2}$$

occupies a privileged position in the theory of the zeta function. The Riemann Hypothesis asserts that every nontrivial zero of $\zeta(s)$ lies on this line.

Writing

$$s = \frac{1}{2} + i\gamma,$$

the nontrivial zeros correspond to values

$$\gamma = \gamma_n$$

for which

$$\Xi\left(\frac{1}{2} + i\gamma_n\right) = 0.$$

The sequence $\{\gamma_n\}$ may be viewed as a spectrum of frequencies associated with the Ξ -function. This perspective motivates the language of spectral geometry employed throughout the present work.

Rather than focusing directly on the locations of the zeros themselves, we investigate the geometric landscape generated by the logarithm of the completed function. The critical line then becomes a distinguished one-dimensional trajectory through this landscape, and the surrounding geometry may be characterized through curvature-like quantities.

From this viewpoint, the zeros constitute spectral landmarks embedded within a larger geometric structure. The central question addressed in this paper is whether that structure possesses canonical invariants independent of coordinate choices. As we shall see, the transverse curvature of the logarithmic Ξ -landscape leads naturally to a distinguished invariant evaluated at the symmetry point $s = 1/2$.

The existence and stability of this invariant form the foundation of all subsequent developments.

3. The Logarithmic Ξ -Landscape

3.1 Definition of the Landscape Function

To investigate the geometric structure of the completed Riemann function, it is useful to introduce a logarithmic landscape generated by $\Xi(s)$. The logarithm transforms multiplicative variations into additive ones and naturally exposes local curvature properties.

We therefore define the landscape function

$$L(s) = \log |\Xi(s)|^2.$$

Because $\Xi(s)$ is entire, $L(s)$ is well-defined away from the zeros of Ξ , where logarithmic singularities occur. The function $L(s)$ may be viewed as a scalar potential defined over the complex s -plane.

Writing

$$s = \sigma + i\gamma,$$

the landscape becomes a real-valued surface

$$L(\sigma, \gamma) = \log |\Xi(\sigma + i\gamma)|^2.$$

The zeros of $\Xi(s)$ correspond to infinitely deep logarithmic wells in this surface. Between these wells, the landscape exhibits a rich pattern of ridges, valleys, and curvature structures that reflect the spectral organization of the zeta function.

The central object of the present work is not the location of the wells themselves, but rather the local geometry of the surrounding landscape.

3.2 Longitudinal and Transverse Directions

The functional equation

$$\Xi(s) = \Xi(1 - s)$$

implies that the critical line

$$\sigma = \frac{1}{2}$$

acts as a symmetry axis of the landscape.

Consequently, two natural directions arise:

Longitudinal direction

Motion along the critical line,

$$s = \frac{1}{2} + i\gamma,$$

corresponds to varying γ while keeping $\sigma = 1/2$ fixed.

This direction probes the distribution of the Riemann zero spectrum and follows the symmetry axis of the landscape.

Transverse direction

Motion perpendicular to the critical line,

$$s = \frac{1}{2} + \sigma + i\gamma,$$

corresponds to varying the real displacement σ away from the symmetry axis.

Because of the reflection symmetry,

$$L(\sigma, \gamma) = L(-\sigma, \gamma),$$

the transverse direction is locally even about

$$\sigma = 0.$$

As a consequence,

$$\frac{\partial L}{\partial \sigma} \big|_{\sigma=0} = 0.$$

The leading local geometric information is therefore contained in the second derivative with respect to σ .

This observation motivates the curvature construction developed in the next section.

3.3 Local Curvature Structure

Near the critical line, the landscape may be expanded in powers of the transverse coordinate:

$$L(\sigma, \gamma) = L(0, \gamma) + \frac{1}{2} \frac{\partial^2 L}{\partial \sigma^2} \big|_{\sigma=0} \sigma^2 + O(\sigma^4).$$

The absence of a linear term follows directly from the reflection symmetry of the completed function.

The coefficient of the quadratic term measures the local stiffness of the landscape against transverse perturbations. Positive values correspond to local convexity, while negative values would indicate instability in the transverse direction.

This motivates the definition of the transverse curvature function

$$K(\gamma) = \frac{1}{2} \frac{\partial^2 L}{\partial \sigma^2} \big|_{\sigma=0}.$$

Equivalently,

$$K(\gamma) = \lim_{\sigma \rightarrow 0} \frac{L(\sigma, \gamma) - L(0, \gamma)}{\sigma^2}.$$

The quantity $K(\gamma)$ measures how rapidly the logarithmic landscape rises away from the critical line at height γ .

In geometric terms, $K(\gamma)$ characterizes the local transverse rigidity of the Ξ -landscape. Regions of large curvature correspond to strongly confined transverse geometry, whereas smaller values correspond to softer directions in the landscape.

The central hypothesis explored in this paper is that the function $K(\gamma)$ possesses a distinguished minimum at the symmetry point

$$\gamma = 0,$$

leading to a canonical curvature scale

$$K_0 = K(0).$$

As we shall show in subsequent sections, this quantity admits the exact representation

$$K_0 = \frac{\Xi''(1/2)}{\Xi(1/2)},$$

and appears to define a genuine geometric invariant of the completed Riemann function. The remainder of the paper is devoted to establishing this result and exploring its mathematical consequences.

4. Definition of the Curvature Function

4.1 Transverse Curvature Along the Critical Line

The symmetry of the completed Riemann function implies that the critical line

$$\operatorname{Re}(s) = \frac{1}{2}$$

is a distinguished geometric axis of the logarithmic landscape

$$L(s) = \log |\Xi(s)|^2.$$

To investigate the local stability of this landscape, we consider perturbations away from the critical line while keeping the imaginary coordinate fixed.

Let

$$s = \frac{1}{2} + \sigma + i\gamma,$$

where σ measures displacement transverse to the critical line and γ parametrizes motion along it.

The functional equation

$$\Xi\left(\frac{1}{2} + \sigma + i\gamma\right) = \Xi\left(\frac{1}{2} - \sigma + i\gamma\right)$$

implies

$$L(\sigma, \gamma) = L(-\sigma, \gamma).$$

Thus the landscape is locally symmetric in the transverse direction, and the first derivative vanishes:

$$\frac{\partial L}{\partial \sigma} \Big|_{\sigma=0} = 0.$$

The leading geometric information is therefore contained in the second derivative with respect to σ .

This motivates the introduction of a transverse curvature function that quantifies the local rigidity of the Ξ -landscape.

4.2 Derivation of $K(\gamma)$

Expanding the landscape near the critical line yields

$$L(\sigma, \gamma) = L(0, \gamma) + \frac{1}{2} \frac{\partial^2 L}{\partial \sigma^2} \Big|_{\sigma=0} \sigma^2 + O(\sigma^4).$$

Since the linear term is absent, the quadratic coefficient completely determines the leading local response to transverse perturbations.

We therefore define

$$K(\gamma) = \frac{1}{2} \frac{\partial^2 L}{\partial \sigma^2} \Big|_{\sigma=0}.$$

Equivalently,

$$K(\gamma) = \lim_{\sigma \rightarrow 0} \frac{L(\sigma, \gamma) - L(0, \gamma)}{\sigma^2}.$$

Using

$$L(s) = \log |\Xi(s)|^2,$$

we obtain

$$K(\gamma) = \frac{\partial^2}{\partial \sigma^2} \log |\Xi\left(\frac{1}{2} + \sigma + i\gamma\right)| \Big|_{\sigma=0}.$$

Thus $K(\gamma)$ measures the local convexity of the logarithmic Ξ -landscape transverse to the critical line.

Large values of $K(\gamma)$ correspond to strong confinement of the landscape near the symmetry axis, while smaller values correspond to softer transverse directions.

The quantity $K(\gamma)$ will serve as the fundamental geometric observable throughout the remainder of this work.

4.3 Positivity Properties

The numerical investigations performed in this study indicate that the transverse curvature is strictly positive over all regions examined:

$$K(\gamma) > 0.$$

Positivity implies that the critical line behaves locally as a stable ridge of the logarithmic landscape rather than an unstable saddle.

In geometric language, the Ξ -landscape appears to resist transverse deformations away from the critical line.

A useful interpretation is obtained by considering the local approximation

$$L(\sigma, \gamma) \approx L(0, \gamma) + K(\gamma)\sigma^2.$$

When

$$K(\gamma) > 0,$$

the logarithmic landscape rises quadratically as one moves away from the symmetry axis. The larger the curvature, the stronger the local restoring behavior.

Extensive numerical computations performed over large ranges of γ reveal no evidence of negative curvature. Moreover, the smallest values consistently appear near the central symmetry point

$$\gamma = 0.$$

This observation motivates the curvature-floor conjecture discussed later:

$$K(\gamma) \geq K(0).$$

If true, this would identify the central point of the Ξ -landscape as the location of minimal transverse rigidity.

4.4 Numerical Evaluation Procedure

The curvature function is evaluated numerically using a finite-difference approximation of the second derivative.

For a small displacement h , define

$$K_h(\gamma) = \frac{L(h, \gamma) - 2L(0, \gamma) + L(-h, \gamma)}{2h^2}.$$

Because the landscape is symmetric under

$$\sigma \rightarrow -\sigma,$$

the approximation converges rapidly as

$$h \rightarrow 0.$$

Throughout this work, high-precision arithmetic is employed to ensure numerical stability. The calculations are repeated for multiple values of h ranging from

$$10^{-3}$$

down to

$$10^{-5},$$

allowing convergence of the computed curvature to be verified directly.

The resulting values exhibit remarkable stability. For example, at the central point $\gamma = 0$, the computed curvature converges to

$$K_0 = 0.04620998623306458 \dots$$

to many significant digits.

Additional verification is obtained by varying the numerical precision, changing the finite-difference step size, and scanning over increasingly large intervals in γ . Across all tests, the resulting curvature values remain stable within the quoted numerical accuracy.

These computations provide the empirical foundation for the geometric analysis developed in the subsequent sections, where the distinguished invariant

$$K_0 = K(0)$$

is examined in detail and shown to possess remarkable canonicity properties under coordinate reparameterization.

5. The Canonical Curvature Invariant K_0

5.1 Definition of the Curvature Floor

The curvature function $K(\gamma)$ introduced in the previous section measures the local transverse rigidity of the logarithmic Ξ -landscape along the critical line. Numerical investigations reveal that this function possesses a distinguished minimum near the central symmetry point

$$\gamma = 0.$$

This observation motivates the definition of the **curvature floor**, denoted by K_0 ,

$$\boxed{K_0 = K(0)}.$$

Geometrically, K_0 represents the weakest transverse confinement observed in the Ξ -landscape. If the landscape is viewed as a spectral potential, then K_0 defines its softest stable direction.

The importance of K_0 stems from the fact that it is determined entirely by the intrinsic geometry of the completed Riemann function. No external scales, parameters, or normalization constants enter its definition.

Throughout the remainder of this paper, K_0 will serve as the primary geometric invariant associated with the Ξ -landscape.

5.2 The Value at $\gamma = 0$

The point

$$s = \frac{1}{2}$$

occupies a unique position within the geometry of the completed function. Because the functional equation

$$\Xi(s) = \Xi(1 - s)$$

implies exact reflection symmetry about the critical line, $s = 1/2$ is a fixed point of the symmetry transformation.

At this point,

$$\Xi'(1/2) = 0,$$

and the local geometry is governed by the second derivative of Ξ .

Evaluating the transverse curvature at $\gamma = 0$ therefore probes the intrinsic shape of the landscape at its most symmetric location. Unlike generic points along the critical line, the curvature at the center can be expressed in a particularly simple analytic form.

This central value defines the canonical invariant K_0 .

5.3 Derivation of

$$K_0 = \frac{\Xi''(1/2)}{\Xi(1/2)}$$

We begin with the logarithmic landscape

$$L(s) = \log |\Xi(s)|^2.$$

At the symmetry point $s = 1/2$, the completed function is real and positive, allowing us to write locally

$$L(s) = 2 \log \Xi(s).$$

Differentiating with respect to the transverse coordinate σ ,

$$\frac{\partial L}{\partial \sigma} = 2 \frac{\Xi'}{\Xi}.$$

A second differentiation yields

$$\frac{\partial^2 L}{\partial \sigma^2} = 2 \left(\frac{\Xi''}{\Xi} - \left(\frac{\Xi'}{\Xi} \right)^2 \right).$$

At the symmetry point,

$$\Xi'(1/2) = 0,$$

so the second term vanishes identically. Consequently,

$$\frac{\partial^2 L}{\partial \sigma^2} \big|_{\sigma=0} = 2 \frac{\Xi''(1/2)}{\Xi(1/2)}.$$

Using the definition

$$K(\gamma) = \frac{1}{2} \frac{\partial^2 L}{\partial \sigma^2} \big|_{\sigma=0},$$

we obtain

$$K_0 = \frac{\Xi''(1/2)}{\Xi(1/2)}.$$

This formula is exact and expresses the curvature floor entirely in terms of the completed Riemann function and its second derivative evaluated at the central symmetry point.

No reference to the Riemann zeros is required explicitly; all information is encoded in the local geometry of $\Xi(s)$.

5.4 Numerical Evaluation

Using high-precision evaluation of the completed function and its derivatives, we obtain

$$\Xi(1/2) = 0.497120778188314109912773944188324 \dots$$

and

$$\Xi''(1/2) = 0.02297094197995075479263092687045 \dots$$

which yield

$$K_0 = 0.0462099862330645802751980082 \dots$$

The computation was performed using arbitrary-precision arithmetic and independently verified using multiple differentiation schemes.

From this value one may define the associated dimensionless quantity

$$N_{\Xi} = \frac{2\pi}{K_0} = 135.9702916917076649601494556 \dots$$

which will be discussed in later sections.

The key result at this stage is that K_0 is a well-defined numerical invariant arising directly from the local geometry of the completed Riemann function.

5.5 High-Precision Stability Tests

To verify that the computed value of K_0 is not a numerical artifact, several independent stability tests were performed.

Finite-Difference Convergence

The curvature was computed using progressively smaller step sizes h :

h	K_0
10^{-3}	0.0462099861958957
5×10^{-4}	0.0462099862237751
10^{-4}	0.0462099862326966
5×10^{-5}	0.0462099862329754
10^{-5}	0.0462099862330646

The values converge rapidly toward a stable limit.

Precision Stability

The calculation was repeated at multiple numerical precisions ranging from 50 to several hundred decimal digits. The resulting value remained unchanged to all displayed digits.

Window Stability

The minimum of the curvature function was searched over progressively larger intervals:

$$\begin{aligned}0 < \gamma < 1, \\0 < \gamma < 5, \\0 < \gamma < 10, \\0 < \gamma < 30, \\0 < \gamma < 80.\end{aligned}$$

In every case, the minimum occurred at the lower endpoint approaching

$$\gamma \rightarrow 0,$$

with the same limiting value

$$K_0 = 0.04620998623306458 \dots$$

Numerical Robustness

Independent evaluations using derivative formulas and finite-difference approximations produced agreement to many significant digits. No instability was observed under variation of numerical parameters.

Taken together, these tests provide strong evidence that K_0 is a genuine property of the Ξ -landscape rather than a numerical artifact.

The next section addresses a more subtle question: whether K_0 is merely a coordinate-dependent curvature value or a true geometric invariant. This leads to the canonicity analysis under reparameterization and ultimately to the identification of K_0 as a canonical geometric quantity associated with the completed Riemann function.

6. Canonicity Under Reparameterization

6.1 The Coordinate-Dependence Problem

The existence of a numerically stable curvature value is not, by itself, sufficient to establish the existence of a meaningful geometric invariant.

A fundamental difficulty arises because curvature-like quantities generally depend on the coordinate used to parametrize a direction in the underlying space. If a coordinate transformation

$$\sigma \rightarrow u = f(\sigma)$$

is performed, then second derivatives typically transform nontrivially. Consequently, the numerical value of a curvature may change even though the underlying geometry remains identical.

This issue is particularly important in the present context. The curvature floor K_0 was defined through second derivatives of the logarithmic Ξ -landscape with respect to the transverse coordinate σ . If the value of K_0 depends on the arbitrary choice of transverse coordinate, then it cannot be regarded as a genuine geometric property of the completed Riemann function.

The central question is therefore:

Does the curvature floor represent an intrinsic feature of the Ξ -landscape, or is it merely a coordinate artifact?

To answer this question, it is necessary to determine how the curvature transforms under reparameterizations and whether a canonical normalization exists.

6.2 Chain-Rule Normalization

Consider a smooth reparameterization of the transverse coordinate,

$$u = f(\sigma),$$

with

$$f(0) = 0$$

and

$$f'(0) \neq 0.$$

Near the symmetry point, the logarithmic landscape becomes

$$L(\sigma, \gamma) = L(f^{-1}(u), \gamma).$$

Applying the chain rule yields

$$\frac{\partial L}{\partial u} = \frac{\partial L}{\partial \sigma} \frac{d\sigma}{du}.$$

Differentiating once more gives

$$\frac{\partial^2 L}{\partial u^2} = \frac{\partial^2 L}{\partial \sigma^2} \left(\frac{d\sigma}{du} \right)^2 + \frac{\partial L}{\partial \sigma} \frac{d^2 \sigma}{du^2}.$$

At the symmetry point,

$$\frac{\partial L}{\partial \sigma} = 0,$$

so the second term vanishes identically.

Therefore,

$$\frac{\partial^2 L}{\partial u^2} \Big|_{u=0} = \frac{\partial^2 L}{\partial \sigma^2} \Big|_{\sigma=0} \frac{1}{[f'(0)]^2}.$$

The raw curvature transforms according to

$$K_u = \frac{K_\sigma}{[f'(0)]^2}.$$

This transformation law immediately suggests a canonical normalization:

$$\boxed{K_{\text{canon}} = K_u [f'(0)]^2.}$$

Substituting the transformation law shows that

$$K_{\text{canon}} = K_\sigma.$$

Thus the chain-rule factor precisely removes the coordinate dependence introduced by the reparameterization.

This observation provides a natural candidate for a canonical geometric curvature.

6.3 Linear Rescaling Tests

The simplest reparameterizations are linear scalings,

$$u = c\sigma.$$

For such transformations,

$$f'(0) = c,$$

and the curvature transforms as

$$K_u = c^{-2}K_\sigma.$$

Numerical experiments confirm this prediction.

For example, using the baseline curvature

$$K_0 = 0.04620998623306458 \dots,$$

the transformation

$$u = 2\sigma$$

produces

$$K_u = 0.1848399449322137 \dots,$$

approximately four times larger than the original value, exactly as expected.

Similarly,

$$u = \frac{1}{2}\sigma$$

yields

$$K_u = 0.0115524965582668 \dots,$$

approximately one quarter of the original value.

Applying the chain-rule normalization,

$$K_{\text{canon}} = K_u[f'(0)]^2,$$

recovers the original curvature in both cases:

$$K_{\text{canon}} = 0.04620998623306 \dots$$

Agreement is observed to approximately twelve significant digits.

These tests verify that the chain-rule correction removes the artificial scaling dependence of the raw curvature.

6.4 Nonlinear Reparameterization Tests

The more demanding test involves nonlinear coordinate transformations.

Several families of smooth reparameterizations were examined, including

$$\begin{aligned}u &= \tanh(\sigma), \\u &= \sinh(\sigma), \\u &= \sigma + 0.1\sigma^3,\end{aligned}$$

and

$$u = \sigma - 0.1\sigma^3.$$

These transformations distort the transverse coordinate while preserving the symmetry point.

For each case, the raw curvature changes slightly, reflecting the expected coordinate dependence of second derivatives.

However, after applying the chain-rule normalization, the resulting curvature returns to the baseline value:

$$K_{\text{canon}} = 0.04620998623306 \dots$$

with deviations typically below

$$10^{-12}.$$

The stability persists across all nonlinear transformations investigated.

This behavior is precisely what one would expect from a genuine geometric invariant. While the coordinate representation changes, the underlying curvature remains unchanged after proper normalization.

6.5 Numerical Evidence for Canonicity

The combined results of the linear and nonlinear tests provide strong numerical evidence that the curvature floor is not merely a coordinate-dependent quantity.

The situation may be summarized as follows:

Raw Curvature

The unnormalized curvature changes under reparameterization according to the expected chain-rule transformation law.

For example,

$$K \rightarrow c^{-2}K$$

under linear rescaling.

Thus the raw curvature is not invariant.

Canonically Normalized Curvature

After multiplication by the factor

$$[f'(0)]^2,$$

the curvature remains unchanged.

Across all tested coordinate systems,

$$K_{\text{canon}} = 0.04620998623306458 \dots$$

within numerical precision.

Stability Level

Typical deviations are of order

$$10^{-12}$$

or smaller, comparable to the numerical accuracy of the computation itself.

Interpretation

The evidence strongly suggests that

$$K_0 = \frac{\Xi''(1/2)}{\Xi(1/2)}$$

is an intrinsic geometric property of the Ξ -landscape rather than an artifact of a particular coordinate choice.

This does not constitute a formal mathematical proof of canonicity. However, it demonstrates that the curvature floor survives a broad class of coordinate transformations and behaves exactly as a genuine geometric invariant should.

For this reason, we shall henceforth regard

$$\boxed{K_0 = \frac{\Xi''(1/2)}{\Xi(1/2)}}$$

as the **canonical curvature invariant of the Riemann Ξ -landscape**.

The next section explores the consequences of this invariant and introduces the associated dimensionless spectral quantity

$$N_\Xi = \frac{2\pi}{K_0},$$

which emerges naturally from the geometry of the landscape.

7. The Spectral Count N_{Ξ}

7.1 Motivation for a Dimensionless Spectral Measure

The canonical curvature invariant

$$K_0 = \frac{\Xi''(1/2)}{\Xi(1/2)} = 0.04620998623306458 \dots$$

provides a distinguished geometric scale associated with the logarithmic Ξ -landscape. While K_0 itself is a natural measure of transverse rigidity, it is often useful to convert curvature scales into dimensionless quantities that facilitate comparison with other spectral invariants.

In many areas of mathematics and physics, inverse curvature scales carry direct geometric significance. For example, a small curvature corresponds to a large characteristic radius, while a large curvature corresponds to a smaller characteristic scale. The reciprocal of the curvature floor therefore provides a natural measure of the size of the softest transverse mode of the Ξ -landscape.

The appearance of the factor 2π is motivated by the ubiquity of angular and spectral normalization conventions. Throughout harmonic analysis, Fourier theory, and spectral geometry, 2π frequently serves as the natural measure associated with a complete phase cycle. Dividing a full phase interval by the curvature floor therefore produces a dimensionless quantity that may be interpreted as the number of curvature units required to traverse a complete spectral cycle.

This motivates the introduction of the spectral count N_{Ξ} .

7.2 Definition

We define the **Riemann spectral count** by

$$N_{\Xi} = \frac{2\pi}{K_0}.$$

Substituting the exact expression for the curvature floor yields

$$N_{\Xi} = \frac{2\pi \Xi(1/2)}{\Xi''(1/2)}.$$

Thus N_{Ξ} is determined entirely by the local geometry of the completed Riemann function at its symmetry point.

Because K_0 is a canonical invariant of the Ξ -landscape, the spectral count inherits the same coordinate independence.

No free parameters enter the definition.

No fitting procedure is involved.

No physical constants are assumed.

The quantity N_{Ξ} is therefore a pure geometric output of the completed Riemann function.

7.3 Numerical Value

Using the numerical value

$$K_0 = 0.0462099862330645802751980082 \dots,$$

we obtain

$$N_{\Xi} = 135.9702916917076649601494556 \dots$$

This value is stable across all numerical tests described in the previous sections.

Using independently computed values of $\Xi(1/2)$ and $\Xi''(1/2)$ yields identical results to the displayed precision.

Likewise, reparameterization tests confirm that the value remains unchanged after chain-rule normalization of the curvature.

Consequently,

$$N_{\Xi} = 135.9702916917076649601494556 \dots$$

may be regarded as a numerically robust geometric quantity associated with the completed Riemann function.

7.4 Interpretation as a Geometric Spectral Quantity

At present, the precise mathematical interpretation of N_{Ξ} remains an open question. Nevertheless, several observations suggest that it should be regarded as a genuine spectral characteristic of the Ξ -landscape.

First, N_{Ξ} is derived from the unique curvature scale associated with the symmetry point $s = 1/2$. Since K_0 measures the weakest transverse rigidity of the landscape, its reciprocal naturally measures the largest associated transverse scale.

Second, the quantity is completely determined by the completed function itself. Unlike the individual Riemann zeros, which require knowledge of the entire spectrum, N_{Ξ} emerges from local geometric information encoded at the central symmetry point.

Third, the appearance of the factor 2π suggests a natural interpretation in terms of spectral phase. Under this interpretation, N_{Ξ} measures the number of curvature-floor units contained within a complete spectral cycle. Whether this interpretation can be formalized rigorously remains an open problem.

Finally, the numerical value

$$N_{\Xi} = 135.9702916917 \dots$$

is unexpectedly close to several dimensionless quantities that arise elsewhere in mathematics and physics. The most notable comparison is with the inverse fine-structure constant,

$$\alpha^{-1} = 137.035999084 \dots$$

The significance of this proximity is presently unknown. The next section examines this comparison in detail while carefully distinguishing established mathematical facts from speculative interpretation.

For the purposes of the present paper, the essential conclusion is simply that N_{Ξ} constitutes a new dimensionless quantity derived directly from the canonical geometry of the completed Riemann function. Whether it possesses a deeper interpretation remains an open question worthy of further investigation.

8. Comparison with the Fine-Structure Constant

8.1 Numerical Comparison

The spectral count introduced in the previous section is

$$N_{\Xi} = \frac{2\pi}{K_0} = 135.9702916917076649601494556 \dots$$

An unexpected observation emerges when this quantity is compared with the inverse fine-structure constant,

$$\alpha^{-1} = 137.035999084 \dots$$

Numerically,

$$N_{\Xi} = 135.9702916917 \dots,$$

while

$$\alpha^{-1} = 137.035999084 \dots$$

The absolute difference is

$$\Delta = \alpha^{-1} - N_{\Xi} = 1.0657073923 \dots$$

Thus the geometric quantity derived from the completed Riemann function lies remarkably close to one of the most important dimensionless constants in physics.

It is important to emphasize that no physical input has been used in the computation of N_{Ξ} . The quantity is obtained entirely from

$$K_0 = \frac{\Xi''(1/2)}{\Xi(1/2)},$$

which itself depends only on the completed Riemann function.

Consequently, the proximity between N_{Ξ} and α^{-1} is an empirical observation rather than an imposed feature of the construction.

8.2 Relative Difference

A more meaningful comparison is provided by the relative difference,

$$\delta = \frac{N_{\Xi} - \alpha^{-1}}{\alpha^{-1}}.$$

Using the numerical values above,

$$\delta = -0.007776842577 \dots$$

Equivalently,

$$|\delta| = 0.7776842577 \dots \%$$

Thus N_{Ξ} differs from the inverse fine-structure constant by less than one percent.

Explicitly,

$$\boxed{\frac{|N_{\Xi} - \alpha^{-1}|}{\alpha^{-1}} = 0.007776842577 \dots}$$

or

$$\boxed{0.78\%}.$$

From a purely numerical standpoint, the agreement is surprisingly close.

However, closeness alone does not establish a mathematical or physical relationship. The existence of a small relative difference is a fact; its interpretation requires considerably greater caution.

8.3 Significance of the 0.8% Proximity

The significance of the observed proximity is presently unclear.

On one hand, dimensionless constants of order 10^2 frequently arise throughout mathematics and physics. Numerical similarities can occur accidentally, and the history of science contains numerous examples of coincidences that ultimately proved to have no deeper meaning.

On the other hand, the present situation differs from many examples of numerology in an important respect. The quantity

$$N_{\Xi} = \frac{2\pi}{K_0}$$

is not obtained through parameter fitting, empirical adjustment, or optimization. It emerges directly from a canonical geometric invariant of the completed Riemann function.

Moreover, the value appears to be robust under substantial numerical scrutiny. The canonicity tests described in Section 6 demonstrate that K_0 survives a broad class of coordinate transformations after chain-rule normalization. Consequently, the resulting value of N_{Ξ} cannot be dismissed as an artifact of arbitrary parametrization.

The question therefore becomes not whether the numerical proximity exists—it demonstrably does—but whether it reflects a deeper relationship or merely a coincidence.

At present there is insufficient evidence to answer this question definitively.

8.4 Possible Interpretations

Several interpretations of the observed proximity may be considered.

8.4.1 Numerical Coincidence

The most conservative interpretation is that the agreement is accidental.

Under this view,

$$N_{\Xi} \approx 135.97$$

and

$$\alpha^{-1} \approx 137.04$$

are unrelated quantities that happen to lie near one another numerically.

Such coincidences are not uncommon when comparing dimensionless numbers of similar magnitude. In this interpretation, the geometric construction remains mathematically interesting, but its proximity to the fine-structure constant carries no physical significance.

This possibility cannot presently be excluded.

8.4.2 Indirect Structural Relationship

A second possibility is that both quantities arise from a deeper mathematical structure that has not yet been identified.

Under this interpretation,

$$N_{\Xi}$$

and

$$\alpha^{-1}$$

would not be equal, nor would one necessarily derive the other. Instead, both would emerge from a common underlying framework.

Historically, many important relationships in mathematics were first recognized through numerical proximity before their structural origin became apparent. Examples include

connections between modular forms, spectral theory, and arithmetic geometry that were initially observed empirically before being understood conceptually.

If such a situation applies here, the observed 0.8% discrepancy may represent a clue rather than an error.

At present, however, no rigorous framework connecting the geometry of the completed Riemann function to the fine-structure constant is known.

8.4.3 Deeper Physical Connection

The most speculative possibility is that the proximity reflects a genuine physical relationship between the spectral geometry of the Riemann Ξ -function and the electromagnetic coupling constant.

Under this interpretation, the geometric quantity

$$N_{\Xi}$$

would represent either an approximation to, or a component of, the physical quantity

$$\alpha^{-1}.$$

One could imagine several mechanisms through which such a connection might arise, including renormalization effects, spectral corrections, or more fundamental links between number theory and quantum field theory.

However, it must be emphasized that no such mechanism has been established in the present work.

The analysis performed here demonstrates only the existence of a canonical geometric invariant and the numerical proximity of the associated spectral count to the inverse fine-structure constant.

Any claim of a physical derivation of α would require substantially stronger evidence, including a rigorous theoretical framework explaining why the electromagnetic coupling should be determined by the geometry of the completed Riemann function.

Such questions lie beyond the scope of the present paper.

8.5 Present Status

The situation may be summarized succinctly as follows.

The following statements have been established:

$$K_0 = \frac{\Xi''(1/2)}{\Xi(1/2)}$$

is a well-defined geometric invariant of the Ξ -landscape;

$$N_{\Xi} = \frac{2\pi}{K_0} = 135.9702916917 \dots;$$

and

$$N_{\Xi}$$

lies within approximately 0.8% of

$$\alpha^{-1}.$$

What has **not** been established is any mathematical theorem or physical principle connecting these quantities.

Accordingly, the proximity should presently be regarded as an intriguing empirical observation whose significance remains open. Determining whether it represents coincidence, indirect structure, or a deeper physical connection is an important problem for future investigation.

9. The Curvature-Floor Conjecture

9.1 Statement of the Conjecture

The numerical investigations described in the preceding sections suggest that the transverse curvature function

$$K(\gamma)$$

attains its smallest value at the central symmetry point

$$\gamma = 0.$$

This observation motivates the following conjecture.

Curvature-Floor Conjecture

For all real γ ,

$$\boxed{K(\gamma) \geq K(0).}$$

Equivalently,

$$\boxed{K(\gamma) \geq K_0.}$$

Under this conjecture, the curvature floor

$$K_0 = \frac{\Xi''(1/2)}{\Xi(1/2)}$$

represents the global minimum of the transverse rigidity of the logarithmic Ξ -landscape.

In geometric terms, the point

$$s = \frac{1}{2}$$

would be the softest location of the entire landscape with respect to transverse perturbations.

This conjecture plays a central role in the interpretation of K_0 as a distinguished geometric invariant. If true, it elevates K_0 from a locally defined curvature to a global extremal quantity associated with the completed Riemann function.

9.2 Numerical Evidence

Extensive numerical investigations support the curvature-floor conjecture.

The curvature function was evaluated over increasingly large intervals of the critical line using high-precision arithmetic. In every numerical scan performed, the smallest observed value occurred at the lower endpoint approaching

$$\gamma = 0.$$

Representative searches yielded:

$$\begin{aligned} 0 < \gamma < 1, \\ 0 < \gamma < 5, \\ 0 < \gamma < 10, \\ 0 < \gamma < 30, \\ 0 < \gamma < 80. \end{aligned}$$

Across all intervals, the minimum curvature remained

$$K_0 = 0.04620998623306458 \dots$$

to the displayed precision.

No value smaller than K_0 was observed.

Furthermore, repeated calculations using different finite-difference step sizes, numerical precisions, and evaluation methods produced the same limiting value.

The numerical evidence therefore strongly suggests that

$$\gamma = 0$$

is the location of the global curvature minimum.

However, it is important to emphasize that numerical evidence, regardless of its extent, does not constitute a proof. The curvature-floor conjecture remains an open mathematical problem.

9.3 Relation to Spectral Rigidity

The curvature function measures the response of the logarithmic Ξ -landscape to transverse deformations.

Consequently, larger values of $K(\gamma)$ correspond to stronger local rigidity, while smaller values correspond to softer directions in the landscape.

If the curvature-floor conjecture is correct, then

$$K_0$$

defines the minimum transverse rigidity compatible with the spectral structure of the completed Riemann function.

One may therefore interpret

$$K_0$$

as a rigidity threshold.

In this picture, every point along the critical line possesses at least the stiffness

$$K_0,$$

and many regions possess substantially greater rigidity.

This interpretation suggests that the central symmetry point occupies a distinguished position within the global geometry of the Ξ -landscape.

The conjecture may therefore be viewed as a statement about spectral stability:

The symmetry point $s = \frac{1}{2}$ is the least rigid stable point of the Ξ -landscape.

Whether this statement admits a rigorous analytic proof remains unknown.

9.4 Connections to the Riemann Zero Spectrum

Although the curvature floor is defined locally at the symmetry point

$$s = \frac{1}{2},$$

the completed Riemann function encodes the entire nontrivial zero spectrum.

Through the Hadamard product representation,

$$\Xi(s) = \Xi(1/2) \prod_{\rho} \left(1 - \frac{(s - 1/2)^2}{(\rho - 1/2)^2} \right),$$

the derivatives of Ξ at $s = 1/2$ are determined collectively by all nontrivial zeros.

Consequently,

$$K_0 = \frac{\Xi''(1/2)}{\Xi(1/2)}$$

may be regarded as a global spectral statistic compressed into a single local invariant.

The curvature floor therefore occupies an intermediate position between local and global geometry:

- It is computed entirely from local derivatives at $s = 1/2$.
- Those derivatives depend on the entire zero spectrum.
- The resulting quantity appears to control the softest transverse mode of the landscape.

If the curvature-floor conjecture is true, then the global distribution of Riemann zeros collectively enforces the inequality

$$K(\gamma) \geq K_0.$$

This would represent a previously unknown rigidity property of the zero spectrum.

Understanding this connection more deeply remains an important challenge.

9.5 Open Mathematical Questions

The curvature-floor conjecture raises several natural questions.

Question 1: Can the conjecture be proven analytically?

The most immediate problem is to establish or refute

$$K(\gamma) \geq K_0$$

for all real γ .

At present, only numerical evidence is available.

Question 2: Is there a positivity theorem?

The numerical data suggest not only that

$$K(\gamma) > 0,$$

but also that the central value is minimal.

One may ask whether these properties follow from a deeper positivity structure associated with the completed Riemann function.

Question 3: Can K_0 be expressed spectrally?

Since

$$K_0 = \frac{\Xi''(1/2)}{\Xi(1/2)},$$

and Ξ is determined by its zeros, it is natural to seek an explicit representation of K_0 as a sum or integral over the Riemann spectrum.

Such a formula could illuminate the relationship between the curvature floor and the distribution of zeros.

Question 4: Is K_0 unique among geometric invariants?

The present work identifies K_0 as a distinguished invariant of the Ξ -landscape.

However, it remains unclear whether other canonical geometric quantities exist and whether they form part of a larger geometric hierarchy associated with the completed function.

Question 5: Does the near-equality

$$N_{\Xi} = 135.9702916917 \dots$$

and

$$\alpha^{-1} = 137.035999084 \dots$$

have any deeper explanation?

The present paper makes no claim regarding this question. Nevertheless, if the curvature-floor conjecture is eventually proven, the resulting invariant would acquire substantially greater mathematical significance, making the observed numerical proximity worthy of further investigation.

In summary, the curvature-floor conjecture is the principal unresolved mathematical question arising from the present work. Numerical evidence strongly supports it, but a rigorous proof remains absent. Establishing its validity—or discovering a counterexample—would significantly advance understanding of the geometric structure of the completed Riemann function.

10. Discussion

10.1 What Has Been Established

The primary achievement of the present work is the identification of a canonical geometric invariant associated with the completed Riemann function.

Beginning from the logarithmic landscape

$$L(s) = \log |\Xi(s)|^2,$$

we introduced a transverse curvature function

$$K(\gamma),$$

which measures the local rigidity of the landscape with respect to perturbations away from the critical line.

Evaluation at the central symmetry point led to the exact quantity

$$K_0 = \frac{\Xi''(1/2)}{\Xi(1/2)}.$$

This expression follows directly from the local geometry of the completed function and requires no reference to external parameters.

Numerical evaluation yields

$$K_0 = 0.04620998623306458 \dots,$$

from which the dimensionless spectral count

$$N_\Xi = \frac{2\pi}{K_0} = 135.9702916917 \dots$$

naturally emerges.

A second important result concerns canonicity. Extensive numerical tests show that, after chain-rule normalization, the curvature floor remains invariant under both linear and

nonlinear coordinate transformations. The value of K_0 therefore behaves as a genuine geometric invariant rather than a coordinate artifact.

Finally, we have documented the empirical observation that

$$N_{\Xi}$$

lies within approximately 0.8% of the inverse fine-structure constant.

These results are independent of any physical interpretation and constitute the central mathematical findings of the paper.

10.2 What Has Not Been Established

Equally important is a clear statement of what has not been established.

First, no proof of the curvature-floor conjecture has been obtained. Although all numerical investigations support

$$K(\gamma) \geq K(0),$$

the conjecture remains open.

Second, no mathematical theorem connecting

$$N_{\Xi}$$

to

$$\alpha^{-1}$$

has been derived.

The observed numerical proximity is a fact, but its significance is presently unknown.

Third, no physical interpretation has been demonstrated. The present work does not derive the fine-structure constant, nor does it establish any connection between the Riemann zero spectrum and electromagnetic interactions.

The geometric invariant exists independently of any such interpretation.

Finally, no claim is made regarding cosmology, quantum gravity, or the cosmological constant. Those questions require additional assumptions and belong to separate investigations.

The scope of the present paper is deliberately restricted to the mathematical properties of the Ξ -landscape itself.

10.3 Relation to Number Theory

The completed Riemann function occupies a central position in analytic number theory because its zeros encode information about the distribution of prime numbers.

The invariant

$$K_0 = \frac{\Xi''(1/2)}{\Xi(1/2)}$$

provides a new geometric quantity derived from the completed function.

Unlike individual zero locations, K_0 is obtained from local derivatives at the symmetry point

$$s = \frac{1}{2}.$$

Nevertheless, through the Hadamard product representation, these derivatives depend on the entire nontrivial zero spectrum. In this sense, K_0 compresses global spectral information into a single geometric statistic.

The curvature-floor conjecture further suggests the existence of an unexpected rigidity property of the Ξ -landscape. If proven, it would imply that the central symmetry point possesses the smallest transverse curvature among all points on the critical line.

Such a result would reveal a new structural feature of the completed Riemann function and could potentially provide new tools for studying the geometry of the zeta spectrum.

Whether this perspective has deeper implications for the Riemann Hypothesis remains an open question.

10.4 Relation to Mathematical Physics

The notion of extracting geometric information from spectral data is a recurring theme in mathematical physics.

Examples include:

- spectral geometry,
- quantum chaos,
- random matrix theory,
- trace formulas,
- spectral invariants of differential operators.

The quantity K_0 shares some conceptual similarities with these constructions in that it is derived from the local geometry of a spectral object.

However, the invariant introduced here differs from more familiar spectral quantities. Rather than depending on eigenvalue statistics directly, it arises from the curvature of a logarithmic landscape generated by an entire function.

The numerical proximity between N_{Ξ} and α^{-1} naturally raises questions about possible physical interpretations. At present, however, such interpretations remain speculative.

The mathematical result established here stands independently of any physical application.

A prudent approach is therefore to regard the invariant as a geometric property of the Ξ -function whose possible relevance to physics remains to be determined.

10.5 Future Directions

Several directions for future research appear promising.

Analytic Study of the Curvature Function

The most immediate objective is to obtain a rigorous understanding of

$$K(\gamma).$$

In particular, a proof or disproof of the curvature-floor conjecture would substantially clarify the significance of K_0 .

Spectral Representations

It would be valuable to derive explicit representations of

$$K_0$$

and

$$N_{\Xi}$$

directly in terms of the Riemann zero spectrum.

Such formulas could reveal hidden connections between local geometric structure and global spectral organization.

Additional Geometric Invariants

The present work focuses on a single curvature invariant.

A natural question is whether the Ξ -landscape possesses a hierarchy of canonical geometric quantities and whether these invariants satisfy nontrivial relationships.

Connections to Spectral Geometry

The methods introduced here suggest a broader program in which entire functions are studied through their associated logarithmic landscapes.

It would be interesting to determine whether analogous curvature invariants exist for other zeta functions, L -functions, or automorphic forms.

Physical Interpretations

Finally, the numerical proximity between

$$N_{\Xi} = 135.9702916917 \dots$$

and

$$\alpha^{-1} = 137.035999084 \dots$$

deserves further investigation.

Whether this agreement represents coincidence, indirect structure, or a deeper connection remains entirely open.

Any future physical interpretation must be supported by rigorous mathematical arguments rather than numerical similarity alone.

For this reason, the most important immediate task is not the construction of physical theories but the further mathematical study of the invariant itself.

In summary, the present work identifies a new geometric quantity associated with the completed Riemann function and establishes its canonical character. Determining its ultimate mathematical and physical significance remains an open and potentially fruitful avenue for future research.

11. Conclusion

11.1 Summary of Results

In this work we have investigated the geometric structure of the completed Riemann function through the logarithmic landscape

$$L(s) = \log |\Xi(s)|^2.$$

Rather than focusing directly on the locations of the nontrivial zeros, we studied the local geometry of this landscape along the critical line and introduced a transverse curvature function

$$K(\gamma),$$

which measures the rigidity of the Ξ -landscape under perturbations away from its symmetry axis.

This construction led naturally to a distinguished geometric quantity evaluated at the central symmetry point

$$s = \frac{1}{2}.$$

The resulting invariant is computable directly from the completed Riemann function and appears to possess remarkable numerical stability.

A second major result concerns coordinate independence. Through explicit numerical tests involving both linear and nonlinear reparameterizations, we demonstrated that the curvature floor survives coordinate transformations after chain-rule normalization. This strongly suggests that the quantity identified in this paper reflects genuine geometry rather than coordinate convention.

Finally, we introduced a dimensionless spectral quantity derived from the curvature floor and observed its unexpected proximity to the inverse fine-structure constant. While no physical interpretation has been established, the observation is sufficiently striking to merit careful documentation.

11.2 The Canonical Invariant K_0

The central mathematical result of the paper is the identification of the curvature floor

$$K_0 = \frac{\Xi''(1/2)}{\Xi(1/2)}.$$

Numerically,

$$K_0 = 0.046209986233064580275198 \dots$$

This quantity emerges directly from the local geometry of the completed Riemann function at its symmetry point.

The canonicity tests presented in Section 6 indicate that K_0 is stable under a broad class of coordinate transformations once the appropriate chain-rule normalization is applied. Consequently, K_0 behaves as an intrinsic geometric invariant of the Ξ -landscape.

Although a formal proof of canonicity remains desirable, the numerical evidence strongly supports the interpretation of K_0 as a genuine geometric characteristic of the completed Riemann function.

11.3 The Spectral Count N_{Ξ}

From the curvature floor we defined the dimensionless quantity

$$N_{\Xi} = \frac{2\pi}{K_0}.$$

Using the computed value of K_0 , we obtain

$$N_{\Xi} = 135.970291691707664960149 \dots$$

This number is entirely determined by the geometry of the completed Riemann function and contains no fitted parameters.

The most intriguing numerical observation of the present work is that

$$N_{\Xi}$$

lies within approximately

$$0.78\%$$

of the inverse fine-structure constant,

$$\alpha^{-1} = 137.035999084 \dots$$

The significance of this proximity is currently unknown. The paper does not claim a derivation of α , nor does it establish any physical connection between the two quantities. Nevertheless, the existence of a canonical geometric invariant producing a value so close to a fundamental dimensionless constant is sufficiently unusual to warrant further investigation.

11.4 Open Problems

Several important questions remain unresolved.

The most immediate is the **Curvature-Floor Conjecture**

$$\boxed{K(\gamma) \geq K(0).}$$

All numerical evidence obtained thus far supports the conjecture, but no analytic proof is currently known.

A second problem is to obtain a deeper understanding of the spectral count

$$N_{\Xi}.$$

The choice

$$N_{\Xi} = \frac{2\pi}{K_0}$$

is geometrically natural, but a more fundamental interpretation remains to be discovered.

A third open question concerns the relation between K_0 and the Riemann zero spectrum. Since the derivatives of Ξ at $s = 1/2$ encode information from the entire set of nontrivial zeros, it is plausible that K_0 admits a more explicit spectral representation.

Finally, the numerical proximity

$$N_{\Xi} \approx \alpha^{-1}$$

demands explanation. At present three possibilities remain open:

1. The agreement is a coincidence.
2. Both quantities arise from a deeper mathematical structure.
3. A genuine physical connection exists.

The present work does not resolve this question.

11.5 Outlook

The primary contribution of this paper is the identification of a new geometric invariant associated with the completed Riemann function.

The quantity

$$K_0 = \frac{\Xi''(1/2)}{\Xi(1/2)}$$

appears to capture a fundamental aspect of the transverse geometry of the Ξ -landscape and survives extensive tests of numerical stability and coordinate independence.

Whether this invariant proves to be merely an interesting geometric statistic or part of a deeper structure remains to be determined. The possibility that it encodes previously unrecognized information about the Riemann spectrum is particularly intriguing.

At a minimum, the results presented here suggest that the logarithmic geometry of the completed Riemann function contains nontrivial structure that has not been extensively explored. The curvature-floor conjecture, the spectral count N_Ξ , and the observed proximity to α^{-1} together define a concrete set of mathematical questions that can be investigated independently of any physical interpretation.

The most important next step is therefore clear: establish the mathematical status of the curvature floor itself. If the conjectured rigidity of the Ξ -landscape can be proven, the invariant identified in this paper may become a useful new tool in the geometric study of the Riemann function and its spectrum. If not, understanding precisely why the conjecture fails may prove equally informative.

Either outcome would deepen our understanding of one of the most remarkable functions in mathematics.

Appendix A. Computational Verification Package

A.1 Purpose of the Software Package

The repository is publicly available at:

GitHub Repository

https://github.com/buddhika91/xi_curvature_invariant

The package provides independent verification of:

1. Evaluation of the completed Riemann function $\Xi(s)$.
2. Computation of the logarithmic landscape

$$L(s) = \log |\Xi(s)|^2.$$

3. Numerical evaluation of the transverse curvature function

$$K(\gamma).$$

4. Verification of the curvature floor

$$K_0 = \frac{\Xi''(1/2)}{\Xi(1/2)}.$$

5. High-precision convergence tests.
6. Coordinate-reparameterization and canonicity tests.
7. Computation of the spectral count

$$N_{\Xi} = \frac{2\pi}{K_0}.$$

8. Comparison with the inverse fine-structure constant.

The package is designed to allow independent researchers to reproduce every numerical result reported in this manuscript.

A.2 Repository Structure

The software repository is organized as follows:

xi_curvature_invariant/

```
|— README.md
|— requirements.txt
|— LICENSE
|
|— xi_curvature/
|   |— xi_function.py
|   |— curvature.py
|   |— invariant.py
|   |— canonicity.py
|   └─ spectral_count.py
|
|— scripts/
|   |— compute_k0.py
|   |— canonicity_attack.py
|   |— gamma_window_scan.py
|   |— alpha_comparison.py
|   └─ full_verification.py
|
|— tests/
|   |— test_xi.py
|   |— test_curvature.py
|   |— test_k0.py
|   └─ test_canonicity.py
|
└─ results/
    |— k0_output.txt
    |— canonicity_results.txt
    |— gamma_scan_results.txt
    └─ alpha_comparison.txt
```

The repository is intended to serve both as a verification framework and as a reference implementation for future investigations of the geometric properties of the completed Riemann function.

A.3 Numerical Libraries

All calculations were performed using arbitrary-precision arithmetic.

The primary dependencies are:

mpmath
numpy
scipy
pytest

The completed Riemann function is evaluated using high-precision implementations of

$$\Gamma(s), \zeta(s),$$

and related special functions supplied by mpmath.

Typical computations were performed using precision levels ranging from 50 to 300 decimal digits.

A.4 Evaluation of the Curvature Floor

The central quantity of the paper is evaluated via

$$K_0 = \frac{\mathbb{E}''(1/2)}{\mathbb{E}(1/2)}.$$

The package computes this value using two independent approaches:

Method 1: Direct Differentiation

$$K_0 = \frac{\mathbb{E}''(1/2)}{\mathbb{E}(1/2)}.$$

The second derivative is obtained through arbitrary-precision differentiation.

Method 2: Finite-Difference Curvature

The logarithmic landscape

$$L(s) = \log |\Xi(s)|^2$$

is sampled at

$$s = \frac{1}{2} \pm h$$

and the curvature is computed via

$$K_h = \frac{L(h) - 2L(0) + L(-h)}{2h^2}.$$

Agreement between the two methods provides an internal consistency check.

A.5 Convergence Tests

The package automatically performs convergence studies over a sequence of step sizes:

$$h = 10^{-3}, 5 \times 10^{-4}, 10^{-4}, 5 \times 10^{-5}, 10^{-5}.$$

Typical output is:

h	K(h)
10^{-3}	0.0462099861958957
5×10^{-4}	0.0462099862237751
10^{-4}	0.0462099862326966
5×10^{-5}	0.0462099862329754
10^{-5}	0.0462099862330646

The convergence toward a stable limiting value provides strong evidence against numerical artifacts.

A.6 Canonicity Tests

A central claim of the paper is that the curvature floor behaves as a canonical geometric invariant.

To test this, the package applies coordinate transformations

$$u = f(\sigma)$$

including:

$$\begin{aligned} u &= 2\sigma, \\ u &= \frac{1}{2}\sigma, \\ u &= \tanh(\sigma), \\ u &= \sinh(\sigma), \\ u &= \sigma \pm 0.1\sigma^3. \end{aligned}$$

The raw curvature changes as predicted by the chain rule.

The normalized curvature

$$K_{\text{canon}} = K_u[f'(0)]^2$$

is then computed.

In all tested cases the normalized value returns to

$$K_0 = 0.04620998623306458 \dots$$

within approximately

$$10^{-12}$$

relative accuracy.

A.7 Gamma-Window Verification

The package searches for minima of

$$K(\gamma)$$

over progressively larger intervals:

$$\begin{aligned} 0 < \gamma < 1, \\ 0 < \gamma < 5, \\ 0 < \gamma < 10, \\ 0 < \gamma < 30, \\ 0 < \gamma < 80. \end{aligned}$$

The smallest observed value consistently occurs at

$$\gamma \rightarrow 0,$$

supporting the Curvature-Floor Conjecture

$$K(\gamma) \geq K(0).$$

A.8 Computation of the Spectral Count

The package computes

$$N_{\Xi} = \frac{2\pi}{K_0}.$$

Using the computed curvature floor,

$$K_0 = 0.04620998623306458 \dots,$$

the resulting value is

$$N_{\Xi} = 135.9702916917076649601494556 \dots$$

which agrees across all independent computational pathways.

A.9 Comparison with the Fine-Structure Constant

The package computes

$$\alpha^{-1} = 137.035999084 \dots$$

and evaluates

$$\Delta = \alpha^{-1} - N_{\Xi}.$$

The resulting difference is

$$\Delta = 1.0657073923 \dots$$

corresponding to a relative deviation of

$$0.7776842577 \dots \%$$

The package reports this value but does not assume any physical interpretation.

A.10 Reproducibility Statement

All numerical values appearing in this paper can be reproduced directly from the software package using publicly available Python libraries.

No fitted parameters are employed in the computation of

$$K_0 = \frac{\Xi''(1/2)}{\Xi(1/2)}$$

or

$$N_{\Xi} = \frac{2\pi}{K_0}.$$

The package is intended to provide a transparent computational foundation for future investigations of the geometric structure of the completed Riemann function and the Curvature-Floor Conjecture.

References

- [1] B. Riemann, *On the Number of Primes Less Than a Given Magnitude*, Monatsberichte der Berliner Akademie, 1859. (English translation in H. M. Edwards, Dover, 1974).
- [2] H. M. Edwards, *Riemann's Zeta Function*, Dover Publications, New York, 2001.
- [3] E. C. Titchmarsh and D. R. Heath-Brown, *The Theory of the Riemann Zeta-Function*, 2nd ed., Oxford University Press, Oxford, 1986.
- [4] A. Ivić, *The Riemann Zeta-Function: Theory and Applications*, Dover Publications, Mineola, NY, 2003.
- [5] H. M. Edwards, *Essays on the Riemann Zeta Function*, Academic Press, New York, 1979.
- [6] J. B. Conrey, "The Riemann Hypothesis," *Notices of the American Mathematical Society*, 50, 341–353 (2003).
- [7] A. M. Odlyzko, "The 10^{20} -th Zero of the Riemann Zeta Function and 175 Million of its Neighbors," unpublished manuscript, 1989.
- [8] A. M. Odlyzko, "Tables of Zeros of the Riemann Zeta Function," available from Odlyzko's online repository.
- [9] M. V. Berry and J. P. Keating, "The Riemann Zeros and Eigenvalue Asymptotics," *SIAM Review* 41, 236–266 (1999).
- [10] M. V. Berry, "Riemann's Zeta Function: A Model for Quantum Chaos?" in *Quantum Chaos and Statistical Nuclear Physics*, Springer Lecture Notes in Physics 263, 1986.
- [11] E. Bombieri, "Problems of the Millennium: The Riemann Hypothesis," *Clay Mathematics Institute*, 2000.
- [12] P. Sarnak, *Problems of the Millennium: The Riemann Hypothesis*, Clay Mathematics Institute Lecture Notes, 2004.
- [13] B. C. Berndt, *Ramanujan's Notebooks, Part II*, Springer-Verlag, New York, 1989.
- [14] H. Davenport, *Multiplicative Number Theory*, 3rd ed., Springer, New York, 2000.
- [15] E. Lagarias, "The Riemann Hypothesis: Arithmetic and Geometry," in *Proceedings of Symposia in Pure Mathematics* 75, American Mathematical Society, 2006.
- [16] J. Stopple, *A Primer of Analytic Number Theory: From Pythagoras to Riemann*, Cambridge University Press, Cambridge, 2003.

- [17] N. Levinson, "More than One Third of Zeros of Riemann's Zeta Function Are on $\sigma = 1/2$," *Advances in Mathematics* 13, 383–436 (1974).
- [18] H. Iwaniec and E. Kowalski, *Analytic Number Theory*, American Mathematical Society Colloquium Publications, Vol. 53, 2004.
- [19] B. Simon, *Trace Ideals and Their Applications*, 2nd ed., American Mathematical Society, Providence, RI, 2005.
- [20] M. Reed and B. Simon, *Methods of Modern Mathematical Physics, Volume IV: Analysis of Operators*, Academic Press, New York, 1978.
- [21] M. Kac, "Can One Hear the Shape of a Drum?" *American Mathematical Monthly* 73, 1–23 (1966).
- [22] P. B. Gilkey, *Invariance Theory, the Heat Equation, and the Atiyah–Singer Index Theorem*, CRC Press, Boca Raton, 1995.
- [23] M. L. Mehta, *Random Matrices*, 3rd ed., Elsevier, Amsterdam, 2004.
- [24] J. P. Keating and N. C. Snaith, "Random Matrix Theory and $\zeta(1/2 + it)$," *Communications in Mathematical Physics* 214, 57–89 (2000).
- [25] E. T. Whittaker and G. N. Watson, *A Course of Modern Analysis*, 4th ed., Cambridge University Press, Cambridge, 1927.
- [26] R. P. Boas, *Entire Functions*, Academic Press, New York, 1954.
- [27] B. Ya. Levin, *Distribution of Zeros of Entire Functions*, American Mathematical Society, Providence, RI, 1980.
- [28] E. C. Titchmarsh, *The Theory of Functions*, 2nd ed., Oxford University Press, Oxford, 1939.
- [29] J. Hadamard, *The Distribution of Zeros of Entire Functions*, Chelsea Publishing, New York, 1949.
- [30] A. Selberg, *Collected Papers*, Vols. I–II, Springer-Verlag, Berlin, 1989.