
Task-Conditioned Reliability under Latent Markov Confounding

Anonymous Author
Anonymous Institution

Abstract

Sequential machine-learning systems often operate under persistent but partially observed context, where uncertainty about latent regimes can propagate into downstream failure. We study how uncertainty in inferred latent context propagates into downstream reliability. For a latent Markov confounding model, we identify two distinct sources of degradation: imperfect latent-state recovery and weak task conditioning. Under explicit regularity conditions, we derive a graceful-degradation bound whose leading term scales as $\varepsilon_\gamma/\lambda_{\min}(\mathbf{J})$, where $\varepsilon_\gamma \equiv \mathbb{E}\|\boldsymbol{\gamma}_t - \mathbf{H}_t\|_1$ measures population posterior state error and $\lambda_{\min}(\mathbf{J})$ measures downstream conditioning. We further show that posterior entropy provides an observable multiplicative surrogate for this difficulty under two-sided spectral comparability, and develop spectral regularization with purged block cross-fitting for weak-overlap sequential estimation. Beyond causal estimation, we evaluate HMM, GRU, Transformer, and state-space representations under disentangled temporal and observation shifts. Across 15 independent benchmark worlds, latent-state classification accuracy alone does not fully explain downstream reliability, while posterior log-loss provides additional explanatory power after controlling for accuracy. Real longitudinal sensor data from four Indian cities reveal complementary failure mechanisms: state uncertainty is more informative in well-conditioned settings, whereas task conditioning becomes informative under severe overlap limitations. A synthetic financial stress test further illustrates an out-of-sample adaptive abstention policy.

Together, these results suggest that reliable sequential ML should assess uncertainty in latent context jointly with the geometry of the downstream task.

1 INTRODUCTION

Sequential machine learning systems frequently operate in environments governed by persistent, partially observed context—such as unobserved physical, macroeconomic, or operational regimes. A widespread design assumption in such systems is that predictive confidence or calibration over the latent state is sufficient for downstream reliability. However, this assumption fails: downstream reliability can depend critically on the *interaction* between latent-state uncertainty and downstream task geometry. When the downstream task is ill-conditioned—such as estimating causal effects under collinear treatments or weak overlap—even small posterior proxy error is catastrophically amplified, causing severe downstream failure despite seemingly confident latent representations.

Continuous-treatment causal inference in atmospheric systems provides a natural laboratory: estimating the causal effect of variation in the observed NO₂ exposure proxy on ambient PM_{2.5} (Dominici et al., 2014) (translating this observational exposure effect to an emissions policy requires structural transport assumptions). Observational time series in this domain violate two foundational assumptions:

1. **Persistent Latent Regime Confounding:** Planetary boundary-layer (PBL) dynamics transition between discrete thermodynamic regimes (e.g., severe nocturnal inversion vs. advective convective clearance). These latent states jointly govern precursor emission dispersion (T_t) and ambient particulate concentration (Y_t).
2. **Imperfect Regime Overlap & Temporal Dependence:** Latent states are never directly observed; auxiliary meteorological proxies Z_t exhibit substantial overlap, making posterior state recovery uncertain ($\varepsilon_\gamma > 0$), while autoregressive persis-

tence breaks cross-sectional exchangeability (Chernozhukov et al., 2018), leaking nuisance errors across temporal folds.

When standard Double Machine Learning (Chernozhukov et al., 2018) or tree-based plug-in estimators are applied naively to environmental time series without accounting for latent regimes, the resulting causal estimands can exhibit severe bias and, in our benchmark, sign inversions—falsely suggesting that precursor reductions exacerbate ambient pollution.

Related Work. Standard DML (Chernozhukov et al., 2018; Foster and Syrgkanis, 2023; Nie and Wager, 2021) relies on observed confounding and i.i.d. exchangeability. In time-series DML, Ciganovic et al. (2026) adapt DML to macroeconomic series via Reverse Cross-Fitting; however, their framework requires strict stationarity and time-reversibility, precluding persistent regime-switching physical systems. In latent-state causal inference, Clouth et al. (2024) examine latent Markov models using parametric g-computation, while Huang et al. (2023) study conditionally stationary time series; neither addresses semiparametric Neyman-orthogonal score stability under weak regime overlap. Crucially, recent advances at the frontier of causal representation learning (Anonymous, 2026) treat weak overlap not as an ignorable nuisance, but as an explicit statistical design problem. OR-DML provides the temporal and latent-regime analogue to this paradigm.

Contributions. We formulate the bridge between latent uncertainty, task conditioning, and sequential reliability:

- **Failure Decomposition (Theorems 3–4):** We formalize latent regime confounding and prove that downstream estimation suffers two distinct degradation mechanisms: imperfect latent recovery (ε_γ) and weak task conditioning ($\lambda_{\min}(\mathbf{J})$).
- **Principled Difficulty Measure (Proposition 5):** We formulate causal difficulty $\mathcal{D}_{\text{causal}} = \frac{\varepsilon_\gamma}{\lambda_{\min}(\mathbf{J})}$ and establish rolling entropy $\mathcal{D}_{\text{operational}}(t) = \frac{H(\gamma_t)}{\lambda_{\min}(\mathbf{J}_t)}$ as an observable multiplicative surrogate under two-sided spectral comparability.
- **Adaptive Sequential Response (Theorems 6–7):** We derive leading-order risk bounds under long-run covariance approximations and develop data-driven empirical risk-surrogate selection $\hat{\lambda}^*$ with purged block cross-fitting for weak-overlap sequential estimation.
- **Multi-Domain Empirical Investigation:** Across controlled multi-world benchmarks, > 14,000 hours of Indian atmospheric sensor data, and a synthetic financial macro stress test, we find

empirical instances of the predicted separation between latent representation accuracy, posterior calibration, and downstream task reliability, demonstrating that task geometry governs failure bottlenecks.

2 STRUCTURAL CAUSAL MODEL & IDENTIFICATION

Let $(\Omega, \mathcal{F}, P)$ be a probability space. Consider a continuous-time atmospheric process observed at discrete equidistant intervals $t \in \{1, \dots, N\}$.

2.1 Structural Equations

We formalize the atmospheric data-generating process via the following Structural Causal Model (SCM):

$$S_t \sim \text{Markov}(\Pi, \pi_0), \quad S_t \in \{1, \dots, K\}, \quad (1)$$

$$Z_t = \mu_Z(S_t) + \xi_t, \quad (2)$$

$$X_t = \phi(X_{t-1}, S_t) + \epsilon_t, \quad (3)$$

$$T_t = m(X_t, S_t) + V_t, \quad \mathbb{E}[V_t | X_t, S_t] = 0, \quad (4)$$

$$Y_t = \theta(S_t)T_t + g(X_t, S_t) + U_t, \quad (5)$$

with $\mathbb{E}[U_t | X_t, S_t, T_t] = 0$, where:

- S_t is an unobserved discrete Markov regime with transition matrix $\Pi_{jk} = P(S_t = k | S_{t-1} = j)$ and stationary distribution π^* .
- $Z_t \in \mathbb{R}^q$ denotes observed auxiliary physical state indicators (e.g., surface wind velocity vectors, thermal vertical gradients, atmospheric pressure).
- $X_t \in \mathbb{R}^p$ is a vector of observed weather confounders and temporal lags.
- $T_t \in \mathbb{R}$ is the continuous treatment (precursor emission proxy, e.g., NO_2).
- $Y_t \in \mathbb{R}$ is the continuous outcome (ambient pollutant, e.g., $\text{PM}_{2.5}$).
- $\theta(S_t) = \sum_{k=1}^K \theta_k^* \mathbb{I}(S_t = k)$ represents the regime-specific causal effect.

2.2 Identification Under Latent Regimes

Our objective is to identify and estimate the regime-specific marginal causal effects:

$$\theta_k^* = \frac{\partial \mathbb{E}[Y_t | \text{do}(T_t = u), S_t = k]}{\partial u}, \quad k \in \{1, \dots, K\}, \quad (6)$$

as well as the stationary Average Treatment Effect (ATE):

$$\theta_{\text{ATE}}^* = \sum_{k=1}^K \pi_k^* \theta_k^*. \quad (7)$$

Assumption 1 (Regime-Conditional Unconfoundedness). Conditional on observed covariates X_t and latent




Figure 1: Causal Directed Acyclic Graph (DAG). Latent regime S_t acts as a common cause for Z_t , X_t , treatment T_t , and outcome Y_t . Omitting S_t opens non-causal confounding pathways.

regime S_t , potential outcomes are independent of treatment assignment:

$$Y_t(u) \perp T_t \mid (X_t, S_t), \quad \forall u \in \text{supp}(T_t). \quad (8)$$

Assumption 2 (Overlap & Positivity). There exist constants $c_1, c_2 > 0$ such that for all $k \in \{1, \dots, K\}$ and almost everywhere on $\text{supp}(X_t)$:

$$c_1 \leq \text{Var}(T_t \mid X_t, S_t = k) \leq c_2. \quad (9)$$

Assumption 3 (Structural Innovation). The outcome innovation $\varepsilon_t \equiv Y_t - \mathbb{E}[Y_t \mid \mathcal{F}_t]$, where $\mathcal{F}_t \equiv \sigma(X_t, T_t, Z_{1:N})$, satisfies $\mathbb{E}[\varepsilon_t \mid \mathcal{F}_t] = 0$ identically. Equivalently, $Y_t - \sum_{k=1}^K \theta_k^* \gamma_{tk}^* T_t - G(X_t, \gamma_t^*)$ has conditional mean zero given $(X_t, T_t, Z_{1:N})$. *Discussion.* This assumption is implied by the SCM (1)–(5) together with $\mathbb{E}[U_t \mid X_t, S_t, T_t] = 0$ and the law of iterated expectations. It ensures the martingale innovation property underpinning the asymptotic normality proof (Theorem 7). Unlike oracle regime separation, Assumption 3 requires only that the SCM correctly specifies the conditional mean structure of Y_t , not that regimes are identified with probability 1.

Assumption 4 (Nuisance Estimation and Mixing Rates). The data process (Y_t, T_t, X_t, Z_t) is strictly stationary and geometrically α -mixing with $\alpha(m) \leq C_\alpha \exp(-c_\alpha m)$ for $C_\alpha, c_\alpha > 0$. Nuisance estimators satisfy $\|\hat{\mu}_Y - \mu_Y\|_{L_2(P)} \cdot \|\hat{\mu}_D - \mu_D\|_{L_2(P)} = o_P(N^{-1/2})$, with individual convergence rates $o_P(N^{-1/4})$.

3 FAILURE OF STANDARD DML: THE OMITTED REGIME BIAS

We now state our first main theoretical result, characterizing the exact failure mode of standard cross-sectional DML.

Theorem 1 (Decomposition of Omitted Regime Bias). Suppose the data are generated according to SCM (1)–(5). Let $\hat{\theta}_{\text{naive}}$ be the standard cross-sectional DML estimator (Chernozhukov et al., 2018) that conditions only on observed covariates X_t , omitting S_t . Define the naive cross-sectional residuals $\tilde{T}_t = T_t - \mathbb{E}[T_t \mid X_t]$ and $\tilde{Y}_t = Y_t - \mathbb{E}[Y_t \mid X_t]$.

Then, as $N \rightarrow \infty$, $\hat{\theta}_{\text{naive}}$ converges in probability to:

$$\hat{\theta}_{\text{naive}} \xrightarrow{P} \sum_{k=1}^K \omega_k \theta_k^* + \mathcal{B}_{\text{heterogeneity}} + \mathcal{B}_{\text{confounding}}, \quad (10)$$

where $\omega_k = \frac{\mathbb{E}[\mathbb{I}(S_t=k)\tilde{T}_t^2]}{\mathbb{E}[\tilde{T}_t^2]}$ with $\sum_k \omega_k = 1$, and the bias components are:

$$\mathcal{B}_{\text{heterogeneity}} = \frac{\sum_{k=1}^K \theta_k^* \mathbb{E}[\bar{m}(X_t) \Delta m_k(X_t) \pi_k(X_t)]}{\mathbb{E}[\tilde{T}_t^2]}, \quad (11)$$

$$\mathcal{B}_{\text{confounding}} = \frac{\sum_{k=1}^K \mathbb{E}[\pi_k(X_t) \Delta m_k(X_t) \Delta g_k(X_t)]}{\mathbb{E}[\tilde{T}_t^2]}, \quad (12)$$

with $\pi_k(X_t) \equiv P(S_t = k \mid X_t)$, $\bar{m}(X_t) \equiv \mathbb{E}[T_t \mid X_t]$, $\Delta m_k(X_t) \equiv \mathbb{E}[T_t \mid X_t, S_t = k] - \bar{m}(X_t)$, and $\Delta g_k(X_t) \equiv g(X_t, k) - \mathbb{E}[g(X_t, S_t) \mid X_t]$.

Proof. Under Neyman orthogonality and consistency of cross-fitted nuisance estimators on X_t , $\hat{\theta}_{\text{naive}} \xrightarrow{P} \frac{\mathbb{E}[\tilde{T}_t \tilde{Y}_t]}{\mathbb{E}[\tilde{T}_t^2]}$. By SCM (4), $T_t = m(X_t, S_t) + V_t$ with $\mathbb{E}[V_t \mid X_t, S_t] = 0$. Thus $\tilde{T}_t = \sum_{k=1}^K \mathbb{I}(S_t = k) \Delta m_k(X_t) + V_t$. Similarly, $\bar{\ell}(X_t) \equiv \mathbb{E}[Y_t \mid X_t] = \sum_{k=1}^K \pi_k(X_t) [\theta_k^* m_k(X_t) + g_k(X_t)]$. Substituting Y_t from (5) and writing $T_t = \tilde{T}_t + \bar{m}(X_t)$, the outcome residual decomposes into:

$$\begin{aligned} \tilde{Y}_t &= \sum_{k=1}^K \theta_k^* \mathbb{I}(S_t = k) \tilde{T}_t + \Delta g(X_t, S_t) + U_t \\ &\quad + \sum_{k=1}^K \theta_k^* [\mathbb{I}(S_t = k) \bar{m}(X_t) - \pi_k(X_t) m_k(X_t)]. \end{aligned}$$

Multiplying by $\tilde{T}_t$ and taking expectations conditional on (X_t, S_t) : First, $\mathbb{E}[\tilde{T}_t U_t] = 0$ by Assumption 1. Second, $\mathbb{E}[\tilde{T}_t \sum_k \theta_k^* \mathbb{I}(S_t = k) \tilde{T}_t] = \sum_k \theta_k^* \mathbb{E}[\mathbb{I}(S_t = k) \tilde{T}_t^2]$. Third, since $\mathbb{E}[\tilde{T}_t \mathbb{I}(S_t = k) \mid X_t] = \Delta m_k(X_t) \pi_k(X_t)$ and $\mathbb{E}[\tilde{T}_t \mid X_t] = 0$, the second term yields $\sum_k \theta_k^* \bar{m}(X_t) \Delta m_k(X_t) \pi_k(X_t)$. Fourth, $\mathbb{E}[\tilde{T}_t \Delta g(X_t, S_t) \mid X_t] = \sum_k \pi_k(X_t) \Delta m_k(X_t) \Delta g_k(X_t)$. Dividing by $\mathbb{E}[\tilde{T}_t^2]$ yields (11) and (12). $\square$

Corollary 2 (Sign Inversion & Bias Amplification). When latent stagnation induces strong positive confounding ($\Delta m_k(X_t) \Delta g_k(X_t) \gg 0$), but residual treatment variance $\mathbb{E}[\tilde{T}_t^2]$ is attenuated due to conditioning

on X_t , the omitted confounding bias $\mathcal{B}_{\text{confounding}}$ undergoes severe bias amplification. In coastal airsheds with negative covariance phases, $\mathcal{B}_{\text{confounding}} < -\sum_k \omega_k \theta_k^*$, driving $\text{plim}(\hat{\theta}_{\text{naive}}) < 0$ despite $\theta_k^* > 0$.

4 METHODOLOGY: OVERLAP-AWARE REGIME DML

To overcome the failure of standard cross-sectional DML, we introduce **Overlap-Aware Regime Double Machine Learning (OR-DML)**.

4.1 Latent Regime Formulation & Observability Diagnostics

Let $Z_t \in \mathbb{R}^q$ denote exogenous meteorological state trajectories (strictly pre-treatment, avoiding outcome contamination). We model the latent atmospheric state $S_t \in \{1, \dots, K\}$ via a Hidden Markov Model with transition matrix Π and emission parameters ϕ . Depending on the regulatory horizon, state posteriors fall into three operational modes:

1. **Retrospective Smoothed Posteriors (Offline):** $\gamma_{tk}^{\text{smooth}} \equiv P(S_t = k \mid Z_{1:N})$ for retrospective environmental epidemiology and structural causal auditing;
2. **Training-Fitted Forward Filtered Posteriors:** $\gamma_{tk}^{\text{filter}} \equiv P(S_t = k \mid Z_{1:t}, \hat{\phi})$ where global transition/emission dynamics $\hat{\phi}$ are calibrated offline, but sequential state updates proceed strictly causally on past-to-present observations $Z_{1:t}$; or
3. **Expanding-Window Online Filter:** where both parameter estimation and forward filtering update causally without access to future measurements.

Our empirical and benchmark implementations evaluate Mode 1 (offline smoothing) and Mode 2 (training-fitted forward filtering). We define the mean posterior entropy as a direct diagnostic of latent state observability:

$$\bar{H}(\gamma) \equiv -\frac{1}{N} \sum_{t=1}^N \sum_{k=1}^K \gamma_{tk} \ln(\gamma_{tk} + \epsilon). \quad (13)$$

A Priori Diagnostics: Standardized separation $\Delta_Z < 0.5$ or entropy $\bar{H} \rightarrow \ln K$ signals non-identification (Theorem 3; rules in Appendix B.3).

4.2 State-Weighted Coupled System & Spectral Regularization

To avoid heuristic decoupled regressions that suffer cross-talk bias under regime overlap, OR-DML esti-

mates regime-specific conditional expectations via state-weighted regressions targeting population expectations $m_k^\gamma(x) \equiv \frac{\mathbb{E}[\gamma_{tk} T_t \mid X_t = x]}{\mathbb{E}[\gamma_{tk} \mid X_t = x]}$ and $\mu_k^\gamma(x) \equiv \frac{\mathbb{E}[\gamma_{tk} Y_t \mid X_t = x]}{\mathbb{E}[\gamma_{tk} \mid X_t = x]}$:

$$\hat{m}_k(X) = \arg \min_m \sum_{t \in \mathcal{T}_b} \gamma_{tk} (T_t - m(X_t))^2, \quad (14)$$

$$\hat{\mu}_k(X) = \arg \min_\mu \sum_{t \in \mathcal{T}_b} \gamma_{tk} (Y_t - \mu(X_t))^2. \quad (15)$$

On evaluation fold $\mathcal{B}_b$, we form centered residuals $\tilde{T}_{tk} = T_t - \hat{m}_k(X_t)$ and $\tilde{Y}_{tk} = Y_t - \hat{\mu}_k(X_t)$. The empirical coupled Gram matrix $\hat{\mathbf{J}} \in \mathbb{R}^{K \times K}$ and score vector $\hat{\mathbf{S}} \in \mathbb{R}^K$ are constructed as:

$$\hat{J}_{jk} = \frac{1}{N} \sum_{t=1}^N \gamma_{tj} \gamma_{tk} \tilde{T}_{tj} \tilde{T}_{tk}, \quad \hat{S}_k = \frac{1}{N} \sum_{t=1}^N \gamma_{tk} \tilde{T}_{tk} \tilde{Y}_{tk}. \quad (16)$$

When regime separation is weak or treatments are collinear, $\lambda_{\min}(\hat{\mathbf{J}}) \rightarrow 0$, rendering the unregularized solve $\hat{\mathbf{J}}^{-1} \hat{\mathbf{S}}$ ill-conditioned. OR-DML introduces the **Spectrally Regularized Estimator**:

$$\hat{\theta}_\lambda = (\hat{\mathbf{J}} + \lambda \mathbf{I})^{-1} \hat{\mathbf{S}}, \quad (17)$$

where $\lambda \geq 0$ trades off regularization bias against collinearity variance. *Scalability & Non-Identifiability:* Inverting $\hat{\mathbf{J}}$ is $\mathcal{O}(K^3)$ with sample complexity $N = \tilde{\Omega}\left(\frac{K \log(K/\delta)}{\min_k (\pi_k^*)^2 \lambda_{\min}^2 \rho_{\text{gap}}}\right)$ under α -mixing (Appendix C.2); when $\kappa(\hat{\mathbf{J}}) > 10^4$ or $\text{CI}(\lambda_{\min})$ covers 0, systems should diagnose non-identifiability and abstain. We distinguish three causal estimand targets: (i) the true stationary population ATE $\theta_{\text{PATE}}^* = \pi^{*\top} \theta^*$, (ii) the estimated stationary ATE $\hat{\theta}_{\text{PATE}} = \hat{\pi}(\hat{\Lambda})^\top \hat{\theta}_\lambda$, and (iii) the empirical sample occupation ATE $\hat{\theta}_{\text{SATE}} = \bar{\gamma}_N^\top \hat{\theta}_\lambda$, where $\bar{\gamma}_N \equiv \frac{1}{N} \sum_{t=1}^N \gamma_t$.

4.3 Purged Block Cross-Fitting with Bartlett Embargo

To ensure validity under temporal dependence, we partition observations into B contiguous blocks. To eliminate cross-fold leakage under α -mixing, each test block $\mathcal{B}_b$ is surrounded by an embargo buffer of width τ^* (Algorithm 1). We determine τ^* from the data via the Bartlett significance threshold:

$$\tau^* = \max \left\{ h \geq 1 : \max_j |\hat{\rho}_j(h)| \geq \frac{1.96}{\sqrt{N}} \right\} + 1. \quad (18)$$

5 THEORETICAL ANALYSIS

We now present the theoretical foundation of OR-DML, formalizing the two distinct failure modes: proxy uncertainty and causal ill-conditioning.




Figure 2: Visualizing the Frisch-Waugh Singularity and Omitted Regime Bias. (a) As residual treatment variance $\mathbb{E}[\tilde{T}_t^2] \rightarrow 0$ via partialling-out X_t , omitted regime bias $\mathcal{B}_{\text{confounding}}$ explodes hyperbolically in Standard DML, while RC-DML remains strictly unbiased ($\mathcal{B}_{\text{regime}} \equiv 0$). (b) Monte Carlo empirical bias across increasing explanatory power $R^2(T | X)$, verifying theoretical bias amplification.

Table 1: The Latent-Confounding Difficulty Frontier: 500 Monte Carlo Replications across Separation Regimes $\Delta_Z \in [0.2, 4.0]$ ($N = 1, 200$, True $\theta_{\text{ATE}}^* = 1.5278$). Nominal coverage target is 95.0% for both Sample ATE (SATE) and Population ATE (PATE).

Δ_Z	Estimator	Mean $\hat{\theta}$	Abs Bias	RMSE	SATE Cov	PATE Cov	$\lambda_{\min}(\mathbf{J})$	$\kappa(\mathbf{J})$	$\hat{\varepsilon}_\gamma$
0.2 (Severe Overlap)	Oracle DML	1.5327	0.0558	0.0703	96.0%	94.0%	0.4485	1.26	0.0000
	Standard DML	9.9833	8.4556	8.4562	0.0%	0.0%	4.9339	1.00	—
	Unreg. Coupled DML	10.1344	8.6066	8.6077	0.0%	0.0%	0.6495	6.67	0.8327
	Spectral OR-DML (Ours)	10.1236	8.5958	8.5969	0.0%	0.0%	0.6495	6.67	0.8327
0.5 (Weak Separation)	Oracle DML	1.5225	0.0552	0.0713	92.0%	95.0%	0.4480	1.28	0.0000
	Standard DML	9.9565	8.4287	8.4296	0.0%	0.0%	4.9560	1.00	—
	Unreg. Coupled DML	10.3373	8.8095	8.8138	0.0%	0.0%	0.7680	3.23	0.3879
	Spectral OR-DML (Ours)	10.3252	8.7974	8.8017	0.0%	0.0%	0.7680	3.23	0.3879
1.0 (Moderate)	Oracle DML	1.5274	0.0572	0.0735	92.0%	92.0%	0.4515	1.26	0.0000
	Standard DML	9.9608	8.4331	8.4340	0.0%	0.0%	4.9535	1.00	—
	Unreg. Coupled DML	5.7832	4.2555	4.3823	0.0%	0.0%	0.5294	67.01	0.1049
	Spectral OR-DML (Ours)	5.7759	4.2481	4.3757	0.0%	0.0%	0.5294	67.01	0.1049
2.0 (Clean Separation)	Oracle DML	1.5289	0.0521	0.0658	93.0%	97.0%	0.4536	1.26	0.0000
	Standard DML	9.9596	8.4318	8.4327	0.0%	0.0%	4.9539	1.00	—
	Unreg. Coupled DML	2.0034	0.5020	1.9036	89.0%	93.0%	0.4321	690.83	0.0501
	Spectral OR-DML (Ours)	2.0014	0.5012	1.9048	87.0%	92.0%	0.4321	690.83	0.0501
	Filtered OR-DML (Ours)	2.0238	0.5203	1.9062	87.0%	90.0%	0.4329	859.89	0.0502
4.0 (Isolated Regimes)	Oracle DML	1.5261	0.0540	0.0670	96.0%	95.0%	0.4394	1.31	0.0000
	Standard DML	9.9787	8.4509	8.4518	0.0%	0.0%	4.9123	1.00	—
	Unreg. Coupled DML	1.6957	0.2236	1.2113	94.0%	93.0%	0.4299	30.34	0.0199
	Spectral OR-DML (Ours)	1.6932	0.2233	1.2102	94.0%	93.0%	0.4299	30.34	0.0199
	Filtered OR-DML (Ours)	1.6932	0.2233	1.2100	94.0%	93.0%	0.4299	30.52	0.0198

Theorem 3 (Non-Identification Under Unconstrained Overlap). *Suppose the latent state is unobserved, and auxiliary proxies Z_t satisfy $P(Z_t | S_t = j) = P(Z_t | S_t = k)$ for all j, k . Then the regime-specific causal effect vector $\theta^* = (\theta_1^*, \dots, \theta_K^*)^\top$ is not point-identified from the joint distribution of observables (Y_t, T_t, X_t, Z_t) : there exist structural causal models P_1 and P_2 satisfying Assumptions 1–3 with identical joint observable laws $P_1(Y_{1:N}, T_{1:N}, X_{1:N}, Z_{1:N}) \equiv P_2(Y_{1:N}, T_{1:N}, X_{1:N}, Z_{1:N})$ for all $N \geq 1$, yet possessing strictly distinct, non-permutation causal effect vectors $\theta^{(1)} \neq \theta^{(2)}$.*

Theorem 3 formally proves via an explicit constructive counterexample (Appendix A) that predictive state inference alone cannot guarantee causal identification

without explicit separation or proxy error bounds.

Theorem 4 (Graceful Degradation Error Bound). *Let $\mathbf{H}_t = \mathbf{e}_{S_t} \in \{0, 1\}^K$ denote the unobserved true regime indicator, and let $\gamma_t \in \Delta^K$ be an estimated posterior state proxy with average L_1 proxy error $\varepsilon_\gamma \equiv \mathbb{E}\|\gamma_t - \mathbf{H}_t\|_1$. Suppose $\|\theta^*\|_2 \leq M_\theta < \infty$, $\sup_t |\tilde{T}_{tk}| \leq M_T < \infty$ almost surely, and regime functions satisfy $\max_{j,k} |g(x, j) - g(x, k)| \leq L_g < \infty$ and $\max_{j,k} |m(x, j) - m(x, k)| \leq L_m < \infty$. If $\lambda_{\min}(\mathbf{J}) > 0$, the unregularized coupled estimator $\hat{\theta}_\gamma = \hat{\mathbf{J}}^{-1} \hat{\mathbf{S}}$ constructed from state-weighted nuisances satisfies:*

$$\|\hat{\theta}_\gamma - \theta^*\|_2 \leq \frac{C \cdot \varepsilon_\gamma}{\lambda_{\min}(\mathbf{J})} + \mathcal{O}_P(N^{-1/2}), \quad (19)$$

where $C = L_g M_T + M_\theta M_T^2 + M_\theta M_T L_m < \infty$.




Figure 3: The Latent-Confounding Difficulty Frontier across 500 Monte Carlo Replications. (a) Absolute causal bias $|\hat{\theta} - \theta^*|$ vs. regime separation Δ_Z . (b) Empirical RMSE. (c) Empirical 95% CI coverage vs. nominal 95% target. (d) Factorial calibration: Absolute bias $|\hat{\theta} - \theta^*|$ vs. empirical difficulty scaling $\hat{\varepsilon}_{\gamma,N}/\lambda_{\min}(\mathbf{J})$, validating the theoretical degradation scaling (Theorem 4) across 15 factorial grid cells, highlighting Adversarial Corners A (high $\hat{\varepsilon}_{\gamma}$, healthy $\lambda_{\min}$) and B (low $\hat{\varepsilon}_{\gamma}$, weak $\lambda_{\min}$).

Algorithm 1 Overlap-Aware Regime DML (OR-DML)

Require: Series $\{Y_t, T_t, X_t, Z_t\}_{t=1}^N$, blocks B , embargo τ^* , spectral penalty $\lambda \geq 0$.

- 1: Infer posteriors $\gamma_t = P(S_t | Z_{1:N})$ (or causal forward filter $P(S_t | Z_{1:t})$).
- 2: **for** block $b = 1$ to B **do**
- 3: Set test block $\mathcal{B}_b$; training set $\mathcal{T}_b = \{1..N\} \setminus (\mathcal{B}_b \pm \tau^*)$.
- 4: **for** regime $k = 1$ to K **do**
- 5: Fit state-weighted nuisances $\hat{m}_k^{(-b)}(X)$ and $\hat{\mu}_k^{(-b)}(X)$ on $\mathcal{T}_b$ with weights γ_{tk} .
- 6: On $\mathcal{B}_b$, form residuals $\tilde{T}_{tk} = T_t - \hat{m}_k(X_t)$ and $\tilde{Y}_{tk} = Y_t - \hat{\mu}_k(X_t)$.
- 7: **end for**
- 8: **end for**
- 9: Compute Gram $\hat{J}_{jk} = \frac{1}{N} \sum_t \gamma_{tj} \gamma_{tk} \tilde{T}_{tj} \tilde{T}_{tk}$ and score $\hat{S}_k = \frac{1}{N} \sum_t \gamma_{tk} \tilde{T}_{tk} \tilde{Y}_{tk}$.
- 10: Compute $\hat{\theta}_\lambda = (\hat{\mathbf{J}} + \lambda \mathbf{I})^{-1} \hat{\mathbf{S}}$ and $\hat{\theta}_{\text{SATE}} = \bar{\gamma}^\top \hat{\theta}_\lambda$, $\hat{\theta}_{\text{PATE}} = \hat{\pi}^\top \hat{\theta}_\lambda$.
- 11: **return** Estimates $\hat{\theta}_\lambda, \hat{\theta}_{\text{SATE}}, \hat{\theta}_{\text{PATE}}$ and diagnostics $(\lambda_{\min}, \kappa, \bar{H})$.

Theorem 4 replaces heuristic orthogonality claims with a rigorous bound connecting *posterior proxy error* ε_γ and *Jacobian conditioning* $\lambda_{\min}(\mathbf{J})$. While the score is Neyman-orthogonal to continuous nuisances (μ_k, m_k) , it is *not* orthogonal to γ_t ; latent proxy uncertainty carries an irremediable statistical price.

The Difficulty Frontier and Operational Surrogate Bridge. Theorem 4 demonstrates that the leading asymptotic bias scales with the theoretical ratio:

$$\mathcal{D}_{\text{causal}} \equiv \frac{\varepsilon_\gamma}{\lambda_{\min}(\mathbf{J})}, \quad (20)$$

where $\varepsilon_\gamma \equiv \mathbb{E} \|\gamma_t - \mathbf{H}_t\|_1$ is the population posterior state error, estimated in finite samples by $\hat{\varepsilon}_{\gamma,N} \equiv \frac{1}{N} \sum_{t=1}^N \|\gamma_t - \mathbf{H}_t\|_1$. Because true states are unobserved at runtime, deployed systems cannot compute ε_γ or $\hat{\varepsilon}_{\gamma,N}$ directly. We bridge unobservable causal difficulty to the computable operational index $\mathcal{D}_{\text{operational}}(t) \equiv H(\gamma_t)/\lambda_{\min}(\mathbf{J}_t)$ via the following re-

sult under spectral comparability:

Assumption 5 (Two-Sided Spectral Comparability). There exist constants $0 < \underline{c}_J \leq \bar{c}_J < \infty$ such that rolling Gram matrices $\mathbf{J}_t$ satisfy $\underline{c}_J \leq \frac{\lambda_{\min}(\mathbf{J}_t)}{\lambda_{\min}(\mathbf{J})} \leq \bar{c}_J$ almost surely for all evaluation windows $t \geq w$.

Proposition 5 (Surrogate Entropy-Difficulty Bridge). For causal forward-filtered posteriors $\gamma_t = P(S_t \in \cdot | \mathcal{F}_t)$ with filtration $\mathcal{F}_t = \sigma(Z_{1:t})$, the posterior error satisfies $\mathbb{E}[\|\gamma_t - \mathbf{H}_t\|_1 | \mathcal{F}_t] = 2 \text{Gini}(\gamma_t) \leq C_K H(\gamma_t)$, where $C_K \leq 1/\ln 2 \approx 1.4427$ for $K = 2$. We define the difficulty hierarchy:

$$\mathcal{D}_{\text{causal}} \equiv \frac{\varepsilon_\gamma}{\lambda_{\min}(\mathbf{J})}, \quad \mathcal{D}_{\text{global}}^H(t) \equiv \frac{H(\gamma_t)}{\lambda_{\min}(\mathbf{J})}, \quad (21)$$

$$\mathcal{D}_{\text{operational}}(t) \equiv \frac{H(\gamma_t)}{\lambda_{\min}(\mathbf{J}_t)}. \quad (22)$$

- (i) Causal difficulty is unconditionally bounded in population by global entropy: $\mathcal{D}_{\text{causal}} \leq C_K \mathbb{E}[\mathcal{D}_{\text{global}}^H(t)]$.
- (ii) Under Assumption 5, the deployable index satisfies:

$$\mathcal{D}_{\text{causal}} \leq C_K \bar{c}_J \mathbb{E}[\mathcal{D}_{\text{operational}}(t)], \quad (23)$$

$$\mathbb{E}[\mathcal{D}_{\text{operational}}(t)] \leq \underline{c}_J^{-1} \mathbb{E}[\mathcal{D}_{\text{global}}^H(t)]. \quad (24)$$

The empirical sample surrogate $\hat{\mathcal{D}}_{\text{causal},N} \equiv \hat{\varepsilon}_{\gamma,N}/\lambda_{\min}(\mathbf{J})$ converges to $\mathcal{D}_{\text{causal}}$ with $\mathcal{O}_P(N^{-1/2})$ error.

Proposition 5 (proof in Appendix B.2) mathematically grounds the deployable monitoring engine across three levels of observability: unobservable causal difficulty $\mathcal{D}_{\text{causal}} \rightarrow$ observable operational surrogate $\mathcal{D}_{\text{operational}}(t) \rightarrow$ realized downstream loss L_{t+h} . We emphasize that (23) establishes $\mathcal{D}_{\text{operational}}(t)$ as an observable surrogate upper bound for $\mathcal{D}_{\text{causal}}$, not a two-sided equivalence. The entropy bridge holds algebraically for retrospective smoothed posteriors $\gamma_t = P(S_t \in \cdot | Z_{1:N})$ as well, though with retrospective rather than prospective interpretation.

Theorem 6 (Leading-Order Risk Bound under Long-Run Covariance Approximation). Let $\hat{\theta}_\lambda = (\hat{\mathbf{J}} +$

$\lambda \mathbf{I})^{-1} \hat{\mathbf{S}}$ be the spectrally regularized OR-DML estimator, and let $\boldsymbol{\theta}_\lambda = (\mathbf{J} + \lambda \mathbf{I})^{-1} \mathbf{S}$. Under the conditions of Theorem 4, the regularized estimand satisfies the exact deterministic identity:

$$\boldsymbol{\theta}_\lambda - \boldsymbol{\theta}^* = (\mathbf{J} + \lambda \mathbf{I})^{-1} (\mathbf{b}_\gamma - \lambda \boldsymbol{\theta}^*), \quad (25)$$

where $\|\mathbf{b}_\gamma\|_2 \leq C\varepsilon_\gamma$. Furthermore, under the long-run score covariance approximation $\boldsymbol{\Omega} = \lim_{N \rightarrow \infty} \text{Var}(N^{-1/2} \sum_t \boldsymbol{\Psi}_t)$, the expected squared estimation error satisfies the leading-order risk bound:

$$\mathbb{E} \|\hat{\boldsymbol{\theta}}_\lambda - \boldsymbol{\theta}^*\|_2^2 \leq \frac{(C\varepsilon_\gamma + \lambda M_\theta)^2 + N^{-1} \text{Tr}(\boldsymbol{\Omega})}{(\lambda_{\min}(\mathbf{J}) + \lambda)^2} + \mathcal{O}(N^{-1}), \quad (26)$$

which accounts for the leading-order stochastic expansion $\mathbb{E} \|\hat{\boldsymbol{\theta}}_\lambda - \boldsymbol{\theta}_\lambda\|_2^2 = \frac{1}{N} \text{Tr}((\mathbf{J} + \lambda \mathbf{I})^{-1} \boldsymbol{\Omega} (\mathbf{J} + \lambda \mathbf{I})^{-1}) + o(N^{-1})$ and deterministic shrinkage bias (Appendix C), sharpening to $\mathcal{O}(N^{-3/2})$ under the local parameter regime $\varepsilon_\gamma + \lambda = \mathcal{O}(N^{-1/2})$. When overlap is ill-conditioned ($\lambda_{\min}(\mathbf{J}) \approx 0$), choosing $\lambda > 0$ reduces the variance upper bound monotonically in λ . Let $R_{\text{bound}}(\lambda) \equiv \frac{(C\varepsilon_\gamma + \lambda M_\theta)^2 + N^{-1} \text{Tr}(\boldsymbol{\Omega})}{(\lambda_{\min}(\mathbf{J}) + \lambda)^2}$ denote the theoretical risk bound, with minimizer $\lambda_{\text{bound}}^* \equiv \arg \min_{\lambda \geq 0} R_{\text{bound}}(\lambda)$. In practice, we evaluate the empirical plug-in surrogate $\hat{R}(\lambda) \equiv \left(\frac{C\varepsilon_\gamma + \lambda \|\hat{\boldsymbol{\theta}}_0\|_2}{\lambda_{\min}(\mathbf{J}) + \lambda} \right)^2 + \frac{\text{Tr}(\hat{\boldsymbol{\Omega}})}{N(\lambda_{\min}(\mathbf{J}) + \lambda)^2}$, yielding the data-driven adaptive parameter $\hat{\lambda}_{\text{emp}}^* = \arg \min_{\lambda \geq 0} \hat{R}(\lambda)$ (Appendix C.1). In our ill-conditioned benchmark ($\lambda_{\min} \approx 0.033$), empirical minimum MSE occurs near $\hat{\lambda}_{\text{empirical}}^* \approx 0.20$ (MSE = 0.0321, a 25.9 $\times$ empirical risk reduction in the benchmark DGP from $\lambda = 0.0$), while $\lambda = 0.10$ achieves an approximately 23.0 $\times$ empirical reduction (MSE = 0.0362 vs. 0.8312) with peak coverage (85.0%). We distinguish λ_{bound}^* and $\hat{\lambda}_{\text{emp}}^*$ from the true infeasible risk minimizer $\lambda_{\text{true}}^* = \arg \min_\lambda \mathbb{E} \|\hat{\boldsymbol{\theta}}_\lambda - \boldsymbol{\theta}^*\|_2^2$.

Theorem 7 (Asymptotic Normality Under Dependent Purged Cross-Fitting). *Under Assumptions 1–4, with embargo buffer $\tau^* \geq C \log N$, the regularized OR-DML estimator under retrospective smoothing or training-fitted forward filtering satisfies:*

$$\sqrt{N} (\hat{\boldsymbol{\theta}}_\lambda - \boldsymbol{\theta}^* - \text{Bias}(\lambda)) \xrightarrow{d} \mathcal{N}(\mathbf{0}, \boldsymbol{\Sigma}_\lambda), \quad (27)$$

where $\boldsymbol{\Sigma}_\lambda = (\mathbf{J} + \lambda \mathbf{I})^{-1} \boldsymbol{\Omega} (\mathbf{J} + \lambda \mathbf{I})^{-1}$. When $\lambda = o(N^{-1/2})$, $\sqrt{N} \text{Bias}(\lambda) \rightarrow \mathbf{0}$, recovering centered asymptotic normality.

Proposition 8 (Joint Asymptotic Distribution for Sample and Population ATE). *The sample ATE estimator $\hat{\theta}_{\text{SATE}} = \bar{\gamma}^\top \hat{\boldsymbol{\theta}}_\lambda$ has conditional asymptotic variance $\sigma_{\text{SATE}}^2 = \bar{\gamma}^\top \boldsymbol{\Sigma}_\lambda \bar{\gamma}$. The population ATE estimator $\hat{\theta}_{\text{PATE}} = \hat{\boldsymbol{\pi}}^\top \hat{\boldsymbol{\theta}}_\lambda$, where $\hat{\boldsymbol{\pi}}$ is the estimated stationary distribution of the Markov transition matrix $\hat{\mathbf{\Pi}}$, satisfies*

the joint delta expansion:

$$\begin{aligned} \sqrt{N} (\hat{\theta}_{\text{PATE}} - \theta_{\text{PATE}}^*) &= \boldsymbol{\pi}^{*\top} \sqrt{N} (\hat{\boldsymbol{\theta}}_\lambda - \boldsymbol{\theta}^*) \\ &\quad + \boldsymbol{\theta}^{*\top} \sqrt{N} (\hat{\boldsymbol{\pi}} - \boldsymbol{\pi}^*) + o_P(1), \end{aligned} \quad (28)$$

yielding asymptotic variance:

$$\sigma_{\text{PATE}}^2 = \boldsymbol{\pi}^{*\top} \boldsymbol{\Sigma}_\lambda \boldsymbol{\pi}^* + \boldsymbol{\theta}^{*\top} \boldsymbol{\Sigma}_\pi \boldsymbol{\theta}^* + 2\boldsymbol{\pi}^{*\top} \boldsymbol{\Sigma}_{\theta\pi} \boldsymbol{\theta}^*, \quad (29)$$

where $\boldsymbol{\Sigma}_\pi = \mathbf{D}_\pi \mathbf{Z} + \mathbf{Z}^\top \mathbf{D}_\pi - \mathbf{D}_\pi - \boldsymbol{\pi}^* \boldsymbol{\pi}^{*\top}$ with fundamental matrix $\mathbf{Z} = (\mathbf{I} - \boldsymbol{\Pi} + \mathbf{1} \boldsymbol{\pi}^{*\top})^{-1}$, and $\boldsymbol{\Sigma}_{\theta\pi} \equiv \lim_{N \rightarrow \infty} \text{Cov}(\sqrt{N}(\hat{\boldsymbol{\theta}}_\lambda - \boldsymbol{\theta}^*), \sqrt{N}(\hat{\boldsymbol{\pi}} - \boldsymbol{\pi}^*))$. The cross-covariance vanishes when the DML score influence is orthogonal to the HMM score direction: $\mathbb{E}[\boldsymbol{\phi}_\theta(W_t) \mathbf{s}_\Lambda(W_t)^\top] = \mathbf{0}$; in general, $\boldsymbol{\Sigma}_{\theta\pi}$ is estimated via the joint HAC sandwich estimator. Under stagnation ($\rho \approx 0.88$), regime occupation variance dominates conditional estimation variance by $\frac{1+\rho}{1-\rho} \approx 15.7\times$.

6 THE LATENT-CONFOUNDING DIFFICULTY FRONTIER

To test the theoretical predictions of Theorems 3–6, we conduct a comprehensive Monte Carlo evaluation across 500 replications, charting the causal estimation error across the **Latent-Confounding Difficulty Frontier**.

6.1 Experimental Design

We simulate dynamic atmospheric time series ($N = 1, 200$) with a two-state Markov chain $S_t \in \{0, 1\}$ (persistence $\rho = 0.88$). Observed meteorological controls $X_t \in \mathbb{R}^3$ follow an autoregressive AR(1) process ($\rho = 0.65$). Treatment assignment is confounded by both X_t and S_t : $T_t = 2.0(1 - S_t) + 6.0S_t + 0.8X_{t1} - 0.5X_{t2} + V_t$. Regime treatment effects are $\theta_0^* = 0.75$ and $\theta_1^* = 2.50$, yielding a true population ATE $\theta_{\text{ATE}}^* = 1.5278$.

The difficulty of latent regime recovery is governed continuously by the separation parameter $\Delta_Z \in \{0.2, 0.5, 1.0, 2.0, 4.0\}$ in exogenous observations $Z_t \sim \mathcal{N}(\pm \Delta_Z \boldsymbol{\mu}, \boldsymbol{\Sigma})$. At $\Delta_Z = 0.2$, state overlap is severe and regimes are near-unidentified ($\bar{H}(\boldsymbol{\gamma}) = 0.434$); at $\Delta_Z = 4.0$, regimes are cleanly separated ($\bar{H}(\boldsymbol{\gamma}) = 0.014$). We evaluate 100 independent Monte Carlo draws per grid point (500 total).

6.2 Difficulty Frontier Findings

Table 1 and Figure 3 reveal three key statistical phenomena:

1. **Persistence of Naive Confounding:** In our benchmark DGP, Standard and Block DML exhibit approximately constant omitted-regime bias

Table 2: Empirical Treatment Elasticities on Clean Longitudinal Ground-Sensor Data across Four Major Indian Megacities (14,000 Complete Hourly Observations, February 2025 – June 2026). Treatment $T_t = \text{NO}_2$, Outcome $Y_t = \text{PM}_{2.5}$.

City	Estimator	Regime	Effect $\hat{\theta}$	Std Error	95% CI	p-value	$\lambda_{\min}(\hat{J})$ [κ]
Delhi ($N = 4,640$)	Standard DML	Pooled	+0.4696	0.0443	[+0.3828, +0.5564]	< 0.0001	251.3 [1.0]
	Block DML	Pooled	+0.5815	0.0468	[+0.4898, +0.6733]	< 0.0001	274.5 [1.0]
	Spectral OR-DML	Regime 1	+0.7038	0.0513	[+0.6032, +0.8044]	< 0.0001	111.8 [1.58]
	Spectral OR-DML	Regime 2	+0.1281	0.0743	[-0.0174, +0.2737]	0.0845	111.8 [1.58]
	Spectral OR-DML	Overall ATE	+0.3742	0.0642	[+0.2484, +0.5000]	< 0.0001	111.8 [1.58]
	Filtered OR-DML	Overall ATE	+0.3745	0.0650	[+0.2471, +0.5019]	< 0.0001	116.3 [1.46]
Mumbai ($N = 4,180$)	Standard DML	Pooled	+2.6397	4.4187	[-6.0208, +11.3003]	0.5502	0.65 [1.0]
	Block DML	Pooled	+1.8174	4.1361	[-6.2893, +9.9242]	0.6604	0.68 [1.0]
	Spectral OR-DML	Regime 1	-0.0294	8.7189	[-17.1183, +17.0596]	0.9973	0.32 [1.09]
	Spectral OR-DML	Regime 2	+3.3036	1.4800	[+0.4027, +6.2045]	0.0256	0.32 [1.09]
	Spectral OR-DML	Overall ATE	+1.5926	4.3636	[-6.9600, +10.1451]	0.7151	0.32 [1.09]
	Filtered OR-DML	Overall ATE	+1.3247	4.2654	[-7.0355, +9.6850]	0.7561	0.30 [1.13]
Bengaluru ($N = 4,093$)	Standard DML	Pooled	-0.0011	0.0598	[-0.1184, +0.1161]	0.9847	11.96 [1.0]
	Block DML	Pooled	-0.0124	0.0583	[-0.1267, +0.1019]	0.8318	12.23 [1.0]
	Spectral OR-DML	Regime 1	+0.0916	0.1551	[-0.2123, +0.3955]	0.5546	4.22 [1.72]
	Spectral OR-DML	Regime 2	-0.1528	0.0689	[-0.2878, -0.0178]	0.0266	4.22 [1.72]
	Spectral OR-DML	Overall ATE	-0.0258	0.0790	[-0.1806, +0.1290]	0.7436	4.22 [1.72]
	Filtered OR-DML	Overall ATE	-0.0211	0.0828	[-0.1834, +0.1413]	0.7993	3.79 [2.02]
Kolkata ($N = 1,209$)	Standard DML	Pooled	+0.5969	2.1183	[-3.5550, +4.7487]	0.7781	0.005 [1.0]
	Block DML	Pooled	-0.4728	1.8772	[-4.1522, +3.2066]	0.8012	0.006 [1.0]
	Spectral OR-DML	Regime 1	+0.6789	1.4037	[-2.0724, +3.4303]	0.6286	0.0025 [1.44]
	Spectral OR-DML	Regime 2	-1.0655	2.6684	[-6.2955, +4.1645]	0.6897	0.0025 [1.44]
	Spectral OR-DML	Overall ATE	-0.0012	1.2995	[-2.5481, +2.5458]	0.9993	0.0025 [1.44]
	Filtered OR-DML	Overall ATE	+0.1777	1.2312	[-2.2355, +2.5909]	0.8853	0.0022 [1.63]

(+8.44, RMSE 8.45) and 0.0% coverage across the entire Δ_Z grid, illustrating Theorem 1. Partialling-out observed controls X_t fails because the latent Markov state S_t remains omitted.

2. **Continuous Graceful Degradation:** In accordance with Theorem 4, OR-DML exhibits continuous graceful degradation. At $\Delta_Z = 0.2$, posterior overlap is high ($\bar{H} = 0.434$), confirming Theorem 3’s non-identification. As Δ_Z increases to 1.0, 2.0, and 4.0, absolute bias shrinks by 98.1% (from 8.59 $\rightarrow$ 4.25 $\rightarrow$ 0.50 $\rightarrow$ 0.16), and coverage rises to 93.0% (with average 95% CI width narrowing from 1.323 at $\Delta_Z = 1.0$ to 0.287 at $\Delta_Z = 4.0$).
3. **Filtering vs. Smoothing:** Causal forward filtering (γ_t^{filter}) tracks retrospective smoothing (γ_t^{smooth}) closely across all separated regimes ($\Delta_Z \geq 2.0$, bias 0.2233 vs 0.2233), establishing that training-fitted forward filtering incurs virtually zero efficiency loss once latent state dynamics are informed by past physical history.

7 EMPIRICAL SENSOR NETWORK STRESS TEST

We evaluate OR-DML on our verified multi-season dataset (14,122 hourly records across Delhi, Mumbai, Bengaluru, and Kolkata, Feb 2025 – Jun 2026; < 0.7% forward-fill). Results appear in Table 2 (profiles in Appendix G, Table 3).

Conditioning Diagnostics & Physical Interpretation: Delhi Basin ($\lambda_{\min} = 111.8$): Clean separation and strong residual variation. OR-DML decomposes pooled causal effect ($+0.47 \pm 0.04$) into Regime 1 ($+0.70 \pm 0.05$, winter inversion) and Regime 2 ($+0.13 \pm 0.07$, convective clearance). **Mumbai Coastal ($\lambda_{\min} = 0.32$):** Marine breeze reversals induce severe treatment collinearity ($\lambda_{\min} = 0.32$). Standard DML

is unstable ($+2.64 \pm 4.42$), while OR-DML isolates the significant stagnation regime effect ($+3.30 \pm 1.48$, $p = 0.026$) from clearance (-0.03 ± 8.72). **Bengaluru Plateau ($\lambda_{\min} = 4.22$):** Standard DML estimates zero effect (-0.001 ± 0.060). OR-DML isolates daytime convective dispersion ($+0.092$) from nocturnal inversion (-0.153 ± 0.069 , $p = 0.027$). **Reliability Contrast:** In Bengaluru, healthy conditioning ($\lambda_{\min} = 4.22$) renders state uncertainty informative ($\text{AUC}(H) = 0.674$), whereas Mumbai’s ill-conditioning ($\lambda_{\min} = 0.32$) blinds raw entropy ($\text{AUC}(H) = 0.486$), requiring coupled index $\mathcal{D}_t$ ($\text{AUC} = 0.531\text{--}0.564$; Table 9). **Kolkata Airshed ($\lambda_{\min} = 0.0025$):** Demonstrates entropy vs. conditioning divergence: despite modest entropy ($\bar{H} = 0.083$), treatments are collinear. Unregularized DML explodes ($\text{SE} = 1.11$), while spectral regularization stabilizes variance ($\text{SE} \rightarrow 0.060$ at $\lambda = 0.10$; Appendix G).

7.1 Dynamic Exposure-Response Curves

Figure 4 (Appendix G) depicts dynamic projections $Y_{t+h} = \theta_h(S_t)T_t + g_h(X_t, S_t) + U_{t+h}$ for $h \in [0, 24]$ hours. Delhi impacts peak at $h = 1$ hour ($+0.72$, 12-hour decay), while Mumbai and Kolkata exhibit diurnal breeze reversals, confirming that static estimators miss multi-hour lag dynamics.

8 CONCLUSION

We investigated how latent-context uncertainty interacts with task geometry to govern sequential ML reliability. Reliable sequential systems must assess latent uncertainty jointly with downstream task geometry, rather than treating state confidence alone as a proxy for trust. OR-DML provides a tractable causal realization of this principle, with experiments confirming its broader validity across sequential learning tasks.

References

- Susan Athey, Julie Tibshirani, and Stefan Wager. Generalized random forests. *The Annals of Statistics*, 47(2):1148–1178, 2019.
- Jushan Bai. Panel data models with interactive fixed effects. *Econometrica*, 77(4):1229–1279, 2009.
- Zineb Banna, Florence Merlevède, and Magda Peligrad. On the matrix Bernstein inequality for α -mixing sequences. *Bernoulli*, 22(4):2073–2098, 2016.
- Peter J Bickel, Chris AJ Klaassen, Ya’acov Ritov, and Jon A Wellner. *Efficient and adaptive estimation for semiparametric models*. Johns Hopkins University Press, 1993.
- Stéphane Bonhomme and Elena Manresa. Grouped patterns of heterogeneity in panel data. *Econometrica*, 83(3):1147–1184, 2015.
- Victor Chernozhukov, Denis Chetverikov, Mert Demirer, Esther Duflo, Christian Hansen, Whitney Newey, and James Robins. Double/debiased machine learning for treatment and structural parameters. *The Econometrics Journal*, 21(1):C1–C68, 2018.
- Victor Chernozhukov, Whitney K. Newey, and Rahul Singh. Automatic debiased machine learning of causal and structural effects. *Econometrica*, 90(3):967–1027, 2022.
- Milos Ciganovic, Federico D’Amario, and Massimiliano Tancioni. Double machine learning for time series. *The Econometrics Journal*, 2026. arXiv:2603.10999.
- Fanny Clouth, Duco Veen, and Maurits Kaptein. Causal inference for latent Markov models. *Psychometrika*, 89(2):481–512, 2024.
- Francesca Dominici, Michael Greenstone, and Cass R Sunstein. Particulate matter matters. *Science*, 344(6181):257–259, 2014.
- Dylan J. Foster and Vasilis Syrgkanis. Orthogonal statistical learning. *The Annals of Statistics*, 51(3):879–908, 2023.
- James D Hamilton. A new approach to the economic analysis of nonstationary time series and the business cycle. *Econometrica*, 57(2):357–384, 1989.
- Biwei Huang, Kun Zhang, and Clark Glymour. Nonstationary causal structure learning via state-dependent conditional independence. *Journal of Machine Learning Research*, 24(120):1–48, 2023.
- Ildar A Ibragimov. Some limit theorems for stationary processes. *Theory of Probability & Its Applications*, 7(4):349–382, 1962.
- John G Kemeny and J Laurie Snell. *Finite Markov Chains*. Springer-Verlag, 1976.
- Marcos López de Prado. *Advances in Financial Machine Learning*. John Wiley & Sons, 2018.
- Xinkun Nie and Stefan Wager. Quasi-oracle estimation of heterogeneous treatment effects. *Biometrika*, 108(2):299–319, 2021.
- Anonymous Authors. Weak-overlap-aware deconfounding representations in causal inference. In *Proceedings of the 29th International Conference on Artificial Intelligence and Statistics (AISTATS)*, 2026.
- Alexandre B Tsybakov. *Introduction to Nonparametric Estimation*. Springer Series in Statistics. Springer, New York, NY, 2009.

AI Use Statement

In this work, we used generative AI tools (large language models) for the following **tasks requiring disclosure**: helping develop the theoretical model and conceptual framework of OR-DML; formulating mathematical claims (Theorems 1–5, Proposition 6); providing critical ingredients for proving mathematical claims and assisting in the writing of proofs (Appendices A–G); implementing methods and software code in Python (`src/or_dml.py`, `src/synthetic_dgp_benchmark.py`, `src/empirical_evaluation.py`, `src/data_pipeline_clean.py`); cleaning and reformatting atmospheric datasets; and supporting literature synthesis in the Related Work section. We have not used generative AI tools to generate synthetic datasets autonomously or to interpret empirical results without author verification. Additionally, we used generative AI tools for the following **tasks with recommended disclosure**: drafting parts of the paper, editing the manuscript to improve readability, suggesting section structures, and formatting bibliographic references.

We have reviewed all AI-assisted work as follows: all mathematical proofs were verified line-by-line from first principles by the authors; all Python code was executed, tested, and verified against independent simulation baselines with known ground-truth parameters; and all empirical numbers in tables and figures were cross-verified directly against the generated `.csv` artifacts in `reports/`. We take full responsibility for the final content of this work, including all text, claims, and artifacts produced with the aid of generative AI.

CHECKLIST

1. For all models and algorithms presented, check if you include:
 - (a) A clear description of the algorithm, mapping, or model. **[Yes]** See Section 3 and Algorithm 1.
 - (b) An analysis of the properties and complexity. **[Yes]** See Theorems 1–5 and Proposition 6.
 - (c) (Optional) Anonymized source code, with specification of all dependencies, including external libraries. **[Yes]** Code included in the supplementary material.
2. For any theoretical claim, check if you include:
 - (a) Statements of the full set of assumptions of all theoretical results. **[Yes]** Assumptions 1–4 are stated in Section 2 and Section 4.
 - (b) Complete proofs of all theoretical results. **[Yes]** Full proofs in Appendices A–F.
 - (c) Clear explanations of any assumptions. **[Yes]** Detailed physical and statistical justifications accompany each assumption.
3. For all figures and tables that present empirical results, check if you include:
 - (a) The code, data, and instructions needed to reproduce the main experimental results. **[Yes]** Scripts in `src/` and reproduction instructions in `README.md`.
 - (b) All the training details (e.g., data splits, hyperparameters, how they were selected). **[Yes]** Section 3 and Section 5 describe block splitting, Bartlett embargo buffer τ^* , and HMM configuration.
 - (c) A clear definition of the specific measure or statistics and error bars (e.g., confidence intervals, p-values). **[Yes]** Table 1 reports empirical bias, RMSE, and 95% coverage over 500 replications; Table 2 reports HAC standard errors and p-values.
 - (d) A description of the computing infrastructure used. **[Yes]** Experiments were run on a standard 16-core workstation; details in the supplementary material.
4. If you are using existing assets (e.g., code, data, models) or curating/releasing new assets:
 - (a) If your work uses existing assets, did you cite the creators? **[Yes]** CPCB ground stations and Open-Meteo reanalysis are cited.
 - (b) Did you mention the license of the assets? **[Yes]** Open public government data terms noted.
 - (c) Did you include any new assets either in the supplemental material or as a URL? **[Yes]** Cleaned longitudinal benchmark code and data bundled in supplementary archive.
 - (d) Did you discuss whether and how consent was obtained from people whose data you’re using/curating? **[Not Applicable]** Aggregate ambient physical air measurements only.
 - (e) Did you discuss whether the data you are using/curating contains personally identifiable information or offensive content? **[Not Applicable]** Contains only environmental measurements.
5. If you used crowdsourcing or conducted research with human subjects:
 - (a) Did you include the full text of instructions given to participants and screenshots? **[Not Applicable]**
 - (b) Did you describe any potential participant risks, with links to Institutional Review Board (IRB) approvals if applicable? **[Not Applicable]**
 - (c) Did you include the estimated hourly wage paid to participants and the total amount spent on participant compensation? **[Not Applicable]**

MATHEMATICAL APPENDIX: COMPLETE PROOFS AND DERIVATIONS

A PROOF OF THEOREM 1: NON-IDENTIFICATION UNDER UNCONSTRAINED OVERLAP

To prove substantive causal non-identification, we constructively exhibit two data-generating distributions, P_1 and P_2 , that are structurally well-formed under the SCM (1)–(5) and satisfy all Assumptions 1–3, yet possess strictly distinct, non-permutation causal effect vectors $\theta^{(1)} \neq \theta^{(2)}$ while inducing identically equal joint distributions over all observable variables:

$$P_1(Y_{1:N}, T_{1:N}, X_{1:N}, Z_{1:N}) \equiv P_2(Y_{1:N}, T_{1:N}, X_{1:N}, Z_{1:N}) \quad \forall N \geq 1. \quad (30)$$

Construction of Substantive Observational Equivalence: Consider a two-state Markov chain ($K = 2$) with symmetric transition probability $P_{01} = P_{10} = p \in (0, 0.5)$, yielding stationary distribution $\pi_0^* = \pi_1^* = 0.5$. Let auxiliary meteorological proxies $Z_t \in \mathbb{R}$ satisfy complete overlap: $Z_t | S_t = 0 \sim \mathcal{N}(0, 1)$ and $Z_t | S_t = 1 \sim \mathcal{N}(0, 1)$ identically, so $Z_t \perp S_t$. Let continuous observed confounders $X_t \in \mathbb{R}$ have marginal density $f_X(x)$ independent of S_t . Let discrete binary treatment $T_t \in \{-1, +1\}$ have equal probability $P(T_t = 1) = P(T_t = -1) = 0.5$, distributed completely independently of (S_t, X_t, Z_t) , which explicitly guarantees $T_t \perp S_t | X_t$ and $T_t \perp (S_t, Z_t, X_t)$. Thus, Assumption 1 (Conditional Unconfoundedness) holds trivially with constant propensity score $e(X_t, S_t) \equiv 0.5 \in [\eta, 1 - \eta]$ satisfying full overlap everywhere. Baseline treatment assignment satisfies $m(X_t, S_t) \equiv 0$ and $V_t = T_t$ with $\mathbb{E}[V_t | X_t, S_t] = 0$.

Now specify two distinct structural outcome models:

1. **Model P_1 (Homogeneous Causal Effect):** Set $\theta^{(1)} = (\theta_0^{(1)}, \theta_1^{(1)})^\top = (1, 1)^\top$ (so the treatment effect is identical across regimes, $\theta_0^{(1)} - \theta_1^{(1)} = 0$), baseline function $g^{(1)}(X_t, S_t) \equiv 0$, and structural outcome innovation $U_t^{(1)} \in \{-1, +1\}$ with $P(U_t^{(1)} = 1) = P(U_t^{(1)} = -1) = 0.5$, distributed independently of $(T_t, X_t, S_t, Z_{1:N})$. Then $\mathbb{E}[U_t^{(1)} | X_t, S_t, T_t] = 0$. Under Model P_1 , the outcome is $Y_t = 1 \cdot T_t + U_t^{(1)}$. The conditional distribution of Y_t given T_t evaluates to:

$$P_1(Y_t = 2 | T_t = 1) = 0.5, \quad P_1(Y_t = 0 | T_t = 1) = 0.5, \quad (31)$$

$$P_1(Y_t = -2 | T_t = -1) = 0.5, \quad P_1(Y_t = 0 | T_t = -1) = 0.5. \quad (32)$$

2. **Model P_2 (Heterogeneous Causal Effect):** Set $\theta^{(2)} = (\theta_0^{(2)}, \theta_1^{(2)})^\top = (2, 0)^\top$ (so the treatment effect is strongly heterogeneous, $\theta_0^{(2)} - \theta_1^{(2)} = 2$), baseline function $g^{(2)}(X_t, S_t) \equiv 0$, and structural error $U_t^{(2)} \equiv 0$ almost surely (trivially satisfying $\mathbb{E}[U_t^{(2)} | X_t, S_t, T_t] = 0$). Under Model P_2 , the outcome is $Y_t = \theta(S_t)T_t$:

- In Regime 0 ($S_t = 0$): $Y_t = 2T_t$, so $Y_t = 2$ if $T_t = 1$ and $Y_t = -2$ if $T_t = -1$.
- In Regime 1 ($S_t = 1$): $Y_t = 0 \cdot T_t = 0$ for all $T_t \in \{-1, +1\}$.

Marginalizing over the unobserved state S_t with prior $\pi_0^* = \pi_1^* = 0.5$:

$$P_2(Y_t = 2 | T_t = 1) = \pi_0^*(1) + \pi_1^*(0) = 0.5, \quad P_2(Y_t = 0 | T_t = 1) = \pi_0^*(0) + \pi_1^*(1) = 0.5, \quad (33)$$

$$P_2(Y_t = -2 | T_t = -1) = \pi_0^*(1) + \pi_1^*(0) = 0.5, \quad P_2(Y_t = 0 | T_t = -1) = \pi_0^*(0) + \pi_1^*(1) = 0.5. \quad (34)$$

Comparing the conditional distributions:

$$P_1(Y_t | T_t, X_t, Z_t) \equiv P_2(Y_t | T_t, X_t, Z_t) \quad \forall (y, t, x, z) \in \text{supp}(Y, T, X, Z). \quad (35)$$

Because the marginal distributions of (T_t, X_t, Z_t) and the temporal Markov transition laws are identical in both models, the full joint likelihoods over all observables are strictly identical for any sample size N :

$$P_1(Y_{1:N}, T_{1:N}, X_{1:N}, Z_{1:N}) \equiv P_2(Y_{1:N}, T_{1:N}, X_{1:N}, Z_{1:N}). \quad (36)$$

Yet the structural causal effects satisfy:

$$\theta^{(1)} = \begin{pmatrix} 1 \\ 1 \end{pmatrix} \neq \begin{pmatrix} 2 \\ 0 \end{pmatrix} = \theta^{(2)}. \quad (37)$$

Crucially, $\theta^{(2)}$ is *not* a permutation of $\theta^{(1)}$. In Model P_1 , the inter-regime causal contrast is $\theta_0^{(1)} - \theta_1^{(1)} = 0$, whereas in Model P_2 , $\theta_0^{(2)} - \theta_1^{(2)} = 2$. No estimation procedure conditioning solely on observable data (Y, T, X, Z)

can distinguish whether the data-generating mechanism is homogeneous (P_1) or heterogeneous (P_2) without informative proxies Z_t that bound the proxy error ε_γ . This establishes substantive non-identification. ■

B PROOF OF THEOREM 2: GRACEFUL DEGRADATION AND THE DIFFICULTY FRONTIER

Let $\mathbf{H}_t = \mathbf{e}_{S_t} \in \{0, 1\}^K$ denote the unobserved true regime indicator vector. Under the true SCM (1)–(5), the oracle regime nuisance functions are:

$$m_k^*(X_t) \equiv \mathbb{E}[T_t | X_t, S_t = k], \quad \mu_k^*(X_t) \equiv \mathbb{E}[Y_t | X_t, S_t = k] = \theta_k^* m_k^*(X_t) + g(X_t, k). \quad (38)$$

The oracle centered residuals are $\tilde{T}_{tk}^* \equiv T_t - m_k^*(X_t)$ and $\tilde{Y}_{tk}^* \equiv Y_t - \mu_k^*(X_t) = \theta_k^* \tilde{T}_{tk}^* + U_t$. Under imperfect regime overlap, the empirical cross-fitted learners $\hat{m}_k$ and $\hat{\mu}_k$ minimize the posterior-weighted empirical risk $\sum_{t \in \mathcal{T}_b} \gamma_{tk} (T_t - m(X_t))^2$. Their exact population targets are the proxy-weighted conditional expectations:

$$m_k^\gamma(X_t) \equiv \frac{\mathbb{E}[\gamma_{tk} T_t | X_t]}{\mathbb{E}[\gamma_{tk} | X_t]}, \quad \mu_k^\gamma(X_t) \equiv \frac{\mathbb{E}[\gamma_{tk} Y_t | X_t]}{\mathbb{E}[\gamma_{tk} | X_t]}. \quad (39)$$

The empirical coupled Gram matrix is $\hat{J} = \frac{1}{N} \sum_t \gamma_{tj} \gamma_{tk} \tilde{T}_{tj} \tilde{T}_{tk}$ and the empirical score vector is $\hat{S}_k = \frac{1}{N} \sum_t \gamma_{tk} \tilde{T}_{tk} \tilde{Y}_{tk}$. The corresponding population objects are $\mathbf{J} \in \mathbb{R}^{K \times K}$ with $J_{jk} \equiv \mathbb{E}[\gamma_{tj} \gamma_{tk} \tilde{T}_{tj}^* \tilde{T}_{tk}^*]$ and $\mathbf{S} \in \mathbb{R}^K$ with $S_k \equiv \mathbb{E}[\gamma_{tk} \tilde{T}_{tk}^* \tilde{Y}_{tk}^*]$.

Neyman Orthogonality vs. Proxy Sensitivity: Consider the state-weighted score for regime k as a functional of candidate nuisances (m_k, μ_k) :

$$\psi_k(W_t; \theta_k, m_k, \mu_k) = \gamma_{tk} (T_t - m_k(X_t)) (Y_t - \mu_k(X_t)). \quad (40)$$

Evaluating the Gateaux derivative with respect to m_k in direction Δm at the oracle target (m_k^*, μ_k^*) :

$$\partial_m \mathbb{E}[\psi_k(W_t; \theta_k^*, m_k^*, \mu_k^*)][\Delta m] = -\mathbb{E}[\gamma_{tk} \Delta m(X_t) (Y_t - \mu_k^*(X_t))] = -\mathbb{E}[\gamma_{tk} \Delta m(X_t) U_t] = 0, \quad (41)$$

since $\mathbb{E}[U_t | X_t, S_t, Z_{1:N}] = 0$ by Assumption 1. Similarly, the Gateaux derivative with respect to μ_k in direction $\Delta \mu$ is:

$$\partial_\mu \mathbb{E}[\psi_k(W_t; \theta_k^*, m_k^*, \mu_k^*)][\Delta \mu] = -\mathbb{E}[\gamma_{tk} (T_t - m_k^*(X_t)) \Delta \mu(X_t)] = -\mathbb{E}[\gamma_{tk} \mathbb{E}[\tilde{T}_{tk}^* | X_t, Z_{1:N}] \Delta \mu(X_t)] = 0. \quad (42)$$

Furthermore, evaluating at the proxy population target $(m_k^\gamma, \mu_k^\gamma)$, $\mathbb{E}[\gamma_{tk} (T_t - m_k^\gamma(X_t)) | X_t] = 0$ holds identically by construction. Therefore, Neyman orthogonality eliminates first-order sensitivity to errors in estimating (m_k, μ_k) , ensuring that nuisance estimation errors contribute only a second-order term $\mathcal{O}_P(\|\hat{m}_k - m_k^\gamma\|_{L_2} \|\hat{\mu}_k - \mu_k^\gamma\|_{L_2}) = \mathcal{O}_P(N^{-1/2})$ under Assumption 4. However, the score ψ_k depends linearly on the posterior state weight γ_{tk} . Because γ_{tk} proxies the unobserved indicator $H_{tk} = \mathbb{I}(S_t = k)$, Neyman orthogonality with respect to nuisance regressions does *not* eliminate proxy estimation error $\mathbf{H}_t - \gamma_t$, which generates a first-order bias.

Score Expansion and Bias Bound (Fix A): Expanding $Y_t = \sum_{j=1}^K H_{tj} [\theta_j^* T_t + g(X_t, j)] + U_t$:

$$\tilde{Y}_{tk}^* = Y_t - \mu_k^*(X_t) = \sum_{j=1}^K H_{tj} \theta_j^* \tilde{T}_{tj}^* + \sum_{j=1}^K H_{tj} [\theta_j^* (m_j^*(X_t) - m_k^*(X_t)) + g(X_t, j) - g(X_t, k)] + U_t. \quad (43)$$

Substituting into the population score $S_k = \mathbb{E}[\gamma_{tk} \tilde{T}_{tk}^* \tilde{Y}_{tk}^*]$:

$$\begin{aligned} S_k &= \sum_{j=1}^K \theta_j^* \mathbb{E}[\gamma_{tk} H_{tj} \tilde{T}_{tk}^* \tilde{T}_{tj}^*] + \mathbb{E} \left[\gamma_{tk} \tilde{T}_{tk}^* \sum_{j=1}^K H_{tj} (\theta_j^* (m_j^*(X_t) - m_k^*(X_t)) + g(X_t, j) - g(X_t, k)) \right] \\ &= \sum_{j=1}^K \theta_j^* \mathbb{E}[\gamma_{tk} \gamma_{tj} \tilde{T}_{tk}^* \tilde{T}_{tj}^*] + \sum_{j=1}^K \theta_j^* \mathbb{E}[\gamma_{tk} (H_{tj} - \gamma_{tj}) \tilde{T}_{tk}^* \tilde{T}_{tj}^*] + \mathbb{E}[\gamma_{tk} \tilde{T}_{tk}^* \Delta g_{tk}] \\ &= (\mathbf{J} \boldsymbol{\theta}^*)_k + b_{\gamma, k}, \end{aligned} \quad (44)$$

where $b_{\gamma, k} \equiv \sum_{j=1}^K \theta_j^* \mathbb{E}[\gamma_{tk} (H_{tj} - \gamma_{tj}) \tilde{T}_{tk}^* \tilde{T}_{tj}^*] + \mathbb{E}[\gamma_{tk} \tilde{T}_{tk}^* \sum_{j=1}^K (H_{tj} - \gamma_{tj}) (\theta_j^* \Delta m_{jk}(X_t) + \Delta g_{jk}(X_t))]$. Under the bounded residual support condition of Theorem 4 ($\sup_t |\tilde{T}_{tk}^*| \leq M_T$ a.s., $\sup_{j,k,x} |\Delta m_{jk}(x)| \leq L_m$, and $\sup_{j,k,x} |\Delta g_{jk}(x)| \leq L_g$), we bound the expectation directly without requiring Hölder's inequality on unbounded

fourth moments:

$$\begin{aligned}
 |b_{\gamma,k}| &\leq \sum_{j=1}^K |\theta_j^*| \mathbb{E} \left[\gamma_{tk} |\tilde{T}_{tk}^*| |\tilde{T}_{tj}^*| |H_{tj} - \gamma_{tj}| \right] + \sum_{j=1}^K \mathbb{E} \left[\gamma_{tk} |\tilde{T}_{tk}^*| |H_{tj} - \gamma_{tj}| (|\theta_j^*| L_m + L_g) \right] \\
 &\leq M_\theta M_T^2 \mathbb{E} \left[\sum_{j=1}^K |H_{tj} - \gamma_{tj}| \right] + M_T (M_\theta L_m + L_g) \mathbb{E} \left[\sum_{j=1}^K |H_{tj} - \gamma_{tj}| \right] \\
 &= (M_\theta M_T^2 + M_\theta M_T L_m + L_g M_T) \mathbb{E} \|\mathbf{H}_t - \boldsymbol{\gamma}_t\|_1 = C \cdot \varepsilon_\gamma,
 \end{aligned} \tag{45}$$

where $C = L_g M_T + M_\theta M_T^2 + M_\theta M_T L_m < \infty$. Since $\lambda_{\min}(\mathbf{J}) > 0$, the unregularized estimand satisfies:

$$\|\boldsymbol{\theta}_\gamma - \boldsymbol{\theta}^*\|_2 = \|\mathbf{J}^{-1} \mathbf{b}_\gamma\|_2 \leq \|\mathbf{J}^{-1}\|_{\text{op}} \|\mathbf{b}_\gamma\|_2 \leq \frac{C \cdot \varepsilon_\gamma}{\lambda_{\min}(\mathbf{J})}. \tag{46}$$

Combining with purged cross-fitting nuisance estimation error of order $\mathcal{O}_P(N^{-1/2})$ yields $\|\hat{\boldsymbol{\theta}}_\gamma - \boldsymbol{\theta}^*\|_2 \leq \frac{C \varepsilon_\gamma}{\lambda_{\min}(\mathbf{J})} + \mathcal{O}_P(N^{-1/2})$, completing the proof of Theorem 4. ■

B.1 The Latent-Confounding Difficulty Frontier $\mathcal{D}_{\text{causal}}$

Theorem 4 demonstrates that the upper bound on structural causal estimation error decomposes into:

$$\|\hat{\boldsymbol{\theta}}_\gamma - \boldsymbol{\theta}^*\|_2 \leq C \cdot \mathcal{D}_{\text{causal}} + \mathcal{O}_P(N^{-1/2}), \quad \text{where} \quad \mathcal{D}_{\text{causal}} \equiv \frac{\varepsilon_\gamma}{\lambda_{\min}(\mathbf{J})}. \tag{47}$$

We highlight several structural properties of the difficulty index $\mathcal{D}_{\text{causal}}$:

1. **Dimensionless Product:** Both population proxy error ε_γ and Gram conditioning $\lambda_{\min}(\mathbf{J})$ interact multiplicatively in governing downstream estimation risk. When either proxy recovery is poor ($\varepsilon_\gamma \rightarrow 1$) or overlap is weak ($\lambda_{\min} \rightarrow 0$), estimation risk escalates sharply.
2. **Asymmetric Sensitivity:** In our factorial simulations (Table 6), reducing $\lambda_{\min}$ by an order of magnitude produces an equivalent variance and bias inflation to a tenfold increase in state proxy error ε_γ .

B.2 Proof of Proposition 4: Surrogate Entropy-Difficulty Bridge

Let $\gamma_t \equiv P(S_t \in \cdot \mid \mathcal{F}_t) \in \Delta^K$ be the Bayesian posterior state distribution with respect to filtration $\mathcal{F}_t \equiv \sigma(Z_{1:t})$. Under proper Bayesian calibration, conditional on $\mathcal{F}_t$, the true latent indicator $\mathbf{H}_t \equiv \mathbf{e}_{S_t}$ is distributed as a categorical random variable taking value $\mathbf{e}_k$ with probability γ_{tk} .

Step 1: Expected L_1 Error as Gini Impurity. Evaluating the conditional expectation of the L_1 error:

$$\begin{aligned}
 \mathbb{E} [\|\gamma_t - \mathbf{H}_t\|_1 \mid \mathcal{F}_t] &= \sum_{k=1}^K \gamma_{tk} \|\gamma_t - \mathbf{e}_k\|_1 = \sum_{k=1}^K \gamma_{tk} \left((1 - \gamma_{tk}) + \sum_{j \neq k} \gamma_{tj} \right) \\
 &= \sum_{k=1}^K \gamma_{tk} \cdot 2(1 - \gamma_{tk}) = 2 \left(1 - \sum_{k=1}^K \gamma_{tk}^2 \right) = 2 \text{Gini}(\gamma_t).
 \end{aligned} \tag{48}$$

Step 2: Entropy Bound on Gini Impurity. By the standard concave inequality $\ln(x) \leq x - 1$, for any $x > 0$:

$$H(\gamma_t) = - \sum_{k=1}^K \gamma_{tk} \ln \gamma_{tk} = \sum_{k=1}^K \gamma_{tk} \ln \left(\frac{1}{\gamma_{tk}} \right) \geq \sum_{k=1}^K \gamma_{tk} (1 - \gamma_{tk}) = \text{Gini}(\gamma_t). \tag{49}$$

Furthermore, for any binary regime partition ($K = 2$) with $\gamma_{t1} = p$ and $\gamma_{t2} = 1 - p$, $2 \text{Gini}(\gamma_t) = 4p(1 - p)$. Comparing to binary Shannon entropy $H(p) = -p \ln p - (1 - p) \ln(1 - p)$, the ratio $\frac{4p(1-p)}{H(p)}$ achieves its maximum at $p = 1/2$, where $\frac{4(1/4)}{\ln 2} = \frac{1}{\ln 2} \approx 1.4427$. Hence for all $p \in [0, 1]$:

$$2 \text{Gini}(\gamma_t) \leq C_K H(\gamma_t), \quad \text{where} \quad C_K \leq \frac{1}{\ln 2} \approx 1.4427. \tag{50}$$

Step 3: Population and Observable Bridge. Taking the population expectation of Step 1 and applying Step 2:

$$\varepsilon_\gamma \equiv \mathbb{E} [\mathbb{E} [\|\gamma_t - \mathbf{H}_t\|_1 \mid \mathcal{F}_t]] = 2 \mathbb{E} [\text{Gini}(\gamma_t)] \leq C_K \mathbb{E} [H(\gamma_t)]. \tag{51}$$

Dividing by the global minimal singular value $\lambda_{\min}(\mathbf{J})$:

$$\mathcal{D}_{\text{causal}} \equiv \frac{\varepsilon_\gamma}{\lambda_{\min}(\mathbf{J})} \leq C_K \mathbb{E} \left[\frac{H(\gamma_t)}{\lambda_{\min}(\mathbf{J})} \right] = C_K \mathbb{E}[\mathcal{D}_{\text{global}}^H(t)]. \quad (52)$$

Furthermore, under Assumption 5 (Two-Sided Spectral Comparability), the rolling Gram matrix satisfies $\lambda_{\min}(\mathbf{J}_t) \leq \bar{c}_J \lambda_{\min}(\mathbf{J})$ almost surely for all evaluation windows $t \geq w$, which implies $\frac{1}{\lambda_{\min}(\mathbf{J})} \leq \frac{\bar{c}_J}{\lambda_{\min}(\mathbf{J}_t)}$. Pointwise substitution yields:

$$\mathcal{D}_{\text{causal}} \leq C_K \bar{c}_J \mathbb{E} \left[\frac{H(\gamma_t)}{\lambda_{\min}(\mathbf{J}_t)} \right] = C_K \bar{c}_J \mathbb{E}[\mathcal{D}_{\text{operational}}(t)], \quad (53)$$

and $\mathbb{E}[\mathcal{D}_{\text{operational}}(t)] \leq \bar{c}_J^{-1} \mathbb{E}[\mathcal{D}_{\text{global}}^H(t)]$. In finite samples, the ergodic theorem and central limit theorem for stationary α -mixing chains ensure that $\hat{\varepsilon}_{\gamma,N} \equiv \frac{1}{N} \sum_{t=1}^N \|\gamma_t - \mathbf{H}_t\|_1 = \varepsilon_\gamma + \mathcal{O}_P(N^{-1/2})$, so the sample surrogate $\hat{\mathcal{D}}_{\text{causal},N} \equiv \hat{\varepsilon}_{\gamma,N} / \lambda_{\min}(\mathbf{J}) = \mathcal{D}_{\text{causal}} + \mathcal{O}_P(N^{-1/2})$. We emphasize that these bounds establish $\mathcal{D}_{\text{operational}}(t)$ as an observable surrogate upper bound for $\mathcal{D}_{\text{causal}}$, not a two-sided equivalence. Across our multi-season ground-sensor streams, the empirical ratio $\lambda_{\min}(\mathbf{J}_t) / \lambda_{\min}(\mathbf{J})$ has 95th percentile in $[1.06, 1.72]$ and maximum ≤ 7.35 , empirically confirming that $\bar{c}_J$ is finite and bounded. ■

B.3 A Priori Proxy Diagnostics and Separation Screening

Before fitting OR-DML or downstream prediction heads, practitioners should assess proxy informativeness on auxiliary measurements $Z_t \in \mathbb{R}^q$. Let $\hat{\boldsymbol{\mu}}_{Z,k} \equiv \frac{1}{N_k} \sum_{t:\hat{S}_t=k} Z_t$ and $\hat{\boldsymbol{\Sigma}}_{Z,k}$ denote empirical cluster moments. We define the standardized separation index:

$$\Delta_Z \equiv \min_{j \neq k} \frac{\|\hat{\boldsymbol{\mu}}_{Z,j} - \hat{\boldsymbol{\mu}}_{Z,k}\|_2}{\sqrt{\frac{1}{2}(\text{Tr}(\hat{\boldsymbol{\Sigma}}_{Z,j}) + \text{Tr}(\hat{\boldsymbol{\Sigma}}_{Z,k}))/q}}. \quad (54)$$

When $\Delta_Z < 0.5$ (or silhouette score < 0.15), the overlap between regime emission densities $P(Z_t | S_t = j)$ and $P(Z_t | S_t = k)$ is severe, driving posterior entropy $\bar{H}(\gamma) \rightarrow \ln K$. By Theorem 3, point identification fails identically in the limit $\Delta_Z \rightarrow 0$. Deployed pipelines should enforce an automated screening gate: if $\Delta_Z < 0.5$ or mean entropy exceeds $0.80 \ln K$, the system must flag non-identifiability and refuse to emit point causal effects without external structural constraints.

C PROOF OF THEOREM 3: SPECTRAL REGULARIZATION BIAS-VARIANCE FRONTIER

Let $\hat{\boldsymbol{\theta}}_\lambda = (\hat{\mathbf{J}} + \lambda \mathbf{I})^{-1} \hat{\mathbf{S}}$, and define the population regularized estimand $\boldsymbol{\theta}_\lambda \equiv (\mathbf{J} + \lambda \mathbf{I})^{-1} \mathbf{S}$. We decompose the estimation error around the regularized population estimand $\boldsymbol{\theta}_\lambda$ via the exact algebraic identity:

$$\mathbb{E} \|\hat{\boldsymbol{\theta}}_\lambda - \boldsymbol{\theta}^*\|_2^2 = \|\boldsymbol{\theta}_\lambda - \boldsymbol{\theta}^*\|_2^2 + \mathbb{E} \|\hat{\boldsymbol{\theta}}_\lambda - \boldsymbol{\theta}_\lambda\|_2^2 + 2\langle \mathbb{E}[\hat{\boldsymbol{\theta}}_\lambda - \boldsymbol{\theta}_\lambda], \boldsymbol{\theta}_\lambda - \boldsymbol{\theta}^* \rangle, \quad (55)$$

which accounts explicitly for the cross term between stochastic estimation error and deterministic regularization bias. For the deterministic bias, substituting $\mathbf{S} = \mathbf{J}\boldsymbol{\theta}^* + \mathbf{b}_\gamma$:

$$\begin{aligned} \boldsymbol{\theta}_\lambda - \boldsymbol{\theta}^* &= (\mathbf{J} + \lambda \mathbf{I})^{-1} (\mathbf{S} - (\mathbf{J} + \lambda \mathbf{I})\boldsymbol{\theta}^*) \\ &= (\mathbf{J} + \lambda \mathbf{I})^{-1} (\mathbf{S} - \mathbf{J}\boldsymbol{\theta}^* - \lambda \boldsymbol{\theta}^*) \\ &= (\mathbf{J} + \lambda \mathbf{I})^{-1} (\mathbf{b}_\gamma - \lambda \boldsymbol{\theta}^*), \end{aligned} \quad (56)$$

which establishes the exact deterministic identity (25). Since $\mathbf{J} \succeq 0$ with minimal eigenvalue $\lambda_{\min}(\mathbf{J}) \geq 0$, the spectral operator norm is $\|(\mathbf{J} + \lambda \mathbf{I})^{-1}\|_{\text{op}} = \frac{1}{\lambda_{\min}(\mathbf{J}) + \lambda}$. Applying the triangle inequality:

$$\|\boldsymbol{\theta}_\lambda - \boldsymbol{\theta}^*\|_2 \leq \|(\mathbf{J} + \lambda \mathbf{I})^{-1}\|_{\text{op}} (\|\mathbf{b}_\gamma\|_2 + \lambda \|\boldsymbol{\theta}^*\|_2) \leq \frac{C\varepsilon_\gamma + \lambda M_\theta}{\lambda_{\min}(\mathbf{J}) + \lambda}. \quad (57)$$

For the stochastic variance, expanding $\hat{\boldsymbol{\theta}}_\lambda - \boldsymbol{\theta}_\lambda$:

$$\hat{\boldsymbol{\theta}}_\lambda - \boldsymbol{\theta}_\lambda = (\mathbf{J} + \lambda \mathbf{I})^{-1} \left(\hat{\mathbf{S}} - \mathbf{S} - (\hat{\mathbf{J}} - \mathbf{J})\boldsymbol{\theta}_\lambda \right) + o_p(N^{-1/2}). \quad (58)$$

By Theorem 7, the score empirical process satisfies $\text{Var} \left(\frac{1}{\sqrt{N}} \sum_{t=1}^N \boldsymbol{\Psi}_t \right) \rightarrow \boldsymbol{\Omega}$. Thus, under the long-run score covariance approximation, the stochastic variance contribution admits the leading-order expansion:

$$\mathbb{E} \|\hat{\boldsymbol{\theta}}_\lambda - \boldsymbol{\theta}_\lambda\|_2^2 = \frac{1}{N} \text{Tr} \left((\mathbf{J} + \lambda \mathbf{I})^{-1} \boldsymbol{\Omega} (\mathbf{J} + \lambda \mathbf{I})^{-1} \right) + o(N^{-1}) \leq \frac{\text{Tr}(\boldsymbol{\Omega})}{N(\lambda_{\min}(\mathbf{J}) + \lambda)^2} + o(N^{-1}). \quad (59)$$

To bound the cross term $2\langle \mathbb{E}[\hat{\theta}_\lambda - \theta_\lambda], \theta_\lambda - \theta^* \rangle$ in (55): by Neyman orthogonality of the score function ψ and purged block cross-fitting with Bartlett embargo buffer $\tau^* \geq C \log N$, the finite-sample expectation of the stochastic error satisfies $\|\mathbb{E}[\hat{\theta}_\lambda - \theta_\lambda]\|_2 \leq C_1 N^{-1}$ (Theorem 7). Furthermore, the deterministic regularized bias satisfies $\|\theta_\lambda - \theta^*\|_2 \leq \frac{C\varepsilon_\gamma + \lambda M_\theta}{\lambda_{\min}(\mathbf{J}) + \lambda}$. By Cauchy-Schwarz, the cross term is bounded by:

$$\left| 2\langle \mathbb{E}[\hat{\theta}_\lambda - \theta_\lambda], \theta_\lambda - \theta^* \rangle \right| \leq 2\|\mathbb{E}[\hat{\theta}_\lambda - \theta_\lambda]\|_2 \|\theta_\lambda - \theta^*\|_2 = \mathcal{O}(N^{-1}) \quad (60)$$

in general (since $\|\theta_\lambda - \theta^*\|_2 = \mathcal{O}(1)$ without vanishing-bias rate restrictions), which sharpens to $\mathcal{O}(N^{-3/2})$ under the local parameter regime $\varepsilon_\gamma + \lambda = \mathcal{O}(N^{-1/2})$. Combining the deterministic squared bias bound with the asymptotic long-run score covariance variance bound establishes the leading-order risk bound:

$$\mathbb{E}\|\hat{\theta}_\lambda - \theta^*\|_2^2 \leq \left(\frac{C\varepsilon_\gamma + \lambda M_\theta}{\lambda_{\min}(\mathbf{J}) + \lambda} \right)^2 + \frac{\text{Tr}(\mathbf{\Omega})}{N(\lambda_{\min}(\mathbf{J}) + \lambda)^2} + \mathcal{O}(N^{-1}). \quad (61)$$

Risk Bound Minimizer vs. Empirical Grid Optimum: Differentiating the leading-order risk upper bound $R_{\text{bound}}(\lambda) \equiv \left(\frac{C\varepsilon_\gamma + \lambda M_\theta}{\lambda_{\min}(\mathbf{J}) + \lambda} \right)^2 + \frac{\text{Tr}(\mathbf{\Omega})}{N(\lambda_{\min}(\mathbf{J}) + \lambda)^2}$ with respect to $\lambda \geq 0$, the bound-optimal shrinkage parameter $\lambda_{\text{bound}}^* \equiv \arg \min_{\lambda \geq 0} R_{\text{bound}}(\lambda)$ satisfies:

$$\frac{\partial}{\partial \lambda} \left[\left(\frac{C\varepsilon_\gamma + \lambda M_\theta}{\lambda_{\min}(\mathbf{J}) + \lambda} \right)^2 + \frac{\text{Tr}(\mathbf{\Omega})}{N(\lambda_{\min}(\mathbf{J}) + \lambda)^2} \right] = 0. \quad (62)$$

When the overlap Gram matrix is ill-conditioned ($\lambda_{\min}(\mathbf{J}) \approx 0$), the variance term $\frac{\text{Tr}(\mathbf{\Omega})}{N(\lambda_{\min} + \lambda)^2}$ dominates for small λ , while the deterministic shrinkage bias $(\frac{\lambda M_\theta}{\lambda_{\min} + \lambda})^2$ dominates for large λ . The interior minimizer $\lambda_{\text{bound}}^* > 0$ balances variance reduction against shrinkage bias in the upper bound. We distinguish λ_{bound}^* and the empirical surrogate minimizer $\hat{\lambda}_{\text{emp}}^* = \arg \min_{\lambda} \hat{R}(\lambda)$ from the true infeasible risk minimizer $\lambda_{\text{true}}^* \equiv \arg \min_{\lambda} \mathbb{E}\|\hat{\theta}_\lambda - \theta^*\|_2^2$. In our empirical simulation benchmark across 500 Monte Carlo replications (`reports/or_dml_regularization_frontier.csv`), the discrete grid minimum occurs at $\hat{\lambda}_{\text{empirical}}^* \approx 0.20$ (MSE = 0.0321, representing a $25.9\times$ empirical risk reduction in this benchmark DGP relative to the unregularized estimator $\lambda = 0.0$ where MSE = 0.8312), while $\lambda = 0.10$ achieves MSE = 0.0362 ($23.0\times$ empirical reduction) with peak coverage of 85.0%. We emphasize that Theorem 6 provides the functional upper-bound characterization of the bias-variance trade-off; the exact numerical reduction factors ($23.0\times$ and $25.9\times$) are empirical simulation properties of the benchmark DGP rather than universal theoretical constants. ■

C.1 Derivation of Adaptive Spectral Regularization Parameter and Oracle Bound

We consider the data-driven selection of the spectral shrinkage parameter $\lambda \geq 0$. While classical Tikhonov regularization selects shrinkage via prediction cross-validation, in OR-DML the target is the structural causal parameter vector θ^* , where prediction cross-validation conflates causal identification with prognostic predictive fit. We distinguish two distinct regularization regimes:

1. **Inferential Regularization** ($\lambda = o(N^{-1/2})$): Required by Theorem 7 to guarantee asymptotic unbiasedness $\sqrt{N}\text{Bias}(\lambda) \rightarrow \mathbf{0}$, ensuring centered Gaussian limit laws and valid nominal confidence interval coverage.
2. **Finite-Sample Risk Minimization** ($\lambda_{\text{risk}} = \mathcal{O}(1)$): Minimizes the finite-sample expected ℓ_2 estimation risk $\mathbb{E}\|\hat{\theta}_\lambda - \theta^*\|_2^2$ under severe causal ill-conditioning ($\lambda_{\min}(\mathbf{J}) \approx 0$).

Empirical Risk Surrogate and Convexity: Define the empirical plug-in risk surrogate $\hat{R}(\lambda)$ by substituting empirical estimators $(\hat{\varepsilon}_\gamma, \lambda_{\min}(\hat{\mathbf{J}}), \hat{\mathbf{\Omega}}, \|\hat{\theta}_0\|_2)$ into the risk upper bound of Theorem 6:

$$\hat{R}(\lambda) = \frac{(C\hat{\varepsilon}_\gamma + \lambda\|\hat{\theta}_0\|_2)^2 + N^{-1}\text{Tr}(\hat{\mathbf{\Omega}})}{(\lambda_{\min}(\hat{\mathbf{J}}) + \lambda)^2}. \quad (63)$$

Let $u = \lambda_{\min}(\hat{\mathbf{J}}) + \lambda \geq \lambda_{\min}(\hat{\mathbf{J}}) > 0$. Rewriting $\lambda = u - \lambda_{\min}(\hat{\mathbf{J}})$, the numerator becomes:

$$\left(C\hat{\varepsilon}_\gamma - \lambda_{\min}(\hat{\mathbf{J}})\|\hat{\theta}_0\|_2 + u\|\hat{\theta}_0\|_2 \right)^2 + N^{-1}\text{Tr}(\hat{\mathbf{\Omega}}) = (A + uB)^2 + V, \quad (64)$$

where $A = C\hat{\varepsilon}_\gamma - \lambda_{\min}(\hat{\mathbf{J}})\|\hat{\theta}_0\|_2$, $B = \|\hat{\theta}_0\|_2$, and $V = N^{-1}\text{Tr}(\hat{\mathbf{\Omega}})$. Dividing by u^2 :

$$\hat{R}(u - \lambda_{\min}) = B^2 + \frac{2AB}{u} + \frac{A^2 + V}{u^2}. \quad (65)$$

Differentiating with respect to u :

$$\frac{d\hat{R}}{du} = -\frac{2AB}{u^2} - \frac{2(A^2 + V)}{u^3} = -\frac{2}{u^3} [ABu + (A^2 + V)]. \quad (66)$$

Setting the derivative to zero yields the unique critical point $u^* = \frac{A^2 + V}{|A|B}$ when $A < 0$. When $A \geq 0$ (i.e., $C\hat{\varepsilon}_\gamma \geq \lambda_{\min}(\hat{\mathbf{J}})\|\hat{\boldsymbol{\theta}}_0\|_2$), $\frac{d\hat{R}}{du} < 0$ on $(0, \infty)$, indicating that spectral shrinkage strictly decreases risk until bounded by the constraint $\lambda \geq 0$. Constraining $\lambda \geq 0$, the closed-form interior minimizer is:

$$\hat{\lambda}^* = \max \left(0, \frac{N^{-1}\text{Tr}(\hat{\boldsymbol{\Omega}}) + (C\hat{\varepsilon}_\gamma)^2}{C\hat{\varepsilon}_\gamma\|\hat{\boldsymbol{\theta}}_0\|_2 + \lambda_{\min}(\hat{\mathbf{J}})\|\hat{\boldsymbol{\theta}}_0\|_2^2} - \lambda_{\min}(\hat{\mathbf{J}}) \right). \quad (67)$$

Because $R'(u) = -\frac{2}{u^3} [ABu + (A^2 + V)]$ changes sign strictly from negative to positive at u^* when $A < 0$, u^* is the unique global minimizer on $(0, \infty)$. By uniform convergence of the empirical plug-in surrogate on compact parameter sets $\lambda \in [0, \bar{\lambda}]$, $\sup_{\lambda \in [0, \bar{\lambda}]} |\hat{R}(\lambda) - R_{\text{surr}}(\lambda)| = \mathcal{O}_P(N^{-1/2})$. Applying M-estimation consistency to the unique minimum of $\hat{R}(\lambda)$ yields the adaptive risk-surrogate selection guarantee:

$$R_{\text{surr}}(\hat{\lambda}^*) \leq \inf_{\lambda \geq 0} R_{\text{surr}}(\lambda) + \mathcal{O}_P(N^{-1/2}), \quad (68)$$

establishing that data-driven spectral shrinkage $\hat{\lambda}^*$ adaptively tracks the optimal surrogate risk trade-off without prior knowledge of the true latent proxy error. ■

C.2 State Space Scalability and Matrix Concentration for $K > 2$

While our primary experiments study binary regimes ($K = 2$), the OR-DML coupled estimator generalizes directly to arbitrary finite state dimensions $K \geq 2$. Here we formalize the computational and statistical scalability of the coupled Gram matrix $\hat{\mathbf{J}} \in \mathbb{R}^{K \times K}$.

Computational Complexity: Constructing $\hat{\mathbf{J}}$ and score $\hat{\mathbf{S}}$ requires evaluating K nuisance pairs across B folds, taking $\mathcal{O}(BK \cdot \text{Cost}(\text{Nuisance}))$ operations. Forming $\hat{\mathbf{J}}$ takes $\mathcal{O}(NK^2)$ operations, and solving $(\hat{\mathbf{J}} + \lambda \mathbf{I})^{-1} \hat{\mathbf{S}}$ via Cholesky factorization takes $\mathcal{O}(K^3)$ operations. For moderate regime counts ($K \leq 20$), the linear solve takes < 1 ms and is computationally negligible compared to nuisance estimation.

Statistical Sample Complexity under α -Mixing: The statistical rate is governed by the concentration of the random matrix $\hat{\mathbf{J}} = \frac{1}{N} \sum_{t=1}^N \boldsymbol{\Gamma}_t \tilde{\mathbf{T}}_t \tilde{\mathbf{T}}_t^\top \boldsymbol{\Gamma}_t$ around its expectation $\mathbf{J}$. Let the latent Markov chain have spectral gap $\rho_{\text{gap}} \equiv 1 - |\lambda_2(\boldsymbol{\Pi})| > 0$. By the Matrix Bernstein Inequality for geometrically α -mixing processes (Banna et al., 2016), with probability at least $1 - \delta$:

$$\|\hat{\mathbf{J}} - \mathbf{J}\|_{\text{op}} \leq C_J \sqrt{\frac{K \log(K/\delta)}{N \cdot \rho_{\text{gap}}}} + \frac{C'_J K \log(K/\delta)}{N \cdot \rho_{\text{gap}}}. \quad (69)$$

By Weyl's theorem, $|\lambda_{\min}(\hat{\mathbf{J}}) - \lambda_{\min}(\mathbf{J})| \leq \|\hat{\mathbf{J}} - \mathbf{J}\|_{\text{op}}$. Furthermore, the population minimal eigenvalue satisfies $\lambda_{\min}(\mathbf{J}) \leq \min_{1 \leq k \leq K} \pi_k^* \mathbb{E}[\tilde{T}_{tk}^2]$. Therefore, to ensure that the empirical Gram matrix remains non-singular ($\lambda_{\min}(\hat{\mathbf{J}}) \geq \frac{1}{2} \lambda_{\min}(\mathbf{J}) > 0$) with high probability, the required sample size scales as:

$$N = \tilde{\Omega} \left(\frac{K \log(K/\delta)}{\min_k (\pi_k^*)^2 \lambda_{\min}(\mathbf{J})^2 \rho_{\text{gap}}} \right). \quad (70)$$

This bound reveals that state-space scaling is limited not by linear algebra ($\mathcal{O}(K^3)$), but by the rarity of the least frequent regime ($\pi_{\min}^*$) and task ill-conditioning ($\lambda_{\min}$).

Non-Identifiability vs. Shrinkage: When Systems Must Abstain: Spectral shrinkage $\hat{\boldsymbol{\theta}}_\lambda = (\hat{\mathbf{J}} + \lambda \mathbf{I})^{-1} \hat{\mathbf{S}}$ stabilizes variance under ill-conditioned overlap ($\lambda_{\min} \approx 0$). However, when data lack identifying variation entirely—diagnosed by condition number $\kappa(\hat{\mathbf{J}}) > 10^4$ or a moving-block bootstrap 95% confidence interval for $\lambda_{\min}(\hat{\mathbf{J}})$ that covers zero—the structural parameter $\boldsymbol{\theta}^*$ is non-identified (Theorem 3). In this regime, heavy shrinkage $\lambda \rightarrow \infty$ artificially forces $\|\hat{\boldsymbol{\theta}}_\lambda\| \rightarrow 0$ without recovering structural effects. Deployed decision pipelines should treat $\kappa(\hat{\mathbf{J}}) > 10^4$ as a structural abstention trigger rather than masking non-identifiability with excessive shrinkage.

D PROOF OF THEOREM 4: DECOMPOSITION OF OMITTED REGIME BIAS

Consider the Structural Causal Model (1)–(5):

$$\begin{aligned} T_t &= m(X_t, S_t) + V_t, \quad \mathbb{E}[V_t | X_t, S_t] = 0, \\ Y_t &= \sum_{k=1}^K \theta_k^* \mathbb{I}(S_t = k) T_t + g(X_t, S_t) + U_t, \quad \mathbb{E}[U_t | X_t, S_t, T_t] = 0. \end{aligned}$$

Define the unconditioned expectations: $\bar{m}(X_t) \equiv \mathbb{E}[T_t | X_t] = \sum_{k=1}^K \pi_k(X_t) m_k(X_t)$ and $\bar{g}(X_t) \equiv \mathbb{E}[g(X_t, S_t) | X_t] = \sum_{k=1}^K \pi_k(X_t) g_k(X_t)$, where $\pi_k(X_t) \equiv P(S_t = k | X_t)$. Residuals from cross-fitting on X_t alone satisfy $\tilde{T}_t = T_t - \bar{m}(X_t) = \sum_{k=1}^K \mathbb{I}(S_t = k) \Delta m_k(X_t) + V_t$, where $\Delta m_k(X_t) = m_k(X_t) - \bar{m}(X_t)$. The outcome residual is:

$$\tilde{Y}_t = Y_t - \mathbb{E}[Y_t | X_t] = \sum_{k=1}^K \theta_k^* \mathbb{I}(S_t = k) T_t - \sum_{k=1}^K \theta_k^* \pi_k(X_t) m_k(X_t) + g(X_t, S_t) - \bar{g}(X_t) + U_t. \quad (71)$$

Writing $T_t = \tilde{T}_t + \bar{m}(X_t)$, the treatment interaction terms decompose as:

$$\sum_{k=1}^K \theta_k^* \mathbb{I}(S_t = k) T_t - \sum_{k=1}^K \theta_k^* \pi_k(X_t) m_k(X_t) = \sum_{k=1}^K \theta_k^* \mathbb{I}(S_t = k) \tilde{T}_t + \sum_{k=1}^K \theta_k^* [\mathbb{I}(S_t = k) \bar{m}(X_t) - \pi_k(X_t) m_k(X_t)]. \quad (72)$$

Taking the inner product with $\tilde{T}_t$ and evaluating expectations: (i) By Assumption 1, $\mathbb{E}[\tilde{T}_t U_t] = 0$. (ii) $\mathbb{E}[\tilde{T}_t \sum_k \theta_k^* \mathbb{I}(S_t = k) \tilde{T}_t] = \sum_{k=1}^K \theta_k^* \mathbb{E}[\mathbb{I}(S_t = k) \tilde{T}_t^2] = \mathbb{E}[\tilde{T}_t^2] \sum_{k=1}^K \omega_k \theta_k^*$, where $\omega_k = \frac{\mathbb{E}[\mathbb{I}(S_t = k) \tilde{T}_t^2]}{\mathbb{E}[\tilde{T}_t^2]}$ with $\sum_k \omega_k = 1$. (iii) Since $\mathbb{E}[\tilde{T}_t | X_t] = 0$ and $\mathbb{E}[\tilde{T}_t \mathbb{I}(S_t = k) | X_t] = \Delta m_k(X_t) \pi_k(X_t)$, the mean adjustment yields $\sum_{k=1}^K \theta_k^* \mathbb{E}[\bar{m}(X_t) \Delta m_k(X_t) \pi_k(X_t)]$. (iv) The confounding component evaluates to $\mathbb{E}[\tilde{T}_t (g(X_t, S_t) - \bar{g}(X_t))] = \sum_{k=1}^K \mathbb{E}[\pi_k(X_t) \Delta m_k(X_t) \Delta g_k(X_t)]$. Dividing by $\mathbb{E}[\tilde{T}_t^2]$ yields (11) and (12). ■

E PROOF OF THEOREM 5: ASYMPTOTIC NORMALITY UNDER DEPENDENT PURGED CROSS-FITTING

The empirical score expansion around true θ^* and true nuisances η^* is:

$$\sqrt{N} \left(\hat{\theta}_\lambda - \theta^* - \text{Bias}(\lambda) \right) = (\hat{\mathbf{J}} + \lambda \mathbf{I})^{-1} \frac{1}{\sqrt{N}} \sum_{t=1}^N \Psi(W_t; \theta^*, \hat{\eta}^{(-b(t))}). \quad (73)$$

With empirical Bartlett embargo window $\tau^* \geq C \log N$, the α -mixing dependence between training set $\mathcal{T}_b$ and test fold $\mathcal{B}_b$ satisfies:

$$\alpha(\tau^*) \leq M \exp(-c\tau^*) \leq MN^{-cC} = o(N^{-1}). \quad (74)$$

Consequently, cross-fold covariance is asymptotically negligible, and nuisance estimation errors satisfy:

$$\|\hat{\mu}_Y - \mu_Y^*\|_{L_2} \cdot \|\hat{\mu}_D - \mu_D^*\|_{L_2} = o_P(N^{-1/2}). \quad (75)$$

By the Ibragimov (1962) central limit theorem for strictly stationary, α -mixing processes satisfying $\sum_m \alpha(m)^{\delta/(2+\delta)} < \infty$:

$$\frac{1}{\sqrt{N}} \sum_{t=1}^N \Psi(W_t; \theta^*, \eta^*) \xrightarrow{d} \mathcal{N}(\mathbf{0}, \mathbf{\Omega}), \quad (76)$$

where $\mathbf{\Omega} = \lim_{N \rightarrow \infty} \text{Var} \left(N^{-1/2} \sum_{t=1}^N \Psi_t \right)$. Applying Slutsky's theorem with $\hat{\mathbf{J}} \xrightarrow{P} \mathbf{J}$:

$$\sqrt{N} \left(\hat{\theta}_\lambda - \theta^* - \text{Bias}(\lambda) \right) \xrightarrow{d} \mathcal{N}(\mathbf{0}, \mathbf{\Sigma}_\lambda), \quad (77)$$

where $\mathbf{\Sigma}_\lambda = (\mathbf{J} + \lambda \mathbf{I})^{-1} \mathbf{\Omega} (\mathbf{J} + \lambda \mathbf{I})^{-1}$. When $\lambda = o(N^{-1/2})$, $\sqrt{N} \text{Bias}(\lambda) \rightarrow \mathbf{0}$, completing the proof.

Operational Modes of Forward Filtering. In offline retrospective smoothing, $\gamma_{ik}^{\text{smooth}} = P(S_t = k | Z_{1:N})$. In real-time decision making, three distinct forward modes exist: (1) *Oracle Filter*: $P(S_t = k | Z_{1:t}, \phi^*)$ using the true HMM parameters; (2) *Training-Fitted Forward Filter*: $P(S_t = k | Z_{1:t}, \hat{\phi})$ where parameters $\hat{\phi}$ are fitted on a historical calibration window and state inference updates strictly forward on past-to-present observations $Z_{1:t}$;

and (3) *Expanding-Window Online Filter*: where both parameters $\hat{\phi}_t$ and state updates proceed sequentially. Our implementations evaluate Mode 1 and Mode 2, where Mode 2 enforces causal forward state filtering without future observation leakage. ■

E.1 Observed-Data Orthogonal Score under Estimated Latent Dynamics

In semiparametric causal inference (Chernozhukov et al., 2018; Bickel et al., 1993), when nuisance parameters consist of both non-parametric outcome/propensity regressions $\eta = (m, \mu)$ and finite-dimensional parameters Λ governing latent state transitions and emissions, one must characterize the sensitivity of the causal estimator to the first-order estimation error in $\hat{\Lambda}$.

Let the state-weighted regime score vector be:

$$\psi_{\text{regime}}(W_t; \theta, \eta, \Lambda) \equiv \sum_{k=1}^K \gamma_{tk}(\Lambda) (T_t - m_k(X_t)) (Y_t - \mu_k(X_t) - \theta_k(T_t - m_k(X_t))). \quad (78)$$

In a hidden Markov model, the latent parameters Λ are estimated via sequence-level maximum likelihood with score function $s_{\Lambda, N}(Z_{1:N}) = \nabla_{\Lambda} \log p(Z_{1:N}; \Lambda)$. Evaluating the Gateaux derivative with respect to Λ at the oracle targets $(\theta^*, \eta^*, \Lambda^*)$:

$$\mathbf{A}_{\Lambda} \equiv \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{t=1}^N \sum_{k=1}^K \mathbb{E} \left[\frac{\partial \gamma_{tk}(\Lambda^*)}{\partial \Lambda^{\top}} (T_t - m_k^*(X_t)) (Y_t - \mu_k^*(X_t) - \theta_k^*(T_t - m_k^*(X_t))) \right]. \quad (79)$$

Importantly, $\mathbf{A}_{\Lambda}$ does not vanish pointwise across regimes. While Assumption 3 guarantees that the structural innovation has zero mean conditional on the true state, $\mathbb{E}[Y_t - \mu_k^*(X_t) - \theta_k^*(T_t - m_k^*(X_t)) \mid X_t, T_t, S_t = k] = 0$, conditioning on observed history $Z_{1:N}$ alone yields:

$$\mathbb{E}[Y_t - \mu_k^*(X_t) - \theta_k^*(T_t - m_k^*(X_t)) \mid X_t, T_t, Z_{1:N}] = \sum_{j \neq k} \gamma_{tj}(\Lambda^*) [(\mu_j^*(X_t) - \mu_k^*(X_t)) + (\theta_j^* - \theta_k^*)(T_t - m_j^*(X_t))]. \quad (80)$$

This residual is non-zero whenever regimes differ in baseline potential outcomes ($\mu_j^* \neq \mu_k^*$) or treatment responsiveness ($\theta_j^* \neq \theta_k^*$).

Under purged block cross-fitting (Algorithm 1), this first-order variation is controlled via two complementary mechanisms:

1. **Decoupling via Purged Embargo Buffer:** The parameters $\hat{\Lambda}$ are estimated on historical calibration blocks or out-of-fold data separated by an embargo buffer $\tau^* \geq C \log N$. By the α -mixing decay of the underlying process (Assumption 4), the covariance between evaluation fold innovations and the parameter estimation error $\hat{\Lambda} - \Lambda^*$ decays as $O(N^{-C\alpha})$.
2. **Stacked Joint HAC Variance Representation:** When $\hat{\Lambda}$ is estimated at parametric rate $\sqrt{N}(\hat{\Lambda} - \Lambda^*) = O_p(1)$, the joint asymptotic expansion takes the stacked M-estimation form:

$$\sqrt{N} \begin{pmatrix} \hat{\theta}_{\lambda} - \theta^* - \text{Bias}(\lambda) \\ \hat{\Lambda} - \Lambda^* \end{pmatrix} \xrightarrow{d} \mathcal{N} \left(\mathbf{0}, \begin{bmatrix} \Sigma_{\lambda} & \Sigma_{\theta\Lambda} \\ \Sigma_{\theta\Lambda}^{\top} & \mathcal{I}_{\Lambda}^{-1} \end{bmatrix} \right), \quad (81)$$

where $\Sigma_{\theta\Lambda}$ is consistently captured by the block Newey-West HAC covariance estimator. In well-separated regimes ($\Delta_Z \rightarrow \infty$), posterior uncertainty vanishes ($\gamma_{tk} \rightarrow \mathbb{I}(S_t = k)$) and $\partial \gamma_{tk} / \partial \Lambda \rightarrow \mathbf{0}$ exponentially fast, rendering $\|\mathbf{A}_{\Lambda}\| \leq C e^{-c\Delta_Z^2}$ and recovering asymptotic orthogonality.

F PROOF OF PROPOSITION 6: JOINT DELTA-METHOD INFERENCE FOR POPULATION ATE

We distinguish three distinct regime weighting vectors:

1. **True Stationary Distribution π^* :** The invariant measure of the underlying Markov transition kernel Π^* , satisfying $\pi^{*\top} \Pi^* = \pi^{*\top}$.
2. **Estimated HMM Stationary Distribution $\hat{\pi}(\hat{\Lambda})$:** The stationary vector derived from the estimated HMM transition matrix $\hat{\Pi}$, where $\hat{\Lambda}$ denotes the vector of estimated HMM parameters.

3. Sample Posterior Occupation Distribution $\bar{\gamma}_N$: The empirical average of posterior smoothed regime weights across the observed sample: $\bar{\gamma}_N \equiv \frac{1}{N} \sum_{t=1}^N \gamma_t$.

The sample Average Treatment Effect estimator is $\hat{\theta}_{\text{SATE}} = \sum_{k=1}^K \bar{\gamma}_{N,k} \hat{\theta}_{\lambda,k} = \bar{\gamma}_N^\top \hat{\theta}_\lambda$. Conditional on the realized sample regime occupancy, its asymptotic variance is governed solely by parameter estimation uncertainty:

$$\sigma_{\text{SATE}}^2 = \bar{\gamma}_N^\top \Sigma_\lambda \bar{\gamma}_N. \quad (82)$$

For the population estimand $\theta_{\text{PATE}}^* = \pi^{*\top} \theta^*$, the plug-in estimator is $\hat{\theta}_{\text{PATE}} = \hat{\pi}^\top \hat{\theta}_\lambda$. Expanding $\hat{\theta}_{\text{PATE}}$ around (π^*, θ^*) via the multivariate delta method:

$$\sqrt{N} (\hat{\theta}_{\text{PATE}} - \theta_{\text{PATE}}^*) = \pi^{*\top} \sqrt{N} (\hat{\theta}_\lambda - \theta^*) + \theta^{*\top} \sqrt{N} (\hat{\pi} - \pi^*) + o_P(1). \quad (83)$$

The asymptotic variance of $\hat{\theta}_{\text{PATE}}$ is given by the quadratic form of the joint limiting distribution:

$$\sigma_{\text{PATE}}^2 = \lim_{N \rightarrow \infty} \text{Var} \left(\sqrt{N} (\hat{\theta}_{\text{PATE}} - \theta_{\text{PATE}}^*) \right) = \pi^{*\top} \Sigma_\lambda \pi^* + \theta^{*\top} \Sigma_\pi \theta^* + 2\pi^{*\top} \Sigma_{\theta\pi} \theta^*, \quad (84)$$

where $\Sigma_\lambda \equiv \lim_{N \rightarrow \infty} N \text{Var}(\hat{\theta}_\lambda)$, $\Sigma_\pi \equiv \lim_{N \rightarrow \infty} N \text{Var}(\hat{\pi})$, and $\Sigma_{\theta\pi} \equiv \lim_{N \rightarrow \infty} N \text{Cov}(\hat{\theta}_\lambda, \hat{\pi})$.

Cross-Covariance Analysis and HMM Score Orthogonality: When both $\hat{\theta}_\lambda$ and $\hat{\pi}$ are estimated from the same longitudinal record, they share the estimated HMM parameter vector $\hat{\Lambda}$. Let $\phi_\theta(W_t) \equiv (\mathbf{J} + \lambda \mathbf{I})^{-1} \Psi_t$ denote the uncentered DML influence function and $\mathbf{s}_\Lambda(Z_t)$ denote the score function of the HMM parameters. The asymptotic representation of $\hat{\pi}$ satisfies:

$$\sqrt{N} (\hat{\pi} - \pi^*) = \nabla_\Lambda \pi^* \mathcal{I}_\Lambda^{-1} \frac{1}{\sqrt{N}} \sum_{t=1}^N \mathbf{s}_\Lambda(Z_t) + o_P(1), \quad (85)$$

where $\mathcal{I}_\Lambda$ is the Fisher information matrix of the HMM. Consequently, the asymptotic cross-covariance evaluates to:

$$\Sigma_{\theta\pi} = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{t=1}^N \sum_{s=1}^N \mathbb{E} [\phi_\theta(W_t) \mathbf{s}_\Lambda(Z_s)^\top] \mathcal{I}_\Lambda^{-1} (\nabla_\Lambda \pi^*)^\top. \quad (86)$$

Therefore, $\Sigma_{\theta\pi} = \mathbf{0}$ if and only if the causal score innovations and the latent Markov score satisfy asymptotic cross-orthogonality: $\mathbb{E}[\phi_\theta(W_t) \mathbf{s}_\Lambda(Z_s)^\top] = \mathbf{0}$ for all lag intervals $|t - s| \geq 0$. While this orthogonality holds under structural innovation independence where $\mathbb{E}[\Psi_t | Z_{1:N}] = \mathbf{0}$, in empirical settings where unmodeled feedback exists between emissions and treatment errors, $\Sigma_{\theta\pi}$ may be non-zero. In practice, σ_{PATE}^2 is consistently estimated by computing a joint Newey-West HAC covariance over the stacked $(2K)$ -dimensional influence sequence $(\phi_\theta(W_t)^\top, \phi_\pi(Z_t)^\top)^\top$.

Stationary Distribution Variance for Markov Chains: For an irreducible, aperiodic Markov chain with transition matrix Π^* and stationary distribution π^* , the fundamental matrix (Kemeny and Snell, 1976) is $\mathbf{Z} \equiv (\mathbf{I} - \Pi^* + \mathbf{1}\pi^{*\top})^{-1}$. The asymptotic covariance matrix of the sample occupation distribution satisfies:

$$\Sigma_\pi = \mathbf{D}_\pi \mathbf{Z} + \mathbf{Z}^\top \mathbf{D}_\pi - \mathbf{D}_\pi - \pi^* \pi^{*\top}, \quad (87)$$

where $\mathbf{D}_\pi = \text{diag}(\pi^*)$. For a two-state Markov chain ($K = 2$) with persistence parameter $\rho = 1 - P_{01} - P_{10} = \lambda_2(\Pi^*)$:

$$\text{Var}(\bar{S}_N) = \frac{\pi_0^* \pi_1^*}{N} \left(\frac{1 + \rho}{1 - \rho} \right) + o(N^{-1}). \quad (88)$$

When $\Delta\theta^* = \theta_1^* - \theta_0^* \neq 0$, the sampling variation in regime occupancy inflates the estimation variance by $\frac{1+\rho}{1-\rho} \approx 15.7\times$ under persistent stagnation ($\rho \approx 0.88$). ■

G EMPIRICAL SENSITIVITY ANALYSES AND DATA PROVENANCE

G.1 Meteorological Regime Characteristics across Megacities

Table 3 details the empirical regime profiles uncovered by the Gaussian HMM across all four megacities. Across all airsheds, the inferred regimes correspond directly to distinct synoptic meteorological states: cooler, high-humidity, low-wind nocturnal/winter inversion states versus warmer, lower-humidity daytime convective dispersion states.

Table 3: Meteorological Regime Profiling across Four Indian Megacities from Ground-Sensor Observations.

City	Regime	Share	Persistence	Wind (m/s)	Temp (°C)	RH (%)
Delhi	Regime 1	42.7%	0.989	5.98	19.63	61.4%
Delhi	Regime 2	57.3%	0.992	7.52	30.93	56.3%
Mumbai	Regime 1	41.7%	0.789	11.49	26.69	83.2%
Mumbai	Regime 2	58.3%	0.857	16.48	28.39	77.9%
Bengaluru	Regime 1	52.2%	0.843	10.83	20.77	84.6%
Bengaluru	Regime 2	47.8%	0.828	11.22	27.67	45.3%
Kolkata	Regime 1	60.7%	0.871	7.32	26.73	85.3%
Kolkata	Regime 2	39.3%	0.810	7.89	32.65	51.2%




Figure 4: Dynamic Exposure-Response Curves $\hat{\theta}_h$ across Horizons $h \in [0, 24]$ Hours for Delhi, Mumbai, Bengaluru, and Kolkata. Shaded bands represent 95% HAC confidence intervals. Regime stratification reveals substantial structural heterogeneity in dynamic pollutant accumulation over multi-hour horizons.

G.2 Regularization Sensitivity in Ill-Conditioned Regimes (Kolkata Airshed)

To verify the theoretical behavior established in Theorem 6 under real-world conditions, Table 4 reports the empirical sensitivity of OR-DML across a fine 13-point regularization grid $\lambda \in [0.0, 1.0]$ in the ill-conditioned Kolkata airshed ($\lambda_{\min} = 0.0024$).

Table 4: Regularization Sensitivity Grid for Kolkata Airshed ($\lambda_{\min} = 0.0024$, $N = 1,209$). Standard errors decrease monotonically from 1.1071 at $\lambda = 0.0$ to 0.0063 at $\lambda = 1.0$, stabilizing estimation variance while shrinking point estimates toward zero.

λ	$\hat{\theta}_{\text{overall}}$	SE	95% CI	$\lambda_{\min}(\hat{\mathbf{J}} + \lambda \mathbf{I})$	$\kappa(\hat{\mathbf{J}} + \lambda \mathbf{I})$
0.0000	-0.5570	1.1071	[-2.7269, +1.6130]	0.0049	1.28
0.0001	-0.5485	1.0878	[-2.6807, +1.5836]	0.0050	1.28
0.0005	-0.5172	1.0170	[-2.5105, +1.4760]	0.0054	1.28
0.0010	-0.4828	0.9406	[-2.3263, +1.3607]	0.0059	1.28
0.0024	-0.4068	0.7776	[-1.9309, +1.1173]	0.0073	1.28
0.0050	-0.3147	0.5890	[-1.4691, +0.8398]	0.0099	1.28
0.0100	-0.2191	0.4022	[-1.0074, +0.5691]	0.0149	1.28
0.0200	-0.1363	0.2463	[-0.6191, +0.3464]	0.0249	1.28
0.0500	-0.0639	0.1140	[-0.2872, +0.1595]	0.0549	1.28
0.1000	-0.0339	0.0601	[-0.1517, +0.0840]	0.1049	1.28
0.2000	-0.0175	0.0309	[-0.0781, +0.0432]	0.2049	1.28
0.5000	-0.0071	0.0126	[-0.0318, +0.0175]	0.5049	1.28
1.0000	-0.0036	0.0063	[-0.0160, +0.0088]	1.0049	1.28

As predicted by Theorem 6, when $\lambda \rightarrow 0$, standard errors explode to 1.1071 due to near-singularity in residual treatment variation. Adding spectral regularization shifts the minimal singular value from 0.0049 to 0.1049 at $\lambda = 0.10$, reducing standard errors by an order of magnitude ($1.1071 \rightarrow 0.0601$) while shrinking point estimates conservatively toward zero. This confirms that spectral regularization acts as a principled safety mechanism against catastrophic variance blowup in ill-conditioned environments.

G.3 Data Imputation Sensitivity Analysis (Strictly Observed vs. Cleaned)

To verify that our empirical findings are not artifacts of missing-data handling, we compare OR-DML causal estimates obtained on the full multi-season cleaned dataset (which uses causal forward-fill on isolated gaps $< 0.7\%$ of hours) against a strictly observed dataset where every row containing any imputed value is entirely deleted.

Table 5: Sensitivity Analysis Comparing Full Dataset ($< 0.7\%$ Causal Forward-Fill) vs. Strictly Observed Dataset (Rows with Any Imputed Entry Deleted) Across Four Megacities. Point estimates and standard errors exhibit identical signs and overlapping confidence bands.

City	N_{Full}	$\hat{\theta}_{\text{Full}}$	SE_{Full}	N_{Strict}	$\hat{\theta}_{\text{Strict}}$	$\text{SE}_{\text{Strict}}$
Delhi	4,640	+0.3634	0.0460	4,602	+0.5366	0.0674
Mumbai	4,180	-5.7671	2.1367	4,146	-10.9323	2.6792
Bengaluru	4,093	+0.0046	0.0101	4,084	+0.0202	0.0112
Kolkata	1,209	-0.5563	1.1055	1,206	-0.1924	1.0148

As documented in Table 5:

- In Delhi, deleting the 38 forward-filled hours (0.8%) yields $\hat{\theta}_{\text{Strict}} = +0.5366 \pm 0.0674$, matching the positive elasticity observed under the full series ($+0.3634 \pm 0.0460$).
- In Mumbai, deleting 34 hours (0.8%) retains the strong negative dispersion elasticity (-10.9323 ± 2.6792 vs. -5.7671 ± 2.1367).
- In Bengaluru, deleting 9 hours (0.2%) preserves the near-zero net effect ($+0.0202 \pm 0.0112$ vs. $+0.0046 \pm 0.0101$).
- In Kolkata, deleting 3 hours (0.2%) preserves identical negative directionality with overlapping confidence intervals (-0.1924 ± 1.0148 vs. -0.5563 ± 1.1055).

These results confirm that our conclusions are robust to missing data and do not depend on the causal forward-fill protocol.

H LATENTREGIMEBENCH: MULTI-DIMENSIONAL BENCHMARK RESULTS

To systematically evaluate machine learning under persistent, partially observed, and shifting latent regimes, we conduct comprehensive evaluations on **LatentRegimeBench** across five diagnostic configurations ($N = 1,200$, 30 Monte Carlo replications each) and four representation architectures (Table 6 and Table 7, with summary visualizations in Figure 5).




Figure 5: Universal LatentRegimeBench Evaluation. (a) Calibration of causal estimation bias against theoretical degradation ratio $\mathcal{D} = \bar{\varepsilon}_\gamma / \lambda_{\min}(\mathbf{J})$. (b) Decision-time abstention safety comparing unconstrained point MSE against uncertainty-gated MSE. (c) Representation zoo out-of-distribution (OOD) generalization across HMM, GRU, Causal Transformer, and Linear SSM.

Table 6: LatentRegimeBench: Multi-Dimensional Stress Test across 5 Strategic Difficulty Configurations (30 Replications each). In well-conditioned regimes (C1, C3), OR-DML achieves a $> 5.5\times$ bias reduction over Standard DML, while under severe proxy error (C2, C4) or persistence inversion (C5), degradation tracks the difficulty ratio $\mathcal{D}$.

Configuration	Δ_Z	Δ_T	ρ	$\mathcal{D}$	OR-DML Bias	Naive Bias	Advantage
Ideal (High Obs, High Overlap)	3.5	5.0	0.85	0.001	0.2165	1.2061	$5.57\times$
Adversarial A (Low Obs, High Overlap)	0.3	5.0	0.85	0.709	1.1950	1.2061	$1.01\times$
Adversarial B (High Obs, Weak Overlap)	3.5	0.4	0.85	0.001	0.2498	1.3886	$5.56\times$
Challenging (Low Obs, Weak Overlap)	0.5	0.5	0.90	2.748	1.3237	1.3862	$1.05\times$
High-Persistence Inversion	2.0	2.0	0.96	0.159	1.5934	1.3175	$0.83\times$

Representation Accuracy vs. Downstream Task Reliability. To evaluate whether latent sequence representations translate to reliable downstream prediction under distribution shift, we fit a downstream Ridge

Table 7: Latent Representation Zoo under Disentangled Distribution Shifts ($N = 2,000$ per evaluation stream, 5 independent random seeds; mean $\pm$ std). Shift A isolates transition dynamics ($\rho = 0.85 \rightarrow 0.96$); Shift B isolates observation noise ($\Delta_Z = 3.5 \rightarrow 1.0$); Shift C compounds both perturbations. While deep neural encoders achieve competitive latent recovery under compound shift, Gaussian HMM maintains the lowest compound calibration error (ECE = 0.0397 ± 0.0242).

Architecture	In-Dist F_1	Shift A F_1	Shift B F_1	Shift C F_1	ΔF_1 Gap	Shift C ECE
Gaussian HMM (Causal Filter)	0.9986 ± 0.0004	1.0000 ± 0.0000	0.8770 ± 0.0114	0.9170 ± 0.0118	$+0.0817 \pm 0.0115$	0.0397 ± 0.0242
Neural GRU Encoder	0.9997 ± 0.0007	0.9997 ± 0.0004	0.8426 ± 0.0128	0.8481 ± 0.0136	$+0.1516 \pm 0.0136$	0.1047 ± 0.0129
Causal Transformer Encoder	0.9997 ± 0.0004	1.0000 ± 0.0000	0.8328 ± 0.0125	0.8372 ± 0.0105	$+0.1625 \pm 0.0108$	0.1561 ± 0.0136
Linear State Space Model (SSM)	0.9995 ± 0.0007	1.0000 ± 0.0000	0.8563 ± 0.0132	0.8580 ± 0.0118	$+0.1415 \pm 0.0119$	0.0558 ± 0.0159

predictor $f_m : (X_t, T_t, \gamma_t^{(m)}) \mapsto Y_t$ strictly on the training partition ($N = 1,200$) and freeze it. In addition, nuisance models $\hat{m}_k(X_t) = \mathbb{E}[T_t | X_t, S_t = k]$ are estimated on training data to construct residualized treatments $\tilde{T}_{tk} = T_t - \hat{m}_k(X_t)$. We compute the exact coupled regime Gram matrix $\mathbf{J}_t = \frac{1}{w} \sum_{s=t-w+1}^t \begin{pmatrix} \gamma_{s0}^2 \tilde{T}_{s0}^2 & \gamma_{s0} \gamma_{s1} \tilde{T}_{s0} \tilde{T}_{s1} \\ \gamma_{s0} \gamma_{s1} \tilde{T}_{s0} \tilde{T}_{s1} & \gamma_{s1}^2 \tilde{T}_{s1}^2 \end{pmatrix}$ across rolling evaluation windows ($w = 48$) and extract its exact minimum eigenvalue $\lambda_{\min}(\mathbf{J}_t)$. On held-out Shift C ($N = 1,200$, 5 independent random seeds), we split into a calibration partition ($N_{\text{cal}} = 400$) to determine the extreme failure loss threshold $q_{0.90}$ once and freeze it for out-of-sample test evaluation, preventing threshold re-definition during bootstrap resamples. We evaluate future lead loss $L_{t+1} = (Y_{t+1} - \hat{Y}_{t+1})^2$ against three distinct signals: (i) State Posterior Entropy $H_t = H(\gamma_t)$, (ii) Task Conditioning alone $1/\lambda_{\min}(\mathbf{J}_t)$, and (iii) Coupled Task Difficulty $\mathcal{D}_t = H_t/\lambda_{\min}(\mathbf{J}_t)$. To quantify estimation uncertainty over temporally dependent observations, intervals are computed via Moving-Block Bootstrap (MBB, $B = 1,000$ resamples with block length $L = 24$ hours), serving as an empirical uncertainty quantification procedure rather than asserting independent sample generation. Sensitivity analysis across block lengths $L \in \{12, 24, 48, 72\}$ confirms that MBB 95% CIs are remarkably stable (HMM CI width 0.142–0.147 across all L ; `reports/representation_zoo_mbb_sensitivity.csv`).

Offline Representation Diagnostics vs. Online Deployable Signals: We distinguish two operational regimes forming a three-level **Reliability Triangle**: (1) *State Identification* (F_1): Can the encoder identify the latent state? (2) *Probabilistic Trustworthiness* (ECE, NLL): Are posterior probabilities well-calibrated? These are retrospective representation diagnostics evaluated on labeled benchmark streams. (3) *Downstream Reliability* (ROC-AUC, $\mathcal{D}_t$): Can the system anticipate downstream failure? At online deployment, true states S_t are unobserved, precluding ECE or NLL computation; monitoring must rely on prospective signals ($H_t, \lambda_{\min}(\mathbf{J}_t), \mathcal{D}_t$).

As shown in Table 8, deep neural sequence models (GRU, Causal Transformer, Linear SSM) achieve competitive latent recovery ($F_1 \approx 0.84$ – 0.86), yet their uncalibrated softmax posteriors produce near-chance downstream failure detection (ROC-AUC ≈ 0.515 – 0.549). In contrast, Gaussian HMM causal filtering achieves well-calibrated posterior states (ECE = 0.0397) and provides statistically informative failure discrimination in this benchmark (ROC-AUC = **0.6153 ± 0.0246** , MBB 95% CI [0.540, 0.683], PR-AUC = 0.1300 ± 0.0169). This discrepancy stems from two structural failure mechanisms: (i) neural sequence models trained under cross-entropy maximize margin rather than probability calibration, yielding overconfident softmax distributions under distribution shift that blind raw entropy alarms; and (ii) standard sequence models are entirely task-geometry agnostic, omitting the downstream Gram matrix $\mathbf{J}_t$. When downstream tasks experience collinearity, naive models collapse regardless of latent classification accuracy, whereas coupled difficulty $\mathcal{D}_t$ anticipates failure by integrating task conditioning. To evaluate whether calibration provides independent explanatory power beyond classification accuracy within the same benchmark environment, we conduct a multi-world fixed-effects hierarchical regression across $D = 15$ independent benchmark worlds ($N = 60$ observations; `reports/representation_zoo_hierarchical_regression.csv`): $\text{ReliabilityAUC}_{d,m} = \alpha_d + \beta_1 F_{1,d,m} + \beta_2 \text{Calibration}_{d,m} + \epsilon_{d,m}$. In Model 1 (F_1 alone), classification F_1 is strongly positively associated with reliability ($\beta_1 = +0.9817 \pm 0.2011, p = 0.0002, R_{\text{within}}^2 = 0.4313$), confirming that state identification is essential. However, in Model 3 ($F_1 + \text{NLL}$), posterior log-loss NLL provides statistically significant incremental explanatory power after controlling for F_1 ($\beta_{\text{NLL}} = -0.0653 \pm 0.0295, p = 0.0442$), increasing within-world explanatory power by 9.24 percentage points ($\Delta R_{\text{within}}^2 = 0.0924$, raising $R_{\text{within}}^2 \rightarrow 0.5237$). In Model 2 ($F_1 + \text{ECE}$), calibration ECE exhibits a consistent negative association and increases within-world fit ($\Delta R_{\text{within}}^2 = 0.0647$, raising $R_{\text{within}}^2 \rightarrow 0.4960$). Thus, classification accuracy explains substantial reliability variation, but calibrated confidence provides incremental explanatory power. Furthermore, sensitivity analysis over numerical eigenvalue floors $\epsilon \in [10^{-6}, 10^{-3}]$ confirms identical AUCs (0.6153), demonstrating invariance to numerical conditioning floors.

Crucially, we identify three empirically distinct operating regimes across our benchmark and case studies:

1. **Regime I (State Uncertainty Dominates):** In well-conditioned airsheds with ample residual variation (Bengaluru Plateau, $\lambda_{\min} = 4.22$), downstream failure is governed by thermodynamic state transitions, making raw state posterior entropy H_t the dominant anticipator (AUC = 0.6737).
2. **Regime II (Task Conditioning Dominates):** In coastal airsheds where marine breeze reversals induce severe treatment collinearity (Mumbai Coastal, $\lambda_{\min} = 0.32$; Table 9), raw state entropy acts as a coin flip (AUC = 0.4855), while coupled task difficulty $\mathcal{D}_t$ incorporates task conditioning $1/\lambda_{\min}(\mathbf{J}_t)$ to restore failure anticipation (AUC = 0.5309–0.5640).
3. **Regime III (Mixed / Joint Limitation):** Under compound distribution shifts (Shift C), neither state uncertainty nor task conditioning alone is universally sufficient; on Shift C with Gaussian HMM, state entropy alone achieves AUC = 0.6209 while coupled difficulty achieves 0.6153. The coupled index $\mathcal{D}_t$ acts as a conservative, task-conditioned surrogate rather than universally dominating entropy, demonstrating that sequential reliability is governed jointly by both mechanisms.

Table 8: Representation Accuracy vs. Downstream Reliability: Evaluating Task-Conditioned Difficulty $\mathcal{D}_t$ across Sequence Encoders on a Genuinely Learned Downstream Model under Compound Distribution Shift (Shift C, 5 random seeds; mean $\pm$ seed std, with 95% Moving-Block Bootstrap CIs [$L = 24$]). High classification F_1 does not guarantee downstream reliability: uncalibrated softmax posteriors in deep neural encoders yield near-chance downstream failure detection (ROC-AUC ≈ 0.515 –0.549), whereas the Gaussian HMM causal filter achieves well-calibrated failure anticipation (ROC-AUC = **0.6153** $\pm$ **0.0246**, MBB 95% CI [0.540, 0.683]).

Architecture	Shift C F_1	ROC-AUC $\mathcal{D}$	MBB 95% CI	Baseline H AUC	Cond AUC	PR-AUC $\mathcal{D}$	MDA $\rho(\mathcal{D}, L)$
Gaussian HMM (Causal Filter)	0.9170 $\pm$ 0.012	0.6153 $\pm$ 0.025	[0.540, 0.683]	0.6209 $\pm$ 0.021	0.5003 $\pm$ 0.046	0.1300 $\pm$ 0.017	+0.0656 $\pm$ 0.050
Neural GRU Encoder	0.8481 $\pm$ 0.014	0.5228 $\pm$ 0.036	[0.446, 0.600]	0.5217 $\pm$ 0.031	0.5048 $\pm$ 0.035	0.0987 $\pm$ 0.019	−0.0032 $\pm$ 0.040
Causal Transformer Encoder	0.8372 $\pm$ 0.011	0.5150 $\pm$ 0.055	[0.438, 0.589]	0.5173 $\pm$ 0.063	0.4965 $\pm$ 0.013	0.1044 $\pm$ 0.027	+0.0009 $\pm$ 0.056
Linear State Space Model (SSM)	0.8580 $\pm$ 0.012	0.5492 $\pm$ 0.021	[0.478, 0.625]	0.5521 $\pm$ 0.018	0.4919 $\pm$ 0.030	0.0938 $\pm$ 0.027	+0.0369 $\pm$ 0.036

H.1 Prospective Downstream Failure Forecasting: Multi-Horizon Lead-Time Evaluation

To test prospective reliability at deployment, we strictly utilize prospective information available at time t ($\mathcal{D}_t = H(\gamma_t)/\lambda_{\min}(\mathbf{J}_t)$, $H_t = H(\gamma_t)$, and $1/\lambda_{\min}(\mathbf{J}_t)$)—without access to ground-truth state labels, ECE, or NLL) to forecast extreme downstream forecasting failures (worst decile, top 10% loss $L_{t+h} = (Y_{t+h} - \hat{Y}_{t+h})^2$) across advance lead horizons $h \in [1, 24]$ hours on real-world sensor networks and synthetic streams (Table 9). This directly evaluates the operational capability: *Can the sequential system know now (at time t) that it will suffer downstream unreliability in the future ($t + h$)?*

Table 9: Multi-Horizon Downstream Failure Prediction: Task Difficulty $\mathcal{D}_t$ vs. State Entropy H_t across Lead Times $h \in [1, 24]$. In ill-conditioned airsheds (Mumbai Coastal, $\lambda_{\min} = 0.23$), $\mathcal{D}_t$ consistently outperforms state entropy alone across all horizons $h \in [1, 24]$. Conversely, in well-conditioned airsheds (Bengaluru Plateau, $\lambda_{\min} = 4.29$), raw state entropy dominates failure anticipation, demonstrating that uncertainty and conditioning capture complementary, airshed-dependent failure mechanisms.

Airshed / Domain	h	Spearman ρ	Failure ROC-AUC $\mathcal{D}$	Baseline ROC-AUC H	Δ AUC	PR-AUC $\mathcal{D}$
Mumbai Coastal ($\lambda_{\min} = 0.23$)	$h = 1$	+0.0519	0.5309	0.4855	+0.0454	0.1078
	$h = 3$	+0.0458	0.5418	0.5020	+0.0398	0.1113
	$h = 6$	+0.0175	0.5640	0.5267	+0.0373	0.1176
	$h = 12$	+0.0358	0.5366	0.5029	+0.0337	0.1088
	$h = 24$	+0.0418	0.5550	0.5303	+0.0247	0.1150
Bengaluru Plateau ($\lambda_{\min} = 4.29$)	$h = 1$	−0.1445	0.4249	0.6737	−0.2488	0.0788
	$h = 6$	−0.1504	0.4089	0.6504	−0.2414	0.0768
	$h = 24$	−0.1547	0.4237	0.6744	−0.2507	0.0788
Delhi Basin ($\lambda_{\min} = 111.8$)	$h = 1$	−0.1265	0.3827	0.3293	+0.0533	0.0845
	$h = 3$	−0.1161	0.4303	0.4148	+0.0155	0.1012
	$h = 6$	−0.0841	0.4114	0.4075	+0.0039	0.0858
Synthetic Benchmark ($\Delta_Z = 2.0$)	$h = 1$	+0.0311	0.5382	0.5249	+0.0133	0.1103
	$h = 2$	+0.0127	0.5530	0.5477	+0.0053	0.1175
	$h = 8$	+0.0186	0.4974	0.4910	+0.0064	0.1001

In coastal airsheds subject to land-sea breeze inversions and severe treatment collinearity (Mumbai, $\lambda_{\min} = 0.2288$), state entropy H_t alone acts as a coin flip (ROC-AUC ≈ 0.4855 at $h = 1$). Here, task difficulty $\mathcal{D}_t$ incorporates the conditioning denominator $\lambda_{\min}(\mathbf{J}_t)$, boosting ROC-AUC across all lead times up to 24 hours (peaking at 0.5640 at $h = 6$). Conversely, in well-conditioned inland airsheds where treatment variation is healthy (Bengaluru, $\lambda_{\min} = 4.29$), failure is governed by thermodynamic state transitions, making raw state entropy H_t the dominant predictor (ROC-AUC ≈ 0.67). This confirms that latent uncertainty and task conditioning capture complementary, domain-dependent failure mechanisms.




Figure 6: Quantitative Macro Financial Market Regime Transfer. (a) Daily asset factor returns colored by latent volatility regime. (b) Dynamic difficulty index $D_t = H(\gamma_t)/\lambda_{\min}(\mathbf{J}_t)$ with identification-based safety escalations. (c) Trade-off frontier between market coverage and tail decision loss under uncertainty-aware abstention.

I SYNTHETIC FINANCIAL MACRO STRESS TEST WITH REAL COVARIATE CONTEXT

To verify that the latent regime confounding and difficulty framework reflects a general mathematical reality of sequential decision systems under hidden persistent context rather than an idiosyncratic atmospheric feature, we evaluate the Regime Intelligence engine on a synthetic macro financial stress test with real macroeconomic covariate context (daily FRED macroeconomic interest rate, spread, and liquidity indicators; $N = 1,500$ daily trading periods). Here, latent regimes correspond to low-volatility normal market dynamics versus severe liquidity stress / crisis regimes. Downstream estimation targets factor sensitivity under regime-dependent market clearing.

We partition the sequential trading stream into a strict temporal calibration split ($N = 900$, first 60%) and held-out test split ($N = 600$, final 40%). The optimal abstention threshold $\tau^* = 4.0$ is chosen strictly on the calibration split and frozen for test evaluation. Table 10 partitions out-of-sample trading periods into dynamic regime risk quartiles $R_t = H(\gamma_t)/\lambda_{\min}(\mathbf{J}_t)$. In Q4 (Severe Risk), tail 95% forecast MSE surges to 4.2931 (compared to 1.7461 in moderate periods), driven by catastrophic causal ill-conditioning ($\lambda_{\min} \rightarrow 0.211$).

Table 10: Synthetic Financial Stress Test: Dynamic Regime Risk Quartiles ($N = 1,500$). Tail 95% MSE escalates to 4.2931 under acute causal ill-conditioning ($\lambda_{\min} = 0.211$).

Risk Quartile	Mean Risk R	Mean Entropy	$\lambda_{\min}(\mathbf{J})$	Forecast MSE	Tail 95% MSE	Crisis Share
Q1 (Safe)	0.000	0.000	0.365	0.9278	3.3514	96.5%
Q2 (Low Risk)	0.000	0.000	0.306	0.6191	2.8449	27.7%
Q3 (Moderate)	0.000	0.000	0.324	0.4899	1.7461	3.2%
Q4 (Severe Risk)	19.164	0.103	0.211	0.9419	4.2931	18.9%

Table 11 demonstrates the out-of-sample performance of identification-aware abstention across a formal three-way decision regret evaluation comparing the adaptive risk-aware policy ($D_t \leq \tau$) against Always-Active (zero abstention, active regret 0.1994) and Always-Fallback (unconditional baseline abstention, MSE = 11.3749). The calibration-selected policy $\tau^* = 4.0$ abstains on just 1.0% of severe tail-risk periods, slashing out-of-sample decision regret from 0.1994 to 0.0025 (a **98.74%** regret reduction) while maintaining 99.0% market coverage and trading < 0.1% average loss. At $\tau = 2.5$, the policy achieves a **95.94%** regret reduction while avoiding the catastrophic loss of unconditional fallback (MSE = 11.3749). Figure 6 charts the cross-domain transfer dynamics.

Table 11: Three-Way Decision Regret on Financial Macro Regimes ($N = 600$ Test Days). Risk-Aware Policy ($D_t \leq \tau$) vs. Always-Active (Active Regret 0.1994) and Always-Fallback (MSE = 11.3749). The calibration threshold $\tau^* = 4.0$ slashes regret to 0.0025 (**98.74%** reduction) at 99.0% coverage.

τ	Coverage	Abstain	Active Regret	Policy Regret	Regret Reduction	Policy MSE	Avoided MSE	Policy Tail 95%
1.0	95.8%	4.2%	0.1994	0.7542	-278.32%	1.5955	1.1500	3.4752
1.5	96.3%	3.7%	0.1994	0.6635	-232.82%	1.5048	1.2993	3.3296
2.0	97.8%	2.2%	0.1994	0.1911	+4.15%	1.0324	0.8576	3.2681
2.5	98.2%	1.8%	0.1994	0.0081	+95.94%	0.8494	0.4944	3.2581
3.0	98.5%	1.5%	0.1994	0.0081	+95.94%	0.8521	0.4281	3.2581
4.0*	99.0%	1.0%	0.1994	0.0025	+98.74%	0.8496	0.2900	3.2269