

Integration is a linear functional

week 1 calculation

Ed Bueler

UAF Math 617 Functional Analysis

Spring 2026

welcome to functional analysis!

plan

each week starts with a concrete calculation, then the rest of the week will cover the theory supporting that calculation

- I will motivate functional analysis as the theory behind *partial differential equations*, *Fourier series*, and the *finite element method*
- most of the theory will be from the textbook, K. Saxe, *Beginning Functional Analysis*
 - when in doubt, assume that *material on the exams* comes from the theory and the textbook, not the more numerical/calculational parts
- this is an *experiment* . . . I have taught functional analysis before but not this way

Outline

- 1 continuity and integration
- 2 integration is a linear functional
- 3 numerical integration
- 4 theory this week

continuity and $C([a, b])$

- continuity and integration are fundamental ideas we need from analysis

Definition (continuity)

- a function $f : [a, b] \rightarrow \mathbb{R}$ is *continuous at* $x \in [a, b]$ if for all $\epsilon > 0$ there is $\delta > 0$ so that

$$|y - x| < \delta \text{ and } y \in [a, b] \implies |f(y) - f(x)| < \epsilon$$

- a function $f : [a, b] \rightarrow \mathbb{R}$ is *continuous* if it is continuous at each $x \in [a, b]$
- picture:

Definition

$$C([a, b]) = \{f : [a, b] \rightarrow \mathbb{R} : f \text{ is continuous}\}$$

- for example $a = 0 < 1 = b$: $C([0, 1])$
- note $C([0, 1])$ is an infinite set because it contains distinct elements $1, x^1, x^2, \dots$
- picture:

$C([0, 1])$ is vector space

Theorem

$V = C([0, 1])$ is a (real) vector space

Proof.

If $f, g \in V$ then $h(x) = f(x) + g(x) = g(x) + f(x)$ is continuous, so V is closed under addition. If $\alpha \in \mathbb{R}$ then $\alpha f(x)$ is continuous, so V is closed under scalar multiplication. The zero function is a continuous function, and $f(x) + 0 = f(x)$, so V has an additive identity. Also $1f(x) = f(x)$, so the scalar identity works. And distribution rules are true: $\alpha(f + g) = \alpha f + \alpha g$, $\alpha(\beta f) = (\alpha\beta)f$. $\square$

- “linear space” is a synonym for vector space
- wait . . . what is the definition of a *vector space*?

Definition (Riemann's integral)

for a mesh $a = x_0 < x_1 < \cdots < x_j < \cdots < x_n = b$, and evaluation points $x_j^* \in [x_{j-1}, x_j]$, define

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{j=1}^n f(x_j^*) (x_j - x_{j-1})$$

- this is *not* the only definition of an integral
- the limit process for an integral is more subtle than the limit process for continuity or derivatives
- we will get to Lebesgue's integral later; we will use it a lot, but only lightly cover its details

Theorem

if $f : [a, b] \rightarrow \mathbb{R}$ is continuous then $\int_a^b f(x) dx \in \mathbb{R}$ is well-defined

integration example

- picture:
- example: compute $\int_0^3 x \, dx$ from the definition

Theorem

if $f : [a, b] \rightarrow \mathbb{R}$ is continuous and if $F : [a, b] \rightarrow \mathbb{R}$ satisfies $F'(x) = f(x)$ then

$$\int_a^b f(x) dx = F(b) - F(a)$$

- example: redo integral on last slide

more integration examples

- example: compute $\int_0^1 x^k \frac{1}{1+x^2} dx$ for $k = 0, 1, 2$

claim: for $k \in \mathbb{Z}^+ = \{0, 1, 2, \dots\}$, the values $c_k = \int_0^1 x^k \frac{1}{1+x^2} dx$ are

$$c_k = \frac{\pi}{4}, \frac{\ln(2)}{2}, 1 - \frac{\pi}{4}, \dots = 0.78539, 0.34657, 0.21460, 0.15342, 0.11873, 0.09657, \dots$$

Outline

1. continuity and integration
2. integration is a linear functional
3. numerical integration
4. theory this week

linear functionals on $C([0, 1])$

Definition (linear functional)

a function $\ell : C([0, 1]) \rightarrow \mathbb{R}$ is a *linear functional* if

$$\ell[\alpha f + \beta g] = \alpha \ell[f] + \beta \ell[g]$$

for all $\alpha, \beta \in \mathbb{R}$ and $f, g \in C([0, 1])$

- example: $\ell[f] = f(0.4)$
- example?: $\ell[f] = 4f$
- example: $\ell[f] = 4f(0) + 2f(1)$
- example?: $\ell[f] = f'(0.4)$

integration against a weight

- example: suppose $w \in C([0, 1])$ is fixed. then

$$\ell[f] = \int_0^1 f(x)w(x) dx$$

is a linear functional on $C([0, 1])$

Proof.



- example?: $\ell[f] = \int_0^1 f(x) dx$
- example?: $\ell[f] = \int_0^x f(t) dt$

example

example. let $w(x) = \frac{1}{1+x^2}$. consider the linear functional

$$\ell[f] = \int_0^1 f(x) \frac{1}{1+x^2} dx$$

what are its values?

$$f(x) = x^k:$$

$$f(x) = \sin(x): \quad = 0.3217935447, \text{ says claudé}$$

$$f(x) = \sin(k\pi x):$$

questions about linear functionals

regularity question

we don't need f and w to be continuous for this to be well-defined:

$$\ell[f] = \int_0^1 f(x)w(x) dx$$

how general can these functions be? can they be discontinuous? completely arbitrary?

recovery question

suppose we think ℓ has a formula $\ell[f] = \int_0^1 f(x)w(x) dx$, but we don't know $w(x)$. can we recover $w(x)$ from the values $\ell[f]$ for various f ?

conceptual question

consider the space V' of all linear functionals on $V = C([0, 1])$. is it a vector space? (yes!) how do we *concretely* understand this abstractly-defined space? is each $\ell \in V'$ represented by a $w(x)$?

recovery question 1

- consider:

$$\ell[f] = \int_0^1 f(x)w(x) dx$$

- suppose $w(x)$ is a degree 4 polynomial, but we don't know which one.
how to recover $w(x)$?

recovery question 2

- consider:

$$\ell[f] = \int_0^1 f(x)w(x) dx$$

- suppose $w(x) = 1/(1 + x^2)$ in fact, but we don't know that. how to recover $w(x)$?

try $f(x) = x^k$?:

try $f(x) = \sin(k\pi x)$?:

recovery question 3

- consider:

$$\ell[f] = \int_0^1 f(x)w(x) dx$$

- suppose $w(x) = 1/(1 + x^2)$ in fact, but we don't know that. how to recover $w(x)$?

try hat functions on a uniform mesh:

Outline

1. continuity and integration
2. integration is a linear functional
3. numerical integration
4. theory this week

numerical integration rules

- **trapezoid rule** for m equal intervals on $[0, 1]$:

$$\int_0^1 f(x) dx \approx T_m(f) = \frac{h}{2} \left(f(x_0) + 2f(x_1) + \cdots + 2f(x_{m-1}) + f(x_m) \right)$$

where $h = 1/m$ and $x_j = jh$

- **Simpson's rule** for *even* m intervals, and same h, x_j :

$$\int_0^1 f(x) dx \approx S_m(f) = \frac{h}{3} \left(f(x_0) + 4f(x_1) + 2f(x_2) + 4f(x_3) + 2f(x_4) \right. \\ \left. + \cdots + 4f(x_{m-1}) + f(x_m) \right)$$

- ... there are many more rules, and many are better, but they have the same flavor ...

numerical integration is a linear functional

- suppose we list values of f in a column vector:
- then

$$f \approx \begin{bmatrix} f(x_0) \\ f(x_1) \\ \vdots \\ f(x_m) \end{bmatrix}$$

$$T_m(f) =$$

$$S_m(f) =$$

linear functionals in finite dimensions

- suppose we have a real, finite-dimensional vector space V
- suppose we have a basis for V : $\mathcal{B} = \{b_1, \dots, b_m\} \subset V$
- write $f \in V$ in this basis: for $f_j \in \mathbb{R}$,

$$f = f_1 b_1 + \dots f_m b_m$$

- the coefficients are usually written in a column vector:

$$(f)_{\mathcal{B}} = \begin{bmatrix} f_1 \\ f_2 \\ \vdots \\ f_m \end{bmatrix}$$

Theorem

if $\ell : V \rightarrow \mathbb{R}$ is a linear functional then there are values $\ell_j, j = 1, \dots, m$, so that

$$\ell[f] = [\ell_1 \quad \ell_2 \quad \dots \quad \ell_m] \begin{bmatrix} f_1 \\ f_2 \\ \vdots \\ f_m \end{bmatrix} = \sum_{j=1}^m \ell_j f_j$$

- in other words, we usually represent linear functions by *row vectors* when we are representing the vectors themselves by *column vectors*
- then $\ell[f]$ is computed by usual rules of matrix-matrix multiplication

Outline

1. continuity and integration
2. integration is a linear functional
3. numerical integration
4. theory this week

theory on Wednesday and Friday

definitions:

- metric space
- norms
- normed vector space
- inner products
- sequence spaces ℓ^p , and norms $\|(x_n)\|_p$ and $\|(x_n)\|_\infty$

= Chapter 1 in Saxe

Assignment 1 posted on bueler.github.io/fa