

 Lagunita is retiring and will shut down at 12 noon Pacific Time on March 31, 2020. A few courses may be open for self-enrollment for a limited time. We will continue to offer courses on other online learning platforms; visit <http://online.stanford.edu>.

Course > Introduction > Contents > Contents for Inference: Relationships $C \rightarrow Q$

 Bookmark this page

Contents for Inference: Relationships $C \rightarrow Q$

Case $C \rightarrow Q$

- Inference for Relationships Introduction
- Case $C \rightarrow Q$: Overview
- Case $C \rightarrow Q$: Independent and Matched Samples

Two Independent Samples

- Two Independent Samples: Overview
- Two Independent Samples: Hypotheses
- Two Independent Samples: Conditions and Two-Samples t-test
- Two Independent Samples: How to Read Statistical Software Output
- Learn By Doing Activity
- Two Independent Samples: Finding the p-value
- Two Independent Samples: Example
- Learn By Doing Activity
- Two Independent Samples: Confidence Interval
- Two Independent Samples: Summary

Matched Pairs

- Matched Pairs: Overview
- Matched Pairs: Idea Behind the Paired t-test
- Matched Pairs: Hypotheses

- Matched Pairs: Conditions and Paired t-test
- Matched Pairs: Finding the p-value
- Matched Pairs: Example
- Learn By Doing Activity
- Matched Pairs: Confidence Interval
- Matched Pairs: Summary

ANOVA

- ANOVA: Overview
- ANOVA: Hypotheses
- ANOVA: The Idea Behind the F-Test
- ANOVA: Conditions and F-test
- ANOVA: Finding the p-value
- ANOVA: Example
- Learn By Doing Activity
- ANOVA: Summary

Case $C \rightarrow Q$ Summary

- Summary (Case $C \rightarrow Q$)

Open Learning Initiative [🔗](#)



[🔗](#) Unless otherwise noted this work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License [🔗](#).

© All Rights Reserved