🛕 Lagunita is retiring and will shut down at 12 noon Pacific Time on March 31, 2020. A few courses may be open for selfenrollment for a limited time. We will continue to offer courses on other online learning platforms; visit http://online.stanford.edu.

Course > Inference: Hypothesis Testing for the Population Mean > Type I and Type II Errors > Wrap-Up (Type I and Type II Errors)

☐ Bookmark this page

Wrap-Up (Type I and Type II Errors)

It is important to remember to take into consideration the possibility of the occurrence of Type I or Type II error, when drawing conclusions from hypothesis tests. Thus:

- Whenever there is a failure to reject the null hypothesis, it is possible that a Type II error has occurred. Thus, it is concluded from the results of the study, that there are no significant differences, even though, in reality, there are significant differences.
- Whenever the null hypothesis is rejected, it is possible that a Type I error has been committed. That is, it is concluded from the results of the study, that there are significant differences, when, in reality, there are no differences.

However, it is not possible to know when either Type I or Type II error has occurred.

	We reject the H ₀ (accept H _a)	We fail to reject the H ₀ (not enough evidence to accept H _s)
H ₀ is true	Type I error	Correct Decision
H ₀ is false (H _a is true)	Correct Decision	Type II error

Open Learning Initiative

● Unless otherwise noted this work is licensed under a Creative Commons Attribution-

NonCommercial-ShareAlike 4.0 International License ...

© All Rights Reserved