

**Clinical Trial ID:**

**NCT00000424**

**Title:**

**Tidal Lavage in Knee Osteoarthritis**

**Summary:**

**This study compared the effects of tidal lavage (washing out) of the knee joint and an**

**imitation lavage procedure in people with knee osteoarthritis. In tidal lavage, the**

**doctor flushes out a knee joint with repeated injections of a mild salt solution, done**

**under local anesthesia. Study participants had to meet standard criteria for diagnosis of**

**osteoarthritis but could have low, medium, or high severity of x-ray changes indicating**

**knee osteoarthritis. We performed the lavage procedure once, and did quarterly followups**

**for 1 year. We permitted patients to use some other osteoarthritis treatments during the**

**study, such as non-narcotic pain relievers, nonsteroidal anti-inflammatory drugs, and**

**physical therapy.**

**Detailed Description:**

**This study was a randomized, double-blind, sham procedure-controlled evaluation of tidal**

lavage as a treatment for knee osteoarthritis. We achieved blinding by raising a drape during the study procedure and mimicking the sensory aspects of tidal lavage, but not flushing the knee, in the sham lavage group. We gave all study participants subcutaneous anesthesia with lidocaine and then bupivacaine and then attempted aspiration of the knee, removing up to 5 ml of synovial fluid for examination and then injecting the knee with 20 ml of bupivacaine.

In people who received the sham treatment, we placed a 16-gauge catheter in the lateral suprapatellar position just to the knee joint capsule and infused small volumes (less than 5 ml per "exchange" x 20) of saline into the subcutaneous tissue while manipulating the anterior knee to mimic efforts to shift fluid within the knee during the "aspiration" phase of each exchange. Patients could see the supply bag of sterile saline but could not see the waste bag of this closed drainage system.

For tidal lavage patients, we made the catheter puncture in the lateral suprapatellar

**pouch and repeatedly distended the knee with 30-50 ml of sterile saline, then aspirated**

**this volume (about 20 exchanges for a total of 1 liter of flush solution).**

**At the end of the procedure, we told the patients that their knee might be swollen due to**

**retained saline, and to expect this additional swelling to resolve over 24-48 hours,**

**during which time they were to minimize activity. The person who did the procedure (the**

**principal investigator) then left the room. The study nurse, who was not present for the**

**procedure and was blinded to the procedure's identity, asked the patients which treatment**

**they thought they received (tidal lavage or sham lavage). We scheduled patient followups**

**with this study nurse for 3, 6, and 12 months, and patients completed questionnaires at**

**each visit. Questionnaires were both arthritis-specific (WOMAC) and global (Quality of**

**Well-Being).**

**Eligibility Criteria:**

**Inclusion Criteria:**

- Knee pain attributed to osteoarthritis for at least 1 year.**

- **Meet American College of Rheumatology clinical or clinical plus x-ray criteria for knee osteoarthritis**
- **Have at least a moderate pain rating on at least one of the five Western Ontario-McMaster University Osteoarthritis Index (WOMAC) scales**

**Exclusion Criteria:**

- **Significant conditions of the spine, hips, or feet that affect the ability to walk**
- **Significant medical conditions that affect the ability to walk and function**
- **Inflammatory arthritis, such as rheumatoid arthritis or gout**
- **Degenerative arthritis secondary to other conditions, such as hemochromatosis, Wilson's disease, or ochronosis**
- **Current significant soft tissue rheumatism such as fibromyalgia, anserine bursitis, or trochanteric bursitis**

**Gender:**

**All**

**Minimum Age:**

**40 Years**

**Maximum Age:**

**N/A**

**Phase:**

**Phase 2**

**Conditions:**

**- Osteoarthritis, Knee**

**Interventions:**

**- Tidal lavage vs. sham lavage of the knee**

**Locations:**

**- Indiana University School of Medicine, Indianapolis, Indiana**