Euler's Cabinet of Curiosities

Physics fidelity report: how each of the 21 exhibits is simulated, and how it was tested against theory

Bullet Physics Playground (bpp), demo/WyomingWill

Every exhibit run afresh on 15 October 2026, and run again after the fixes described in section 2

21 exhibits	**252** checks against theory, tabulated
14 agree	**7** agree, with a numerical or open caveat

Contents

1. Summary **3**

2. What was found, and what was done **5**

2.1 Fixed: the simulations and their checks . . . 5

2.2 Fixed: the documentation . . . 5

2.3 Still open . . . 6

3. How the gallery is simulated and checked **7**

4. Wing A: Turning **8**

4.1 Gömböc variety . . . 8

4.2 Bille and the polyhedra . . . 11

4.3 Galileo's ramp . . . 14

4.4 The double cone . . . 16

4.5 The oloid and the sphericon . . . 18

4.6 Euler's disk . . . 20

4.7 The rattleback . . . 22

4.8 The spinning eggs . . . 24

4.9 The tippe top . . . 27

4.10 The Dzhanibekov effect . . . 30

4.11 The gyroscope . . . 32

4.12 The bicycle wheel . . . 34

5. Wing B: Swinging **38**

5.1 Newton's cradle . . . 38

5.2 Huygens' clocks . . . 41

5.3 Kapitza's pendulum . . . 42

5.4 The double pendulum . . . 45

5.5 Pendulums with a twist . . . 47

5.6 The chain fountain . . . 49

5.7 The falling Slinky . . . 53

5.8 The Brazil-nut effect . . . 55

5.9 The solar system . . . 58

1. Summary

Euler's Cabinet of Curiosities is a gallery of physics exhibits in the Bullet Physics Playground. Its point is to be faithful to the physics: each exhibit should do what the real thing does, for the reason the real thing does it, and say how it knows. This report collects, exhibit by exhibit, how each one is simulated and how it was tested against theory, with the numbers.

Every simulated number here comes from a run made for this report, on 15 October 2026, or from the exhibit's README, and the exhibit sections say which. Most theory values are worked out by the exhibits themselves from their own parameters; where a theory value, a fit or a comparison was worked out for this report instead (for example the Dzhanibekov flip period, the gyroscope's formula values, the fits to Galileo's ramp), the section says so. Each exhibit was run headless, from its own folder, with the start pause off (HOLD_S=0, which changes nothing in the physics; see section 3), and its console output was read against its README and its code.

A first pass of this report found several exhibits with real problems (section 2.1 lists them) and many places where the documentation said something the code didn't do. Those were fixed the same day, and every exhibit that changed was run again: the exhibit sections describe the gallery as it now is, and section 2 lists what was found, what was done about it, and what is still open.

Wing A: Turning

Exhibit	Checks	Headline result	Status
Gömböc variety	9	One stable point: Gömböc-C ends upright in 11 of 11 random drops; righting from a headstand 8.75 and 13.03 s against the earlier study's 8.8 and 13.2 s. Sloan shapes 0 of 16 upright in one run, 3 of 9 in a seeded re-run (study: 9%; small samples, and the bodies knock into each other).	**Agrees**
Bille and the polyhedra	11	Bille ends on its one stable face in 27 of 27 drops and 6 of 6 set-downs, by the predicted face paths (a quick crossing of a face is now logged, in brackets).	**Agrees**
Galileo's ramp	10	Finish times within -0.013% of sqrt(2d(1+k)/(g sin θ)); distances in equal times 1 : 2.998 : 4.997 (odd numbers; fitted for this report); first-order convergence in step size.	**Agrees**
The double cone	9	Rolls up or down exactly as the rail geometry predicts on all three tracks; centre's drop within 1 µm of the exact geometry; energy loss 0.1-0.3% (numerical).	**Agrees**
The oloid and the sphericon	8	Oloid's centre stays within 4.2426-4.5000 cm, the roller and sphericon level at 4.2426 cm, exactly as theory says; side-to-side weaves 3.6, 5.1 and 5.8 cm, now stated correctly.	**Agrees**
Euler's disk	12	Precession rate within 0.08-0.18% of theory from 25° down to 1°; with losses off, energy steady to the eighth figure. Its finite-time end follows its rolling-resistance law (now said), not Moffatt's air.	**Agrees**
The rattleback	11	Spun the wrong way it reverses (0.67 s), the right way never; the reference model's growth rates (8.9/s and 1.7/s at 1 turn a second) and the rolling-resistance window now stated as measured. New README.	**Agrees**
The spinning eggs	17	The left egg rises at 0.30 s and stands at 0.67-0.68 s at every step rate from 2400 to 38,400 (was 0.50 s to never); its rise from 60° to 30° within 8% of Moffatt and Shimomura (2% started in balance); Jellett's quantity changes by 0.8-1.7%; energy budgets converge with step. Open: how long it stays standing depends on the step.	**Agrees, with caveats**
The tippe top	11	Inversion criterion right for all three tops; centre of mass rises 1.057 to 2.545 cm as theory says; stem threshold about 80 rad/s, within the sleeping-top rule's bounds. Open: the green top's spin collapses after 40 s without rolling resistance.	**Agrees, with caveats**
The Dzhanibekov effect	12	Flip period 3.3129 s against 3.3131 s from the exact elliptic-function solution (-0.01%), 36 flips in 120 s. Energy and angular momentum are held by projection, and the README now says so. New README.	**Agrees**

Exhibit	Checks	Headline result	Status
The gyroscope	13	Precession 2.58 rad/s against m g r/(I ω) = 2.59 at the spin it has, and to 0.0-0.5% with the tilt term as it sags; README now right (2.56 rad/s at 100 turns/s). Rocked 1.5° it slides further than aimed for, cause not separated.	**Agrees, with caveats**
The bicycle wheel	8	Stool's turning matches angular-momentum conservation exactly with friction off; L kept to 0.093% over 120 s. A guard now reports and recovers from a NaN blow-up, whose cause (seen once) is still unknown.	**Agrees, with caveats**

Wing B: Swinging

Exhibit	Checks	Headline result	Status
Newton's cradle	13	First strike the same at every step rate from 1200 to 9600 steps/s (was: a third of the energy lost at 9600); momentum to 0.00%; ball 4 gets 13.6% (Herrmann & Seitz: ~12%); the row settles over minutes, as real cradles do.	**Agrees**
Huygens' clocks	8	All three starts lock in anti-phase at 180.0°, at 0.9985-0.9988 Hz against 0.9984 Hz from theory; at 9600 steps/s the same to 1e-4 Hz (one start, 45 s).	**Agrees**
Kapitza's pendulum	9	Stability threshold 0.783 cm bracketed (0.76 falls, 0.80 stands); rocking period 1.133 s against 1.135 s for its 7.6° swing (-0.2%), now at a converged 19,200 steps/s (was 4800, 1.7% short).	**Agrees**
The double pendulum	16	Energy to 2e-11; half-step twin 64 fm off at 1 s; Lyapunov 3.42/s; normal modes to 0.000% and 0.135%; Floquet window 0.728-1.356 cm. The throw-over time of the too-hard chain (not the fact) isn't converged, and the README says so.	**Agrees, with caveats**
Pendulums with a twist	12	No flip below 70.53°; at μ = 3 the hidden integral stays 0.7500000000 (wanders at 2.8 and 3.2); spring's growth 0.526/s against 0.527 (Floquet).	**Agrees**
The chain fountain	21	Independent of the display speed; empties at every bend. With the push the falling chain carries, Biggins and Warner's relations hold. Run with every link simulated, the chain still passes none of the pile's push on (so that loss is the model's, not the shortcut's), but its fountain is lower: the exhibit's shortcut makes it 1.2x (30°) to 2x (60-90°) too tall. A faithful mode costs 5-8x per frame.	**Agrees, with caveats**
The falling Slinky	13	Bottom stays put 0.2038 s against sqrt(M/3k) = 0.2018 s: 0.6% from having 86 separate coils (an exact 86-coil chain waits 0.2030 s) and 0.4% from Bullet's step, both now explained in the README.	**Agrees**
The Brazil-nut effect	12	Qualitative: the big bead rises and the convection runs the right way. Floor push 0.18 of the bed's weight against 0.20 rigid. Lift-off at 46-48° against 19.5° for a rigid bed is mostly Bullet's soft contacts (stiffer: 28-30°), not converged.	**Agrees, with caveats**
The solar system	17	Planets within 5 km of JPL's DE421 after 10 years, the Moon 39 km; Mercury's perihelion 42.981″/century against 42.981″; rotation checks hold. The checks now stop at their exact dates, and give the same lines at any rate.	**Agrees**

2. What was found, and what was done

A first pass of this report, earlier on 15 October 2026, ran every exhibit afresh and read its output against its README and code. What it found was fixed the same day, and the exhibits that changed were run again; the numbers in sections 4 and 5 are from those runs. First the fixes to simulations and their checks, then to the documentation, then what is still open.

2.1 Fixed: the simulations and their checks

- **Newton's cradle.** At 9600 physics steps a second the first strike lost a third of its energy. The cause: before each step, only contacts that would still be squeezed at the step's end if the balls moved freely were worked out, so a contact that was squeezed but already opening was skipped, and Bullet let the balls pass through, losing the stored compression. Now any contact squeezed now or touching within the step is worked out, and each ball is handed back on its hinge's circle, turned with its threads. The first strike is the same at every step rate from 1200 to 9600. The fast merging of the row (in about 15 s) was the same energy leak: now, with no damping the row never merges, and with the default damping it settles over minutes (its relative swing halves in about 80 s), as real cradles do (Hutzler et al.: 1-2 minutes).
- **The chain fountain.** Links were woken and frozen once per drawn frame, so the physics depended on the display speed. All of that now runs per physics step, with a fixed 1/1800 s step; the output is identical at 0.25x, 0.5x and 1x. The 60° stall: links woken ahead of the pick-up had free ends and were dragged off, until the whole coil was loose in the jar and the hanging chain rested on the frozen links below; the last woken link is now held to the unmoved coil, and landed links stop colliding. It empties at every bend. A theory check against Biggins and Warner (2014) is now printed, and the README's height table is re-measured. A second pass found the coil laid with joints bent past their limits (304 of 1054 at 20°, plus upright links between layers), which Bullet snapped back when woken, kicking the chain: that, not the stiffness, was why the stiff chain's fountain was higher. The coil is now laid within the limits, the speed is the rate links land, and the pile's push can be measured directly (CF_PUSH=1).
- **The spinning eggs.** From 3° the left egg sat at the edge where friction holds its contact still, and the step's small errors decided when it broke free: 0.50 s at 4800 steps a second, 2.45 s at 19,200, never at 28,800 and 38,400. It now starts at 6°, twice that threshold, and rises at 0.30 s at every step rate; a new console check compares its rise with Moffatt and Shimomura's rate.
- **The solar system.** The yearly and monthly checks ran on the first frame after their date, and at high rates reached back to it inaccurately. The integration now stops at each check's exact date, runs the check, and goes on; and a second rate dependence in the moons' stepping was found and fixed. At 5, 60 and 100 days a second the check lines are now identical; at 100 days a second, Mimas on 2026-02-01 went from 4688 km to 41, Mercury in 2036 from 1220 km to 1. The README's samples are fresh.
- **Kapitza's pendulum.** At its old default of 4800 steps a second the rocking period was 1.7% short of converged, and that error happened to cancel the large-swing correction. The default is now 19,200 steps a second (converged to 0.1%), the console also compares with the period for the actual swing (1.133 s against 1.135 s), and a restart that misreported the size of the rocking (8.0° instead of 7.6°) is fixed.
- **The gömböcs and the Billes.** Their attendant's lift ran during the start pause, so with the pause on their output differed. The lift now waits for the pause to end, and the pause changes nothing in them either (checked with fixed seeds).
- **The bicycle wheel.** A guard now checks every body's state each frame; if anything is not a number it prints the full state, the time and the move under way, and restarts the run. Pressing R mid-turn no longer prints a stray event.

2.2 Fixed: the documentation

- **The gyroscope.** Precession at 100 turns a second now 2.56 rad/s (was 2.67, not what its own parameters give); off-wire times re-measured (rocked 1°: 62.0 s; 1.5°: 49.0 s; 2°: 12.7 s; 3°: 5.9 s; still wire 55.0 s, or 48.2 s on a rough foot); the turns-a-second it comes off at stated from runs (38, 45, 52).
- **The oloid.** The weaves are now 3.6, 5.1 and 5.8 cm (was '4 to 6 cm' for all three).
- **The rattleback and the Dzhanibekov effect.** Each has a README now. The rattleback's rolling-resistance window and growth rates, and the Dzhanibekov header's figures for Bullet's own modes (-31.9% energy, -17.5% angular momentum), are as measured; both run from the repository root.
- **The tippe top.** Rolling resistance never stops the red top turning over (0-300 μm); what it does do is now stated. The stem threshold is about 80 rad/s; the README's example is a fresh run.
- **The falling Slinky.** Its 1% offset is explained: 0.6% from 86 separate coils (an exact chain of 86 point masses waits 0.2030 s; its excess over the continuum's 0.2018 s halves as the coils double) and 0.4% from Bullet's step. On the stairs, nearly one walk in two goes wrong at the normal speed (12 of 21 clean).
- **Euler's disk, the spinning eggs, the double pendulum, Brazil nut.** Euler's disk says its end follows its rolling-resistance law, not Moffatt's; the eggs' README gives the energy budget by step size, egg by egg; the double pendulum's says the throw-over time isn't converged and the Lyapunov comparison is an order-of-magnitude one; the Brazil-nut's says its late lift-off is mostly soft contacts.
- **Comments and help text.** Huygens' clocks (the in-phase mode is slower), Kapitza (env options), double pendulum and pendulums with a twist (stale figures), chain fountain help, Gömböc comments and reconstruction notes (resting tilt, where a second basin first appears; the reconstruction README notes the findings PDF's own inconsistency), Bille (its step depends on the speed setting).

2.3 Still open

- **The bicycle wheel.** The cause of the NaN blow-up seen once in the GUI is unknown; the guard will leave a full diagnostic if it happens again.
- **The chain fountain.** With every link simulated, rigid capsule links lifted from a pile still pass none of the pile's push (0.20-0.49 m v^2) into the falling chain; why, and whether real bead chains do better (Biggins and Warner's h2 = 0.14 h1 implies an effective α of about 0.12), is open. The exhibit's shortcut (10 links woken ahead, the last held to the unmoved coil) keeps the chain off the rim and makes the fountain 1.2x (30°) to 2x (60-90°) taller than the full simulation (8.6, 6.3, 5.5 cm at 30°, 60°, 90°). Four live top layers (CF_WAKE=layer) reproduce the full chain at 30° and 60° but cost 21 ms a frame at quarter speed against 4.4 ms now, and none matches at 90°; the default is unchanged and the README says the fountain reads up to twice too tall.
- **The tippe top.** With no rolling resistance, the green top's spin collapses from 25.6 to 3.2 turns a second between 38 and 46 s, far faster than its spin friction allows. Not investigated.
- **The spinning eggs.** Once standing, how long the left egg stays up depends on the step (it starts to fall back at 22.3, 20.6 and 17.6 s at 4800, 9600 and 19,200 steps a second): the same slow first-order tip, which a second-order rolling contact should remove.
- **The gyroscope.** Rocked 1.5°, the tightrope gyroscope wobbles wider (9-17°) and slides out to 14-15 cm against the 9 cm aimed for; the wobble pushing the foot along is the likely cause, not yet separated from the slope's push.
- **The Brazil-nut effect.** The lift-off phase isn't converged even at the stiffest contacts tried (28-30° at 9600 steps/s with 60 iterations, against 19.5° for a rigid bed); stiffer runs are too slow to show live.
- **The double pendulum.** The too-hard chain's throw-over time isn't converged (3.96 s at 4800 steps/s, 4.46 s at 9600); only that it is thrown over is.
- **The falling Slinky.** Kept at 4800 steps a second for speed, so its 0.4% step error is documented rather than removed.

3. How the gallery is simulated and checked

How the exhibits are simulated

Bullet (the Bullet Physics library, through bpp) carries the rigid bodies, joints and ordinary contacts. Where Bullet's contacts can't do the physics faithfully, the exhibit works it out itself and hands the result to Bullet each step: Newton's cradle integrates Hertz's law between the balls in quarter-microsecond sub-steps; the oloid and the spinning eggs work out their own contacts; the tippe top and the rattleback advance their spin by Euler's equations (RK4) with Bullet's gyroscopic term off; the gyroscope applies its own gyroscopic torque (its rotor is a number, a gyrostat, not a body spun at 100 turns a second). Some exhibits don't use Bullet's dynamics at all, and Bullet only draws them: Euler's disk integrates the exact rolling equations; the double pendulum, pendulums with a twist and the solar system integrate their exact equations of motion by Bulirsch-Stoer (Gragg's modified midpoint, extrapolated, tenth order), with RK4 for the shaken pendulums and Wisdom-Holman for the moons. Each exhibit's section gives its integrator, its step and its parameters.

What 'tested against theory' means here

Most exhibits print, as they run, the quantities theory predicts, next to what the simulation did: periods, rates, thresholds, conserved quantities, positions against an ephemeris. The theory values are worked out by the exhibit from its own parameters (so a change of slider changes both), or, for the solar system, taken from JPL. Some don't, or only partly: the rattleback's console is qualitative (its numbers are in a separate reference model), and the Brazil-nut effect is mostly qualitative; for these, and wherever else it helped, the comparison was made for this report and is marked so. Where possible a check is also run at a different step size, to see that the agreement isn't a numerical accident; the sections say where that was done and where it wasn't.

The hall and the exhibits on their own

Each exhibit runs in its own physics world inside the hall, and the hall only runs the one you're at. Every exhibit's console output in the hall is compared, line for line, with its output when run on its own, and they are identical. This was checked for each exhibit as it was added, and again, for this report, for every exhibit whose console changed in the fixes (Newton's cradle, the chain fountain, the solar system, Kapitza's pendulum, the Brazil-nut effect, the bicycle wheel, and the exhibits whose text changed). The exceptions are the Gömböc and Bille drops, which are seeded from the clock and so differ from run to run anywhere.

The start pause

Each exhibit stands still for half a second whenever it starts, so you see how it begins. During the pause bpp's own time step is zero and the exhibit's step is skipped, so the pause changes nothing in the physics: with the pause on, the output is the output without it, delayed, line for line. This was checked for all twenty exhibits that have it when it was added (at 2 s), and at 0.5 s for every exhibit changed in the fixes, including the gömböcs and the Billes with fixed seeds (their attendant's lift now waits for the pause to end). The runs in this report were made with it off.

Rendering and sound

Bodies the physics moves are drawn where the physics has them. Some things are drawn without being simulated, and the sections say so: the chain fountain's coil in the jar until it is picked up and its links once they land, the falling Slinky's pile once it has landed (laid to rest), the cradle's threads, the bicycle figure's arms, the cords of the swinging Atwood machines and springs (drawn as beads). The solar system's maps of each surface are drawn, not photographs (their markings placed where they are on the real body; the turning itself is the IAU's model, and is checked), and the gyroscope's tilt gauge turns twenty times as far as the wire does. Newton's cradle's clack is computed from the same Hertz strike, as the sound a jerked steel sphere sends into the air.

4. Wing A: Turning

4.1 Gömböc variety

Agrees with theory. One stable point: Gömböc-C ends upright in 11 of 11 random drops; righting from a headstand 8.75 and 13.03 s against the earlier study's 8.8 and 13.2 s. Sloan shapes 0 of 16 upright in one run, 3 of 9 in a seeded re-run (study: 9%; small samples, and the bodies knock into each other).
Folder: demo/WyomingWill/Gomboc/

What it shows

A Gömböc is a convex body of uniform density with exactly one stable and one unstable equilibrium, so however it is set down it rolls back to the same upright pose. The table drops up to 12 bodies at random orientations, cycling through three shapes: the real Gömböc (Gomboc-C, at 9 cm and at 75% and 125% size) and two of M. L. Sloan's analytic Gömböcs (β = 0.07 and 0.08). The lesson is that having one resting point in theory is not enough. Gomboc-C rights itself every time, while the gently sloped Sloan shapes usually stall on their sides under ordinary rolling resistance.

Model

Bullet handles contact. Each body is a `btConvexHullShape`:

- **Gomboc-C:** the convex hull of a 1.2 mm tessellation of the GOMBOC900 CAD B-spline surfaces (13,476 points, per gomboc-drop-c.lua).
- **Sloan shapes:** hull points of $r^4 = 1 + 4\beta \sin\varphi \cos(\theta - 5\varphi)$.
- **Margin:** 0.04 cm, scaled with size.
- **Mass and inertia:** exact (CAD or analytic) at resin density 1.2 g/cm³. That gives 323 g at 9 cm, 136 g at 75%, 630 g at 125%, and 407 g / 401 g for β = 0.07 / 0.08.

The script steps the world itself at 1200 steps per simulated second (20 substeps of 1/1200 s per 1/60 s frame), using Bullet's semi-implicit Euler. Before every substep it applies two losses as constant moments at the contact:

- rolling resistance $M = \delta \cdot m \cdot g$, with lever arm δ = 150 µm by default;
- spin friction of 0.7 rad/s².

Other settings: friction 0.5 (Bullet multiplies it by the table's 0.8), restitution 0.1 (table 0.2), air drag 0, no taps. The tray is flat with 8° ramps at the edges and a rim. Status thresholds: "Upright" means within 15° of the resting normal for 2 s. "At rest" means less than 0.2° of turn and 0.02 cm of movement over 2 s. Five seconds after a body comes to rest it is lifted away and the next shape is dropped (GV_REDROP).

Theory checks

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Gomboc-C rests only on its one stable point	1 stable + 1 unstable equilibrium (CAD surface checked, code comment)	random drops: 11 of 11 runs that came to rest were upright, tilt 0.4–7.9°; 60 s tally: 8 Gomboc-C runs, 0 not upright	100% upright
Gomboc-C from a headstand (balanced on the unstable point, 1–3° nudge)	must leave the unstable point and right itself	3 of 3 at rest upright: 0.3°, 4.0°, 12.5° (the last propped against a neighbour)	consistent
Sloan β 0.07 / 0.08 self-righting at 150 µm rolling resistance	one basin (mathematically monostatic); findings PDF Table 5b: 14 of 157 (9%) upright	drops: 13 of 13 runs at rest NOT upright (18–142°); headstand 0 of 3 (71–138°)	0/16 vs 9% expected (about 1.4 of 16); small sample

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Residual tilt of Gomboc-C at rest	0° ideal; with a constant rolling moment it stops short. gomboc-drop-c.lua comment, corrected 15 Oct 2026: "0.3-8 deg ... (7.9 deg the most seen, on the 75% one in Gomboc Variety: the smaller, the further off)" (was "0.3-7 deg")	report run 0.3–7.9° (excluding the propped 12.5°); seeded re-run 15 Oct 2026: 0.3–4.6°	within the corrected range
Time to right from a headstand, 75% / 100% / 125%	findings PDF Table 6: 8.8 / 13.2 / 14.3 s	8.75 / 13.03 / 2.73 s (125% was knocked by a neighbour)	75% and 100% within 0.2 s
Mass at 100 / 75 / 125%	V·ρ = 268.851 cm³ × 1.2 g/cm³ = 322.6 g; × s³: 136.1 / 630.1 g	323 / 136 / 630 g	exact
Hull's false resting faces (Gomboc-C)	ideally none	code comment: 34 hull resting faces, all within 16° of true rest, deepest needs a 0.53 µm lift	— (documented)
Height of balancing point above resting point	findings Table 5: Gomboc-C 11.3 mm; Sloan β 0.07 / 0.08: 4.8 / 5.1 mm	— (documented)	—
Energy drift with all losses off (step-size check)	no gain	gomboc-drop-c.lua comment: at 1200 steps/s "energy stays level, with no steady gain"; 2400 "no better"; 600 also no steady gain	documented, not rerun

The 60-second tally printed by this run (time to rest, median and range): Gomboc-C 2 runs, 1.2 s (1.2–6.4). At 125%, 3 runs, 17.2 s (6.4–22.0). At 75%, 3 runs, 8.3 s (5.0–11.0). β 0.07: 5 runs, 3.1 s, 5 not upright. β 0.08: 5 runs, 5.7 s, 5 not upright.

Run used (random seed from the clock; one run each, so the numbers vary)

```
cd demo/WyomingWill/Gomboc && HOLD_S=0 stdbuf -oL timeout 3000 bpp -f gomboc-variety.lua -n 4200   # 70 s, 6 bodies,
    redrop 5
HOLD_S=0 GV_START=headstand GV_REDROP=0 timeout 3000 bpp -f gomboc-variety.lua -n 3600             # headstands
```

(A 9000-frame run was started but stopped: on the shared 2-core machine it ran at well under 1/5 of real time.)

```
  Orange  at rest after  17.15 s, 0.5 deg off upright -- Orange (Gomboc-C at 125%)
  Yellow  at rest after   8.30 s, 7.9 deg off upright -- Yellow (Gomboc-C at 75%)
  Purple  at rest after   2.25 s, 133.6 deg off upright  ** NOT UPRIGHT ** -- Purple (Sloan beta 0.07)
=== 60 s: 18 finished runs ===
    Gomboc-C at 125%             3 runs, time to rest median  17.2 s (6.4-22.0)
    Gomboc-C at 75%              3 runs, time to rest median   8.3 s (5.0-11.0)
    Sloan beta 0.07              5 runs, time to rest median   3.1 s (1.9-11.1), 5 not upright
    Sloan beta 0.08              5 runs, time to rest median   5.7 s (1.9-13.1), 5 not upright
  Yellow  upright after   8.75 s -- Yellow (Gomboc-C at 75%)
  Red     upright after  13.03 s -- Red (Gomboc-C)
  Red     at rest after  16.05 s, 0.3 deg off upright -- Red (Gomboc-C)
  Green   at rest after  16.22 s, 137.5 deg off upright  ** NOT UPRIGHT ** -- Green (Sloan beta 0.07)
```

Seeded re-run after the fixes (15 Oct 2026):

```
GV_SEED=3 GV_COUNT=4 GV_REDROP=1 HOLD_S=0 bpp -f gomboc-variety.lua -n 3700
=== 60 s: 13 finished runs ===
    Gomboc-C                     3 runs, time to rest median  12.8 s (11.8-14.0)
    Gomboc-C at 125%             2 runs, time to rest median   6.4 s (2.6-10.3)
    Gomboc-C at 75%              2 runs, time to rest median   5.1 s (3.0-7.1)
    Sloan beta 0.07              3 runs, time to rest median   3.5 s (1.6-5.1), 2 not upright
    Sloan beta 0.08              3 runs, time to rest median  10.4 s (7.2-20.3), 1 not upright
```

In the 60 s tally 1 of 3 β 0.07 and 2 of 3 β 0.08 runs ended upright; over the whole run (every body that came to rest, first report each) 1 of 4 β 0.07 and 2 of 5 β 0.08. So in this seeded run the Sloan shapes sometimes righted themselves, unlike the report's 0 of 16. 3 of 9 is more than the PDF's 9% (Table 5b) would make likely (a 4% chance), 0 of 16 less (22%); the samples are small and bodies here knock into each other, so neither run tests the 9% well. One β 0.07 body (Green, run 1) was re-reported at rest five times (4.4 to 51.6 s, creep) and so never entered the tally.

Pause and hall checks after the lift fix (15 Oct 2026): with `GV_SEED=3 GV_COUNT=4 GV_REDROP=1`, HOLD_S=0 for 900 frames is line for line a prefix of the default half-second pause for 930 frames (31 lines, lifts and redrops inside the window); hall against standalone (cmp2.sh, 1200 frames): IDENTICAL (32 run lines).

Known departures and honest caveats

- Fixed 15 Oct 2026: the attendant's lift ran during the start pause (a `holdTick` that called `liftStep`), so with the pause on the output differed. `holdTick` is removed; the lift waits for the pause to end and the pause changes nothing (checks above).
- **Rolling resistance and spin friction are script-side models.** They are constant moments, not Bullet physics. δ = 150 µm and 0.7 rad/s² were chosen, not measured. The findings PDF (Table 5b) shows that the outcome depends strongly on δ: without rolling resistance nothing ever settles, Gomboc-C included.
- **Gomboc-C stops a few degrees short of upright** (here up to 7.9°). The constant moment holds it wherever gravity's restoring lever is smaller than δ, and its valley is flat. The code says this is real behaviour.
- **The Sloan shapes are not strictly convex at these β.** They roll on their convex hull. The findings PDF cites arXiv:2604.17120, which finds strict convexity only below β ≈ 0.036. The hull gains a second basin at higher β. Re-checked 15 Oct 2026 by re-running the reconstruction's basin scan (scanb.py's method, form 1, 25000 surface points) at finer β: deepest extra basin 5.9 µm at 0.08 (the hull facets' own noise is ~5 µm), 11.9 µm at 0.085, 25.3 at 0.09, 43.0 at 0.095, 66.7 at 0.10 (the PDF's Table 5 gives 67 at 0.10 and none at 0.08). The lua comment ("first found at beta 0.085") was consistent with this; it now gives the depths and notes that the PDF rounds the limit to "above β ≈ 0.09". The scan result is recorded in variety-reconstruction/README.txt. The PDF was not edited.
- **The β 0.08 resting point:** the lua says its lowest point is ~35° from the formula's minimum; computed 15 Oct 2026 from the `down` and `top` vectors in gomboc-meshes/beta1-08.lua: 34.6° (β 0.07: 4.6°). The findings PDF §4 says the Sloan shapes' "true resting orientation lies 4–5° from the formula's minimum", which holds for β 0.07 only; its own §5 says the balancing point sits ~35° off the top on β 0.08. This inconsistency inside the PDF is now noted in variety-reconstruction/README.txt (there is no Gomboc README.md). The PDF was not edited.
- **Re-reported rests (creep).** Several bodies were reported "at rest" again and again: Yellow #3 (125%) five times between 19.6 and 32.4 s, and one β 0.07 three times. Each time the body had moved more than 0.2° or 0.2 mm within 2 s after being declared at rest, which matches the contact-solver creep admitted in gomboc-drop-c.lua ("up to ~0.3 deg and ~0.7 mm per minute"). The tally uses the last of these times.
- Fixed on 15 Oct 2026, tally median: with an even number of runs the "median" used to print the lower middle value; it is now the mean of the two middle values, in gomboc-variety.lua and gomboc-drop-c.lua. Seeded re-run: "Gomboc-C at 125% 2 runs, time to rest median 6.4 s (2.6-10.3)" = mean of 2.58 and 10.27 s.
- **Collisions between bodies** (props, knocks) affect individual results. The console flags them as "touching".
- Fixed on 15 Oct 2026: variety-reconstruction/README.txt listed meshes beta1-06 and beta2-04, which are not in gomboc-meshes/. It now names the two kept (beta1-07, beta1-08), says gen.py's __main__ makes a different larger-β set, and that the kept files' `down` is the lowest face of the hull's one basin, not the formula minimum gen.py writes. Sloan's form 2 is left out by design.
- Statistics are from one short run. The Sloan upright fraction (0/16) is compatible with the 9% in the findings PDF, given the small sample.

References

- Kiss Attila, GOMBOC900 SolidWorks CAD model (2013), as cited in code.
- M. L. Sloan, "An Analytical Gomboc", arXiv:2306.14914 (2023); MathWorld, "Gömböc".

- "Sloan's Analytical Gömböc at Published β: A Strict-Convexity-Constrained Reanalysis", arXiv:2604.17120 (2026), as cited in the findings PDF.
- P. L. Várkonyi, G. Domokos, "Mono-monostatic bodies: the answer to Arnold's question", Math. Intelligencer 28 (2006).
- G. Domokos, P. L. Várkonyi, "Geometry and self-righting of turtles", Proc. R. Soc. B 275 (2008).
- W. M. Spears with Claude, "One resting point is not enough" (gomboc-findings.pdf, 9 Oct 2026, working notes).

4.2 Bille and the polyhedra

Agrees with theory. Bille ends on its one stable face in 27 of 27 drops and 6 of 6 set-downs, by the predicted face paths (a quick crossing of a face is now logged, in brackets).
Folder: demo/WyomingWill/Bille/

What it shows

These are bodies that rest on only one face because of where their mass is, not because of their shape alone.

- **Bille** is the monostable tetrahedron of Almadi, Dawson & Domokos (2025): a hollow carbon-tube frame with a dense tungsten-carbide wedge, so that set down on any face it tips, face to face, onto face D. It has one stable face and two unstable corners.
- **The spiral polyhedra** (after Domokos & Kovács 2023) have 21, 26 and 37 corners, each with equal weights at the corners. Each has one stable face (the base) and one unstable vertex (the apex). Their bands of side faces tip one onto the next: … → S2 → S1 → base.

Bodies are dropped at random orientations or set down on a wrong face, and the console prints the faces each one passed through.

Model

Bullet handles contact through each body's convex hull: the 4 vertices for Bille, the corner points for the polyhedra. Mass and principal inertia are taken from the data files:

- **Bille at size 1:** 120 g, I = (0.219, 1.225, 1.321) kg·cm². The collision margin is the real tube radius, 0.05 cm.
- **Polyhedra:** scaled 4.5× to 40 cm tall, with mass scaling as size³.
- **Per-colour sizes:** 1.0, 0.6, 1.3, 0.8, 1.15 and 0.7.

The losses are applied by the script before every physics step, as constant moments: rolling resistance with lever arm δ = 50 µm (not scaled with size) and spin friction of 0.7 rad/s². Other settings: friction 0.5 (× table 0.8), restitution 0.1, no air drag.

Time step. Requested: 1200 steps per simulated second (slider), but at least one step per frame. Actual, from the code `NSTEPS = max(1, floor(steps·speed/60 + 0.5))`:

- At the default 1/50 slow motion (speed 0.02), NSTEPS rounds to 1, so the step is 1/3000 s.
- At speed 0.1 it is 1/1200 s.

Detection thresholds. A face counts as "lain on" when its normal is within 2° of straight down for 0.03 s. "At rest" means less than 0.2° and 0.03 cm of movement over 0.4 s. An edge counts as "rolled on" when exactly two of its corners are within 0.15% of the body's size of the table.

Theory checks

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Bille rests only on face D (random drops)	monostable on D (paper; reconstruction puts the COM 0.81 mm inside the face-D loading zone, which is at most 1.48 mm deep: README.txt)	27 of 27 drops on D (6 at speed 0.02, 21 at speed 0.1); tallies "16 runs, 16 on the stable face"	100%
Bille set down on a wrong face	ends on D	6 of 6 on D	100%
Falling pattern from C	C → D (quasi-static rule)	2 of 2: C → D, over edge AB (= C∩D)	matches
Falling pattern from A	A → D	1 of 1: A → D, over edge BC (= A∩D)	matches
Falling pattern from B	B → A → D	report run: 3 of 3 logged as "B → D" (edges CD = B∩A then BC = A∩D). Since 15 Oct 2026 a face crossed in under 0.03 s is logged in brackets: re-run (BD_SEED=5 BD_BODIES=1 BD_COUNT=3 BD_START=setdown) logs Green "B -> (A) -> D; edges it rolled on: CD, BC"	path now matches in the face log too
Polyhedra rest only on the base (drops)	one stable face	21-corner 4/4 (incl. one in the mixed run), 26-corner 2/2, 37-corner 4/4	10/10
Polyhedra set down on a band	S_n → ... → S1 → base	37-corner: S6→S5→S4→S3→S2→S1 (then rocks S2/S1/base); 26-corner from S4: S4 → base; 21-corner from S1: S1 → base	all end on base; order never reversed except brief rocking (e.g. 26-corner drop S1→S2→S1→base)
Bille time to rest after drop	header: a single face-to-face tip takes ~1/20 s	median 0.39–0.41 s (tallies), range 0.36–0.69 s	—
Polyhedra time to rest (40 cm scale)	findings Table 6 at 9 cm: 4.3 / 2.3 / 2.5 s (21/26/37)	21: 11.15–16.22 s; 26: 4.09–5.16 s; 37: 4.33–6.01 s	not comparable directly (scaled 4.5×, sizes 0.6–1.3)
Masses	120 g × s³; polyhedra 109 / 142 / 201 g at 9 cm (findings) × (4.5 s)³	Bille 120 / 264 / 183 g at s = 1 / 1.3 / 1.15; 21-corner 9951 g at s=1 (109.2 g at 9 cm), 26: 28362 g (141.7 g), 37: 27878 g (201.1 g)	exact
Earlier statistics (README)	—	code header: Bille "re-dropped 346 times, always ended on face D"; findings: polyhedra 116/117 on base (26-corner: 37/38)	documented

Run used (random seeds from the clock; one run each)

```
cd demo/WyomingWill/Bille && HOLD_S=0 timeout 1500 bpp -f bille-drop.lua -n 9000                    # speed 0.02, 3
    s simulated
HOLD_S=0 BD_SPEED=0.1 bpp -f bille-drop.lua -n 9000                                                 # 15 s simulated
HOLD_S=0 BD_START=setdown BD_COUNT=6 BD_REDROP=0 bpp ... -n 3600   (speed 0.02, and again with BD_SPEED=0.1)
HOLD_S=0 BD_SPEED=0.1 BD_BODIES=2 BD_COUNT=6 BD_REDROP=1 bpp ... -n 12000                           # polyhedra only
```

Re-run after the 15 Oct 2026 logging change (seeded):

```
BD_SEED=5 BD_BODIES=1 BD_COUNT=3 BD_START=setdown BD_REDROP=0 HOLD_S=0 bpp -f bille-drop.lua -n 2500
  Green   set down on B; predicted: B -> A -> D
  Yellow  at rest on face D after  0.23 s; faces it lay on: A -> D; edges it rolled on: BC -- Yellow (Bille)
  Red     at rest on face D after  0.25 s; faces it lay on: A -> D; edges it rolled on: BC, AB -- Red (Bille)
  Green   at rest on face D after  0.34 s; faces it lay on: B -> (A) -> D; edges it rolled on: CD, BC -- Green (Bille)
BD_SEED=5 BD_BODIES=1 BD_COUNT=3 BD_REDROP=1 HOLD_S=0 bpp -f bille-drop.lua -n 4000
=== 1 s: 3 finished runs ===
    Bille                                3 runs, 3 on the stable face, time to rest median 0.33 s; paths: D x3
```

(Bille seeds from the clock unless BD_SEED is set; seeded, it is checked for hall identity below. The tally's new median was not exercised with an even count in this short run; the formula is (t[floor((n+1)/2)] + t[floor(n/2)+1]) / 2.)

```
  Yellow  set down on B; predicted: B -> A -> D
  Yellow  at rest on face D after  0.45 s; faces it lay on: B -> D; edges it rolled on: CD, BC -- Yellow (Bille)
  Red     at rest on face D after  0.19 s; faces it lay on: C -> D; edges it rolled on: AB -- Red (Bille)
  Orange  at rest on face D after  0.21 s; faces it lay on: A -> D; edges it rolled on: BC -- Orange (Bille)
=== 12 s: 16 finished runs ===
    Bille                                16 runs, 16 on the stable face, time to rest median 0.41 s; paths: D x16
  Purple  set down on S6; predicted: S6 -> S5 -> S4 -> S3 -> S2 -> S1 -> base
  Purple  at rest on face base after  5.04 s; faces it lay on: S6 -> S5 -> S4 -> S3 -> S2 -> S1 -> S2 -> S1 -> base
    -> S1 -> base -- Purple (37-corner)
    21-corner polyhedron (ideal)           3 runs, 3 on the stable face, time to rest median 14.98 s; paths: S1 ->
    base x2, S3 -> S2 -> S1 -> base -> S1 -> base x1
    37-corner polyhedron (shell + weights)   4 runs, 4 on the stable face, time to rest median 4.58 s; ...
```

Pause and hall checks after the lift fix (15 Oct 2026): with `BD_SEED=5 BD_BODIES=1 BD_COUNT=3 BD_REDROP=1`, HOLD_S=0 for 3000 frames is a prefix of the default pause for 3030 frames (36 lines); hall against standalone (cmp2.sh, 1500 frames): IDENTICAL (12 run lines). With a fixed seed the hall-identity check applies like any other exhibit's.

Known departures and honest caveats

- Fixed 15 Oct 2026: the attendant's lift ran during the start pause (a `holdTick` that called `liftStep`), so with the pause on the output differed. `holdTick` is removed; the lift waits for the pause to end and the pause changes nothing (checks above).
- Fixed on 15 Oct 2026: the tally's "median" with an even number of runs printed the lower middle value; it is now the mean of the two middle values (same fix in ../Gomboc).
- **The Bille shape is a reconstruction, not the authors' drawing.** The paper gives vertex coordinates only in a figure. The shape was fitted to the published longest edge (500 mm), volume (668.624 cm³) and the four loading-zone volumes, all matched exactly. The mirror image fits equally well. The WC wedge mass design is the author's own, not the paper's.
- **Bille has a tiny design margin.** Its centre of mass sits 0.81 mm inside a zone at most 1.48 mm deep, and "a 0.1 g glue blob at a far vertex moves it ~0.25 mm" (README.txt).
- **The 21-corner body is the paper's idealisation** (point masses, weightless skin). The findings PDF says it is "essentially a mathematical object". Domokos & Kovács' Table 1 centre-of-mass values could not be reproduced from the stated construction, so the shapes were refitted (opt21/optnk). The 26- and 37-corner bodies are refitted for a 0.5 mm polycarbonate shell with tungsten weights.
- **Logged paths are dynamic, not quasi-static.** Bodies dropped or set down carry momentum and can cross a face in under the 0.03 s that counts as lying on it, or rock back up a band (S1 → S2). Fixed on 15 Oct 2026: such a quickly crossed face is no longer dropped from the log; it is printed in brackets, e.g. "B -> (A) -> D"; a bracketed face later lain on properly is replaced by the plain name.
- **Rolling resistance and spin friction are script-side constant moments** (δ = 50 µm chosen, not measured).
- **Time step depends on speed.** The effective step is 1/3000 s at the default speed and 1/1200 s at speed 0.1, despite the "steps" slider reading 1200 (there is always at least one step per frame). Documented on 15 Oct 2026 in the lua header and the slider's description ("at least one a frame (at 0.02 speed: 3000 a second)"); not changed, since a step independent of speed would need fractional steps per frame. No convergence study is documented for this exhibit.
- **Not fully tested.** The polyhedra were tested in only 10 drops and 3 set-downs here, and the 21-corner body takes 11–16 s to settle at this scale. All results come from one short session.

References

- G. Almadi, R. J. MacG. Dawson, G. Domokos, "Building a monostable tetrahedron", arXiv:2506.19244 (2025); Quanta Magazine, 2025-06-25.
- G. Domokos, G. Almadi, K. Regős, R. Dawson, "On Equilibria of Tetrahedra", Math. Intelligencer (2023).
- G. Domokos, F. Kovács, "Conway's spiral and a discrete Gömböc with 21 point masses", Amer. Math. Monthly 130 (2023); arXiv:2103.13727.

- S. Bozóki, G. Domokos, F. Kovács, K. Regős, Int. J. Solids Struct. (2021) and Eur. J. Oper. Res. 310 (2023), as cited in the findings PDF.
- W. M. Spears with Claude, "One resting point is not enough" (../Gomboc/gomboc-findings.pdf, 2026).

4.3 Galileo's ramp

Agrees with theory. Finish times within -0.013% of sqrt(2d(1+k)/(g sin θ)); distances in equal times 1 : 2.998 : 4.997 (odd numbers; fitted for this report); first-order convergence in step size.
Folder: demo/WyomingWill/GalileoRamp/

What it shows

Five rollers are let go together on a 5-degree ramp: a wooden ball, a small steel ball, a wooden disc, a brass hoop and a can of soup. The order at the finish line depends only on how much of each body's mass sits out at its rim ($k = I/(m\ r^2)$), not on its mass or size. The soup can wins because the liquid inside hardly turns, the two balls tie, and the hoop comes last.

Model

Bullet does the contacts. The balls are `btSphereShape`. The disc, hoop and can also collide as spheres of their own radius, but they are held to their lanes (linear factor (1,1,0), angular factor (0,0,1)) and given their own inertia tensors. The README says Bullet's cylinders, touching along a line, "jitter and lose 2% of their speed", which is why they are not used. The racers (code and README):

- wooden ball: r = 2.5 cm, 0.7 g/cm^3, 45.8 g, k = 0.4
- steel ball: r = 1.0 cm, 7.85 g/cm^3, 32.9 g, k = 0.4
- wooden disc: r = 2.5 cm, 2 cm thick, 27.5 g, k = 0.5
- brass hoop: 3 mm thick, 1.5 cm wide, 56.5 g; $k = (r^2 + r_i^2)/(2\ r^2) = 0.887$
- can: 40 g of tin plus 330 g of soup. Only the tin counts towards the spin inertia, which gives k = 0.094.

Physics settings: friction 0.8 on both ramp and bodies, restitution 0, damping 0, g = 981 cm/s^2, 30 solver iterations, 2400 steps per simulated second by default (GR_STEPS). Bullet's integrator is its usual semi-implicit (symplectic) Euler. The ramp is 120 cm long, and the finish line is 100 cm below the start line. The script interpolates each finish crossing to within one physics step.

Theory checks

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Soup time to 100 cm	t = sqrt(2d(1+k)/(g sinθ)), k=0.094: 1.5994 s	1.5992 s	-0.013%
Wooden ball time	k=0.4: 1.8097 s	1.8095 s	-0.011%
Steel ball time (ties wooden ball)	k=0.4: 1.8097 s	1.8095 s	-0.011%; ties ball to 0.1 ms
Disc time	k=0.5: 1.8732 s	1.8730 s	-0.011%
Hoop time	k=0.887: 2.1011 s	2.1009 s	-0.010%
Speed at line, soup / ball / disc / hoop	v = sqrt(2 d g sinθ/(1+k)): 125.0 / 110.5 / 106.8 / 95.2 cm/s	125.1 / 110.5 / 106.8 / 95.2 cm/s	≤0.1 cm/s
Acceleration (least-squares fit of d = ½at² + v0 t + d0 to GR_LOG, 0.2 s to finish)	a = g sinθ/(1+k): ball 61.07, disc 57.00, hoop 45.30, soup 78.18 cm/s²	61.071 / 56.999 / 45.305 / 78.182 cm/s²	<0.01%
Galileo's odd-number rule (distance in successive 0.5 s intervals)	1 : 3 : 5	1 : 2.998 : 4.997 (all five racers)	0.1%
Step-size convergence, soup time error	error → 0 as dt → 0	600 st/s -0.052%; 2400 -0.013%; 9600 -0.003%	error ∝ dt (first order, ~4× per 4× steps)

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Finish order	soup < balls (tie) < disc < hoop	Soup, Ball, Steel, Disc, Hoop	matches

The theory values for time and speed are printed by the script. The fit and odd-number rows I computed from the GR_LOG CSV of this run. Times in the README are identical to this run's, to every printed digit.

Run used

```
cd demo/WyomingWill/GalileoRamp && HOLD_S=0 timeout 1500 bpp -f galileo-ramp.lua -n 600            # 2400 steps/s
HOLD_S=0 GR_STEPS=600  ... -n 600 ;  HOLD_S=0 GR_STEPS=9600 ... -n 600                            # convergence
HOLD_S=0 GR_LOG=gr.csv ... -n 150                                                                   # distance log
```

```
  Soup   the can of soup         370.0 g, k = I/(m r^2) = 0.094: a = g sin(theta)/(1 + k) =  78.2 cm/s^2; theory
    1.599 s
  Hoop   the brass hoop           56.5 g, k = I/(m r^2) = 0.887: a = g sin(theta)/(1 + k) =  45.3 cm/s^2; theory
    2.101 s
  Soup   1st across the line at 1.5992 s (theory 1.5994, -0.013%), at 125.1 cm/s (theory 125.0)
  Ball   2nd across the line at 1.8095 s (theory 1.8097, -0.011%), at 110.5 cm/s (theory 110.5)
  Steel  3rd across the line at 1.8095 s (theory 1.8097, -0.011%), at 110.5 cm/s (theory 110.5)
  Disc   4th across the line at 1.8730 s (theory 1.8732, -0.011%), at 106.8 cm/s (theory 106.8)
  Hoop   5th across the line at 2.1009 s (theory 2.1011, -0.010%), at 95.2 cm/s (theory 95.2)
GR_STEPS=600:  Soup   1st across the line at 1.5986 s (theory 1.5994, -0.052%) ...
GR_STEPS=9600: Soup   1st across the line at 1.5994 s (theory 1.5994, -0.003%) ...
```

Known departures and honest caveats

- The disc, hoop and can are spheres to the collision detector and are locked to planar motion with spin about their axes only. That is exact for an ideal line contact, but the model cannot show wobble, steering or edge effects. Collision shape and inertia are chosen separately, on purpose.
- The can is an idealisation: the soup is treated as not turning at all, so it counts only towards translation. Real soup is viscous and slowly spins up, so a real can would sit somewhere between k = 0.094 and that of a solid cylinder (0.5).
- All times are about 0.2 ms early (-0.01%). The error halves roughly with each doubling of the step rate (first-order integrator). At the default 2400 steps/s it is far below anything visible.
- Friction 0.8 was chosen so that nothing slips. At 5° the friction needed is small ($\mu \geq k \tan\theta/(1+k)$, at most 0.041 for the hoop), so this is comfortable. Rolling resistance and air drag are absent: the ramp is ideal.
- The odd-number rule and acceleration fit are not printed by the exhibit. I computed them from the optional log. In the log, the ball passes the 114 cm end of the ramp before t = 2.0 s, so only the first three intervals (up to 1.5 s) were used.
- The README's console excerpt matches this run exactly.

References

- Galileo Galilei, inclined-plane experiments (named in the exhibit title; no specific citation in code or README).
- Standard rolling-without-slipping result $a = g \sin\theta/(1 + I/(m r^2))$ (stated in code and README; no source cited).

4.4 The double cone

Agrees with theory. Rolls up or down exactly as the rail geometry predicts on all three tracks; centre's drop within 1 µm of the exact geometry; energy loss 0.1-0.3% (numerical).
Folder: demo/WyomingWill/DoubleCone/

What it shows

Two cones are joined base to base and set on a pair of rails that rise and spread apart in a V. Let go at the low, narrow end, the double cone rolls up to the high, wide end, so it seems to roll uphill. It does not really: the spreading rails meet it nearer its points, where it is thinner, so its centre of mass sinks faster than the rails rise. Two control tracks sit beside it. On the same rails a plain cylinder rolls down. On rails rising 5 cm instead of 1.5 cm (too steep), the same cone rolls down.

Model

Bullet does the contacts:

- **Double cone:** a `btCompoundShape` of two `btConeShapeZ` (R = 5 cm, H = 12 cm each), collision margin 0.01 cm, 400 g, solid. Inertia is I_axis = 3/10 m R^2 = 3.0 kg·cm^2 and I_perp = m(3/20 R^2 + H^2/10).
- **Cylinder:** a `btCylinderShapeZ` with r = 3 cm, length 26 cm, 400 g.
- **Rails:** 8 mm brass rods (radius 0.4 cm), 50 cm long, 2.5 cm from the centre line at the low end and 9.5 cm at the high end, so tan φ = 0.14 per side. They rise 1.5 cm (tan θ = 0.03) at the back and middle tracks and 5 cm (tan θ = 0.10) at the front. For collision, each rail is a compound of 10 cm capsules laid end to end. The README says a single 54 cm cylinder gave an "invisible bump halfway up", and 3 cm pieces lost "a few per cent" of energy at the joins.
- **Settings:** friction 0.9, restitution 0, damping 0, 30 solver iterations, 2400 steps per simulated second (DC_STEPS), integrated by Bullet's semi-implicit Euler.

The script itself works out the exact predicted centre height at every x. It finds by bisection the height at which the rail's centre line just clears a cone enlarged by (ρ + margin)/cos(taper), allowing for the off-centre contact point on sloping rails. It also tracks the total energy, kinetic (translational plus rotational) plus potential.

Theory checks

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Rolls up or down (back)	rolls up iff tan θ < tan φ·R/H; 0.030 < 0.14·5/12 = 0.0583	rolls up 36.8 cm to the high end	correct
Rolls up or down (front)	0.100 > 0.0583, so rolls down	rolls down 23.9 cm	correct
Back cone: centre drop while rolling up 36.8 cm	exact geometry (script): 1.045 cm; thin-rail estimate (tan φ R/H − tan θ)·Δx = 0.0283·36.8 = 1.04 cm (my arithmetic)	1.045 cm down	0.000 cm (exact); 0.2% (estimate)
Front cone: centre drop rolling down 23.9 cm	geometry 0.994 cm; estimate (0.100 − 0.0583)·23.9 = 1.00 cm	0.994 cm	exact; 0.6% (estimate)
Cylinder: centre drop rolling down 31.8 cm	geometry 0.955 cm; tan θ·Δx = 0.954 cm	0.955 cm	exact
Largest deviation from predicted centre height along the run	0	back 0.0000 cm, front 0.0001 cm, middle 0.0001 cm (4 cycles)	≤1 µm
End-to-end centre change (48 cm), printed prediction	back −1.36 cm, cylinder +1.44 cm, front +2.00 cm	(prediction only)	—
Energy conservation (rolling, no slip)	ΔE = 0	loss as % of centre-fall energy, over 4 cycles: back 0.14–0.17%, front 0.09–0.12%, middle 0.20–0.26%	≤0.26% lost

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Step-size check (DC_STEPS=600)	—	deviation ≤0.0004 cm; energy loss back 0.42–0.50%, front 0.48–0.57%, middle 0.62–0.65%	loss about 3× larger at 4× the step

Run used

```
cd demo/WyomingWill/DoubleCone && HOLD_S=0 timeout 1500 bpp -f double-cone.lua -n 1200     # 2400 steps/s, 4 cycles
HOLD_S=0 DC_STEPS=600 timeout 1500 bpp -f double-cone.lua -n 1200                          # step-size check
```

```
  Back   rails rise 1.5 cm over 50 (tan 0.030); the cone's centre would go down by 1.36 cm from end to end:
         tan(theta) 0.030 < tan(phi) R/H = 0.140 5.000 / 12.000 = 0.0583, so it rolls UP the rails (its centre going
    down)
  Middle a cylinder on the same rails: its centre rises 1.44 cm with them, so it rolls down
  Front  rails rise 5.0 cm over 50 (tan 0.100); the cone's centre would go up by 2.00 cm from end to end:
  Front  nears the low, narrow end at 1.55 s, having rolled 23.9 cm down the rails (the rails falling 2.39 cm): its
    centre went down 0.994 cm (the geometry: 0.994),
         off the geometry by 0.0001 cm at worst on the way; it lost 0.12% of the energy its centre's fall gave it
  Middle nears the low, narrow end at 1.80 s, having rolled 31.8 cm down the rails (the rails falling 0.95 cm): its
    centre went down 0.955 cm (the geometry: 0.955),
         off the geometry by 0.0001 cm at worst on the way; it lost 0.20% of the energy its centre's fall gave it
  Back   nears the high, wide end at 2.32 s, having rolled 36.8 cm up the rails (the rails rising 1.10 cm): its
    centre went down 1.045 cm (the geometry: 1.045),
         off the geometry by 0.0000 cm at worst on the way; it lost 0.14% of the energy its centre's fall gave it
```

The README's back-cone line (2.32 s, 36.8 cm, 1.045 cm, 0.0000 cm, 0.14%) matches this run's first cycle exactly. The README's "0.1 to 0.3%" energy loss matches the 0.09–0.26% range seen here.

Known departures and honest caveats

- The "geometry" check is the script's own exact contact geometry, which includes the rail thickness and Bullet's collision margin. It shows that Bullet holds the body on the ideal surface; it is not an independent physical measurement. The simple thin-rail formula agrees to within 0.6%.
- Energy is not exactly conserved. 0.1–0.3% of the energy released by the centre's fall is lost, and the loss roughly triples at 600 steps/s, so it is numerical (contact and solver dissipation), not physical rolling resistance. The losses also vary slightly from cycle to cycle (e.g. back 0.14–0.17%), because each restart begins from Bullet's leftover contact state.
- The rails were segmented into 10 cm capsules specifically to avoid Bullet contact artefacts: a bump with one long cylinder, energy loss at joins with 3 cm pieces (README). This was tuned, not derived.
- Friction 0.9 is high, chosen so that nothing slips. Rolling resistance and air drag are absent.
- The checks are reported only between 6.5 cm from each end (away from the stops). Behaviour at the stops is not checked.
- Only the outcome (direction and centre descent) is checked. Times to reach the end (e.g. 2.32 s) are not compared with an analytic prediction.

References

- The 18th-century "mechanical paradox" lecture-room instrument (mentioned in code and README; no specific author or paper cited).

4.5 The oloid and the sphericon

Agrees with theory. Oloid's centre stays within 4.2426-4.5000 cm, the roller and sphericon level at 4.2426 cm, exactly as theory says; side-to-side weaves 3.6, 5.1 and 5.8 cm, now stated correctly.
Folder: demo/WyomingWill/Oloid/

What it shows

Three wooden rollers, each the convex hull of two circular arcs (radius 6 cm), are pushed along a level board at 15 cm/s. None is round, yet each rolls and lays its whole surface on the board while weaving from side to side. The oloid's (Schatz) centre of mass rises and falls a little as it rolls. The two-circle roller's (Stewart) and the sphericon's stay at one height, R/√2.

Model

Each roller is a Bullet rigid body (mass and principal moments from `reconstruction/gen-meshes.py`). Bullet's own collision is switched off (`collides=false`), and Bullet's gravity and gyroscopic term are off too (`setFlags(1)`). Before every physics step the script does three things. (1) It applies gravity (g = 981 cm/s²). (2) It advances the spin with Euler's torque-free equations, using RK4 in the body frame. (3) It finds the exact contact, the lowest point of each of the two arcs, from the analytic arc support function. Normal and Coulomb-friction impulses (μ = 0.6, a 2D friction cone) are solved together by sequential impulses, 8 iterations, with Baumgarte bias 0.2 on penetration. Rolling resistance is a moment δ·N with δ = 20 μm (`DELTA = 0.002` cm), applied against the rolling but not about the board normal. Bullet then only integrates positions and orientations. The run uses 2400 physics steps/s (`OL_STEPS`) and no damping. Theory for the height range is computed when the script loads: it scans 720 × 180 "down" directions and bisects for directions where both arcs reach equally far. Masses (this run, matching README): oloid 461.5 g, two-circle 496.2 g, sphericon 316.7 g (wood 0.7 g/cm³). Setting-down yaws (−14.7°, 12.0°, −25.6°) were "found by trying" so each rolls straight on average.

Theory checks

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Oloid COM height range while rolling	between R/√2 and 3R/4 = 4.2426 to 4.5000 cm (script's own scan of the shape)	4.2426 to 4.5000 cm	exact to 4 decimals (0.1 μm)
Two-circle roller COM height	constant, 4.2426 cm (scan; Stewart roller)	4.2426 to 4.2426 cm	constant to 4 decimals
Sphericon COM height	constant R/√2 = 4.2426 cm	4.2426 to 4.2426 cm	constant to 4 decimals
Energy loss besides rolling resistance (should be 0 for rolling without slipping), 2400 steps/s	0	sphericon 0.229 %, two-circle 0.681 %, oloid 0.227 % of push KE	README: "0.2 to 0.7 %" — matches
Same, 9600 steps/s (step-size convergence)	0; README: "about a fifth" of the 2400 value	0.046 %, 0.126 %, 0.054 %	ratios 0.20, 0.19, 0.24 — matches README
Share of push KE taken by rolling resistance (δ = 20 μm)	(bookkeeping, no theory value)	6.6 %, 18.9 %, 15.9 %	—
Side-to-side weave	README/header now: two-circle 3.6, oloid 5.1, sphericon 5.8 cm; help text "3.6 to 5.8 cm" (fixed 15 Oct 2026; was "4 to 6 cm")	sphericon 5.8, two-circle 3.6, oloid 5.1 cm (same on every push in a 25 s run)	matches
Oloid volume	3.0524 R³ (README, from gen-meshes.py)	not re-run	README only

Run used

```
cd demo/WyomingWill/Oloid && HOLD_S=0 timeout 1500 bpp -f oloid.lua -n 720
cd demo/WyomingWill/Oloid && HOLD_S=0 OL_STEPS=9600 timeout 1500 bpp -f oloid.lua -n 720
cd demo/WyomingWill/Oloid && HOLD_S=0 bpp -f oloid.lua -n 1500      # 15 Oct 2026 re-run, weave figures, identical to
    the above
```

Default (2400 steps/s):

```
  Oloid     461.5 g; lying along a line on the board, its centre is between 4.2426 and 4.5000 cm up (theory)
  Sphericon reaches the far end at 3.93 s, 108.3 cm along the board, weaving 5.8 cm side to side:
            its centre 4.2426 to 4.2426 cm up (theory: 4.2426 to 4.2426);
            energy: of what the push gave it, 6.6% went to rolling resistance, and 0.229% otherwise (rolling without
    slipping, it should lose none)
  2-circle  reaches the far end at 6.83 s, 109.6 cm along the board, weaving 3.6 cm side to side:
            energy: of what the push gave it, 18.9% went to rolling resistance, and 0.681% otherwise (rolling without
    slipping, it should lose none)
  Oloid     reaches the far end at 9.02 s, 108.8 cm along the board, weaving 5.1 cm side to side:
            its centre 4.2426 to 4.5000 cm up (theory: 4.2426 to 4.5000);
```

OL_STEPS=9600:

```
            energy: of what the push gave it, 6.6% went to rolling resistance, and 0.046% otherwise (rolling without
    slipping, it should lose none)
            energy: of what the push gave it, 18.9% went to rolling resistance, and 0.126% otherwise (rolling without
    slipping, it should lose none)
            energy: of what the push gave it, 15.9% went to rolling resistance, and 0.054% otherwise (rolling without
    slipping, it should lose none)
```

The README's sample console block matches this default run line for line (times 3.93/9.02 s, distances, 0.229 %).

Known departures and honest caveats

- The height check partly tests the script against itself. The "theory" range comes from the same arc-support function the contact uses, scanned numerically, not from a closed-form derivation. It does agree with the closed-form values $R/\sqrt{2} = 4.2426$ and $3R/4 = 4.5000$.
- The min/max height is recorded once per frame (60 Hz), not every physics step.
- The contact model is the script's own sequential-impulse solver with Baumgarte stabilisation, not exact rolling constraints. The residual energy loss of 0.05-0.7 % is step-size error and shrinks about 5× from 2400 to 9600 steps/s.
- The rolling resistance (20 µm) is a chosen parameter. It takes 7-19 % of the push energy over the ~110 cm run.
- The setting-down yaw angles (12-26° off the axle) were tuned by hand so the rollers run straight on average.
- The board is level and the rollers are pushed. README: on a slope, past about 40 cm/s, one arc's normal force nearly vanishes and they skid. That limit was not tested here.
- Fixed 15 Oct 2026: README, header comment and help text said all three weave "4 to 6 cm"; the two-circle roller weaves 3.6 cm. They now give each body's measured weave (3.6 / 5.1 / 5.8 cm; help text 3.6 to 5.8 cm). The Cabinet's placard and catalogue text for this exhibit do not mention the weave. Text only; physics and console unchanged (hall vs standalone IDENTICAL, 900 frames).
- The oloid's centre height is not perfectly constant (it rises and falls by 0.26 cm). A real oloid on a slope therefore rolls unevenly; the exhibit shows this only as the pen trace.

References (as cited in README and code)

- Paul Schatz, the oloid, 1929.
- A. T. Stewart, the two-circle roller, 1966 (no title or journal given).
- Sphericon (no source cited).
- Mass properties: `reconstruction/gen-meshes.py` (NumPy/SciPy).

4.6 Euler's disk

Agrees with theory. Precession rate within 0.08-0.18% of theory from 25° down to 1°; with losses off, energy steady to the eighth figure. Its finite-time end follows its rolling-resistance law (now said), not Moffatt's air.
Folder: demo/WyomingWill/EulersDisk/

What it shows

A heavy steel disc is set rolling on its edge on a mirror at a 30° lean. It rolls round in place and leans further over as it loses energy. The flatter it gets, the faster its axis goes round, until it suddenly stops, flat. This is Joseph Bendik's 1990s toy; a spinning coin does the same.

Model

Bullet is not used for the dynamics. The README says Bullet's contacts "added energy about as fast as the rolling resistance took it". The script integrates the exact rolling-without-slipping equations: Newton's and Euler's equations for the disc, with the contact force solved each instant so that the contact point is stationary: $(I + m(r\cdot r - r\, r^{T}))\ \dot{\omega} = -\omega\times I\omega + m\, r\times(g\cdot up - \omega\times\dot{r}) +$ moments. Integration is RK4 at 9600 steps/s (`ED_STEPS`), with Gram-Schmidt re-orthonormalisation and the centre height reset from the contact geometry every step. The disc is steel (ρ = 7.85 g/cm³), a = 3.8 cm radius, 1.2 cm thick, rolling on the edge of its bottom face (h = 0.6 cm). Run values: m = 427 g, I1 = 1.594, I3 = 3.085 kg cm². Dissipation has two parts: (1) rolling resistance, a moment $\delta\cdot m\cdot g$ against the level part of ω, with δ = 30 µm (`ED_DELTA`); (2) a drag of 10 /s on the centre's sideways motion (`ED_DRAG`; README: "the air, and the mirror's grip not being perfect"). The disc is set going at α = 30° with the steady-precession Ω for that lean, 7.47 turns/s. It counts as stopped when α < 0.3°.

Theory checks

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Precession Ω at 24.92°	$\Omega^2 = m\, g\, a\,(a\cos\alpha - h\sin\alpha) / (\sin\alpha\,(I1\, a\cos\alpha - I3\, h\sin\alpha))$ → 8.06 turns/s	8.04	−0.18 %
Ω at 19.88°	same → 8.88	8.87	−0.14 %
Ω at 14.98°	same → 10.11	10.10	−0.11 %
Ω at 9.99°	same → 12.25	12.24	−0.09 %
Ω at 5.97°	same → 15.73	15.72	−0.08 %
Ω at 4.00°	same → 19.15	19.14	−0.08 %
Ω at 1.98°	same → 27.14	27.11	−0.09 %
Ω at 0.99°	same → 38.33	38.29	−0.11 %
Rate of fall of √α near the end	$\delta\sqrt{(m\, g\, a / I1) / (3a)}$ = 0.00832 /s	0.00818 /s (line fit to samples with α < 2°)	−1.7 %
Finite-time singularity constant	$\Omega\,(t0 - t) = 3a/\delta$ = 3800 (t0 from the fit: 90.6 s)	3861 (0.32°), 3864 (0.69°), 3867 (1.20°), 3872 (1.84°)	+1.6 to +1.9 %
Energy conservation with no losses (ED_DELTA=0, ED_DRAG=0, 19.8 s)	E constant	2523.0883 kg cm²/s² at all 77 logged samples	no change at the logged 8 significant figures (< 4e-8 relative), with or without the drag; README now says "to the eighth figure over 20 s"
Stopping lean / time	(not theory) README: 0.30°, 81.73 s, 67 turns/s	0.30°, 81.73 s, 67 turns/s	identical to README

Run used

```
cd demo/WyomingWill/EulersDisk && HOLD_S=0 timeout 1500 bpp -f eulers-disk.lua -n 5400
cd demo/WyomingWill/EulersDisk && HOLD_S=0 ED_DELTA=0 ED_DRAG=0 ED_LOG=disk_e.csv timeout 1500 bpp -f eulers-disk.lua
    -n 1200
```

```
--- set going: leaning 30 degrees, going round 7.47 times a second; rolling resistance 30 microns ---
  the disc: 427 g, 7.6 cm across, 1.2 cm thick; I1 1.594, I3 3.085 kg cm^2
  t=  8.72 s  leaning 24.92 degrees: going round   8.04 times a second (theory, going round in place:   8.06; -0.18%)
  t= 51.17 s  leaning  5.97 degrees: going round  15.72 times a second (theory, going round in place:  15.73; -0.08%)
  t= 74.63 s  leaning  0.99 degrees: going round  38.29 times a second (theory, going round in place:  38.33; -0.11%)
  t= 81.73 s  it stops, flat, leaning 0.30 degrees, going round 67 times a second
              sqrt(alpha) falling at 0.00818 a second (theory, near the end: delta sqrt(m g a / I1) / (3 a) = 0.00832);
              so it would lie flat at t0 = 90.6 s, going round infinitely fast
              t= 77.17 s, leaning 0.69 degrees: W (t0 - t) = 3864 (theory: 3 a / delta = 3800)
              t= 68.63 s, leaning 1.84 degrees: W (t0 - t) = 3872 (theory: 3 a / delta = 3800)
```

Every number in the README's console sample (10.10/10.11, 15.72/15.73, 38.29/38.33, 81.73 s, 0.00818 vs 0.00832, t0 = 90.6 s, 3864 vs 3800) was reproduced exactly.

Known departures and honest caveats

- **The dissipation law is a modelling choice.** It is a constant rolling-resistance moment δ·m·g. That gives √α falling linearly, so $\alpha \propto (t0 - t)^2$ and $\Omega \propto 1/(t0 - t)$. Moffatt's finite-time-singularity analysis (Nature 2000) uses viscous dissipation in the air layer instead, giving $\alpha \propto (t0 - t)^{1/3}$. The exhibit's "theory" is the theory of its own dissipation model, so it cannot discriminate between real-world dissipation mechanisms. Since 15 Oct 2026 the README and lua header say this plainly: the finite-time end comes from the chosen rolling-resistance law, Moffatt's air law would give $\alpha \propto (t0 - t)^{1/3}$, $\Omega \propto (t0 - t)^{-1/6}$, and the exhibit can't tell them apart (README also cites Easwar, Rouyer and Menon 2002, vacuum experiments).
- The near-end law is a small-angle asymptote. The 1.7 % gaps (√α rate, Ω(t0 − t)) are consistent with the finite angles of the fit window (α < 2°) and with the extrapolated t0. Note t0 = 90.6 s, while the disc actually stops at 81.73 s.
- The extra sideways drag (10 /s) was added to keep the centre still. Without it, the README says, the centre drifts into mm-sized circles and Ω runs up to 20 % below the in-place theory. The 0.1 % agreement therefore depends partly on this tuned drag.
- The disc never slips and never bounces. The README admits the real toy stops around 0.3° when bouncing and slipping take over, which this model does not include. The 0.3° stop is an imposed threshold, not an emergent result.
- Fixed on 15 Oct 2026: the README and lua header said "430 g"; they now say 427 g, as the console prints.
- The energy test (no losses): re-run 15 Oct 2026 with ED_DELTA=0 alone and with ED_DELTA=0 ED_DRAG=0, 1200 frames (19.8 s): 2523.0883 kg cm²/s² at all 77 logged samples in both. The README and header used to say "a few parts in a million"; they now say "to the eighth figure over 20 s".

References

- Joseph Bendik, Euler's Disk toy (1990s), as named in README.
- H. K. Moffatt (2000), Nature: Euler's disk and its finite-time singularity (cited in README and lua header since 15 Oct 2026, as the contrasting air-viscosity law).
- K. Easwar, F. Rouyer, N. Menon (2002), vacuum experiments on the spinning disc (README, since 15 Oct 2026; journal not given there).

4.7 The rattleback

Agrees with theory. Spun the wrong way it reverses (0.67 s), the right way never; the reference model's growth rates (8.9/s and 1.7/s at 1 turn a second) and the rolling-resistance window now stated as measured. New README.
Folder: demo/WyomingWill/Rattleback/

What it shows

A rattleback (celt) is a boat-shaped body whose mass axes are turned a few degrees from the lines of its hull. Spun one way, it spins smoothly. Spun the other way, it starts to rock, stops, and spins back. Tapped on the end from rest, it starts spinning its preferred way. Three identical rattlebacks are shown: left, spun clockwise (the wrong way); middle, spun anticlockwise (its own way); right, "restless", spun clockwise on an almost lossless table, so it reverses more than once.

Model

The body is a solid half-ellipsoid with half-axes 6 × 1.5 × 1.2 cm, plastic at 1.2 g/cm³, with the hull turned 8° about the vertical from the principal axes. From `rattleback-data.lua`: m = 27.14 g, I = (0.01454, 0.20765, 0.19775) kg cm². Collision uses Bullet's own contact (unlike the other three exhibits here): a btConvexHullShape of 3073 surface points, 0.1 mm margin, contact-processing threshold 0.001 cm, friction 0.6, restitution 0. Bullet's gyroscopic term is off (`setFlags(0)`). Instead, Euler's torque-free equations are advanced with RK4 before every Bullet step. Rolling resistance is a moment δ·m·g against the horizontal part of ω about the contact, which is taken as directly below the centre of mass. Defaults: δ = 50 μm (left and middle) and 5 μm (restless). The run uses 1200 steps/s, no damping. Initial conditions: 1 turn/s about the vertical plus a 1 rad/s "hand wobble" about the long axis; the tap is 3 rad/s of pitch. Documentation: README.md (written 15 Oct 2026: what it shows, keys, model, theory and console, the reference model and its checks, files, headless env vars), the lua header, and `reference/` (rb_ref.py, an independent rolling-without-slipping ODE model; rb_scan.py; rattleback-test.lua).

Theory checks

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Mass properties	rb_ref.py: m = (2/3)π A B C ρ; I from the half-ellipsoid formulas	data file 0.0271434 kg; I = 0.0145353, 0.2076467, 0.1977530; rb_ref gives 0.02714336, 0.01453527, 0.20764671, 0.19775295	identical
Linear growth rate, wrong way (clockwise)	eigenvalues of linearised rb_ref (recomputed 15 Oct 2026); header/README now "8.9/s at 1 turn/s; 8.5/s at 6 rad/s" (fixed; header said 8.5 at 1 turn/s)	8.493 /s at 6 rad/s; 8.882 /s at 2π rad/s (1 turn/s)	matches
Growth rate, own way (anticlockwise)	code comment / README: 1.7 /s at 1 turn/s (README: 1.66 at 6 rad/s)	1.661 /s at 6 rad/s; 1.728 /s at 1 turn/s	OK
Skew 0 control	no instability	growth 0.0000 /s both ways (rb_ref); rb_ref integration: no reversal in 30 s	OK
Lossless reference reverses both ways	code: "unstable in BOTH directions"	rb_ref, ±6 rad/s, skew 8: first reversal at 4.68 s (+spin) and 1.07 s (−spin); energy drift −1.4e-9 / −1.5e-9	OK
Exhibit: wrong-way spin reverses	expected	Left REVERSED at 0.67 s (rocking up to 7°)	qualitative OK
Exhibit: own-way spin does not reverse	expected (δ = 50 μm damps the weak instability)	Middle never reversed in 60 s	OK
Restless (δ = 5 μm) reverses repeatedly	"again and again"	3 reversals: 0.65, 4.22, 6.23 s; then ~+0.02 turn/s creep to 60 s	qualitative OK; only 3 reversals
Tap from rest spins anticlockwise	expected	all three anticlockwise at 0.03 s; restless later reverses at 3.10 and 5.52 s	OK

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
δ window (left and middle; RB_ROLL, 30 s runs, exhibit start)	header/README now state the measured window (fixed 15 Oct 2026; header said "20-100 µm one-way; at 200 neither way reverses")	0, 10 µm: both reverse repeatedly. 20: Left reverses 3x (0.55, 5.35, 7.62 s). 30: Left 3x (0.52, 3.93, 5.38 s), Middle once at 8.73 s. 40, 50, 75, 100: one-way (Left once at 0.52-0.68 s, Middle never). 150, 300: neither reverses. 200: Left reverses late, 4.57 s	one-way window 40-100 µm; documented as such
Bullet harness vs reference, no losses (rattleback-test.lua, RB_EULER=1, RB_ROLL=0, skew 8, ±6 rad/s, 0.02 rad/s rock)	rb_ref first reversal: 1.07 s (−6, wrong way), 4.68 s (+6)	0.70 s (−6), 3.15 s (+6)	both ways, same order; Bullet reverses ~1/3 sooner

Run used

```
cd demo/WyomingWill/Rattleback && HOLD_S=0 timeout 1500 bpp -f rattleback.lua -n 3600
HOLD_S=0 RB_START=tap ... -n 1800 ;  HOLD_S=0 RB_ROLL={0,20,100,200} ... -n 1200
python3 reference/rb_ref.py (run via runpy, PYTHONDONTWRITEBYTECODE=1); linearised growth: rb_scan-style Jacobian on
    rb_ref
reference/: RB_SKEW={8,0} RB_SPIN={-6,6} RB_EULER=1 RB_ROLL=50 RB_LOG=... bpp -f rattleback-test.lua -n 1200
15 Oct 2026 re-runs: default -n 3600; RB_START=tap -n 1800; RB_ROLL={0,10,20,30,40,75,100,150,200,300} -n 1800;
  rb_ref.py and the growth rates again; rattleback-test.lua as above plus RB_ROLL=0 (skew 8, ±6), -n 600
```

```
  Restless REVERSED at  0.65 s: now spinning anticlockwise (rocking up to 8 deg)
  Left  REVERSED at  0.67 s: now spinning anticlockwise (rocking up to 7 deg)
t=  2.0s  turns/s: Left +0.62  Middle +0.79  Restless +0.71   (+ = anticlockwise)
  Restless REVERSED at  4.22 s: now spinning clockwise (rocking up to 16 deg)
  Restless REVERSED at  6.23 s: now spinning anticlockwise (rocking up to 16 deg)
skew 8 spin +6:  ... first reversal 4.68 s  rock 28.7 pitch 6.2  energy drift -1.4e-09
skew 8 spin -6:  ... first reversal 1.07 s  rock 28.7 pitch 6.1  energy drift -1.5e-09
skew 0 spin +6:  ... first reversal none  rock 0.0 pitch 0.0  energy drift -4.4e-14
(RB_ROLL=200)  Left  REVERSED at  4.57 s: now spinning anticlockwise (rocking up to 5 deg)
```

Known departures and honest caveats

- The theory comparison is **qualitative in the exhibit itself**. The console reports reversals and spin rates, never comparing them with a theory number. The quantitative checks (growth rates, the both-ways instability) live in `reference/rb_ref.py`, an independent rigid-ellipsoid, no-slip, energy-conserving model, not in the exhibit.
- The exhibit uses Bullet's general convex-hull contact, with friction, possible slip and a hull faceted into 3073 points. The reference model assumes no slip on a smooth ellipsoid. The two are not compared directly on reversal time or spin after reversal.
- One-way behaviour is **tuned via rolling resistance**. With no losses, this shape reverses both ways. The real asymmetry depends on table losses, a modelling choice. Fixed 15 Oct 2026: the header's window ("20-100 µm one-way; at 200 neither") is replaced by what runs from the exhibit's start show: one-way for 40-100 µm; at 20 and 30 µm the wrong-way one turns round and back (at 30 the own-way one also reverses once, late); at 150 and 300 neither reverses, but at 200 the wrong-way one still does, late (4.57 s), so above 100 it is near the edge. The window is narrow and not monotone; the default 50 µm sits inside it.
- The rolling-resistance contact point is approximated as directly below the centre of mass.
- After the restless one settles, it keeps a ~0.02 turn/s spin out to 60 s rather than coming to rest. This is likely contact-solver creep.
- Corrected 15 Oct 2026 (this dossier's earlier reading was wrong): in the reference harness (`rattleback-test.lua`, RB_ROLL=50) the body sits on the table from the start (height 0.7600 cm is the rest height plus the 0.01 cm margin). It spins at ±6.00 for ~10 s because the 50 µm rolling resistance at once removes almost all of the 0.02 rad/s starting rock (0.01 deg -> 0.0009 deg at the first frame); the rock then regrows slowly, and skew 8 at −6 rad/s reverses at 11.1 s; +6 does not, skew 0 neither,

in 20 s. With RB_ROLL=0 it reverses both ways (0.70 s and 3.15 s; rb_ref 1.07 and 4.68 s).

- Fixed 15 Oct 2026: the data-file fallback path was `demo/rattleback/`; it is now `demo/WyomingWill/Rattleback/`, and the exhibit runs from the repository root (checked).
- Fixed 15 Oct 2026: the header quoted 8.5 /s "at 1 turn/s"; that is the 6 rad/s value. It now says 8.9 /s at 1 turn/s (8.5 at 6 rad/s). The header's key list also said R spins "left clockwise, right anticlockwise" and T taps "both"; now left and right clockwise, middle anticlockwise, and all three.
- The restless one reverses 3 times (0.65, 4.22, 6.23 s), not "again and again" as the help text says; the header now gives the three times; the help text was left as it is.
- Text and a fallback path only; physics and console unchanged (hall vs standalone IDENTICAL, 900 frames; a fresh 600-frame standalone run matches the start of the earlier 3600-frame run line for line).

References

None are cited in code or reference files. The classical analyses (Walker 1896; Bondi 1986, Proc. R. Soc. A; Garcia & Hubbard 1988, Proc. R. Soc. A) are the standard theory for the spin-direction-dependent instability. They are added here for context and are not used by the exhibit.

4.8 The spinning eggs

Agrees with theory, with a numerical or open caveat. The left egg rises at 0.30 s and stands at 0.67-0.68 s at every step rate from 2400 to 38,400 (was 0.50 s to never); its rise from 60° to 30° within 8% of Moffatt and Shimomura (2% started in balance); Jellett's quantity changes by 0.8-1.7%; energy budgets converge with step. Open: how long it stays standing depends on the step.
Folder: demo/WyomingWill/SpinningEgg/

What it shows

A hard-boiled egg spun fast on its side rises onto its end and spins there like a top. Sliding friction at the table does it, as with the tippe top. Spun faster, it also jumps briefly off the table as it rises. A raw egg spun the same way stays down: its liquid does not turn with the shell, the egg slows quickly, and after a finger stops it, it starts turning again by itself.

Model

Each egg is a Bullet rigid body; Bullet only integrates position and orientation. The egg is a prolate spheroid with half-axes a = 2.85 cm and b = 2.15 cm, mass 58 g, with $A = m(a^2+b^2)/5$ across and $C = 2mb^2/5$ about the long axis. Before every step the script applies gravity and runs Euler's torque-free equations (RK4; Bullet's gyroscopic term off). It then computes the exact contact: the spheroid's lowest point, with a unilateral normal impulse (so the egg can leave the table) and Coulomb friction $\mu = 0.3$ in a 2D cone, by sequential impulses over 6 iterations with Baumgarte bias 0.2. The dish rim is handled the same way. Spin friction is a moment $\kappa \cdot N$ about the vertical with $\kappa = 100$ µm. Rolling resistance is $\delta \cdot N$ with $\delta = 100$ µm. The run uses 4800 steps/s, no damping. Raw egg: a 0.35 mm shell (10 % of the mass) is the rigid body. Potential-flow added inertia $m_f/5 \cdot (a^2-b^2)^2/(a^2+b^2)$ is added to it. The rest of the liquid is a second rotor, $I_liq = 0.1194$ kg cm², coupled to the shell by a linear drag with time constant 1.5 s. Spins: left and right 25 turns/s, middle 40 turns/s. Each egg starts tipped 6° from horizontal (3° until 15 Oct 2026; see caveats), spun about the vertical.

Theory checks (default run unless noted)

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Spin when upright, left 25 t/s (Jellett, Moffatt & Shimomura 2002)	W = W0·A b/(C a) = 26.00 t/s; corrected for the J removed by spin friction and rolling resistance: 25.17	25.31 t/s	J changed +0.83 %
Same, middle 40 t/s (jumps)	41.60; corrected 40.26	40.95	+1.73 %

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Same, κ = 0 (EG_KAPPA=0), left	26.00; corrected 25.85	26.02 (25.98 at EG_STEPS=9600)	+0.81 % (+0.70 %)
Rate of rise, left, 60° → 30° from upright (added 15 Oct 2026)	Moffatt-Shimomura gyroscopic balance, sliding: J dθ/dt = −μ M g h², J from the egg at 60°, κ taking J: 0.240 s	0.259 s, the same at EG_STEPS 2400 /4800/9600/14400/19200/38400	+8.2 %. Started in the balance (EG_BALANCED=1, no initial nutation): 0.246 vs 0.241 s, +2.1 % (+2.4 % at 19200)
Same, middle (jumps)	0.384 s	0.468 s (0.435 at 2400, 0.469-0.470 at 9600-38400)	+22.0 %; EG_BALANCED=1: 0.401 vs 0.385, +4.2 %
Energy budget, left: lost vs friction + landing work	equal	41.17 vs 41.71 μJ	−1.29 %
Energy budget, middle	equal	106.19 vs 107.08 μJ	−0.83 %
Step-size convergence of the energy budget (EG_STEPS 2400/4800/9600/14400/19200/38400; above 9600 env only, beyond the slider)	-> 0	left: −2.56 / −1.29 / −0.69 / −0.50 / −0.40 / −0.30 %; middle: −1.62 / −0.83 / −0.44 / −0.32 / −0.29 / −0.26 %	both converge (fixed 15 Oct 2026; the left one did not before)
Rise onset vs step (same runs)	should not depend on step	left: rising (tilt < 60°) at 0.30 s and standing at 0.67-0.68 s at every step; J kept to +1.05 / +0.83 / +0.72 / +0.69 / +0.67 / +0.64 %; middle 0.52 s and 1.18-1.22 s	converged (fixed 15 Oct 2026; was 0.50 / 0.65 / 0.90 / 2.45 s / never, see caveats)
Left egg after standing up (EG_STEPS 4800/9600/19200, 1800 frames)	should not depend on step	tilt at 3 s: 4.2 / 7.1 / 9.2°; falling back at 22.33 / 20.62 / 17.60 s (spinning 6.4 / 7.6 / 10.0 t/s); back on its side 26.13 / 25.67 / 25.18 s	**not converged** (open; see caveats). README says so
Jump threshold (Shimomura et al. 2006 observed jumps; README: "faster than about 30 t/s")	—	30 t/s: 1 jump (4.0 ms, < 0.01 mm); 32: 8 jumps (≤0.33 mm); 34: 13 jumps (≤0.58 mm)	OK
Jumps at 40 t/s (README: 24 jumps, ≤0.6 mm, 20 ms)	measured real eggs: fractions of a mm, up to a few 10 ms	24 jumps, 0.32-1.08 s, longest 19.8 ms, highest 0.58 mm	matches README
Jumps vs friction at 35 t/s (README: still jumps for μ 0.2-0.5)	—	μ = 0.2: 20 jumps; μ = 0.5: 9 jumps	OK
Raw egg: shared spin of shell + liquid	W0·I_s/(I_s+I_liq) = 5.5 t/s (I_s 0.0336, I_liq 0.1194)	κ = 0 at 1.2 s: shell 5.81, liquid 5.24 (moment-weighted 5.37, my calculation); default κ: 4.44 / 4.73	κ = 0: −2.3 % (−0.9 % from the old 3° start; the 6° egg touches off-centre and sliding friction takes some spin). README says "nearly do", with these numbers
Raw egg re-spin after finger (κ = 0)	liquid's L shared: 3.38 t/s	3.34 t/s (3.36 vs 3.39 at EG_STEPS=9600)	−1.2 % (−0.9 %)
Raw egg re-spin, default κ	README: "about half"	1.41 vs 2.76 t/s	OK
Falls back after slowing (README: ~20 s, ~6 t/s, step-dependent)	—	falling back at 22.33 s at 6.4 t/s; on its side 26.13 s (4800 steps/s)	OK at 4800, but see the not-converged row

Run used

```
cd demo/WyomingWill/SpinningEgg && HOLD_S=0 timeout 1500 bpp -f spinning-egg.lua -n 1800      (also EG_STEPS=9600 /
    19200 with EG_LOG, -n 1800)
HOLD_S=0 EG_STEPS={2400,9600,14400,19200,38400} ... -n 120;  EG_BALANCED=1 (and with EG_STEPS=19200) ... -n 120
HOLD_S=0 EG_KAPPA=0 ... -n 300 (also EG_STEPS=9600);  EG_KAPPA=0 EG_SPIN={20,30} (EG_STEPS 4800/9600/19200) ... -n 300
EG_FAST={30,32,34} ... -n 150;  EG_FAST=35 EG_MU={0.2,0.5} ... -n 240
EG_TILT=3 EG_STEPS={4800,9600,14400,19200,28800,38400} ... -n 300 (the old start); EG_TILT 2-10 and EG_MU/EG_SPIN
    variations to map the hold
```

```
        Jellett's J = 49.87 (A b W0 = 49.93 lying level); standing up it should spin W0 A b / (C a) = 26.00 turns/s
    (x 1.040)
  Left   rising at  0.30 s (tilt 31 deg, spinning 25.2 turns/s)
  Left   standing on its end at  0.67 s (tilt 10 deg, spinning 25.3 turns/s)
        spinning 25.31 turns/s; by Jellett's J it should be 25.17 (J 48.74, from 49.87, 48.33 with spin friction:
    +0.83%)
        energy lost 41.17 microjoules; friction took 41.73, landings -0.02: 41.71 (-1.29% apart)
        it rose from 60 to 30 deg in 0.259 s; by J dtheta/dt = mu M g h^2 (sliding) it should take 0.240 (+8.2%)
  Middle standing on its end at  1.22 s (tilt 2 deg, spinning 41.0 turns/s)
        spinning 40.95 turns/s; by Jellett's J it should be 40.26 (J 78.64, from 79.80, 77.31 with spin friction:
    +1.73%)
        it rose from 60 to 30 deg in 0.468 s; by J dtheta/dt = mu M g h^2 (sliding) it should take 0.384 (+22.0%)
        it jumped 24 times as it rose (0.32 to 1.08 s): longest 19.8 ms, highest 0.58 mm off the table
  Left   falling back at 22.33 s (tilt 16 deg, spinning 6.4 turns/s)
  [EG_BALANCED=1]        it rose from 60 to 30 deg in 0.246 s; by J dtheta/dt = mu M g h^2 (sliding) it should take
    0.241 (+2.1%)
  [EG_STEPS=19200] Left  falling back at 17.60 s (tilt 15 deg, spinning 10.0 turns/s)
  [EG_TILT=3, EG_STEPS=19200] Left   rising at  2.45 s (tilt 23 deg, spinning 24.0 turns/s)   (28800, 38400: never)
```

The default-run lines match the README's console sample digit for digit. Hall and standalone output IDENTICAL (cmp2.sh, 1800 frames, 21 run lines); HOLD_S=0 for 1200 frames is an exact prefix of the default 0.5 s pause for 1230.

Known departures and honest caveats

- The egg is a symmetric spheroid; a real egg is fatter at one end (README admits this).
- Fixed on 15 Oct 2026: the left egg's rise onset was step-dependent (0.50 / 0.65 / 0.90 / 2.45 s at 4800 / 9600 / 14400 / 19200 steps/s). Cause, by experiment: started 3° from level, the egg sat at the edge where Coulomb friction can hold the contact point still. Converged (28800 and 38400 steps/s, the first followed for 20 s) the left egg never slides: it rolls on its side without slipping (contact slip 0 after each step, friction at 50-100 % of μN), drifts to level and slows; it never rises. At coarser steps the step's first-order errors tip it up until it breaks free. The hold is a stick/slip threshold, not a noise-seeded unstable equilibrium: once sliding, the rise is deterministic (Coulomb friction gives a finite rise rate, not exponential growth), and after breaking free the 3° egg rose from 60° to 30° in 0.233-0.241 s at every step that let it go (Moffatt-Shimomura: 0.227-0.239, the J left by then). Measured threshold at μ 0.3, 25 t/s: between 3° and 4° (4°: rises at 0.35 s at every step; 3.5°: 0.38 vs 0.42 s). It moves the expected way: μ 0.2 at 3° rises at 0.50 s at both 4800 and 38400; the 40 t/s egg rises from 1° and 1.5° at both 4800 and 38400. Fix: start at 6°, about twice the threshold, stated in the script and README (EG_TILT sets it). Not tuned to a theory value; the rise check is the rate (60° → 30°) against Moffatt-Shimomura, not the onset. All three eggs start at 6°, so the middle and raw eggs' numbers changed a little (24 jumps instead of 21; raw-egg sharing −2.3 % instead of −0.9 %).
- Still open: up on its end (about 10° tilt) the egg rolls without slipping, and the same first-order step error slowly tips it further up: tilt at 3 s 4.2° / 7.1° / 9.2° at 4800 / 9600 / 19200, so the time it starts falling back is 22.3 / 20.6 / 17.6 s (at 6.4 / 7.6 / 10.0 t/s). Not converged; the README now says so. Likewise at 20 t/s (EG_SPIN=20 EG_KAPPA=0) the egg rises to about 17° identically at every step but then stands (< 15°) at 1.25 s at 4800, 2.13 s at 9600, not within 5 s at 19200. The error comes from enforcing no-slip at the start-of-step lowest point while the contact runs round the egg at the spin rate (a fix to the translation, moving the centre on the exact arc about the contact, changed nothing). Evaluating the contact at the mid-step orientation cut the standing drift about 7x (tilt at 3 s 13.1° at 4800 vs 12.3° at 19200) but at 4800 and below it damped the nutation during the rise (60° → 30° in 0.238 s instead of the converged 0.259), so it was not adopted. A proper second-order rolling contact is still to do.

- The rise-rate check: the left egg rises 8 % slower than Moffatt-Shimomura's leading-order balance; the measured axis of rotation sits 7.65° from vertical at 64.4° tilt against the balance's 7.84° ($\omega = ((n - \Omega \cos\theta) \sin\theta, \theta, \Omega \sin^2\theta + n \cos\theta)$, $C\, n = A\, \Omega \cos\theta$). The gap is the nutation from starting with pure spin about the vertical: started in balance it is 2 %. The middle egg is 22 % slower (4 % from balance); it jumps throughout its rise.
- Jellett's J is only approximately conserved. The script measures a 0.8-1.7 % change and corrects the prediction for the J removed by spin friction (and rolling resistance and the rim; the console calls the sum "spin friction", also with EG_KAPPA=0), so the "theory" value (25.17) already includes a model-side correction. The bare Moffatt-Shimomura prediction is 26.00 t/s. The first console line now gives the egg's actual J at the start (49.87) and A b W0 lying level (49.93); before, it labelled J0 as "A b W0".
- Spin friction κ and rolling resistance δ (100 μm each) are chosen parameters. κ sets how long the egg stays up; δ stops a fallen egg rolling across the dish.
- Contact is a sequential-impulse solver, not an exact unilateral constraint. Landings are perfectly inelastic, so jump heights and durations depend on that solver. The left egg's "landings −0.02" μJ is the normal impulse's work on the small Baumgarte overlap, not a landing.
- The raw egg's liquid is a two-rotor model: added inertia plus a linearly coupled rotor with a time constant of 1.5 s. It is not a fluid simulation, and the 1.5 s is an assumed value.

References (as cited in README)

- H. K. Moffatt and Y. Shimomura, "Spinning eggs — a paradox resolved", Nature 416, 385 (2002).
- Y. Shimomura and colleagues, Proc. R. Soc. A (2006), on spinning eggs jumping; reported in Science News, 23 May 2006, "Well-spun egg also jumps".
- The rise equation $J\, d\theta/dt = -F\, h^2$ and the balance $C\, n = A\, \Omega \cos\theta$ as restated in K. Sasaki, "Spinning eggs — which end will rise?", arXiv:physics/0310163 (eqs. 7, 9-11).

4.9 The tippe top

Agrees with theory, with a numerical or open caveat. Inversion criterion right for all three tops; centre of mass rises 1.057 to 2.545 cm as theory says; stem threshold about 80 rad/s, within the sleeping-top rule's bounds. Open: the green top's spin collapses after 40 s without rolling resistance.
Folder: demo/WyomingWill/TippeTop/

What it shows

A ball with a short stem, its centre of mass a little below the ball's centre. Spun on its ball, stem up, it turns itself over within a few seconds and spins on its stem, raising its centre of mass; sliding friction at the table does the work. Three tops: red, spun fast (40 turns/s), turns over and stands on its stem; blue, the same top spun slowly (15 turns/s), only gets as far as lying on ball and stem; green, with its centre of mass almost at the ball's centre, never turns over. All end stem up again as they slow.

Model

- Top: one Bullet rigid body, 15 g, hollow ball R = 1.5 cm; $I3 = 2/3\ m\ R^2 = 0.0225$ kg cm^2 about the stem, $I1 = 0.9\ I3$ across. Centre of mass a = 0.45 cm below the ball's centre (alpha = a/R = 0.3); green top a = 0.06 cm (alpha = 0.04).
- Contact shape: Bullet compound of a true sphere (the ball) and a capsule (the stem, round tip radius 0.6 cm standing 0.6 cm out). Sliding friction 0.2 (Bullet), restitution 0, contact processing threshold 0.001.
- Bullet's gyroscopic term is off; before every physics step the script advances the body-frame angular velocity by Euler's equations with RK4 (as in the rattleback exhibit).
- Script-applied losses: "spin friction", a deceleration 0.5 rad/s^2 of the spin component about the vertical; rolling resistance, a constant moment delta*m*g against the horizontal angular velocity,

delta = 30 microns, with the matching linear-velocity change about the contact.

- 4800 physics steps/s (80 per 1/60 s frame), no damping, g = 981 cm/s$^{2.}$ Initial state: stem up tilted 12 deg (random azimuth), spin about the stem.

Theory checks

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Inversion possible, red/blue (alpha = 0.3)	1 - alpha < I1/I3 < 1 + alpha: 0.7 < 0.9 < 1.3 (README)	red inverts (on its side 2.5 s, up on its stem 3.5 s); blue reaches 124-127 deg (ball and stem)	qualitative, agrees
Inversion impossible, green (alpha = 0.04)	0.9 outside 0.96-1.04 (README)	max tilt 1 deg (2 s), 0 deg thereafter; never turns over in 120 s	agrees
Centre-of-mass height, start	R - a cos(12 deg) = 1.06 cm	1.057 cm (TT_LOG)	agrees
CoM height standing on stem	stem-tip contact 2.55 cm below the CoM: 2.55 cm	max 2.545 cm, 2.49 cm at 10 s (tilt 166 deg)	agrees (CoM rises ~1.5 cm)
Green top CoM height	R - a = 1.44 cm	1.441 -> 1.440 cm	agrees
Time to turn over (red)	README now: "about 3.5 seconds", example "Left up on its stem at 3.5 s" (fixed 15 Oct 2026; was "about 3", "3.3 s")	on its side 2.5 s, up on its stem 3.5 s (re-run identical)	matches
Spin needed to stand on the stem	README/lua now: "about 80 rad/s or more" with the rule's bounds (fixed 15 Oct 2026; was "about 100"). w > 2 sqrt(I1' m g d)/I3, d = 1.95 cm: 68 rad/s with I1' about the CoM (tip sliding freely); 132 (I about the tip-sphere centre) to 163 rad/s (I about the contact) for a tip that can't slide	TT_LOG: up on its stem with ~91 rad/s; steady (tilt 171-173 deg) while >= 83 rad/s; tilt first below 170 deg at 24.0 s with 81.8 rad/s; onto ball and stem at 38.7 s; blue never above 44 rad/s after 8.3 s, never stood	between the bounds, nearer the slippery one
Red returns stem up	README: sinks back "twenty-odd seconds later"; all three stem up in ~1.5 min	sinking from ~24-26 s; ball+stem 38.7 s, on ball 66.6 s, 0 deg by 92 s; blue 0 deg by 76 s	agrees
Without spin friction	README: red "spins on its stem for ever"	TT_SPINF=0: 175 deg, 14.7 t/s still at 60 s	agrees (60 s run)
Rolling resistance, effect (TT_ROLL sweep, 15 Oct 2026)	README/lua now give the measured effect (fixed; was "100 microns stops the top turning over at all")	red up on its stem at 3.4-3.7 s for every value 0-300 µm; back onto ball and stem at 47.0 s (0), 22.4 (10), 38.7 (30), 37.1 (50), 33.8 (75), 30.0 (100), 23.1 (150), 12.8 s (300). Blue: reaches ball and stem at 30 µm (and 10), stays within 8 deg (2-s reports) at 50-300. At 0 nothing settles: blue rocks 16-65 deg, green's tilt grows from ~40 s to 71 deg at 78 s with its spin collapsing (36 -> 0.4 t/s)	documented
Determinism	(repeat run)	identical event times on a second run	reproducible

No conservation check is printed by the exhibit; Jellett's integral (the quantity conserved during sliding on the ball) was not logged and was not checked here.

Run used

```
cd demo/WyomingWill/TippeTop
HOLD_S=0 bpp -f tippe-top.lua -n 7200           # main run, 120 s
HOLD_S=0 TT_LOG=tt.csv bpp ... -n 600           # CoM heights
HOLD_S=0 TT_ROLL=100 bpp ... -n 1200            HOLD_S=0 TT_SPINF=0 bpp ... -n 3600
# 15 Oct 2026: HOLD_S=0 TT_LOG=tt.csv bpp ... -n 7200 (same events as above);
#   TT_ROLL={0,10,50,75,100} -n 4800; TT_ROLL={150,300} -n 1800
```

```
  Left   on its side at   2.5 s (stem 83 deg from up)
  Left   up on its stem at   3.5 s (stem 156 deg from up)
t=  4.0s  stem from up, spin: Left 161 deg 14.6 t/s | Middle  10 deg 13.9 t/s | Right   1 deg 37.2 t/s
  Middle on its ball and stem at   8.3 s (stem 124 deg from up)
t= 26.0s  stem from up, spin: Left 160 deg 12.6 t/s | Middle 113 deg  5.0 t/s | Right   0 deg 35.4 t/s
  Left   on its ball and stem at  38.7 s (stem 127 deg from up)
  Middle spinning on its ball at  49.5 s (stem 59 deg from up)
  Left   spinning on its ball at  66.6 s (stem 59 deg from up)
t= 92.0s  stem from up, spin: Left   0 deg  1.2 t/s | Middle   0 deg  0.1 t/s | Right   0 deg 30.0 t/s
  Left   up on its stem at   3.4 s (stem 179 deg from up)        [TT_ROLL=100]
```

Known departures and honest caveats

- Several parameters were tuned to make the exhibit work (README): a hollow ball (a solid 2/5 m R^2 ball turned over but never stood up), a short round-ended stem (a long thin one stuck leaning), rolling resistance 30 microns (to settle the tops), spin friction 0.5 rad/s^2 (otherwise the red top never comes down).
- Spin friction and rolling resistance are phenomenological torques applied by the script, not contact physics; contact friction is Bullet's (Coulomb, 0.2).
- Event times depend on the random tilt direction (math.random); two runs here gave identical times, but other builds/Lua versions may differ.
- The stand-on-stem threshold is not a sharp prediction; the stem tip is a 6 mm sphere, and contact friction makes the effective criterion lie between the slippery and fixed-point bounds I estimated (68-132 rad/s).
- Fixed 15 Oct 2026: README and lua said 100 microns of rolling resistance stops the (red) top turning over; it does not, at any value tried up to 300 µm. They now say what it does: settles the tops (with none they rock on), brings the red one down off its stem sooner (not monotonically: 10 µm 22 s), and from 50 µm keeps the slow blue one from turning over. Also fixed: the README's console example (3.3 s -> 3.5 s), "about 3 seconds" -> 3.5, the stem threshold "about 100 rad/s" -> about 80 with the run and the rule's bounds, and the header's env-var list (TT_SPINF, TT_ROLL added).
- Open: with no rolling resistance the green (centred) top, which cannot invert, starts rocking after ~40 s and loses its spin within ~10 s (25.6 -> 3.2 t/s between 38 and 46 s; the spin friction alone would take 0.6 t/s), far faster than the spin friction (0.5 rad/s^2) allows. Not investigated; likely a contact-solver effect that the rolling resistance otherwise damps. The README says it hasn't been looked into.
- Text only; physics and console unchanged (hall vs standalone IDENTICAL, 900 frames).
- The green "centred" top spins on for a long time (27.7 t/s at 120 s), slowed only by spin friction; it is "stem up" throughout.
- The state labels (on its side, on ball and stem, up on its stem) come from tilt and height thresholds held for 0.33 s; reported times lag the actual change by that much.

References

None cited in the README or code (the lua header mentions Bohr and Pauli watching one). The inversion condition 1 - alpha < I1/I3 < 1 + alpha is the standard result from the tippe-top literature, not referenced in the exhibit.

4.10 The Dzhanibekov effect

Agrees with theory. Flip period 3.3129 s against 3.3131 s from the exact elliptic-function solution (-0.01%), 36 flips in 120 s. Energy and angular momentum are held by projection, and the README now says so. New README.
Folder: demo/WyomingWill/Dzhanibekov/

What it shows

Three identical aluminium T-handles shaped like a wing nut float without gravity, each spun at 1 turn/s about a different principal axis with the same 1% extra wobble. The one spun about its intermediate axis (the stem) flips end over end every few seconds, again and again, as Vladimir Dzhanibekov's wing nut did on Salyut 7 in 1985; the two spun about the smallest (bar) and largest (across both) axes just spin steadily (the "tennis racket theorem").

Model

(Facts from the lua header, `dzhanibekov-meshes/t-handle-data.lua`, and README.md, written 15 Oct 2026.)

- Body: solid aluminium (2.7 g/cm^3) T-handle, bar r = 0.5 cm x 10 cm along body x, stem r = 0.8 cm x 4 cm along -y; mass 42.92 g; principal moments I = (0.102132, 0.184989, 0.277521) kg cm^2 about (bar, stem, across). Stem = middle moment.
- Each handle is a Bullet rigid body (convex-hull shape used only for mouse hover; no gravity, no contacts), damping 0. Bullet integrates only the orientation from the angular velocity.
- Bullet's gyroscopic term is off (setFlags(0)). Before every physics step the script advances the body-frame angular velocity by Euler's free-body equations with classical RK4, then projects it back onto the initial kinetic energy and |L| (two Newton steps along the gradients of E and L^2), and writes it back to Bullet.
- 600 physics steps/s by default (10 per 1/60 s frame; slider/DZ_STEPS 120-2400). Initial state: w = W about the chosen axis plus eps*W about the next axis, W = 2pi rad/s, eps = 1%.
- Console: each sign change of the spin component along the chosen axis is a "flip"; every 10 s it prints relative energy and angular-momentum error and the maximum tilt of the spin axis from vertical.
- Companion script `bullet-rotation-test.lua` (I = 1:2:3, w0 = (0.01, 10, 0) rad/s) tests Bullet's own gyroscopic options against Euler's equations.

Theory checks

Exact values computed now from Euler's equations: closed form T = 4K(k)/w_p with w_p = sqrt((I3-I2)(L^2-2E I1)/(I1 I2 I3)), k^2 = (I2-I1)(2E I3-L^2)/((I3-I2)(L^2-2E I1)) (case L^2 > 2E I2, here k^2 = 0.99968), cross-checked by a high-precision ODE integration (scipy DOP853, rtol 1e-11) that also gives first-flip times and axis tilts.

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Time between flips, stem (middle axis)	T/2 = 2K(k)/w_p = 3.3131 s	mean of 35 intervals 3.3129 s (each printed as 3.30 or 3.32, frame-quantised)	-0.01%
Time of first flip	ODE: 1.657 s (1.6566)	1.67 s (reported at frame end, 1/60 s resolution)	within one frame
Flips in 120 s	ODE: 36	36 (last at 117.62 s; ODE 117.615 s)	exact
Bar (smallest axis): steady	ODE: 0 flips, axis tilt max 2.08 deg	0 flips, "axis wandered up to 2 deg"	agrees (console rounds to 1 deg)
Across (largest axis): steady	ODE: 0 flips, axis tilt max 0.60 deg	0 flips, "up to 1 deg"	agrees to the console's rounding
Energy conservation (all three, 120 s)	E constant	worst printed relative error 4.3e-7	~single-precision level
Angular momentum conservation	L constant	worst printed relative error 2.2e-7	~single-precision level

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Flip period claim (lua header)	header now: "3.3129 s between flips (mean of 35, over 2 minutes) against the exact 3.3131 s" (fixed 15 Oct 2026; was "to 0.2%, with energy and angular momentum exact")	3.3129 s (re-run 15 Oct 2026: 36 flips, last 117.62 s)	-0.01%
Bullet gyro OFF, no Euler step (test, 60 s)	header: "nothing flips"	0 flips (exact: 21)	confirms
Bullet "explicit" gyro (flag 2)	header: energy +59%	E +59.1%, L +26.1%; mean flip interval 2.34 s vs exact 2.873 s	confirms
Bullet default "implicit" (flag 8)	header/README now: energy -31.9%, L -17.5%, flips on time (fixed 15 Oct 2026; header said -44% / -25%)	re-run 15 Oct 2026: E -31.9%, L -17.5%; first flip 1.433 s vs exact 1.437 s; same with flag 4	matches
Implicit at 16x steps (9600 Hz)	header now: "-3.3% and -1.7% a minute"	E -3.3%, L -1.7% (re-run)	confirms

Linearised instability rate (for context, not printed by the exhibit): lambda = W sqrt((I2-I1)(I3-I2)/(I1 I3)) = 3.27 /s (e-folding 0.31 s).

Run used

```
cd demo/WyomingWill/Dzhanibekov
HOLD_S=0 DZ_LOG=dz.csv bpp -f dzhanibekov.lua -n 7200
DZ_FLAGS={0,2,4,8} DZ_LOG=brt_N.csv bpp -f bullet-rotation-test.lua -n 3600   (plus DZ_FLAGS=8 DZ_HZ=9600; DZ_FLAGS=0
    DZ_EULER=1)
15 Oct 2026 re-runs: HOLD_S=0 bpp -f dzhanibekov.lua -n 7200; DZ_FLAGS={0,2,4,8} and DZ_FLAGS=8 DZ_HZ=9600, -n 3600
    (all as before)
```

```
--- spin 1.0 turns/s, nudge 1.0% ---
  Left   flipped at   1.67 s  (flip 1)
  Left   flipped at   4.98 s, 3.32 s after the last  (flip 2)
  Left   flipped at   8.28 s, 3.30 s after the last  (flip 3)
t=   10s  Left: flips 3, axis wandered up to 90 deg, energy +8.0e-08, ang. mom. +4.0e-08 | Middle: flips 0, axis
    wandered up to 2 deg, energy +1.3e-07, ang. mom. +6.7e-08 | Right: flips 0, axis wandered up to 1 deg, energy
    +2.3e-07, ang. mom. +1.2e-07
t=   70s  Left: flips 21, axis wandered up to 90 deg, energy -8.3e-08, ang. mom. -4.1e-08 | Middle: flips 0, axis
    wandered up to 2 deg, energy +3.1e-07, ang. mom. +1.5e-07 | Right: flips 0, axis wandered up to 1 deg, energy
    +4.3e-07, ang. mom. +2.2e-07
  Left   flipped at 117.62 s, 3.30 s after the last  (flip 36)
```

The bullet-rotation-test figures come from its CSV logs (the script prints nothing), analysed now.

Known departures and honest caveats

- The rotational dynamics are the script's, not Bullet's: Bullet's own gyroscopic options were found inadequate (above), so Euler's equations are integrated in Lua with RK4 plus an explicit projection onto constant E and |L|. Conservation is therefore enforced, not emergent; the flip period is the real test, and it matches. Since 15 Oct 2026 the header and README say so plainly (the header said energy and L were "exact" and the console "checks" them; the console's E/L figures, at most 4.3e-7 and 2.2e-7, show only single-precision rounding).
- Bullet is single precision: E and L hold only to ~1e-7. The header admits that with a vanishingly small nudge rounding noise starts the wobble sooner. Seen clearly in the test script's own Euler mode (energy-only projection, 0.1% nudge): mean flip interval 2.52 s vs exact 2.873 s (-12%), because the energy noise (about +/-1.5e-6 relative) exceeds the orbit's distance from the separatrix (~2.5e-7). At the exhibit's 1% nudge this does not matter (0.01%).
- Fixed 15 Oct 2026: the header's figures for Bullet's default implicit integrator (-44% energy, -25% L per minute) were not reproducible; re-run gives -31.9% / -17.5% at 600 Hz (same with flag 4), and header and README now quote those.
- No step-size convergence study is reported; only the default 600 steps/s was run here.
- Which axis is "middle" depends on proportions (header: with a 7 cm stem the bar becomes the middle axis); the 4 cm stem is a design choice.

- Fixed 15 Oct 2026: the header's headless-test path and the data-file fallback pointed at `demo/dzhanibekov/`; both now `demo/WyomingWill/Dzhanibekov/`, and the exhibit runs from the repository root (checked: first flip 1.67 s as from the folder).
- Fixed 15 Oct 2026: README.md written (what it shows, keys, theory and console, physics with the Bullet test results, files, env vars).
- Not changed: `bullet-rotation-test.lua`'s own comment says its DZ_EULER mode holds E and L, but the code projects energy only (the README says energy only).
- Text and paths only; physics and console unchanged (hall vs standalone IDENTICAL, 900 frames).

References

None cited beyond the historical note (Dzhanibekov, Salyut 7, 1985) in the lua header. Elliptic-function solution of Euler's equations: standard (e.g. Landau & Lifshitz, Mechanics, sec. 37) — added here for the check, not cited by the exhibit.

4.11 The gyroscope

Agrees with theory, with a numerical or open caveat. Precession 2.58 rad/s against m g r/(I ω) = 2.59 at the spin it has, and to 0.0-0.5% with the tilt term as it sags; README now right (2.56 rad/s at 100 turns/s). Rocked 1.5° it slides further than aimed for, cause not separated.
Folder: demo/WyomingWill/Gyroscope/

What it shows

Three identical toy gyroscopes spun to 100 turns/s. Left: held only by one end of its axle on a pedestal point, it does not fall but precesses with the axle level. Middle: the same, hung by a string. Right: axle upright, standing on a tightrope on a forked foot, balancing like a top on its point; an attendant rocks the wire 1 degree each way so it slides to and fro, upright, about 9 cm each way. As the rotors slow, the left two precess faster and sag; the tightrope one wobbles wider and finally comes off.

Model

- Gyroscope: one Bullet rigid body (hoop 50 g, axle 10 g, rotor 100 g; M = 160 g). Rotor r = 3 cm, I_rotor = 0.65 m r^2 = 0.585 kg cm^2; body moments (code): axial 1.069, in-hoop-plane 0.849, across-hoop 1.333 kg cm^2. Holding balls at r = 6.0 cm from the centre; the foot's wire contact 6.38 cm (PIV + ball + wire radius).
- Rotor spin is a scalar angular momentum h = I_rotor*S inside the body (gyrostat); Bullet never spins a body at 100 turns/s. Bullet's gyroscopic term is off; the script applies -w x (I w + h) as an external torque each step, 2nd-order Adams-Bashforth (README: a one-step torque grows the wobble; a direct velocity update makes it sag steadily).
- Pedestal: btPoint2PointConstraint at the ball's centre; released when the axle sags 40 deg below level. String: a 2 g, 12 cm rigid rod with ball joints at both ends. Wire: kinematic capsule, radius 0.06 cm, 30 cm long; foot = sphere + two capsule prongs; foot-wire friction 0.005 (slides on slopes > 0.3 deg).
- Losses: rotor bearing 1 turn/s per second; pivot friction 0.5e-3 N m against any turning at the pedestal (a quarter at the string's knot). Rocking: wire tipped +/-1 deg about its middle in 0.6 s (cosine ramp), always up on the gyroscope's side, judged 0.3 s ahead, with timing adapted to reach ~9 cm each way.
- 2400 physics steps/s, 30 solver iterations, no damping, g = 981 cm/s^2.

Theory checks (theory values computed now from the code's constants unless marked README)

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Precession rate at 100 turns/s	Omega = m g r / (I_rotor w) = 0.16*981*6/(0.585*628.3) = 2.562 rad/s (2.45 s per turn); README/lua now say the same (fixed 15 Oct 2026; they said 2.67 rad/s, 2.35 s)	2.582 rad/s over 0.5-2 s (mean spin ~99 t/s; theory 2.588); 2.656 rad/s over 0-8 s at mean spin 96 t/s (theory 2.669)	-0.2% / -0.5%

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Turns round in first 8 s (100 -> 92 t/s)	integral of m g r/(I w(t)) = 3.40 turns; README now "3.35 turns ... (theory 3.40; the console rounds it to 3.4)"	pedestal 3.35, string 3.36 (from GY_LOG); console "3.4 times round"	-1.5%
Precession as it sags (pedestal), with tilt term	m g r = Omega h + Omega^2 (I_pivot - I_axial) sin(theta), I_pivot = 7.093, I_axial = 1.069 kg cm^2	8-16 s 2.878 vs 2.877; 16-24 s 3.132 vs 3.132; 24-32 s 3.413 vs 3.430; 32-36 s 3.692 vs 3.685	0.0 to -0.5% (first-order formula alone: -1% to -5%)
Sag with no pivot friction (dip from the wobble)	m g r l / (I_rotor w)^2 (I = 6.6-7.1 about pivot): 2.6-2.8 deg at 100 t/s, 7.3-7.9 at 60, 16.5-17.7 at 40. README: "3 deg at 100, 16 at 40"	GY_PIVF=0: pedestal 8.3 deg at 60 t/s, 17.3 deg at 40 t/s; 0.1-3.4 deg sampled at 92-98 t/s	agrees (within the I range)
Sleeping-top criterion on the wire	I3 w > 2 sqrt(I m g l), l = 6.38 cm, I about the foot 7.36-7.85 kg cm^2: 46.7-48.2 t/s. README: "above 46 (46.7 to 48.2 ...)"	still wire, rough (GY_MU=0.3): off at 52 t/s (48.2 s)	README: 52, matches
Still wire, slippery foot (0.005)	README: off at 45 t/s; with I about the CoM (ice rule) the bound is 15.9-19.9 t/s	off at 45 t/s (55.0 s)	matches README
Rocked 1 deg (default)	README: lasts 62 s, off at 38 t/s	off the wire at 62.0 s, 38 t/s	matches
Rocked 1.5 deg	README/lua now: 49 s, at 51 t/s, shorter than the still wire (fixed 15 Oct 2026; they said 55 s, "as long as on a still wire, or longer")	off at 49.0 s, 51 t/s	matches
Rocked 2 deg	README: thrown over in 13 s	off at 12.7 s	matches
Rocked 3 deg	README: 6 s	off at 5.9 s	matches
Slide range when rocked 1 deg	README: "about 9 cm each way (5 to 12 cm; the first time 14 ...)"	turn points (scratch copy logging the attendant's turn detection, 0-55 s): 13.9 first, then 4.8-12.5 cm	matches
Slide range when rocked 1.5 deg	README: stops 1 to 15 cm out; 14-15 cm in its last 6 s, attendant at its 4 cm limit	turn points 1.2-14.7 cm; 14.1, 13.9, 14.5, 14.7 cm at 42-47 s with the timing offset at its clamp (-4 cm)	matches (documented, not fixed)
Pedestal slides off / string stops	README now: pedestal off "after 37 seconds"; tightrope off "after a minute, at 38 turns a second"; string rotor stops at 100 s	pedestal off at 37.2 s (63 t/s); tightrope off 62.0 s (38 t/s); string spin 0 t/s at 100.0 s	matches

Run used

```
cd demo/WyomingWill/Gyroscope
HOLD_S=0 GY_LOG=gy.csv bpp -f gyroscope.lua -n 7200          # defaults: rock 1 deg
HOLD_S=0 GY_ROCK=0 bpp ... -n 7200     HOLD_S=0 GY_ROCK=0 GY_MU=0.3 bpp ... -n 5400
HOLD_S=0 GY_ROCK=1.5 bpp ... -n 5400   HOLD_S=0 GY_ROCK=2 ... -n 3600   HOLD_S=0 GY_ROCK=3 ... -n 1800
HOLD_S=0 GY_PIVF=0 bpp ... -n 3600
# re-run 15 Oct 2026 (after the text fixes): GY_ROCK = 1, 1.5, 2, 3, 0 with GY_LOG, -n 4800; GY_ROCK=0 GY_MU=0.3 -n
    4200
# 1.5 deg investigation: a scratch copy printing the attendant's turn points and timing offset (rock.late), GY_ROCK=1
    and 1.5, -n 3300
```

```
t=  8.0s  Pedestal below level  9.5 deg,  92 t/s, 3.4 times round | String below level  5.4 deg,  92 t/s, 3.4 times
    round | Tightrope from upright  4.1 deg,  92 t/s,  +0.8 cm along the wire, tipped -0.63 deg
t= 14.0s  ... Tightrope from upright  5.0 deg,  86 t/s,  +9.2 cm along the wire, tipped -1.00 deg
  Pedestal  off the pedestal at  37.2 s (spin 63 turns/s)
  Tightrope off the wire at  62.0 s (spin 38 turns/s)                        [rock 1]
  Tightrope off the wire at  55.0 s (spin 45 turns/s)                        [rock 0]
  Tightrope off the wire at  48.2 s (spin 52 turns/s)                        [rock 0, mu 0.3]
  Tightrope off the wire at  49.0 s (spin 51 turns/s)                        [rock 1.5]
  Tightrope off the wire at  12.7 s (spin 87 turns/s)                        [rock 2]
  Tightrope off the wire at   5.9 s (spin 94 turns/s)                        [rock 3]
t= 60.0s  Pedestal below level 17.3 deg,  40 t/s, 36.6 times round | ...      [pivot friction 0]
```

(The t=14.0 s line is shortened with "..."; precession rates are least-squares fits to the logged axle heading.)

Known departures and honest caveats

- The rotor is not simulated as a spinning body; the gyroscopic torque is applied explicitly (Adams-Bashforth). The README says the more obvious alternatives either grow the wobble or make it sag steadily.
- Sagging comes from a chosen pivot-friction torque (0.5e-3 N m), not from a contact model; the 40 deg release from the pedestal is a scripted rule; the string is a rigid 2 g rod, not a flexible string.
- The foot friction 0.005 ("as if it ran on a little wheel") is chosen so the gyroscope slides on a 1 deg tilt. The attendant is a feedback controller (lead 0.3 s, adaptive timing), not physics.
- Fixed 15 Oct 2026: the precession rate in README and lua header (was 2.67 rad/s, 2.35 s per turn, at 100 t/s; that is the value at about 96 t/s) now reads 2.56 rad/s, 2.45 s, from the stated m = 160 g, r = 6 cm, I = 0.65 m r^2; the "3.4 turns in 8 s" now says 3.35 (theory 3.40, console rounds to 3.4).
- Fixed 15 Oct 2026: rocked 1.5 deg, README and header now give the measured 49.0 s (51 t/s), shorter than the still wire's 55.0 s; only 1 deg (62.0 s) outlasts the still wire. The "about 40"/"about 50" turns/s wording is replaced by the run values: rough still wire 52, slippery still wire 45, rocked 1 deg 38; pedestal off at 37 s.
- Still open, now documented in the README: rocked 1.5 deg the attendant does not hold 9 cm. Its turn points scatter from 1.2 to 14.7 cm (at 1 deg: 4.8-12.5 after a first 13.9). The attendant's correction is an integral on the reach error (gain 0.3, offset clamped at +/-4 cm); at 1.5 deg the same timing error gives 1.5x the distance, and in the last 6 s before it falls it overshoots to 14-15 cm with the offset at its clamp. The wobble is wider (9-17 deg from upright vs 2-8 at 1 deg, 17-28 deg in the last swings); the README says that most likely the wobble pushes the foot along too, and that this has not been separated from the slope's push. Nothing in the physics was changed.
- The sleeping-top threshold is only approximate here: the foot is a sphere sliding on a thin capsule, not a fixed point; the README itself explains the slippery foot shifts the relevant I toward the centre of mass.
- No step-size convergence study is documented; all runs at 2400 steps/s.

References

None cited in the README or code. Formulas used are the textbook gyroscope precession rate and the sleeping-top (Maxwell/Klein-Sommerfeld) stability criterion.

4.12 The bicycle wheel

Agrees with theory, with a numerical or open caveat. Stool's turning matches angular-momentum conservation exactly with friction off; L kept to 0.093% over 120 s. A guard now reports and recovers from a NaN blow-up, whose cause (seen once) is still unknown.
Folder: demo/WyomingWill/BicycleWheel/

What it shows

A wooden lay figure on a swivel stool, feet off the floor, holds a spinning bicycle wheel with its axle upright; the stool is still. Turning the wheel over makes the stool turn the other way; turning it back stops the stool. Pushing the wheel out to arm's length slows the stool, pulling it in speeds it up again (the figure-skater effect). Nothing pushes on the floor: it is angular momentum about the stool's axis being shared between wheel and figure.

Model

Three Bullet rigid bodies (code, README):

- Stool seat + figure: one body, 75 kg, I = 15000 kg cm^2 (1.5 kg m^2) about the stool axis, on a vertical btHingeConstraint to a fixed anchor; bearing friction 0.02 N m (slider `stool friction`, applied by the script as a saturated torque).

- Hands: 1 kg, I = 100 kg cm^2, on a btSliderConstraint straight out in front, motor-driven between 34 cm (close) and 50 cm (out) in 1.5 s (smoothstep path + proportional correction, gain 40).
- Wheel (axle + handles): 3 kg, I = 1150 kg cm^2 about the axle (0.115 kg m^2), 600 across; hinged to the hands along the slide axis; a hinge motor turns it over in 1.2 s.
- The wheel's spin is not a Bullet rotation: it is a scalar angular momentum h = I3·S carried inside the body (a gyrostat). Bullet's gyroscopic term is off (setFlags(0)); the script applies the gyroscopic torque -w x (I w + h) each step, 2nd-order Adams-Bashforth (1.5·T_n - 0.5·T_(n-1)). The hinge reaction to that torque is what turns the stool.
- Wheel bearing friction 0.02 turns/s per second; the lost spin is handed to the axle as torque.
- 1200 physics steps/s (20 per 1/60 s frame), 30 solver iterations, no damping, gravity 981 cm/s^2 (irrelevant to the physics shown).
- The "should" value printed after each move is computed in Lua: (L0 - I3·S·(axle·up)) / (I_P + (m_w+m_hands)·d^2 + I_w,eff + I_hands), with L0 = total angular momentum about the stool axis when the move started.

Theory checks

Spin 4 turns/s: h = 0.115 x 2pi x 4 = 2.890 kg m^2/s. I_close = 1.5 + 4 x 0.34^2 + 0.115 + 0.010 = 2.087 kg m^2; I_out = 1.5 + 4 x 0.50^2 + 0.115 + 0.010 = 2.625 kg m^2 (hand calculation from code constants; README quotes 2.09 and 2.63).

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Stool rate, wheel turned over, held close (no friction)	w = 2h/I_close = 2 x 2.890/2.087 = 2.769 rad/s	2.769 (first round); 2.770 later rounds	0.000 / +0.001 rad/s (<0.04%)
Stool rate, pushed out (no friction)	w = 2h/I_out = 2.202 rad/s	2.202	0.000
Stool rate, pulled back in (no friction)	2.769 rad/s	2.769 (2.770 later)	<0.04%
Stool rate, axle up again (no friction)	0	0.000 (0.001 in rounds 5-6)	<=0.001 rad/s
Angular momentum about stool axis (no friction, 120 s, 6 rounds)	conserved; README: "within 0.1%"	max deviation 0.093%, end +0.050% (from BW_LOG)	within README bound
Stool rate vs console "should", defaults (friction on), 24 moves	console theory from L at move start	turned over 2.719 vs 2.736 ... 1.136 vs 1.152; out 2.099 vs 2.114; in 2.565 vs 2.584; up -0.021 vs -0.018	sim 0.015-0.020 rad/s below (0.6-1.4%), as README predicts (~0.02 rad/s friction during move)
README friction sequence, turned over & close	README: 2.72, 2.46, 2.19 rad/s	2.719, 2.456, 2.192	matches
README frictionless table	README: 2.769 / 2.202 / 2.769 / 0.000	2.769 / 2.202 / 2.769 / +0.000	matches

Run used

```
cd demo/WyomingWill/BicycleWheel
HOLD_S=0 BW_STOOLF=0 BW_WHEELF=0 BW_LOG=bw0.csv bpp -f bicycle-wheel.lua -n 7200   # frictionless
HOLD_S=0 BW_LOG=bw.csv bpp -f bicycle-wheel.lua -n 7200                            # defaults
```

Frictionless:

```
t=  3.7s  wheel turned over (34 cm out): the stool turning at +2.769 rad/s (0.44 turns/s); it should, by its angular
    momentum: +2.769
t=  8.5s  wheel pushed out (50 cm out): the stool turning at +2.202 rad/s (0.35 turns/s); it should, by its angular
    momentum: +2.202
t= 13.0s  wheel pulled in (34 cm out): the stool turning at +2.769 rad/s (0.44 turns/s); it should, by its angular
    momentum: +2.769
t= 17.3s  wheel axle up again (34 cm out): the stool turning at +0.000 rad/s (0.00 turns/s); it should, by its
    angular momentum: +0.000
t=118.1s  wheel turned over (34 cm out): the stool turning at +2.770 rad/s (0.44 turns/s); it should, by its angular
    momentum: +2.770
```

Defaults:

```
t=  3.7s  wheel turned over (34 cm out): the stool turning at +2.719 rad/s (0.43 turns/s); it should, by its angular
    momentum: +2.736
t= 17.3s  wheel axle up again (34 cm out): the stool turning at -0.021 rad/s (0.00 turns/s); it should, by its
    angular momentum: -0.018
t= 22.8s  wheel turned over (34 cm out): the stool turning at +2.456 rad/s (0.39 turns/s); it should, by its angular
    momentum: +2.472
t= 41.8s  wheel turned over (34 cm out): the stool turning at +2.192 rad/s (0.35 turns/s); it should, by its angular
    momentum: +2.208
```

Angular-momentum figures (0.093%, 0.050%) were computed from the L_up column of the BW_LOG csv; no NaN appeared in either 120 s run.

Known departures and honest caveats

- The "theory" column the console prints is computed by the script from the simulated state at the start of each move; it is an internal consistency check (angular momentum bookkeeping), not an independent prediction. The independent hand calculation (2.769, 2.202 rad/s) agrees with it in the frictionless case.
- With friction on (defaults) the stool lags the console "should" by 0.015-0.020 rad/s because friction during the 1.2-1.5 s move is not accounted for (README says so). Total angular momentum falls 60% over 120 s in the default run; that is the stool bearing and wheel bearing, by design (the spin drops from 4.0 to 1.6 turns/s).
- After "axle up again" with friction on, the stool shows -0.021 rad/s rather than 0: the friction-induced residual.
- Mass properties are idealised: the figure+stool is one rigid body with a fixed 1.5 kg m^2; the arms are drawn only (the 1 kg "hands" body stands in for them); the wheel's inertia (0.115 kg m^2) and 600 kg cm^2 transverse are chosen, not derived from the mesh. The wheel's spin is a gyrostat number, not a rotating Bullet body.
- The gyroscopic torque is explicit (Adams-Bashforth); the agreement shown is at the default 1200 steps/s; no step-size study is documented.
- Open issue (user report, not reproduced): a user saw the bicycle wheel go to NaN once in the GUI after visiting the solar system exhibit in the hall (Cabinet). It could not be reproduced headless; the 7200-frame default run on 15 Oct 2026 was NaN-free too. Cause still unknown (the Cabinet prefixes each exhibit's sliders, so the solar system's settings are not shared with this exhibit). Guard added 15 Oct 2026: after every frame the script checks the positions, velocities and angular velocities of the stool, wheel and hands bodies, and the wheel's spin, angle, arm angle and slide distance; if any is not finite it prints one console line ("t=...s ** NOT A NUMBER: the simulation broke down (going by itself, move N, last move: ...; wheel ... cm out, spinning ... rad/s, turned ... rad; steps, speed, step); stool at (...) moving (...) turning (...); wheel ...; hands ... -- starting again") and restarts the run cleanly. Tested by injecting a NaN into the wheel's angular velocity at t = 5 s in a scratch copy: the line was printed, the run restarted, and the restarted run's first report (t = 3.7 s, +2.719 / +2.736) was identical to a fresh start. Also: the bearing-loss term IW3·d/h is now skipped if the step h were ever 0 (it would be 0/0); this does not occur in normal use and is not claimed to be the cause.
- Fixed on 15 Oct 2026: pressing R mid-move used to print a stray "wheel turned over ..." line about 1.7 s into the new run (the cut-off move's pending report, with the restarted clock). restart() now clears the pending report; tested by forcing a restart at t = 2.5 s, mid-turn, in a scratch copy: no stray line.

- Checks after these changes: default 7200-frame run gives console output byte-identical to the previous version of the script (27 lines); hall identity cmp2.sh "the bicycle wheel" 2400 frames: IDENTICAL (8 run lines); pause check 2400/2430 frames: prefix OK (10 lines).

References

None cited in the README or code. The physics is the standard textbook "rotating stool and bicycle wheel" demonstration of angular momentum conservation.

5. Wing B: Swinging

5.1 Newton's cradle

Agrees with theory. First strike the same at every step rate from 1200 to 9600 steps/s (was: a third of the energy lost at 9600); momentum to 0.00%; ball 4 gets 13.6% (Herrmann & Seitz: ~12%); the row settles over minutes, as real cradles do.
Folder: demo/WyomingWill/NewtonsCradle/

What it shows

Five steel balls hang in a row on V-threads. Lift one, two or three and let go: the same number fly out at the far end. Because the balls touch, a strike is a compression pulse through the row rather than a chain of two-ball collisions, so the far ball does not get quite everything; the row breaks up and, over a few minutes, ends up swinging together. With a 0.1 mm gap (key `G`) each strike becomes a clean one-in, one-out sequence.

Model

Balls: rigid spheres, 25 mm diameter, 64 g (code: rho 0.00785 kg/cm^3, 64.2 g computed), centres 15 cm below the thread tops. Each ball is a Bullet rigid body on a `btHingeConstraint` to a static anchor (it swings in the row's plane and turns with its threads); Bullet collisions are off (`collides = false`), solver iterations 40, Bullet damping 0. Bullet steps at 2400 /s (slider 1200-9600). Strikes are done by the script before every physics step: any pair pressed together, or predicted to touch within the step, is integrated with each ball moving along its own swing (the contact force along the line of centres, projected on each ball's swing tangent; inertia along the swing m(1 + 2/5 r^2/L^2), since the ball turns with its threads), with Hertz forces F = K d$^{3/2}$, K = 4/3 E* sqrt(R/2), E* = E / (2(1 - nu^2)), E = 200 GPa, nu = 0.3, times a Hunt-Crossley factor (1 + 1.5 alpha v), alpha = 0.25 x 0.0012 s/cm (a quarter of Hunt-Crossley for steel). Integrator: velocity Verlet, sub-steps <= 0.25 us; each ball is handed back to Bullet on its hinge circle, turned with its threads, with velocity W along its swing tangent and angular rate W/L (so the hinge solver has nothing to correct), placed so Bullet's step ends where the row had it. A squeeze still held at the end of a step is carried into the next step through the positions and worked out there. Knocks with |dv| < 3 cm/s are not counted as strikes. Clack sound is pre-synthesised by `make-sounds.py` (same Hertz strike at 20 ns steps, rigid-sphere dipole radiation with the h2 spherical Hankel function, table image at 0.85, 48 kHz) — not run here because it rewrites `cradle-sounds/`.

Theory checks

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Two-ball contact time at 62.7 cm/s	Hertz: 2.87 (m^2/(R E*^{2} v))$^{1/5}$, m,R reduced: 84.2 us	84.2 us (script's own 2-ball integration)	exact to 0.1 us
Restitution of two-ball strike, damping 0.25	design target 0.982 (README)	0.982 (1.000 with NC_DAMP=0)	matches
Time to first strike	quarter period of physical pendulum, I = m(L^2 + 2/5 r^2), elliptic K, 30 deg: 0.1979 s (I reproduced 0.19789 s independently)	0.1975 s (2400 steps); 0.1977 s (3000, 4800); 0.1978 s (7200, 9600); 0.1967 s (1200)	-0.4 ms = one step (README says so)
Energy in first strike, no damping	conserved: 100%	100.0% (touching); 100.0% (gap)	0.0%
Momentum along row, first strike	conserved	0.00% off	exact
Touching row, no damping: speeds after	README: -4.5, -1.9, -0.9, 8.0, 62.0 cm/s	-4.5, -1.9, -0.9, 8.0, 62.0	identical
Touching row, damping 0.25: speeds after	README: -4.1, -1.6, -0.6, 8.5, 60.4 cm/s; energy 95.3%	-4.1, -1.6, -0.6, 8.5, 60.4; 95.3%	identical

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Gap 0.1 mm, no damping	README: 62.7 out, others < 0.05 cm/s	0.0, -0.0, -0.0, -0.0, 62.7; energy 100.0%	identical
Gap 0.1 mm, damping 0.25	four two-ball strikes at e = 0.982: each striker keeps (1-e)/2 x 62.7 = 0.56 cm/s, out 62.7 x 0.991^4 = 60.5	0.6, 0.6, 0.6, 0.5, 60.5; energy 93.0%	matches
Ball 4's share of incoming momentum	Herrmann & Seitz 1982 five-ball simulation: ~12% (README)	8.5/62.7 = 13.6% (README says 14%)	+1.6 points; same pattern (1-3 back, 4-5 forward)
Row merges to swinging together	Hutzler et al. 2004: 1-2 min on real cradles (README)	1 ball, 30 deg, auto off, 7 min run: together swing 5.2 deg; largest ball's swing about the row's mean falls below 1/2 of it at ~80 s, 1/5 at ~3 min, 1/10 at ~6.5 min (1200 and 9600 steps/s agree to a few s); energy 32% at 15 s, 19% at 1 min, then 16.6% (the together swing). Damping 0: never merges (relative swing ~24 deg after 3 min), energy kept to 1.5%	same scale as real cradles, somewhat slower; set by the contact damping alone (README says so)
Clack peak at listener, 60 cm/s	README (make-sounds.py console): 2.7 Pa, 103 dB, peak ~ $\text{speed}^{1.48}$	not run	README only
Step-size convergence of first strike	should be independent of step	1200, 2400, 3000, 4800, 7200, 9600 steps/s: -4.1 -1.6 -0.6 8.5 60.4-60.5 cm/s, energy 95.3-95.4%; NC_DAMP=0: -4.5 -1.9 -0.9 8.0 62.0, energy 100.0% (100.2% at 1200, over a 1.7 ms window with gravity acting)	converged (fixed 15 Oct 2026)

Run used

```
cd demo/WyomingWill/NewtonsCradle && HOLD_S=0 timeout 1500 ../../../release/bpp -f newtons-cradle.lua -n 3600 | grep
    -v ALSA
  (plus -n 900 / -n 40 runs with NC_DAMP=0; NC_GAP=0.01; NC_GAP=0.01 NC_DAMP=0;
    NC_STEPS=1200,2400,3000,4800,7200,9600 each with
   and without NC_DAMP=0; settling: a scratch copy of the exhibit with the auto-lift off and each ball's angle and
    the total
   energy logged every frame, 1 ball, 25200 frames at 2400/s, 14400 at 9600/s, 7200 at 1200/s, 10800 with NC_DAMP=0)

    two balls meeting at 62.7 cm/s: they touch for 84.2 microseconds (Hertz: 84.2); they part at 0.982 of it
  strike  1 at  0.197 s: speeds before   62.7    0.0    0.0    0.0    0.0
                              after    -4.1   -1.6   -0.6    8.5   60.4 cm/s; energy 95.3%; momentum along the row
    0.00% off ...
                    0.1975 s after letting go; a quarter swing from 30 degrees takes 0.1979 s
  [NC_DAMP=0]  after    -4.5   -1.9   -0.9    8.0   62.0 cm/s; energy 100.0%; momentum along the row 0.00% off
  [NC_GAP=0.01 NC_DAMP=0]  after     0.0   -0.0   -0.0   -0.0   62.7 cm/s; energy 100.0%
  [NC_GAP=0.01]  after     0.6    0.6    0.6    0.5   60.5 cm/s; energy 93.0%
  [1 ball] strike 90 at 13.425 s: ... after    22.8    4.6    5.8    6.9   12.3 cm/s; energy 119.1%; momentum 7.52%
    off (over 14.2 ms)
  [3 balls] strike 240 at 14.775 s: ... after   -20.4  -41.1  -16.5  -43.7  -45.4 cm/s (still not swinging together
    at 15 s)
  [NC_STEPS=9600]  after    -4.1   -1.6   -0.6    8.5   60.4 cm/s; energy 95.3%; momentum along the row 0.00% off
  [NC_STEPS=1200]  after    -4.1   -1.6   -0.6    8.5   60.5 cm/s; energy 95.4%; momentum along the row 0.01% off
```

Known departures and honest caveats

- Fixed 15 Oct 2026: **step-size failure at 9600 steps/s** (first strike kept 66.8% of the energy, ball 4 left at 22.6 cm/s; 6000 steps/s with no damping also failed, 84.4%). Cause: the "touching" test only asked whether a pair would be pressed at the end of the step if it moved freely (gap - closing speed x h < 0). A contact still squeezed but already opening (at 9600/s the pulse straddles steps: ball 4-5 still squeezed 14 um at a step boundary, separating at 23 cm/s) was predicted "open", so the next step skipped the strike and Bullet simply let the balls fly apart through each other, throwing the stored squeeze away. Now a pair is worked out if it is pressed now or will be within the step. A second, smaller error came from the hand-back: balls were moved along x without turning them on their

hinges, and the hinge solver's correction shifted squeezes of ~15 um by 10-50 nm between steps (0.1-0.3% of a strike's energy); and strikes at a swing angle were worked out along x with mass m, then projected by the hinge onto the swing, which made or lost energy by ~angle^2/2 per strike (with damping 0 the energy wandered by +-10% over a minute after the first fix alone); now each ball is handed back on its hinge circle, turned, with velocity along its swing, and the row is worked out along each ball's swing with its inertia m(1 + 2/5 r^2/L^2). The default first strike moved from 60.477 to 60.450 cm/s for ball 5 (shown 60.5 -> 60.4; README updated).

- Fixed 15 Oct 2026: **settling time**. The same lost squeezes, through many later strikes, drained the energy: with NC_DAMP=0 the old code lost 59% of the energy by 5 s and 81% by 20 s, and the row "merged" in ~15 s whether damped or not. That fast merging was the bug. Now with damping 0 energy is kept to 1.5% over 3 min and the row never merges; with the default damping it merges over minutes (table). The README's "Why the balls part" now gives the measured times and says the rate is set by the contact damping alone (a quarter of Hunt-Crossley, chosen for the 0.982 two-ball restitution, not tuned to Hutzler); a real cradle also has air, thread and alignment losses that the model lacks, so the near-agreement with 1-2 min is not a test of the model.
- Late "strikes" in long runs report energy 58-126% and momentum up to ~105% off; these are windows of 10-40 ms of repeated contact in which gravity and the threads act, so the per-strike bookkeeping is not a conservation test there.
- By itself the exhibit relifts every 15 s, so the settling is only seen with `A` off.
- Ball 4 gets 13.6% of the incoming momentum versus Herrmann & Seitz's ~12%: same pattern, not the same number.
- Threads are a hinge, so out-of-plane swinging and thread stretch are absent; the threads are drawn only.
- Sound: README numbers only (not regenerated; files unchanged, md5 checked). Fixed 15 Oct 2026: `make-sounds.py`'s docstring said the loudest file was "four balls at 120 cm/s"; its speeds stop at 100 and the files show clack-n1-v100.wav peaks at 0.95 (n2-v100 0.62, n1-v60 0.56), so it now says "one ball at 100 cm/s".
- Checks after the fix: hall vs standalone (cmp2.sh newton, 1500 frames): IDENTICAL (59 run lines); pause: HOLD_S=0 -n 840 is a prefix of the default -n 870 (30 of 30 lines), and -n 1200 of -n 1260 across the 15 s relift (52 of 52).

References

- H. Hertz (contact law, contact-time formula 2.87(...)$^{1/5}$) — cited by name.
- K. H. Hunt and F. R. E. Crossley (contact damping) — cited by name.
- F. Herrmann and M. Seitz (1982), five-ball simulation of the cradle (README; journal not given).
- E. J. Hinch and S. Saint-Jean (1999), break-up of a line of balls (README).
- S. Hutzler, G. Delaney, D. Weaire and F. MacLeod, "Rocking Newton's cradle", Am. J. Phys. 72 (2004).

5.2 Huygens' clocks

Agrees with theory. All three starts lock in anti-phase at 180.0°, at 0.9985-0.9988 Hz against 0.9984 Hz from theory; at 9600 steps/s the same to 1e-4 Hz (one start, 45 s).
Folder: demo/WyomingWill/HuygensClocks/

What it shows

Two pendulum clocks hang from one beam that can slide a little on a spring. Started out of step (60, then 300, then 120 degrees apart, four minutes each), within about half a minute they swing in exact opposition and the beam stops: Huygens' 1665 "odd kind of sympathy". In opposition their pushes on the beam cancel, so the beam's friction takes no energy; any in-phase part is damped away.

Model

Bullet. The beam (1.5 kg including the clock cases) is a rigid body restricted to slide along x (linear factor (1,0,0), no rotation, no gravity). Each pendulum is one rigid body (brass bob 3.5 cm diameter, 191 g, plus a light aluminium rod; bob centre 25 cm from the pivot) on a `btHingeConstraint` to the beam. Done by the script before every physics step, as velocity changes: the beam's spring (k from 1.5 Hz with the beam alone) and friction (5 kg/s); each pendulum's air drag (a torque proportional to angular rate, Q = 150 alone); and the escapement, an angular-velocity kick at each zero crossing, sized to replace exactly the air loss over half a swing at 8 degrees. Bullet's own damping 0, solver iterations 40, 2400 physics steps/s (Bullet's semi-implicit Euler with the hinge solver). Clock hands advance by each clock's own swing count.

Theory checks

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Anti-phase normal mode (no escapement, no friction)	$w^2 = m\,g\,d / I$ (beam still): 0.9996 Hz	not run as a free mode; used as reference	(my own 3x3 eigen-solve: 0.99964 Hz)
In-phase normal mode	root of det[k - (M+2m)w^2, -2md w^2; -md w^2, mgd - I w^2] = 0: 0.9179 Hz	not compared with a simulated mode	(my eigen-solve: 0.91787 Hz)
Anti-phase at 8 deg amplitude	0.9996 Hz x pi / (2 K(sin(4 deg))): 0.9984 Hz	settled clock frequency 0.9987 Hz (start 60 deg), 0.9988 (300 deg), 0.9985 (120 deg)	+0.03%, +0.04%, +0.01% (README: escapement kicks account for the rest)
Final phase difference	180 deg (anti-phase locking, Bennett et al. 2002)	180.0 deg at t = 60 s for all three starts	exact to 0.1 deg
Beam amplitude decays	beam stops in anti-phase	5.335 -> 0.544 -> 0.062 -> 0.008 mm at 15/30/45/60 s (start 60 deg)	~x10 per 15 s; to 0.008 mm
Time to lock	README: "within a minute"	within 0.5 deg of 180 by 30 s for all three starts	consistent
Step-size check (my run, HC_STEPS=9600)	should not depend on step	161.9 / 179.5 / 180.1 deg; 1.0063 / 1.0005 / 0.9990 Hz; beam 5.338 / 0.543 / 0.062 mm	agrees with 2400 /s to 1e-4 Hz and 0.003 mm
README console listing	README t = 15..60 s lines	identical to this run, digit for digit	matches

No energy or momentum conservation check is performed by the exhibit (the system is driven and damped by design).

Run used

```
cd demo/WyomingWill/HuygensClocks && HOLD_S=0 timeout 1500 ../../../release/bpp -f huygens-clocks.lua -n 3600 | grep
    -v ALSA
  (also HC_PHASE=60, 300, 120 with -n 3700 to reach the t=60 s line; HC_STEPS=9600 -n 3600)

  stopped, they'd swing in opposition at 0.9996 Hz (the beam still), or together at 0.9179 Hz (the beam swinging
    against them)
  in opposition, swinging 8 degrees (a pendulum's slower, the further it swings: the elliptic integral): 0.9984 Hz
  t=   15 s  the clocks 161.9 degrees apart (in opposition: 180); swinging at 1.0063 Hz; the beam moving up to 5.335
    mm
  t=   30 s  the clocks 179.5 degrees apart (in opposition: 180); swinging at 1.0004 Hz; the beam moving up to 0.544
    mm
  t=   45 s  the clocks 180.1 degrees apart (in opposition: 180); swinging at 0.9989 Hz; the beam moving up to 0.062
    mm
  t=   60 s  the clocks 180.0 degrees apart (in opposition: 180); swinging at 0.9987 Hz; the beam moving up to 0.008
    mm
  [HC_PHASE=300] t=   60 s  the clocks 180.0 degrees apart ...; swinging at 0.9988 Hz; the beam moving up to 0.008 mm
  [HC_PHASE=120] t=   60 s  the clocks 180.0 degrees apart ...; swinging at 0.9985 Hz; the beam moving up to 0.004 mm
```

Known departures and honest caveats

- Locks in about 30 s, against Huygens' "within half an hour". The beam parameters (1.5 kg, 1.5 Hz spring, 5 kg/s friction) are chosen for a fast, visible demonstration, not to match Huygens' chairs and beam.
- The escapement is an idealised impulse at the bottom of each swing, sized exactly to the linear air loss at 8 degrees; real escapements (and Bennett et al.'s model) differ. The 0.03% excess of the settled frequency over the free anti-phase mode is put down to the kicks (README), not separately verified.
- Pendulums are confined to the plane (hinges); the beam only slides (no rocking).
- The in-phase mode frequency (0.9179 Hz) is printed but never compared against a simulated in-phase swing.
- Fixed on 15 Oct 2026: the lua header used to call the "together" mode "faster"; it now says "slower", matching its own console value (0.9179 Hz against 0.9996 Hz) and the README. Comment only; the console output is unchanged.
- A third normal mode (mostly the beam on its spring, about 1.63 Hz by my eigen-solve) exists and is not mentioned.
- The 15 s report window's "swinging at" frequency is the last single period of clock 1, so early values (1.0063 Hz) reflect the transient.

References

- C. Huygens, 1665 observation ("an odd kind of sympathy") — README.
- M. Bennett, M. F. Schatz, H. Rockwood and K. Wiesenfeld, Proc. R. Soc. A (2002) — README and lua header (title not given there).

5.3 Kapitza's pendulum

Agrees with theory. Stability threshold 0.783 cm bracketed (0.76 falls, 0.80 stands); rocking period 1.133 s against 1.135 s for its 7.6° swing (-0.2%), now at a converged 19,200 steps/s (was 4800, 1.7% short).
Folder: demo/WyomingWill/Kapitza/

What it shows

An upside-down pendulum stands on a pivot shaken up and down 40 times a second and does not fall; nudged, it rocks slowly about the upright as if gravity pointed up. Left: shaken 1.0 cm, it stays up. Right: shaken 0.6 cm, too little, it falls and ends up hanging. Both are stood up again, tilted 8 degrees, every 15 s.

Model

Bullet. Each pendulum is one rigid body (brass bob 3 cm diameter plus aluminium rod radius 0.25 cm; bob centre 20 cm from the pivot; 130.8 g; centre of mass 19.19 cm from the pivot; inertia from the composite) on a `btHingeConstraint` to a pivot block. The block is a 10^4 kg body moved kinematically by the script before every step: position y = y0 + a cos(w t) and velocity -a w sin(w t), w = 2 pi x 40 Hz. No friction, Bullet damping 0, solver iterations 40, 19200 physics steps/s (480 per shake; raised from 4800 on 15 Oct 2026, see below; steps slider 2400-19200). The slow motion is measured on the angle averaged over each 1/40 s shake; rocking period from upward zero crossings of that average.

Theory checks

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Equivalent length	l = I / (m d): 19.761 cm	(input; independent recomputation 19.7614 cm)	exact
Stability threshold	Kapitza: a w > sqrt(2 g l) = 196.9 cm/s, a > 0.783 cm	1.00 cm stays up; 0.60 cm falls at 0.69 s	qualitatively correct
Threshold bracket (KP_A1=0.80 KP_A2=0.76)	0.80 cm (a w = 201.1) should stand; 0.76 cm (191.0) should fall	0.80 cm "stays up (as it should)"; 0.76 cm falls at 1.39 s	correct on both sides, within +/-3% of threshold
Rocking period, a = 1.0 cm, small-rock theory	2 pi / W, $W^2 = a^2 w^2 / (2 l^2) - g / l$: 1.124 s	1.133 s over 6 rocks, up to 7.6 deg	+0.8% (expected: the rock is 7.6 deg, not small)
Rocking period for its size (new console term)	averaged eq. to next order $\phi'' = -W^2 \phi + c \phi^3$, $c = 2K/3 - g/(6l)$, $K = a^2 w^2/(2 l^2)$; Lindstedt $T = (2\pi/W)(1 + 3 c A^2/(8 W^2))$, A = 7.6 deg: 1.135 s (+0.97%)	1.133 s	-0.2%
Exact (non-averaged) pendulum equation, integrated independently (Python RK4, 400 steps per shake, same shake-averaged measurement)	$\theta'' = (g - a w^2 \cos wt) \sin\theta / l$, from 8 deg: 1.1341 s; from 2 deg: 1.1237 s	1.133 s (8 deg); KP_TILT=2: 1.122 s	-0.1% / -0.15%
Step-size convergence of the rocking period (KP_STEPS)	should converge to the exact equation's 1.134 s	2400: 1.073; 3600: 1.103; 4800: 1.115; 7200: 1.125; 9600: 1.129; 14400: 1.132; 19200: 1.133; 38400: 1.134 s	converged to 0.1% at the new default
Repeatability over 4 restarts (3600 frames)	—	1.133 s and 7.6 deg each time	stable (fixed: see caveats)
Hall identity / pause	identical in the Cabinet; HOLD_S=0 output a prefix of the default-pause output	cmp2.sh "Kapitza" 1500 frames: IDENTICAL (12 run lines); pause check 1500/1530 frames: prefix OK (18 lines)	pass

No energy check (the shaker does work by design).

Run used

```
cd demo/WyomingWill/Kapitza && HOLD_S=0 timeout 1500 ../../../release/bpp -f kapitza.lua -n 3600 | grep -v ALSA
  (also KP_A1=0.80 KP_A2=0.76 -n 1200; KP_TILT=2 -n 600; KP_STEPS=2400..38400 -n 600)

  the pendulum: 130.8 g, its centre 19.19 cm from the pivot, equivalent length l = I/(m d) = 19.761 cm;
  it stays up if a w > sqrt(2 g l) = 196.9 cm/s, i.e. a > 0.783 cm
  Left   shaken 1.00 cm: a w = 251.3 cm/s > 196.9 -- it should stay up, rocking with a period of 2 pi / W = 1.124 s
  Right  shaken 0.60 cm: a w = 150.8 cm/s < 196.9 -- it should fall
  Right  falls over at 0.69 s
  Left   stays up, rocking up to 7.6 degrees either side: a period of 1.133 s over 6 rocks (theory, 2 pi / W: 1.124
    s, +0.8%; for its size: 1.135 s, -0.2%)
  [KP_STEPS=4800, the old default] Left   stays up, rocking up to 7.6 degrees either side: a period of 1.115 s over 6
    rocks (theory, 2 pi / W: 1.124 s, -0.8%; for its size: 1.135 s, -1.7%)
  [KP_TILT=2] Left   stays up, rocking up to 1.9 degrees either side: a period of 1.122 s over 6 rocks (theory, 2 pi
    / W: 1.124 s, -0.2%; for its size: 1.125 s, -0.2%)
  [KP_A1=0.80] Left   stays up (as it should)
  [KP_A2=0.76] Right  falls over at 1.39 s
```

Known departures and honest caveats

- Fixed on 15 Oct 2026, step size: at the old default of 4800 steps/s the rocking period was 1.7% short of the converged value (1.115 s against 1.134 s), and that error happened to cancel the finite-amplitude lengthening, so "-0.8% from 2 pi / W" agreed for the wrong reason. The default is now 19200 steps/s (slider max raised to 19200), where the period is within 0.1% of both the 38400 run and an independent integration of the exact pendulum equation. Cost: about 6 ms of CPU per frame headless.
- Fixed on 15 Oct 2026, comparison: the console now also gives the period for the rock's measured size (Lindstedt correction of the averaged equation, about +1%), so the comparison is no longer between a finite rock and a small-rock formula only. The remaining -0.2% is half the averaging's next terms (the exact equation gives 1.1341 s vs 1.135) and half step error.
- Fixed on 15 Oct 2026, restart bug: after each 15 s restart the first shake-average bin included the previous run's stale bin plus the new 8-deg start angle, so later runs reported "rocking up to 8.0 degrees" (the start tilt) instead of 7.6. Now every run reports 7.6.
- Kapitza's formulas are first-order averages (fast shaking, small rocking); the README says so.
- Near threshold (0.80 cm) the rocking period (4.3 s theory) is too long for the 8 s report window, so only "stays up" is checked, not a period.
- No upper stability limit is checked (Kapitza's averaging cannot see it; the DoublePendulum exhibit's Floquet analysis does that for the two-rod chain).
- The pivot is driven exactly (kinematic), i.e. an ideal shaker; the hinge is frictionless and the motion is planar.
- Fixed on 15 Oct 2026, documentation: the lua header listed a non-existent `KP_LOG=file.csv`; it now lists `KP_A1`, `KP_A2`, as the README does. The README's "131 g" now reads 130.8 g, as the console.

References

- A. Stephenson (1908), first observation — README / lua header.
- P. L. Kapitza (1951), averaged-potential explanation — README / lua header (titles and journals not given).

5.4 The double pendulum

Agrees with theory, with a numerical or open caveat. Energy to 2e-11; half-step twin 64 fm off at 1 s; Lyapunov 3.42/s; normal modes to 0.000% and 0.135%; Floquet window 0.728-1.356 cm. The throw-over time of the too-hard chain (not the fact) isn't converged, and the README says so.
Folder: demo/WyomingWill/DoublePendulum/

What it shows

Three tables. The butterfly: seven double pendulums let go from 120 degrees, each a billionth of a radian further round (bottom bobs 2 Å apart), swing as one for about five seconds and then fly apart: chaos. The two pure swings: the slow normal mode (rods together, lower rod sqrt 2 times as far), the fast one (rods opposed, 2.4 times as often), and a mix of both that never repeats, each with a chart recorder. Upside down: double pendulums on pivots shaken at 40 Hz; 0.6 cm falls, 1.0 cm stands (Kapitza/Acheson), 1.42 cm is thrown over by a parametric sway.

Model

Bullet only draws. The script integrates Lagrange's exact equations for equal point bobs on massless rods. Butterfly (l = 20 cm) and pure swings (l = 12 cm): Bulirsch-Stoer (Gragg modified midpoint with 2, 4, 6, 8, 10 substeps, Neville extrapolation in h^2, tenth order), 240 steps/s. Shaken chains (l = 10 cm): classical RK4, 4800 steps/s (120 per shake), pivot acceleration added to g. No friction anywhere. Extra hidden copies: a "twin" of butterfly pendulum 1 at half step (for a step-halving check) and a "shadow" offset by 1e-8 rad, renormalised every 0.25 s (Benettin's method for the largest Lyapunov exponent; norm in angles and angular rates / sqrt(g/l)). Upright stability: linearised equations integrated over one shake (RK4, 400 steps) to the 4x4 monodromy matrix; Floquet multipliers from its traces; window edges by 40-step bisection. Averaged theory: extra stiffness $(a^2\ w^2 / 2)\ D\ M^{-1}\ D$. Normal-mode amplitude correction: Lindstedt / normal-form average of the quartic Hamiltonian terms.

Theory checks

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Butterfly energy (BS, 30 s)	conserved	max drift 2.1e-11 of m g l over all seven	conserved to 2e-11
Step-halving twin, bottom bob at t = 1 s	should be << 2 Å start spacing	64.4 fm (2 Å / 64.4 fm ~ 3100x)	integration error negligible
Twin reaches 1 cm	later than the seven	8.3 s vs 6.3 s for the neighbour	arithmetic error lags by 2 s
Lyapunov exponent (Benettin, 30 s)	no closed form	3.42 /s, doubling every 0.203 s	—
Separation 2 Å -> 1 cm	ln(1 cm / 2 Å) / lambda = 5.2 s	neighbour reached 1 cm at 6.3 s	observed 1.1 s (21%) later; order-of-magnitude check only
Slow mode, small swings	T = 2 pi / sqrt((g/l)(2 - sqrt 2)) = 0.90796 s (mine 0.907959)	—	—
Slow mode at 7 deg	Lindstedt: $x(1 + 0.0849\ A^2)$: 0.90911 s	0.90911 s over 21 swings	-0.000%
Slow mode shape	lower/upper = +sqrt 2 = 1.4142	1.4091	-0.36%
Fast mode, small swings	2 pi / sqrt((g/l)(2 + sqrt 2)) = 0.37609 s (mine 0.376089)	—	small-swing theory alone 2.7% off
Fast mode at 7 deg	Lindstedt: $x(1 + 1.8526\ A^2)$: 0.38678 s	0.38626 s over 51 swings	-0.135%
Fast mode shape	-sqrt 2 = -1.4142	-1.4112	-0.21%
Mixed swing (first 10 s)	half of each mode, each at its amplitude-corrected period	max deviation 0.1247 deg (1.78% of 7 deg); 5.11 deg with small-swing periods	good with correction

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Lower stability edge, upright chain	averaged: a w > sqrt((2 + sqrt 2) g l): a > 0.728 cm (mine 0.7282); Floquet: 0.728 cm	0.60 cm falls at 0.79 s; 1.00 cm stands	correct
Upper stability edge	Floquet: a < 1.356 cm (averaging cannot see it)	1.42 cm thrown over at 3.96 s (4800/s); 4.46 s (my 9600/s run)	correct; timing step-dependent
Rocking period, a = 1.0 cm	Floquet exact 0.879 s; averaged 0.881 s	0.881 s over 12 rocks, up to 3.8 deg	+0.3% / +0.1% (same at 9600 steps/s)
README console listing	README lines	identical to this run (incl. 1.42 cm at 3.96 s)	matches

Run used

```
cd demo/WyomingWill/DoublePendulum && HOLD_S=0 timeout 1500 ../../../release/bpp -f double-pendulum.lua -n 2400 |
    grep -v ALSA
  (also DP_STEPS=9600 -n 1200 as a step-size check of the shaken chains)

    t=1 s  the bottom bobs: the next one 1.0 nm apart, the farthest 6.3 nm  (the first, stepped twice as finely: 64.4
    fm)
    exactly (Floquet), it stands if 0.728 cm < a < 1.356 cm
    shaken 1.42 cm: the shaking pumps up a sway and throws it over at 3.96 s
    shaken 1.00 cm: it stands, rocking up to 3.8 degrees: a period of 0.881 s over 12 rocks (exact 0.879 s, +0.3%;
    averaged 0.881 s, +0.1%)
  Slow swing: a period of 0.90911 s over 21 swings (theory 0.90911 s for its size, -0.000%; small swings 0.90796 s);
  Fast swing: a period of 0.38626 s over 51 swings (theory 0.38678 s for its size, -0.135%; small swings 0.37609 s);
    to within 0.1247 degrees (1.78% of its 7; with the small-swing periods, 5.11 degrees)
  The butterfly, over 30 s: energy steady in all seven to 2.1e-11 of m g l;
    nearby swings part at 3.42 per second (Lyapunov, Benettin's method): doubling every 0.203 s,
    so 2.0 angstroms grows to 1 cm in about ln(1 cm / 2.0 angstroms) / 3.42 = 5.2 s; the next one got there at 6.3 s
    and the first, stepped twice as finely, only at 8.3 s
  [DP_STEPS=9600] shaken 1.42 cm: the shaking pumps up a sway and throws it over at 4.46 s
```

Known departures and honest caveats

- Idealised: point bobs, massless rods, no friction or air; Bullet plays no part in the physics.
- The Lyapunov estimate is a finite-time value from one 30 s trajectory; the predicted 5.2 s vs observed 6.3 s differ by 21%. Since 15 Oct 2026 the README says so itself: an order-of-magnitude check, not a precise test (the early growth is slower, before the gap aligns with the fastest-growing direction, and the bob gap is not the Benettin norm).
- Fast mode remains 0.135% off after the A^2 correction; the README attributes this to the next order.
- Mode-shape ratios (1.4091, -1.4112) are amplitude ratios of the measured maxima of each rod, slightly off sqrt 2 at 7 deg.
- The 1.42 cm chain's throw-over time is not converged in step size (3.96 s at 4800/s, 4.46 s at 9600/s, re-run 15 Oct 2026): it grows from a tiny seed that the step errors are part of, so only "is thrown over" is robust. Since 15 Oct 2026 the README states this. Its other numbers (0.881 s rocking, 0.79 s fall) were unchanged at 9600/s.
- The rocking period is measured on shake-averaged angles from a 4-degree tilt along the slow mode.
- "the next one 52.6 cm apart, the farthest 25.8 cm" at t = 7 s: "farthest" means pendulum 7 (the largest initial offset), not the maximum separation; after saturation it can be the smaller.
- Fixed on 15 Oct 2026: the lua header used to say energy holds "to about 1e-9" and the twin is "a few femtometres off after a second, a hundred-thousandth of the atom's width"; it now says 2e-11 of m g l and 64 fm, some three thousand times less than 2 Å, as the run gives (2.1e-11, 64.4 fm). The README's "start a hundred thousand times smaller" sentence was corrected the same way. Comments/README only; console output unchanged.

References

- D. J. Acheson (1993), stabilisation of an inverted chain of pendulums by vibration — README / lua header (journal not given).

- P. L. Kapitza (averaging), via the neighbouring Kapitza exhibit — README.
- Benettin's method (Lyapunov exponent), Lindstedt's method / normal form, Floquet theory, Bulirsch-Stoer with Gragg's modified midpoint — cited by method name only.

5.5 Pendulums with a twist

Agrees with theory. No flip below 70.53°; at μ = 3 the hidden integral stays 0.7500000000 (wanders at 2.8 and 3.2); spring's growth 0.526/s against 0.527 (Floquet).
Folder: demo/WyomingWill/PendulumsTwist/

What it shows

Three tables of pendulums that defy intuition. A "flip row" of nine equal-bob double pendulums, released from rest with both rods at the same angle (60 to 120 degrees), shows that energy forbids the lower rod from going over the top below 70.53 degrees, while being *allowed* above that line does not mean it *happens* (72 and 76 degrees never flip). Three swinging Atwood machines (mass ratios mu = 2.8, 3.0, 3.2) look alike, but only at mu = 3 is there a hidden second conserved quantity, shown on a dial whose needle stays still. Two swinging springs show 2:1 autoparametric resonance: tuned bounce:swing = 2:1, the bounce drains into a swing that grows to about 11 degrees and back; tuned 2.4:1 it just bounces.

Model

Every pendulum is integrated by the script from its exact Lagrangian equations of motion; Bullet only draws (all objects are non-colliding, mass-0 display bodies; cords and springs are drawn as beads). Integrator: Bulirsch-Stoer (Gragg modified midpoint with 2, 4, 6, 8, 10 substeps, Richardson-extrapolated to zero step: tenth order), 240 macro-steps per simulated second (4 per 1/60 s frame); the Atwood machines take two half-steps, i.e. 480 per second. No damping or friction anywhere. Units cm, kg, s; g = 981 cm/s^2.

- Flip row: rods l = 7 cm, equal point bobs, angles 60, 66, 70, 72, 76, 80, 90, 100, 120 degrees.
- Atwood: (1 + mu) r'' = r theta'^{2} - g (mu - cos theta), r theta'' = -2 r' theta' - g sin theta; start r = 18 cm, theta = 60 degrees, from rest; zero-size pulleys 8 cm apart.
- Springs: stretched length l = 20 cm at rest, released Z = 2 cm lower with a 0.05-degree swing; k chosen so sqrt(k/m) = 2.0 or 2.4 times sqrt(g/l). The exact (Floquet) growth rate is computed in the script by RK4 (2000 steps) over one bounce period of the linearised swing equation with prescribed r(t).
- Everything restarts every 60 s (PT_CYCLE); the report prints at the end of each cycle.

Theory checks

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Lower-rod flip threshold	E = -3mgl cos a must reach -mgl: cos a <= 1/3, a >= 70.53 deg	Never flipped: 60, 66, 70 (below line); also 72, 76 (above line, allowed but not seen in 60 s). Flipped: 80 (10.86 s), 90 (2.81 s), 100 (1.26 s), 120 (1.82 s)	Consistent: no flip below 70.53 deg
Upper-rod flip threshold	needs +mgl: cos a <= -1/3, a >= 109.47 deg	Only 120 deg flips its upper rod (4.87 s); 100 deg does not	Consistent
Flip-row energy conservation	E constant	max abs(E - E0) = 1.1e-08 of m g l	Conserved (README: 1.1e-08, identical)
Atwood mu = 3 second integral	I = p_xi^2/8 - E xi^2 + 2 xi^4, p_xi = 4 xi'(xi^2 + eta^2), constant only at mu = 3 (Casasayas-Nunes-Tufillaro 1990)	I between 0.7500000000 and 0.7500000000	Constant to 10 printed digits
Same quantity, mu = 2.8	not conserved	I between 0.6151989479 and 1.0500000000	Wanders, as expected

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Same quantity, mu = 3.2	not conserved	I between 0.3572701538 and 0.8542564850	Wanders, as expected
Atwood energy (relative)	E constant	3.1e-13 (2.8), 2.8e-13 (3.0), 2.1e-13 (3.2)	Conserved to round-off
Atwood closest approach to pulley	(no theory; README value)	7.0 / 4.5 / 3.7 cm; no pulley hit	Matches README exactly
Spring 2:1 growth rate	s = (3/4)(Z/l) sqrt(g/l) = 0.525 /s (first order); Floquet multiplier of one bounce: 0.527 /s	fitted from swing peaks 0.2 to 3 deg: 0.526 /s	-0.2% vs Floquet, +0.2% vs first order
Spring 2:1 peak swing	(README: "grows to 11 degrees")	at most 11.4 degrees	Matches README
Spring 2.4:1	Floquet: no growth (multipliers on unit circle)	never grew; at most 0.0 degrees	Exact
Spring energy (relative)	E constant	4.9e-14 (2.0:1), 3.6e-14 (2.4:1)	Conserved to round-off

The threshold angles 70.53 and 109.47, and the theoretical rates 0.525 and 0.527, are computed by the script and printed at each release.

Run used

```
cd demo/WyomingWill/PendulumsTwist && HOLD_S=0 timeout 1500 ../../../release/bpp -f pendulums-twist.lua -n 3620
```

```
    Tuned 2.0:1, the swing grows as exp(s t): s = (3/4)(Z/l) w = 0.525/s to first order, 0.527/s exactly (Floquet);
    flip row, let go at 80 degrees: the lower rod goes over the top at 10.86 s
    the flip row: the lower rod over the top -- 80: 10.9 s, 90: 2.8 s, 100: 1.3 s, 120: 1.8 s (upper 4.9 s); never:
    60, 66, 70, 72, 76 degrees (below 70.53 it can't; above, it may)
      energy steady to 1.1e-08 of m g l
    Atwood, 2.8 times: I between 0.6151989479 and 1.0500000000 (no such constant here); energy steady to 3.1e-13;
    closest to its pulley 7.0 cm
    Atwood, 3.0 times: I between 0.7500000000 and 0.7500000000 (constant, as at 3 it must be); energy steady to
    2.8e-13; closest to its pulley 4.5 cm
    Atwood, 3.2 times: I between 0.3572701538 and 0.8542564850 (no such constant here); energy steady to 2.1e-13;
    closest to its pulley 3.7 cm
    spring, 2.0:1: the swing grew at 0.526/s, from 0.2 to 3 degrees (theory 0.527 exactly, 0.525 to first order); at
    most 11.4 degrees; energy steady to 4.9e-14
    spring, 2.4:1: the swing never grew; at most 0.0 degrees; energy steady to 3.6e-14
```

Every number in the README's console excerpt was reproduced digit for digit by this run.

Known departures and honest caveats

- The README says plainly that the mu = 2.8 and 3.2 Atwood machines are **not visibly chaotic** when released from rest at 60 degrees. The author reports (README; not re-run here) that two copies started 1e-6 rad apart separate at no measurable exponential rate for release angles from 60 to 170 degrees and ratios from 1.3 to 3.2. So the exhibit does not claim to show chaos. What sets mu = 3 apart is only the hidden constant, shown on the dial (needle turns 120 deg per 0.5 change in I).
- The README also states (not re-run here) that from 90 degrees at mu = 2.5 the light weight is whipped into its pulley within 16 s. The script stops integrating a machine if r < 3% of 18 cm. The pulleys have zero size.
- Flip times near the threshold (72, 76 deg "never"; 80 deg at 10.9 s) cover a 60 s window only. "Never" means "not within 60 s", not proven never. The energy line is a necessary condition, not a sufficient one.
- The flip-row energy drift (1.1e-08 of mgl) is about five orders larger than the Atwood and spring drifts; it is still small. Fixed on 15 Oct 2026: the Lua header comment used to say "steady to 1e-10"; it now says "steady to 1e-8 of m g l", matching the console and README (1.1e-08). Comment only; console output unchanged.
- The growth rate is fitted to swing peaks between 0.2 and 3 degrees only, where the linear (exponential) phase holds. The later energy exchange (11.4 deg maximum) is nonlinear and is not compared with theory.

- Idealisations: point bobs, massless rigid rods and inextensible massless cords, frictionless pulleys of zero radius, massless linear springs.

References

- S. Casasayas, A. Nunes, N. Tufillaro (1990): the integrable swinging Atwood machine at mu = 3 (cited in code and README; no journal given).
- Pujol, Tufillaro and group, arXiv:2507.00058: the form of the integral I used here (README).
- Mathieu's equation and Floquet theory, for the parametric growth rate (README; no specific source).

5.6 The chain fountain

Agrees with theory, with a numerical or open caveat. Independent of the display speed; empties at every bend. With the push the falling chain carries, Biggins and Warner's relations hold. Run with every link simulated, the chain still passes none of the pile's push on (so that loss is the model's, not the shortcut's), but its fountain is lower: the exhibit's shortcut makes it 1.2x (30°) to 2x (60-90°) too tall. A faithful mode costs 5-8x per frame.
Folder: demo/WyomingWill/ChainFountain/

What it shows

A long bead chain is pulled over the rim of a jar 1.5 m above the floor. It then siphons out on its own, and rises in an arch above the rim instead of sliding over it. This is the "Mould effect", filmed by Steve Mould in 2013 and explained by Biggins and Warner in 2014. A link picked up by one end rotates and pushes its other end into the pile, and the pile's upward reaction throws the chain up. A rope, with no links, would just slide over.

Model

- Chain: Bullet rigid capsule links, 1.5 cm joint-to-joint, radius 0.3 cm, mass 0.5 g, no damping, no restitution. Neighbours are joined by btConeTwistConstraint with a swing limit equal to the bending angle (default 30 deg; CF_BEND; slider 20-90 deg, was 10-90) and free twist. Friction 0.3, chain on glass and chain on chain.
- Pile and jar: a rigid disc stands in for the coil, lowered one layer (0.65 cm) when the first link of a new layer is woken. The jar is 24 glass slabs (7 cm inside radius, 8 cm wall).
- Initial coil (changed 15 Oct 2026): flat spirals, outside in then inside out, all turning the same way. Every link is 1.5 cm and no joint is laid bent past its limit. The inner radius is max(3 cm, the tightest the chain can turn while tipping down) = 3.1 cm at 30 deg, 4.8 cm at 20. At a layer's end the chain tips down 0.65 cm over 3 links, going on round. 10 layers: 15.8 m (1053 links) at 30 deg, 9.6 m at 20, 16.1 m at 60/90. Waiting links are drawn, not simulated.
- Simulation window: links are woken so that 10 links ahead of the pick-up still lie where they were laid, at most 1200 per simulated second. The last link woken is held at its far end by a cone-twist joint to a fixed body in the next link's place. Links below y = 2.5 cm are frozen: their joint is removed and they become drawn-only, non-colliding meshes. About 130-200 links are active.
- Stepping: exactly 1/1800 s at every display speed. Each frame takes as many whole steps as its share of simulated time. Wake, freeze and measurement run after every step. 20 solver iterations.
- Bullet does all dynamics and contacts. The script does wake, anchor, freeze, the pile disc, and (read-only) measurement.

Theory checks

Biggins and Warner (2014): pile push a m v^2 up on the link being lifted, floor pull b m v^2. Steady flow gives $v^2 = g\,h1/(1 - a - b)$ and $h2 = a\,v^2/g$ (narrow arch). Here b = 0 by construction (frozen links, no joint, no collisions). Fresh runs, 10 layers, CF_SPEED=1, steady window t = 1.00 s to the last coil link woken. v = rate links land x 1.5 cm. "Push" = disc contact force minus the weight of links touching it, divided by m v^2 (CF_PUSH=1). "Carried" (new, CF_PUSH=1) = $-(T - m\,v_i^2 - m\,g\,y)/(m\,v^2)$ on the freely

falling chain 20-100 cm below the pick-up. T is the tension worked out from the links' accelerations, summed up from the free lowest link. A steady B&W chain keeps this equal to a from pile to floor.

Quantity	Theory / source	Simulation (this run)	Agreement
Same run at any display speed	must not depend on frame rate	0.25x with pause = 1x (30 deg); 0.5x = 1x (60 deg, CF_PUSH)	**yes, exact**
Speed bound	ideal chain sqrt(g h1) = 384 cm/s	397 / 450 / 425 / 432 cm/s (30/20/60/90)	faster, as a > 0 requires
Pile push a (direct)	B&W rod model a = (1 - l/(m b^2/4))/2: 1/6 for their chain, 0.31 for these links	0.31 / 0.27 / 0.19 / 0.17; 0.05-0.10 lower (0.22/0.17/0.13/0.12) if every link within 7 mm of the pile is counted as lying on it	push present, same order as theirs
a from speed, 1 - g h1/v^2	= push if nothing else acts	0.06 / 0.27 / 0.18 / 0.21	60, 90: within the push's uncertainty. 30: **no**, the rim holds the chain up with 0.31 m v^2 (20: 0.14)
a from height, g h2/v^2	= push (narrow steady arch)	0.07 / 0.05 / 0.07 / 0.07; h2 = 10.5 / 10.7 / 12.8 / 13.2 cm	**no**: fountain about 1/3 of a v^2/g with the measured push (35 cm at 60 deg)
30 deg: height vs speed	equal	0.07 vs 0.06	agree only by chance (push 0.31)
Energy at landing	lossless chain: 1/(2(1 - a)) of the drop = 60-72%	42 / 44 / 50 / 49%; rms landing speed 352 / 361 / 383 / 380 cm/s, 11-20% below v	**no**: 11-30% of the power lost on the way
a the falling chain carries	= push (steady, lossless)	-0.13 / -0.07 / 0.04 / 0.03	**no**: a third or less of the push reaches the falling chain; below zero where the rim takes more (20, 30)
Stationary pick-up (CF_LAYOUT=flake, CF_JAR=0, 40 layers; 45/60/90 deg)	h2 = a v^2/g	push 0.28/0.23/0.13; h2 18.2/17.3/9.1 cm = 0.36/0.40/0.36 of push v^2/g; spiral on the same table 0.30/0.29 (60/90)	**no**: helps a little (60 deg 17.3 vs 10.6 cm), not the cause
Flake: theory with the carried a (0.10/0.08/0.06)	h2 = a v^2/g; v = sqrt(g h1/(1 - a)); energy 1/(2(1 - a))	h2 ratio 1.01/1.16/0.79; rms landing speed 0.979/0.972/0.962 of the formula; energy 53/51/49% against 56/54/53%	**yes, within 21% (height), 4% (speed), 4 points (energy)**
One thing at a time (flake 60 and spiral 60): steps 3600, 7200; iterations 50, 100; friction 0; joint stops off (CF_STOP=179); 5, 15 woken ahead; links half size (CF_LINK=0.5)		h2/(push v^2/g) 0.25-0.50 (flake), 0.24-0.35 (spiral); energy 43-53%	no model channel restores the fountain
Shortcut vs every link simulated (CF_LIVE=1; 6 layers, 5 runs each at jar 146-154 cm, CF_PUSH=2)	the shortcut should not matter	h2 every link / exhibit's way: 8.6±0.4 / 10.6±1.5 (30), 6.3±0.5 / 10.8±0.6 (60), 5.5±0.4 / 11.2±0.8 (90) cm. Unmoved-pile push 0.49/0.23/0.20 vs 0.20/0.17/0.15. Carried -0.13/-0.19/-0.16 vs -0.01/0.01/0.01. Rim 0.26/0.32/0.28 vs 0.13/0.03/0.03 m v^2. v 399/379/397 vs 420/418/423 cm/s	shortcut **not** the cause of the lost push (every-link chain carries none either), but it **raises the fountain** 1.2x (30) to 2x (60, 90) and keeps the chain off the rim
Reference convergence		every link, 3600 steps/s: h2 8.5, 9.4 (30); 5.7, 6.0 (90). Settled 0.3 s first: 9.6 (30). Landed links left on the floor: 8.5 (30)	within the 5-run spread

Quantity	Theory / source	Simulation (this run)	Agreement
Cheaper match (CF_WAKE=layer)	= every link	4 live layers at 30: v 406±17, h2 8.7±0.6, push 0.48, carried -0.09, rim 0.20; 3 layers match at 60; at 90 neither (v 356-359 vs 397, rim 0.57-0.62 vs 0.28)	matches at 30 and 60
Cost per frame, 0.25x, standalone (Cabinet run, 1200 frames)	16.7 ms at 60 fps	exhibit 4.4 ms (12.8 s); 4 live layers 21 ms (30.3 s); every link 35 ms (46.9 s)	faithful ways not affordable: default unchanged
h2/h1	B&W measured 0.14 (a = 0.12 by their formula with b = 0)	0.070 / 0.071 / 0.085 / 0.088 (spiral), up to 0.13 (flake 45)	fountain lower
Chaos spread (jar 146, 148, 152, 154 cm)		v 414-448 cm/s, push 0.15-0.41, height-a 0.05-0.07	
Step 1/3600 s	converged?	20 deg: v 477, push 0.39, h2 19.8; 30: 437, 0.32, 14.3	**not converged** at 20 deg
Fountain above rim (CF_LOG, t >= 0.5 s), median (IQR), highest; 14 layers in brackets	README	20: 4.7 (3.5-5.8), 11.9 [4.5 (3.0-5.8)]; 30: 5.4 (3.4-8.0), 12.4 [4.1 (1.9-6.7)]; 60: 7.7 (6.0-8.3), 10.1 [6.3]; 90: 7.8 (6.6-8.6), 12.4 [7.3]	README replaced
Stiffer chain, higher fountain	B&W; old README	no trend beyond chaos (medians at 4 jar heights: 6-8 / 5-7 / 4-8 / 5-8 cm)	old trend came from the bad coil
Time to empty	README "about 4 s"	4.25 s (30), 4.00 (60, 90), 2.50 (20, shorter chain)	yes

Run used

```
cd demo/WyomingWill/ChainFountain
CF_PUSH=1 CF_SPEED=1 CF_BEND=<20|30|60|90> CF_LOG=d<b>.csv HOLD_S=0 ../../../release/bpp -f chain-fountain.lua -n 300
CF_PUSH=1 CF_SPEED=1 CF_LAYERS=14 CF_BEND=<b> ... -n 400 ; CF_HEIGHT=<146|148|152|154> ... ; CF_STEPS=3600 ... -n 400
CF_PUSH=1 CF_SPEED=1 CF_JAR=0 CF_LAYOUT=flake CF_LAYERS=40 CF_BEND=<45|60|90> HOLD_S=0 ... -n 400   # plus CF_STEPS,
    CF_ITER, CF_MU=0, CF_STOP=179, CF_AHEAD, CF_LINK=0.5 one at a time
timeout 3000 ../../../release/bpp -f chain-fountain.lua -n 1200        # 0.25x with pause: identical to 1x
bash ../Cabinet/report/cmp2.sh chain ChainFountain chain-fountain.lua 1200 cf   # IDENTICAL (22 lines); CF_PUSH=1:
    IDENTICAL (25); flake 60 CF_PUSH=1 (1300 frames): IDENTICAL (23)
CF_LIVE=1 CF_LAYERS=<6> CF_HEIGHT=<146..154> CF_BEND=<30|60|90> CF_PUSH=2 CF_SPEED=1 HOLD_S=0 ... -n 400   # and the
    same without CF_LIVE, and with CF_WAKE=layer CF_LIVELAYERS=<3|4>
CF_LIVE=1 CF_LAYERS=6 CF_PUSH=2 bash ../Cabinet/report/cmp2.sh chain ... 1100 cfl   # IDENTICAL (21); CF_WAKE=layer
    CF_LIVELAYERS=4 CF_PUSH=2, 1200 frames: IDENTICAL (26)
```

```
t= 4.25s  all of it out of the jar; the fountain stood up to 12.4 cm above the rim
  theory (Biggins and Warner 2014), steady part t=1.00-3.68s; h1 = 150.2 cm, the pick-up above where a link lands
    (the rim is 5.3 cm above the pick-up)
  chain 397 cm/s (the rate links land): an ideal chain (a = b = 0) would go sqrt(g h1) = 384 cm/s and not rise above
    the pick-up
  top of the chain 10.5 cm above the pick-up on average (h2): h2/h1 = 0.070 (their chain: 0.14); b = 0 here, so v =
    sqrt(g (h1 + h2)) = 397 cm/s
  a = g h2/v^2 = 0.07 from the height, 1 - g h1/v^2 = 0.06 from the speed (rod model for one of these links: 0.31)
  the links reach the floor at 352 cm/s (rms), with 42% of the energy of their drop (an ideal chain: 50%; one the
    pile pushes, more)
  the pile's push, measured (its contact forces less the weight lying on it): a = 0.31; the rim holds the chain up
    with 0.31 m v^2
  with that push a steady narrow arch would stand h2 = a v^2/g = 50.4 cm above the pick-up (it stood 10.5)
  what the falling chain carries of it (its tension, from the links' accelerations, 20-100 cm below the pick-up): a =
    -0.13
[flake 60, table] the pile's push ... a = 0.23 ... h2 = a v^2/g = 42.9 cm (it stood 17.3) ... carries of it ... a =
    0.08
[60 deg] a = g h2/v^2 = 0.07 from the height, 1 - g h1/v^2 = 0.18 from the speed ... a = 0.19; the rim ... 0.02 m v^2
```

Pause: CF_SPEED=1, HOLD_S=0 for 300 frames is an exact prefix of the default hold for 330 frames (24 lines; with CF_PUSH=1, 27 lines). Default output (CF_PUSH off) is byte-identical to the committed b241d9c. 0.25x with pause = 1x with CF_PUSH=1; flake 60 at 0.5x = 1x.

Known departures and honest caveats

- Fixed 15 Oct 2026: frame-rate dependence (all work per physics step, fixed 1/1800 s) and the 60-degree stall (last woken link anchored; landed links don't collide).

- Fixed 15 Oct 2026: the coil was laid wrong. The spiral went in to 3 cm at every bending, so at 20 deg 304 of 1054 joints were laid bent past their stop (worst about 29 deg against 20). At every bending each layer began with a 0.65 cm upright link (90 deg bends) and the next layer wound the other way. When woken, the solver kicked these joints straight. The old "stiffer chain, higher fountain" (20 deg median 11.1 cm) came from those kicks: with the fixed coil it is 4.7. Only the lead's 3 corners, woken at t = 0, are still laid past the limit.
- Measurement changes (15 Oct): v is now the rate links land, not the rate they are woken. At 20 deg and 4800 steps/s the waking ran ahead (wake rate 592 cm/s, above the sqrt(2 g h1) = 542 energy bound). New lines: landing speed and energy, and with CF_PUSH=1 the directly measured push. CF_PUSH is off by default because eachContact makes the run 2.6x slower in the Cabinet (34.7 s against 13.2 s for 1200 frames).
- **Located 15 Oct 2026, mechanism still open: the fountain is low because the pile's push doesn't reach the moving chain.** The pile pushes up with a = 0.13-0.31, but the falling chain carries only a third or less of it. A pile whose pick-up stays put (flake on a table) raises the fountain only a little (ratio 0.36-0.40 against 0.29-0.30 for the spiral on the same table). No model channel tried (step x2 and x4, iterations x2.5 and x5, friction 0, joint stops off, link size x0.5, look-ahead 5/15) brings the ratio above 0.50. With the push the chain does carry, the flake obeys all three B&W relations: height within 21%, speed within 4%, energy within 4 points. So the earlier apparent 11-30% energy loss mostly came from using the contact push as a. The gap is in the pick-up zone. The chain is dragged and lifted over several links (still only about 90% of full speed 4-6 cm above the pile, from the inferred tension), not picked up a link at a time. This does not change with step or iterations, so it is not solver convergence. Without joint stops the push itself halves. On the spiral, and with half-size links, the falling chain carries 0.00-0.04 while the fountain stands higher than that would give. So between pile and falling leg those flows are not steady either. Whether real chains also carry only part of the contact push is unknown; B&W's own h2 = 0.14 h1 means a = 0.12 against their rod model's 1/6.
- Not step-converged at 20 deg (1/3600 s: v +6%, h2 +85%). The run is chaotic, with spreads as in the table.
- The waking scheme is not neutral. With 5 links woken ahead instead of 10 (scratch copy with CF_K), v = 476 / 452 cm/s at 30 / 60 deg (+20% / +6%) and a from speed 0.35 / 0.28, against a push of 0.21 / 0.16. The anchor's force is not measured. With 20 the woken links are dragged into a heap (276-446 links moving; at 60 deg the rim carries 1.3 m v^2). So v is known to about 20%.
- **Tested 15 Oct 2026: the wake/anchor shortcut does not lose the push, but it does make the fountain too tall.** With every link simulated, the moving chain still carries none of the pile's push (carried -0.13 to -0.19), so the lost push is how rigid links come off a pile in this model. The full chain drags the top of the pile and rubs the rim (0.26-0.32 m v^2). Its fountain is 8.6 / 6.3 / 5.5 cm against the exhibit's 10.6 / 10.8 / 11.2 at 30 / 60 / 90 deg. The exhibit's fountain is therefore up to 2x taller than the full simulation of the same chain. A cheaper scheme (4 live top layers, no anchor) matches the full chain at 30 deg, but costs 21 ms/frame at quarter speed against 4.4 now (full: 35 ms; budget 16.7). The default is unchanged; the README says so. The flake pile could not be run all-live: on the bare table the live 25-layer stack is dragged off as a mass.
- Experiment options added (env only; defaults unchanged, byte-identical output): CF_LIVE, CF_SETTLE, CF_NOFREEZE, CF_WAKE=layer, CF_LIVELAYERS, CF_PUSH=2; CF_LAYOUT=flake, CF_JAR=0, CF_ITER, CF_MU, CF_AHEAD, CF_STOP, CF_LINK; CF_PUSH=1 now also prints h2 = a v^2/g and the carried a. The flake is for 45 deg and more only (wider turns don't fit), a layer holds about 0.3 m, so a 40-layer pile is 26 cm tall and h1 drifts from about 150 to 125 cm in a run.
- The coil is a rigid disc, not a heap. When the disc drops a layer, the last ~10 woken links of the old layer fall 0.65 cm. The anchor is infinitely strong. The floor is a freeze at 2.5 cm (b = 0 by construction).

References

- S. Mould, video, 2013 (README).
- J. S. Biggins, M. Warner, "Understanding the chain fountain", Proc. R. Soc. A 470 (2014) 20130689, arXiv:1310.4056: $v^2 = g\ h1/(1 - a - b)$, $h2/h1 = a/(1 - a - b)$, rod model $a = (1 - I/(m\ b^2/4))/2 = 1/6$ for three beads per link, measured h2 = 0.14 h1, energy requires a + b <= 1/2.

5.7 The falling Slinky

Agrees with theory. Bottom stays put 0.2038 s against sqrt(M/3k) = 0.2018 s: 0.6% from having 86 separate coils (an exact 86-coil chain waits 0.2030 s) and 0.4% from Bullet's step, both now explained in the README.
Folder: demo/WyomingWill/Slinky/

What it shows

A Slinky hangs stretched under its own weight, with a steel ball hanging level with its bottom. When both are released, the ball falls at once, but the Slinky's bottom hangs motionless until the collapse coming down from the top reaches it. Meanwhile the Slinky's centre of mass falls exactly like the ball. Beside it, the same Slinky walks end over end down seven stairs.

Model

- Falling Slinky: M = 215 g, N = 86 coils (SL_COILS), each a Bullet rigid body of M/N = 2.5 g. The collision shape is a thin square plate, 7 cm across. Closed length is 5.6 cm, stiffness k = 1.76 N/m, with no pre-tension. Neighbours are joined by a btSliderConstraint along the axis, with a lower limit at the closed spacing, no restitution, and rotation locked. This gives inelastic coil-on-coil stops. The script applies the spring force kseg = k(N-1) per gap before every physics step, and only while the gap is stretched. Each coil's initial stretch is set from the weight below it, so nothing moves before release. Bullet steps at 4800 per simulated second with 60 solver iterations and no damping. Coil friction is 0.5 and floor friction 0.6. The display runs at 0.1x real speed (SL_SPEED). The hand releases after a 0.5 s "hold" slider. 0.05 s after landing, the pile is replaced by a frozen closed stack, because loose coils jostle in Bullet.
- Stairs walker: the same M and k, built as 43 capsule pieces of 2 coils each, constrained to the plane (x, y, rotation about z). The two sides of each piece connect to the next piece by pull-only springs of k(42)/2 each, with damping 0.05 N s/m. A shear spring of 10 k(42) keeps pieces in line. Forces are applied by the script as velocity kicks each step, and Bullet handles the contacts (friction 0.6, restitution 0). There are 7 steps, 12 cm rise and 13 cm tread, from 84 cm. It restarts 3 s after coming to rest.

Theory checks

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Time the bottom stays put	t* = sqrt(M / 3k) = 0.2018 s (script)	0.2038 s (86 coils)	+1.0% (README now: 0.2038 s)
Same, the discrete chain itself (new, 15 Oct 2026: reconstruction/slinky-chain.c, point masses + pull-only springs + sticking collisions, step 1e-6 s)	continuum limit 0.2018	43: 0.20412; 86: 0.20296; 172: 0.20238; 344: 0.20208 s (same at 5e-7 s)	+1.15 / +0.58 / +0.29 / +0.14 %: halves as coils double
Bullet vs step size, 86 coils (SL_STEPS, new env)	the chain's 0.2030	4800: 0.2038; 9600: 0.2033; 19200: 0.2031; 38400: 0.2030 s	converges to the chain; the step adds +0.4% at the default
Bullet, 43 / 86 / 172 coils at 4800 steps/s	should approach 0.2018 as N grows (old README)	0.2046 / 0.2038 / 0.2048 s	**does not**: the step error grows with N
Same at 9600 / 19200 steps/s	—	9600: 0.2043 / 0.2033 / 0.2030; 19200: 0.2042 / 0.2031 / 0.2026 s (38400: 43 coils 0.2042, 86 0.2030)	with small steps it does approach, as the chain does

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Detection threshold (1 cm/s vs 10 cm/s, scratch copy)	should not matter	0.2038 s both	not the cause
Hanging stretch	M g / 2k = 59.9 cm	printed length 65.5 cm minus 5.6 cm closed = 59.9 cm	exact (README: "stretches 60 cm")
Ball's fall when the bottom starts moving	g t^2 / 2 at t = 0.2038 s = 20.4 cm	20.4 cm	exact (README: 20 cm)
Bottom stationary before t*	displacement 0	bottom drop 0.000 cm at t = 0.100 and 0.200 s (SL_LOG)	exact
Centre of mass in free fall	y_cm drop = g t^2 / 2 (only gravity acts)	t = 0.1/0.2/0.3/0.4 s: CoM drop 4.915/19.641/44.176/78.521 cm vs ball 4.915/19.640/44.176/78.521 cm; g t^2/2 = 4.905/19.620/44.145/78.480	CoM = ball to 0.001 cm. The 0.01-0.04 cm offset from g t^2/2 is one 1/600 s logging-frame phase
Landing time	(no theory given)	0.492 s after release	(README gives none)
Stairs: time per step, 7 steps	README now: "about 0.4 s a step (2.6 to 3.1 s for a clean walk)"; sample walk 3.03 s	21 walks at native 0.1x speed: clean walks 2.56-3.07 s to the floor	matches the corrected README (the old "a third of a second", 2.64 s sample, was from an earlier run)
Stairs failure rate	old README: "one walk in three or four" goes wrong; README now: "nearly one walk in two"	native 0.1x speed, 21 walks in a row (HOLD_S=0, -n 84000): 12 clean; 4 reached the floor but skipped steps (3 4 5 F; 3 4 5 7 F; 3 4 5 6 F F; 3 4 5 F); 3 stopped on step 7; 2 never started (rest on step 2, no landings). Earlier SL_SPEED=1 sample: 4 of 9 clean	9/21 = 43% imperfect; README corrected

Run used

```
cd demo/WyomingWill/Slinky
SL_LOG=sl.csv HOLD_S=0 timeout 1500 ../../../release/bpp -f falling-slinky.lua -n 9000      # 15 s simulated at 0.1x
SL_COILS=172 HOLD_S=0 timeout 1500 ../../../release/bpp -f falling-slinky.lua -n 600
SL_COILS=43  HOLD_S=0 timeout 1500 ../../../release/bpp -f falling-slinky.lua -n 600
SL_SPEED=1   HOLD_S=0 timeout 2400 ../../../release/bpp -f falling-slinky.lua -n 3600      # 60 s, walk statistics
    (report)
HOLD_S=0 ../../../release/bpp -f falling-slinky.lua -n 84000                                 # 15 Oct 2026: 21 walks
    at 0.1x
SL_STEPS=9600/19200/38400 (SL_COILS=43/86/172) ... -n 400                                    # 15 Oct 2026: step
    study
cc -O2 -o slinky-chain reconstruction/slinky-chain.c -lm; ./slinky-chain {43,86,172,344} 1e-6  # 15 Oct 2026
```

(HOLD_S has no effect here. This exhibit uses its own "hold" slider, 0.5 s.)

```
--- hung up: 215 g, 86 coils, stiffness 1.76 N/m; 65.5 cm long hanging (5.6 cm closed up); the bottom should stay put
    for 0.202 s ---
t= 0.254s  the bottom starts to move, 0.2038 s after letting go (it should, by theory: 0.2018 s); the ball had fallen
    20.4 cm
t= 0.542s  the Slinky lands, 0.492 s after letting go
stairs t= 3.75s  the walking Slinky comes to rest, on the floor: 7 steps down in 3.03 s
stairs t= 3.64s  the walking Slinky comes to rest, on the floor: 7 steps down in 2.98 s
[43 coils]  t= 0.255s  the bottom starts to move, 0.2046 s after letting go (it should, by theory: 0.2018 s); the
    ball had fallen 20.6 cm
[172 coils] t= 0.255s  the bottom starts to move, 0.2048 s after letting go (it should, by theory: 0.2018 s); the
    ball had fallen 20.6 cm
[SL_SPEED=1] stairs t= 2.18s  the walking Slinky comes to rest, on step 6: 5 steps down in 1.40 s
[SL_STEPS=19200] t= 0.253s  the bottom starts to move, 0.2031 s after letting go (it should, by theory: 0.2018 s);
    the ball had fallen 20.2 cm
86 coils, step 1e-06 s: the bottom moves at 0.20296 s (its gap closes at 0.20296 s); sqrt(M/3k) = 0.20179 s
[0.1x, walk 9]  stairs t= 7.25s  the walking Slinky comes to rest, on step 7: 6 steps down in 2.80 s
[0.1x, walk 14] stairs t= 1.28s  the walking Slinky comes to rest, on step 2: 1 steps down in 0.00 s
```

Known departures and honest caveats

- Fixed on 15 Oct 2026, the hang-time explanation: the README used to say more coils bring the wait closer to sqrt(M/3k). At the default 4800 steps/s they do not (43/86/172 coils: 0.2046/0.2038/0.2048 s). Established cause: the +1.0% at 86 coils is +0.6% from the discreteness itself (an exact chain of 86 sticky point masses waits 0.2030 s; the gap halves as coils double) and +0.4% from Bullet's step (0.2038 at 4800 steps/s, 0.2031 at 19200, 0.2030 at 38400). The step error grows with N because the collapse front then crosses each coil in fewer steps, which is why 172 coils are no better at 4800 steps/s. The detection threshold (1 vs 10 cm/s) makes no difference. README and lua header now say this; SL_STEPS was added for headless testing, and reconstruction/slinky-chain.c (the independent chain check the README cites) was added. Default steps not raised: 4800 steps/s already costs ~20 ms per frame here with the 86 coils and the stairs walker.
- The theory is for a continuous spring with **no pre-tension**. The model deliberately has none, although a real Slinky does.
- Coil collisions are perfectly inelastic slider limits, not true contact. After landing, the coils are replaced by a frozen pile, so post-landing dynamics are not physical.
- The walker is a planar, 43-piece reduced model with tuned damping. The README says 0.05 N s/m was chosen because less damping gallops and more stalls. The 12 x 13 cm stairs are the size it "walks best on". Walks are sensitive to tiny differences (the walker is rebuilt identically each time; walks differ through Bullet's internal ordering, so the "statistics" are of that sensitivity). At SL_SPEED=1 landings are detected once per 1/60 s frame instead of per 1/600 s (the physics step, 1/4800 s, is the same), so brief landings can be missed and a clean walk logged as skipping; the README now says to count walks at the native 0.1x speed. Fixed on 15 Oct 2026: the README's "one walk in three or four" is now "nearly one walk in two", with the 21-walk breakdown, and its sample walk listing is replaced by walk 1 of this run.
- The ball is a 50 g Bullet sphere released from a fixed constraint at the same instant.

References

- None cited by name in the code or README. The sqrt(M/3k) result is derived in the README from momentum M g t.

5.8 The Brazil-nut effect

Agrees with theory, with a numerical or open caveat. Qualitative: the big bead rises and the convection runs the right way. Floor push 0.18 of the bed's weight against 0.20 rigid. Lift-off at 46-48° against 19.5° for a rigid bed is mostly Bullet's soft contacts (stiffer: 28-30°), not converged.
Folder: demo/WyomingWill/BrazilNut/

What it shows

Three glass-fronted boxes, each holding a one-bead-thick layer of beads, are shaken vertically at 7 Hz. Shaken gently (peak 0.8 g), the bed never leaves the floor and nothing happens. Shaken hard (3 g), a big bead starting at the bottom rises to the top within seconds and stays there. A small coloured bead in the third box rises too, but then sinks again by a wall and circulates. This shows the convection rolls (up in the middle, down by the walls) that Knight, Jaeger and Nagel identified as the mechanism.

Model

Bullet does everything here; the script adds no custom contact or integration.

- Beads: 140 glass spheres per box (density 0.0025 kg/cm^3), radius 0.5 cm, with a deterministic size spread of +/-5% (BN_POLY = 0.1) so they do not crystallise. Big bead: radius 1.5 cm. All beads are locked to the x-y plane (linear factor (1,1,0), rotation about z only).
- Friction: 0.8 bead-bead, 1.0 walls. Restitution 0.1 for both.
- Box: a 1e4 kg compound body (floor plus two walls, 15 cm wide inside, 24 cm tall) with its own gravity off. The script sets its position and velocity to the sinusoid y = A sin(wt) before every physics step.

The front and back glass are not colliders; the planar constraint stands in for them.

- Shaking: f = 7 Hz, A = Gamma g / w^2, which gives 4.1 mm for Gamma = 0.8 and 15.2 mm for Gamma = 3.
- Stepping: 2400 physics steps per second, 20 solver iterations. Beads settle for 0.5 s before shaking starts. The boxes are re-poured every 90 s.
- Floor push: the floor's push on the bed is measured from how much the heavy box is slowed in each step, binned by phase in 2-degree bins over shakes 5 to 12.

Theory checks (two seeds; the default seed 7 is the README's)

Quantity	Theory (formula / source)	Simulation, seed 7	Seed 8	Agreement
Amplitude for Gamma	A = Gamma g / (2 pi f)^2	4.1 mm (0.8 g), 15.2 mm (3 g)	same	exact (definition)
Gamma = 0.8: bed stays on floor	min floor push = (1 - Gamma) W = 0.20 W (rigid bed)	never left; min push 0.18 W	0.20 W	0.02 W low / exact
Gamma = 3: lift-off phase, middle box	rigid bed: arcsin(1/Gamma) = 19.5 deg (a perfectly inelastic rigid bed, iterated: lands at 1.4 deg, leaves at 19.5)	46 to 48 deg	46 to 48 deg	**Disagrees by about 27 deg; tested 15 Oct 2026: mostly Bullet's contact compliance** (next rows)
Same, right box	19.5 deg	50 to 52 deg	42 to 44 deg	as above
Lift-off phase vs contact stiffness (seed 7; middle, right box)	should tend to 19.5 if the late lift-off is numerical softness	2400 steps/s, 20 iterations (default): 46-48, 50-52; 4800/s: 36-38, 32-34; 100 iterations: 36-38, 38-40; 9600/s + 60 iterations: 28-30, 36-38	moves toward 19.5 as contacts stiffen; not converged at the stiffest tried	
Lift-off phase vs bounce, friction (seed 7)	—	restitution 0: 42-44, 40-42; friction 0.3/0.3: 40-42, 54-56	secondary	
Big bead rises (Brazil-nut effect)	qualitative (Knight et al. 1993: rises with convection)	top at 6.3 s (44 shakes), rose 8.4 cm	21.4 s (149 shakes), rose 10.8 cm (re-run 15 Oct 2026: same)	qualitative yes; timing varies 3x with the pour. At 4800 steps/s (seed 7): 11.9 s (82 shakes); coloured bead 19.3 s
Big bead stays up	qualitative	upper half 77% of the time	91%	yes
Convection, middle box	rise in middle, fall at walls	middle +5.5 mm/s, walls -10.1 mm/s	+4.8 / -9.5 mm/s	correct sign / pattern
Convection, right box	same	+7.8 / -12.9 mm/s	+7.1 / -12.5 mm/s	correct sign / pattern
Coloured bead circulates	up nearer the middle, down nearer the walls	4 round trips; up 3.6 cm, down 4.6 cm from centre (walls at 7.5)	4 trips; 2.6 / 3.4 cm	yes
Coloured bead reaches top	qualitative	11.6 s (81 shakes), rose 8.9 cm	12.7 s (89 shakes), 9.3 cm	yes

The only quantitative theory comparisons are the Gamma < 1 floor push and the rigid-bed lift-off phase. Everything else checks direction, sign and pattern. There is no comparison with a measured rise rate or convection speed, and no conservation check (the system is driven and dissipative).

Run used

```
cd demo/WyomingWill/BrazilNut && HOLD_S=0 timeout 1500 ../../../release/bpp -f brazil-nut.lua -n 3700
cd demo/WyomingWill/BrazilNut && BN_SEED=8 HOLD_S=0 timeout 1500 ../../../release/bpp -f brazil-nut.lua -n 3700
```

```
  Left   Gamma 0.8: the bed never left the floor (the floor's push never less than 0.18 of its weight; a rigid bed's:
    0.20)
  Middle Gamma 3.0: the bed leaves the floor 46 to 48 degrees into the upswing (a rigid bed: arcsin 1/Gamma = 19.5;
  Right  Gamma 3.0: the bed leaves the floor 50 to 52 degrees into the upswing (a rigid bed: arcsin 1/Gamma = 19.5;
  Middle the big bead reaches the top at 6.3 s (44 shakes), having risen 8.4 cm
  Right  the coloured bead reaches the top at 11.6 s (81 shakes), having risen 8.9 cm
  Middle from 5 s to a minute: beads in the middle rising 5.5 mm/s on average, beside the walls falling 10.1 mm/s
  Middle the big bead, since reaching the top, has been in the bed's upper half 77% of the time
  Right  from 5 s to a minute: beads in the middle rising 7.8 mm/s on average, beside the walls falling 12.9 mm/s
  Right  the coloured bead has gone round 4 times: going up it was 3.6 cm from the middle on average, going down 4.6
    cm (the walls are 7.5 cm out)
[seed 8] Middle the big bead reaches the top at 21.4 s (149 shakes), having risen 10.8 cm
```

Seed 7 reproduced every console line in the README exactly. The README omits the "Right ... 50 to 52 degrees" line, but that is not a disagreement.

Console after the 15 Oct 2026 text change (seed 7):

```
  Middle Gamma 3.0: the bed leaves the floor 46 to 48 degrees into the upswing (a rigid bed: arcsin 1/Gamma = 19.5;
         this one presses on longer: Bullet's contacts give a little, and the stiffer they are, the sooner it leaves)
```

Stiffness runs: BN_STEPS=4800; BN_STEPS=9600 (and 60 iterations, scratch copy); 100 iterations (scratch copy); BN_E=0; BN_MUB=0.3 BN_MUW=0.3; each -n 110. BN_STEPS=4800 -n 1500 for the rise. BN_SEED=8 -n 1400.

Pause and hall checks after the console change (15 Oct 2026): hall against standalone (cmp2.sh, 900 frames): IDENTICAL (7 run lines); HOLD_S=0 for 400 frames is a prefix of the default pause for 430 frames (10 lines).

Known departures and honest caveats

- **The parameters were tuned.** The README says that 7 Hz, 3 g and high friction "were found by trying". At 10 Hz and 2.5 g with less friction, the big bead rose "only now and then" (README; not re-run here).
- The lift-off phase is far from the rigid-bed value: about 47 deg against 19.5 deg. The README used to explain it as a springy bed squeezed by its landing, untested. Tested 15 Oct 2026 (scratch copy that also printed the push-vs-phase profile and took a solver-iteration setting): stiffer contacts move the lift-off toward 19.5 (4800 steps/s: 36-38 deg; 100 iterations: 36-38; 9600 steps/s with 60 iterations: 28-30, middle box), so the late lift-off is mostly Bullet's soft, position-corrected contacts, not a property of glass beads. It had not converged at the stiffest setting tried, which is too slow for the exhibit. In the default profile the floor's push is 4-10 W from 0 to 18 deg (the landing; a rigid inelastic bed would land at 1.4 deg), 1.4 W at 30-36 deg, and drops below 0.1 W at 48 deg. The console line, lua header and README now say this ("this one presses on longer: Bullet's contacts give a little, and the stiffer they are, the sooner it leaves"); how much later than 19.5 deg a real bed of glass beads would leave is stated as not tested.
- The rise time depends strongly on the initial packing: 6.3 s for seed 7, 21.4 s for seed 8 (re-run 15 Oct 2026). Since 15 Oct 2026 the README says so (and gives the 4800-steps/s rise, 11.9 s); its table now reads "comes up (within seconds with this pour; it can take twenty)".
- The bed is 2-D (one bead thick) with high wall friction, which drives the wall-shear convection. A real 3-D jar of nuts has other mechanisms (percolation, void filling, inertia) that this setup neither isolates nor excludes.
- The box is a kinematic driver (prescribed position and velocity on a 1e4 kg body), not a dynamic shaker. The floor-push measurement depends on that huge mass ratio.
- Bead sizes come from a deterministic formula, and pouring uses the script's own linear congruential random-number generator, so each pour is identical for a given BN_SEED.

References

- J. B. Knight, H. M. Jaeger, S. R. Nagel, Phys. Rev. Lett. 70, 3728 (1993): the big bead rises with the convection (README).

5.9 The solar system

Agrees with theory. Planets within 5 km of JPL's DE421 after 10 years, the Moon 39 km; Mercury's perihelion 42.981″/century against 42.981″; rotation checks hold. The checks now stop at their exact dates, and give the same lines at any rate.
Folder: demo/WyomingWill/SolarSystem/

What it shows

The Sun, the eight planets, Pluto, five asteroids (Ceres, Pallas, Juno, Vesta, Hygiea) and 21 moons, moving under gravity from 2026-01-01 0h TDB, starting from JPL's states for that day. An orrery shows heliocentric distances square-rooted (directions and ecliptic heights true), and vitrines show each planet with its moons. Every body wears a drawn map and turns by the IAU rotation model, so the Moon's librations and Mercury's 3:2 spin-orbit resonance can be seen. The console compares the integration with JPL every year (planets, Moon) and every month (moons, asteroids).

Model

All gravity is computed in Lua (`solar-engine.lua`). Bullet only places meshes: world gravity is set to 0 and every body's transform is written each frame, so Bullet runs no dynamics. Everything is in ICRF coordinates, AU and days.

- *Main integration*: 16 bodies, all pulling each other (Newton). They are the Sun, Mercury, Venus, Earth, the Moon, Pluto, five asteroids, and the system barycentres of Mars, Jupiter, Saturn, Uranus and Neptune. Two terms are added: the first-order Schwarzschild correction to the Sun's pull, $GM/(c^2 r^3)[(4GM/r - v^2)r + 4(r\cdot v)v]$, and the Earth's J2 acting on the Moon (code comment: the Earth's pole is taken as ICRF z, "near enough"). The step is a fixed 1-day Bulirsch–Stoer step: Gragg modified midpoint with 2, 4, 6, 8 and 10 substeps, Richardson-extrapolated to 10th order, with no error control. Positions between steps come from cubic Hermite interpolation (for display; the checks are taken at whole days, the ends of steps). Alongside runs a massless "Mercury without Einstein" twin, pulled by Newton only.
- *Moon systems* (Mars, Jupiter, Saturn, Uranus, Neptune, Pluto): a Wisdom–Holman drift–kick–drift step. Each moon drifts on its Kepler orbit for half a step. The kick holds the other moons (with the planet's reflex wobble), the planet's J2 and J4, Mars's C22/S22 rotating with Mars, and the Sun's tide (the Sun taken from the main integration). Each system has 16 steps per orbit of its innermost moon (console, in days): Mars 0.020, Jupiter 0.111, Saturn 0.059, Uranus 0.088, Neptune 0.367, Pluto 0.399. Between steps a moon is carried on its Kepler orbit from the last step (less than one step).
- *Checks*: each frame first runs every check whose date falls before the frame's time, with the integration stopped exactly at that date (the yearly and monthly dates are whole days from the start, so the main integrator's 1-day steps land on them), then advances to the frame. The check figures are therefore the same at any SS_RATE (fixed 15 Oct 2026; before, checks ran on the first frame after the date and interpolated back).
- *Perihelion check*: a separate two-body run of Mercury plus the Schwarzschild term, stepped by RK4 at h = 0.01 d. The Laplace–Runge–Lenz vector is averaged over each perihelion-to-perihelion orbit and a straight line is fitted through its direction.
- *Data*: DE421 states and masses for the Sun, planets, Moon and Pluto. Moons and asteroids start from JPL Horizons. Moon-system masses are DE440's. Pluto's GM is 975.43 km^3/s^2, the value Horizons' Charon orbit implies (README: DE421's 977.0 puts Charon 20,000 km off in a year).
- *Rotation*: IAU WGCCRE 2015 pole and prime-meridian models, keeping only the main periodic terms. Mars, its moons and the Moon's terms come from the 2009 report.

Theory checks

Values are from the fixed code (15 Oct 2026); the check lines are identical at SS_RATE=5, 60 and 100. The Phobos, Titania, Miranda and Ceres rates are differences between yearly Horizons lines (2026-01-01 → 2027-01-01 → 2031-01-01).

Quantity	Theory (formula / source)	Simulation (this run)	Agreement
Planets vs DE421 (heliocentric), 2027 / 2031 / 2036	JPL DE421	2027: all 0 km. 2031: Mars 1, Jupiter 2, others 0. 2036: Mercury 1, Earth 1, Mars 1, Jupiter 5, Saturn 1, others 0	≤ 5 km in 10 yr (README "within 5 km") ✓
Moon vs DE421 (geocentric)	DE421	4 / 20 / 39 km (2027 / 2031 / 2036)	39 km ≈ 0.05″ (README)
Mercury without GR, lag behind DE421	needs the Schwarzschild term	49 km (2027), 624 (2031), 1813 (2036); not monotonic (733, 624, 404, 286, 531, 1107, 1813 for 2030–36)	shows that GR is needed; README "49 / 1800 km", and now says the yearly figure is not monotonic ✓
Mercury perihelion advance	$\Delta\varpi = 6\pi GM/(c^2a(1-e^2))$ per orbit = 42.981″/century	42.981″/century (over 3, 20 and 40 orbits)	0.000″ to 3 decimals
Energy (Newtonian)	conserved	max $\lvert\Delta E/E\rvert$ 1.4e-9 (2035)	README "about 1e-9" ✓
Angular momentum	conserved	max $\lvert\Delta L/L\rvert$ 3.0e-10 (2035)	README "3e-10" ✓
Laplace resonance L_I − 3L_E + 2L_G	180° (free libration 0.07°, README)	yearly means 179.87–180.12°, spread 0.15–0.23°; 20-day means 179.65–180.32°	±0.35° (README "179.7–180.3") ✓; these are frame-sampled and move by ~0.01° with SS_RATE
Neptune–Pluto 3:2	P_Pluto/P_Neptune ≈ 1.5; min separation > 17 AU	ratio 1.489–1.515 (osculating); closest 32.04 AU	consistent; 25 yr cannot test the 17 AU bound (README)
Moons, asteroids vs Horizons, 2026-02-01	JPL Horizons	Phobos 83, Io 5, Mimas 41, Titania 102, Titan 0, Ceres 1 km	start-up; README same line identical ✓ (was Io 78, Mimas 518 from a high-rate run)
Same, 2027-01-01	Horizons	Phobos 986, Mimas 575, Titania 858, Miranda 404, Ceres 218, Titan 31 km	README same line identical ✓
Same, 2031-01-01	Horizons	Phobos 4771, Titania 4502, Miranda 2060, Mimas 2794, Ceres 2010, Titan 90, Triton 108, Charon 209 km	README same line identical ✓
Phobos drift	Horizons	≈ 950 km/yr	README "950 km a year" ✓
Titania / Miranda drift	Horizons	≈ 900 / 400 km/yr	README ✓
Earth subsolar point, 2026-01-01 0h TDB	IERS Earth via Skyfield: 23.02 S, 178.87 W	23.01 S, 178.79 W	0.01° lat, 0.08° lon
Moon libration (sub-Earth point)	longitude up to 7.9°; latitude 1.5° + 5.0–5.3° = 6.7–6.8°	each year 7.5–8.0 W to 7.4–8.1 E; 6.8–6.9 S to 6.8 N	✓
Mercury 3:2 hot poles at perihelion	subsolar longitude 0° or 180°	0.0 to 0.2 from 0°/180° (e.g. 2036: −180.0, 0.0, −180.0, 0.0)	≤ 0.2°
WH vs Bulirsch–Stoer prototype (README)	BS reference	Mimas 3 km off after 20 d (README, not rerun)	—

Run used

```
cd demo/WyomingWill/SolarSystem
HOLD_S=0 SS_RATE=60  SS_YEARS=10 timeout 3000 ../../../release/bpp -f solar-system.lua -n 3660
HOLD_S=0 SS_RATE=100 SS_YEARS=10 timeout 3000 ../../../release/bpp -f solar-system.lua -n 2200
HOLD_S=0 SS_RATE=5|60|100 SS_YEARS=4 ... -n 13180|1100|662          # three years, rate comparison
```

All 120 check lines (DE421 blocks and Horizons lines, including the yearly Laplace means) are identical, character for character, at rates 60 and 100 over ten years, and at 5, 60 and 100 over three; the rate-60 lines are also identical to the pre-fix rate-60 run (at 1 d/frame the old sampling already landed on the dates). Only the frame-sampled lines (20-day Laplace means, the Moon's face extremes, Mercury's perihelion longitudes) differ, by 0.01–0.2. Verbatim:

```
the Earth: on 2026-01-01 at 0h (TDB) the Sun is overhead at 23.01 S, 178.79 W (... through Skyfield: 23.02 S, 178.87
    W ...)
the moons: Wisdom-Holman, steps (days): Mars 0.020, Jupiter 0.111, Saturn 0.059, Uranus 0.088, Neptune 0.367, Pluto
    0.399
2026-02-01  against Horizons, km off: Phobos 83, Deimos 4, Io 5, Europa 11, Ganymede 19, Callisto 17, Mimas 41,
    Enceladus 6, ...
2027-01-01  against JPL's DE421, km off, from the Sun: Mercury 0, Venus 0, Earth 0, Mars 0, Jupiter 0, Saturn 0,
    Uranus 0, Neptune 0, Pluto 0;
            Mercury's perihelion turned by Einstein: 42.981" a century, over 3 orbits (theory, ... an orbit: 42.981")
2031-01-01  against Horizons, km off: Phobos 4771, ..., Mimas 2794, ..., Titania 4502, Oberon 937, Triton 108, Charon
    209, Ceres 2010, ...
2036-01-01  against JPL's DE421, km off, from the Sun: Mercury 1, Venus 0, Earth 1, Mars 1, Jupiter 5, Saturn 1,
    Uranus 0, Neptune 0, Pluto 0;
            the Moon, from the Earth: 39 km
            a Mercury without Einstein's correction: 1813 km off
            energy steady to -1.3e-09, angular momentum to -2.5e-10
```

Before the fix, the rate-100 run gave (2026-02-01) Io 945, Mimas 4688, Enceladus 2014 and (2036) Mercury 1220, Venus 11, Moon 122 km; now Io 5, Mimas 41, Enceladus 6 and Mercury 1, Venus 0, Moon 39, as at rate 60.

Checks: hall vs standalone (`SS_RATE=100 SS_YEARS=3 bash ../Cabinet/report/cmp2.sh solar SolarSystem solar-system.lua 700 ss`): IDENTICAL (101 lines). Start pause: HOLD_S=0 for 600 frames is a prefix of the default pause for 630 frames (SS_RATE=100, SS_YEARS=3).

Known departures and honest caveats

- **Fixed 15 Oct 2026: check sampling at high SS_RATE.** The checks used to fire on the first frame after their date. At 100 d/s (1.67 d a frame) the main bodies were then read off the Hermite cubic outside its step (extrapolated up to a day: Mercury 1220 km, the Moon 122 km in 2036) and the moons carried back from their last Wisdom–Holman step by a bare Kepler orbit (Mimas 4,688 km, Io 945 km on 2026-02-01). Now each frame first stops the integration at every check date it passes and runs the check there (`checks(tn)` in solar-system.lua), then goes on to the frame's time; the displayed positions are unchanged. A second rate dependence was removed in `E.advance` (solar-engine.lua): the moons' steps inside the current day are finished before the next day is stepped, so each kick takes the Sun from the same day's interpolant whatever the frame pattern. With both, the check lines agree to the km at rates 5, 60 and 100.
- **Fixed 15 Oct 2026: README sample output.** The 2027 block (Laplace 180.08, spread 0.18, with its 20-day-means line), the 2036 planets (Mercury 1, Earth 1, Mars 1, Jupiter 5, Saturn 1), the Horizons 2026-02-01 / 2027 / 2031 lines, "within 5 km", Phobos 950 km/yr, "20–210 km" and the Mimas wording now come from the fresh run. The README now says the no-Einstein distance is not monotonic year to year (49, 190, 494, 733, 624, 404, 286, 531, 1107, 1813 km for 2027–36): the orbit differs, and the distance depends on where Mercury is in it each Jan 1.
- Still frame-sampled (by design, not checks against theory values): the Laplace 20-day means and yearly means, the Moon's face extremes, Mercury's perihelion longitudes and Neptune–Pluto closest approach are taken once per frame, so they change slightly with SS_RATE (20-day means by ~0.01°, the yearly means agreed to 0.01° in these runs, perihelion longitudes by up to 0.2°; at 1.67 d/frame the perihelion is located to within a frame).
- Between Wisdom–Holman steps a moon's position is its Kepler drift from the last step (no kick); at the check dates this covers under one step (≤ 0.4 d for Pluto, 0.059 d for Saturn) and is the same at every rate.
- Physics left out (README):

- The Moon's 39 km is the Moon's figure, tides and Earth-pole precession; the J2 pole is fixed at ICRF z.
- Phobos (~950 km/yr) is Mars's higher-order gravity field. Adding C22/S22 cut it from 3,200 km/yr, and Deimos from 330 to 50.
- Titania and Miranda (900 and 400 km/yr) are not explained by J2 or older masses.
- Mimas: 575 km after a year, 2,794 after five, growing with a month-to-month wobble of 100–200 km (the README's old "500–2,400, rising and falling" is not what the monthly series shows; README updated). The README attributes it to its 2:1 resonance with Tethys (70-year swing); that cause is not tested here.
- Asteroids (Ceres 2,000 km in 5 yr, ~1″) lack Horizons' 11 extra perturbing asteroids and DE441's barycentre.

- DE421 masses drive the main integration, but DE440 masses drive the moon systems. This mix is deliberate.
- The 1-day Bulirsch–Stoer step has no error control. Pluto–Neptune "> 17 AU" is a known fact, not shown in the 10 years run.
- Rotation models keep only the main periodic terms. Several moons' meridians (Titan, Mimas, Enceladus, Tethys, Dione, Iapetus) and Pluto/Charon are not cross-checked against a second source (README).
- Surface maps are drawn, not photographs. Variable features such as the Red Spot and sunspots are illustrative.
- The README's Lainey L1.2 alternative start (1.5° swing, period 2039 d vs 2070 d theory) was not rerun, since it needs the Horizons files removed.

References (as cited in the code and README)

- JPL DE421 ephemeris (via skyfield-data/jplephem); DE440/DE441 (masses, Horizons' asteroid barycentre); JPL Horizons; JPL satellite solutions.
- Archinal et al. 2018, "Report of the IAU WGCCRE: 2015", Celest. Mech. Dyn. Astr. 130:22; the 2009 WGCCRE report, CMDA 109:101; NAIF pck00008; the hapsira library.
- Wisdom & Holman mapping; Bulirsch–Stoer / Gragg modified midpoint; Lainey's L1.2 theory of the Galilean moons (via astronomy-engine).
- IERS Earth orientation via Skyfield; Natural Earth coastlines.