Problem Set 1

Applied Stats/Quant Methods 1

Due: October 1, 2023

Instructions

- Please show your work! You may lose points by simply writing in the answer. If the problem requires you to execute commands in R, please include the code you used to get your answers. Please also include the .R file that contains your code. If you are not sure if work needs to be shown for a particular problem, please ask.

- Your homework should be submitted electronically on GitHub.

- This problem set is due before 23:59 on Sunday October 1, 2023. No late assignments will be accepted.

- Total available points for this homework is 80.

Question 1 (40 points): Education

A school counselor was curious about the average of IQ of the students in her school and took a random sample of 25 students' IQ scores. The following is the data set:

```
y <- c(105, 69, 86, 100, 82, 111, 104, 110, 87, 108, 87, 90, 94, 113, 112, 98,
    80, 97, 95, 111, 114, 89, 95, 126, 98)
```

1. Find a 90% confidence interval for the average student IQ in the school.

   Answer: The 90% confidence interval for the average student IQ in the school is (93.96 , 102.92)

2. Next, the school counselor was curious whether the average student IQ in her school is higher than the average IQ score (100) among all the schools in the country.

   Using the same sample, conduct the appropriate hypothesis test with $\alpha = 0.05$.

   Answer:

   Null hypothesis: Average student IQ in the school is not higher than the average IQ score (100) among all schools in the country

   Alternative hypothesis: Average student IQ in the school is higher than the average IQ score (100) among all schools in the country

   t.test(y, mu = 100, alternative = "greater")

   One Sample t-test

   data: y t = -0.59574, df = 24, p-value = 0.7215 alternative hypothesis: true mean is greater than 100 95 percent confidence interval: 93.95993 Inf sample estimates: mean of x 98.44

   Fail to reject the null hypothesis

Question 2 (40 points): Political Economy

Researchers are curious about what affects the amount of money communities spend on addressing homelessness. The following variables constitute our data set about social welfare expenditures in the USA.

State	*50 states in US*
Y	*per capita expenditure on shelters/housing assistance in state*
X1	*per capita personal income in state*
X2	*Number of residents per 100,000 that are "financially insecure" in state*
X3	*Number of people per thousand residing in urban areas in state*
Region	*1=Northeast, 2= North Central, 3= South, 4=West*

Explore the `expenditure` data set and import data into `R`.

- Please plot the relationships among *Y, X1, X2,* and *X3*? What are the correlations among them (you just need to describe the graph and the relationships among them)?

  [,1] [,2] [,3] [,4] [,5] [1,] NA 0.5317212 0.4482876 0.4636787 0.06563026 [2,] 0.53172116 NA 0.2056101 0.5952504 -0.21890587 [3,] 0.44828764 0.2056101 NA 0.2210149 0.33380548 [4,] 0.46367873 0.5952504 0.2210149 NA -0.23100105 [5,] 0.06563026 -0.2189059 0.3338055 -0.2310011 NA

-

- Please plot the relationship between *Y* and *Region*? On average, which region has the highest per capita expenditure on housing assistance?

  Answer: Region 4

- Please plot the relationship between *Y* and *X1*? Describe this graph and the relationship. Reproduce the above graph including one more variable *Region* and display different regions with different types of symbols and colors.

-

3

Figure 1: relationships among *Y*, *X1*, *X2*, and *X3*

Figure 2: relationship between *Y* and *Region*

Figure 3: relationship between *Y* and *X1*