

From the three-particle Green's function to the parquet equations, to ADC, and beyond

The Faddeev-decomposition of the three-particle vertex is used to derive a Hermitian effective Hamiltonian of Faddeev-ADC(3) type. Besides the Parquet approximation and the use of a Hartree–Fock Green's function, the derivation is based on two further approximations: couplings between different correlation channels as well as overlaps between different time-orderings are neglected. Faddeev-ADC(3) contains FLEX (first-order resolvent), ADC(2)-X (TDA limit) ADC(3) (TDA limit + first-order backward couplings) as limiting cases. Several self-energy schemes that are derived in Hedin's scheme or in the pp channel as well as PSD-completions (double counting of second-order terms aside) are recovered as a limiting case of Faddeev-ADC(3), and they are reviewed. Numerical results show that Faddeev-ADC(3), as well as all other methods that explicitly include triplet backward couplings to all orders come with very large errors. Switching off the triplets largely overcomes these issues, but only the performance of *GW* is recovered. Replacing the ring- and ladder-CCD amplitudes of Faddeev-ADC(3) with CCD ones, one recovers essentially the performance of CCSD. Coupling correlation channels seems to be decisive for accurate charged excitations.

CONTENTS

I. Introduction	1	2. Pair problems and reducible vertices	18
II. Faddeev Approach to the Self-Energy	2	3. Pair propagators	18
A. Martin–Schwinger Equations	2	4. Three-particle propagators	19
B. Eliminating G_2	2	C. Riccati equations and decoupling conditions	19
C. Faddeev Decomposition	3	1. Comparison to Löwdin downfolding	20
D. Faddeev form of Eq. (13): the ADC–parquet link	3	D. The ADC hierarchy	20
1. The Faddeev Series	3	1. ADC(3) from the first-order Faddeev kernel	20
2. Truncation at fixed order: ADC	4	a. GF3 self-energy	21
3. Own-Channel and No-Overlap Approximations	4	b. From GF3 to ADC(3)	21
III. Parquet in effective-Hamiltonian form	5	I. INTRODUCTION	
A. Resummation of the no-overlap own-channel Faddeev series	5	Via the three-particle Green's function G_3 , two routes to the one-particle self-energy are connected: the parquet equations for the two-particle vertex and the algebraic-diagrammatic construction (ADC). Starting from the Martin–Schwinger hierarchy (Sec. II A), we express the correlation self-energy in terms of a 6-point response function R , which obeys a Dyson equation with a three-particle kernel (Sec. II B). Dropping its 3-body irreducible part leaves the Faddeev kernel, ^{1–3} in which two of the three lines scatter through a two-particle vertex while the third line is a spectator. In the Faddeev form, this vertex is the full parquet vertex F (Sec. II C). ⁴	
B. Pair and three-particle propagators	7	Truncated at fixed order in v , the Faddeev series gives GF n , and Schirmer's matching gives ADC(n) (App. D). The connection becomes useful once the ladders in the pairs are resummed beyond the Tamm–Dancoff ladders contained in ADC (Sec. III).	
1. Amplitude Form of Propagators	7	With the RPA ladders of the first-order Faddeev kernel in every pair, and without overlaps between the ladders of different pairs, a Hermitian effective Hamiltonian of the size of ADC(3) is obtained, that describes the coupling of a single particle to pp and eh phonons. This is essentially the Faddeev random-phase approximation (FRPA) of Barbieri and co-workers. ^{5–8} In FRPA, the couplings are those of ADC(3), $U^{(1)} + U^{(2)}$, added by hand; ⁸ here they follow from the backward amplitudes of the phonons, the ring-CCD (eh) and ladder-CCD (pp) amplitudes of the channels. ⁹ The hierarchy is $\text{ADC}(3) \subset \text{FRPA} \subset \text{parquet}$ for the dynamic self-energy. At fourth order, the crossed channels, the overlaps and the line dressing are added in the 3p2h space.	
C. External vertices and effective Hamiltonian	8		
D. Phonon form	9		
IV. Relations to various schemes to construct electronic self-energies	9		
A. Tamm–Dancoff limit and ADC(3)	9		
B. Connections to (vertex-corrected) <i>GW</i>	9		
1. Double counting	10		
2. <i>GW</i>	10		
3. Σ^{TDHF}	10		
4. <i>G3W2</i> -type vertex corrections	11		
5. PSD-version of <i>G3W2</i> -type self-energies	11		
C. The pp channel	13		
D. FLEX and other combinations of eh and pp channels	13		
V. Numerical results	14		
VI. Conclusions	16		
A. Parquet vertices	17		
B. Pair and three-particle propagators	18		
1. Notation	18		

Three approximations are made by construction. The parquet approximation, $\Lambda = \Lambda^{(1)}$, keeps F exact through third order, since $\Lambda = \Lambda^{(1)} + O(v^4)$, Eq. (A4). With a static HF reference in all lines of the vertex, every diagram of the vertex is a product of instantaneous interactions and G_{HF} lines, whose frequency integrals turn into resolvents of configuration spaces. This makes the effective-Hamiltonian form possible. Finally, the Faddeev approximation affects the self-energy from fifth order on. The orders through which the resulting self-energies are exact are collected in Table I; the constructions derived in these notes are exact through fourth order at most.

TABLE I. Order through which the self-energy is exact, and the highest ADC(n) contained, for different approximations. Λ : fully irreducible two-particle vertex; $\Xi_{3\text{B}}$: 3-body irreducible kernel; SDE: Schwinger–Dyson equation. The second row is the construction of these notes. The first three rows use the parquet approximation, $\Lambda = \Lambda^{(1)}$, Eq. (A4).

Approximation	Eqs.	exact through	contains
Parquet, SDE	(8), (A7)	4th	ADC(4)
Parquet F , Faddeev	(18), (20)	4th	ADC(4)
Parquet $F + \Xi_{3\text{B}}^{(3)}$	(15), (13)	5th	ADC(5)
$\Lambda^{(\leq 4)}$, SDE	(8), (A7)	5th	ADC(5)
$\Lambda^{(\leq 4)} + \Xi_{3\text{B}}^{(\leq 4)}$	(15), (13)	6th	ADC(6)

II. FADDEEV APPROACH TO THE SELF-ENERGY

A. Martin–Schwinger Equations

Let us define the n -particle Green’s function as

$$G_n(1 \dots n; 1' \dots n') = (-i)^n \langle \mathcal{T}_C \hat{\psi}(1) \dots \hat{\psi}(n) \times \hat{\psi}^\dagger(n') \dots \hat{\psi}^\dagger(1') \rangle, \quad (1)$$

with $\hat{\psi}$ in the Heisenberg picture. G_n might be expressed in terms of G_{n-1} and G_{n+1} through the Martin–Schwinger hierarchy,¹⁰

$$\begin{aligned} & [i\partial_{t_1} - h_0(1)] G_n(1 \dots n; 1' \dots n') \\ &= \sum_{j=1}^n (-1)^{j-1} \delta(1j') G_{n-1}(2 \dots n; 1' \dots \widehat{j'} \dots n') \\ &+ (-1)^n i v(1\bar{2}; \bar{3}\bar{4}) G_{n+1}(\bar{3}\bar{4} 2 \dots n; 1' \dots n' \bar{2}^+) . \end{aligned} \quad (2)$$

We define

$$G_0^{-1}(11') = \left[i \frac{\partial}{\partial t_1} - h_0(1) \right] \delta(11'), \quad (3)$$

and its inverse via

$$G_0^{-1}(1\bar{1}) G_0(\bar{1}1') = \delta(11'). \quad (4)$$

Using both relations, G_1 through G_3 become

$$G(11') = G_0(11') - i G_0(1\bar{3}) v(\bar{3}\bar{4}; \bar{5}\bar{6}) G_2(\bar{5}\bar{6}; 1'\bar{4}^+), \quad (5)$$

$$\begin{aligned} G_2(12; 1'2') &= G_0(11') G(22') - G_0(12') G(21') \\ &+ i G_0(1\bar{3}) v(\bar{3}\bar{4}; \bar{5}\bar{6}) \\ &\times G_3(2\bar{5}\bar{6}; 1'2'\bar{4}^+), \end{aligned} \quad (6)$$

$$\begin{aligned} G_3(123; 1'2'3') &= G_0(11') G_2(23; 2'3') \\ &- G_0(12') G_2(23; 1'3') \\ &+ G_0(13') G_2(23; 1'2') \\ &- i G_0(1\bar{4}) v(\bar{4}\bar{5}; \bar{6}\bar{7}) \\ &\times G_4(2\bar{3}\bar{6}\bar{7}; 1'2'3'\bar{5}^+), \end{aligned} \quad (7)$$

Through Eq. (5), we can define the single-particle self-energy according to

$$\Sigma(11') = -iv(1\bar{2}; \bar{3}'\bar{2}') G_2(\bar{3}'\bar{2}'; \bar{3}\bar{2}^+) G^{-1}(\bar{3}1'). \quad (8)$$

B. Eliminating G_2

We can now use Eq. (6) to eliminate the dependence of Eq. (8) on G_2 in favor of G_3 . G_4 might be eliminated as well, and Eq. (8) can be expressed in terms of G_3 and G_5 , or, more generally, in terms of odd-order Green’s functions. Alternatively, combining Eq. (6) with Eq. (7), one can express Eq. (8) in terms of G_2 and G_4 , or, more generally, in terms of even-order Green’s functions.

Here, we take the odd-order route. To do so, following Winter² and Ethofer and Schuck³, we introduce the 6-point response function

$$\begin{aligned} R(123; 1'2'3') &= G_3(123; 1'2'3') \\ &- G_2(23; 3'\bar{4}) G^{-1}(\bar{4}\bar{4}') G_2(\bar{4}'1; 1'2'), \end{aligned} \quad (9)$$

so that the EOM for G_2 becomes

$$\begin{aligned} G_2(12; 1'2') &= G(11') G(22') - G(12') G(21') \\ &+ i G(1\bar{3}) v(\bar{3}\bar{4}; \bar{5}'\bar{6}') R(2\bar{5}'\bar{6}'; 1'2'\bar{4}^+). \end{aligned} \quad (10)$$

Plugging Eq. (10) into Eq. (8) and using Eq. (4) gives

$$\Sigma(11') = \Sigma_{\text{HF}}(11') + \Sigma^c(11') \quad (11)$$

with

$$\Sigma_{\text{HF}}(11') = -i [v(1\bar{2}; 1'\bar{2}') - v(1\bar{2}; \bar{2}'1')] G(\bar{2}'\bar{2}^+), \quad (12)$$

$$\begin{aligned} \Sigma^c(11') &= v(1\bar{2}; \bar{3}'\bar{2}') G(\bar{3}'\bar{4}) v(\bar{4}\bar{5}; \bar{6}'\bar{7}') \\ &\times R(\bar{2}'\bar{6}'\bar{7}'; \bar{3}\bar{2}\bar{5}^+) G^{-1}(\bar{3}1'). \end{aligned} \quad (13)$$

The response function R obeys a Dyson equation with the 2p1h-irreducible three-particle kernel Ξ , whose free propagator is the 2p1h propagator of three independent (dressed) lines,

$$\mathcal{G}(312; 1'2'3') = G(33') [G(11') G(22') - G(12') G(21')] . \quad (14)$$

As in G_3 , all three lines run from the second to the first triple. From here on, the first triple of R is ordered as in Eq. (14), with the line 3 of $\mathcal{G}$ in the first slot. The second triple keeps the order (1'2'3'), so that line 3 runs from the last slot of the second triple to the first slot of the first.

C. Faddeev Decomposition

The kernel Ξ of the Dyson equation for R is split into the Faddeev (pairwise-reducible) part and the 3-body irreducible kernel Ξ_{3B} , which is dropped from now on. In the Faddeev part, one pair of lines scatters through a two-particle vertex while the third line propagates freely.^{1-3,5} Since the two lines of every pair run in the same direction, each pair scatters in its pp channel, with the pp-irreducible vertex Γ^{pp} (exact at first order, see App. A). As in Refs. 2,3, the Dyson equation for R then reads

$$R = \mathcal{G} + \mathcal{K}R, \quad \mathcal{K} = \mathcal{K}_{12}[\Gamma^{\text{pp}}] + \mathcal{K}_{31}[\Gamma^{\text{pp}}] + \mathcal{K}_{32}[\Gamma^{\text{pp}}], \quad (15)$$

where the pair operators act, for any 4-point Y and any χ antisymmetric in its first pair, as

$$\begin{aligned} [\mathcal{K}_{12}[Y]\chi](312; 1'2'3') &= -\frac{1}{4}[G(1\bar{1})G(2\bar{2}) - G(1\bar{2})G(2\bar{1})] \\ &\quad \times Y(\bar{1}\bar{2}; \bar{1}'\bar{2}')\chi(3\bar{1}\bar{2}'; 1'2'3'), \\ [\mathcal{K}_{31}[Y]\chi](312; 1'2'3') &= G(3\bar{3})G(1\bar{1})Y(\bar{3}\bar{1}; \bar{1}'\bar{3}') \\ &\quad \times \chi(\bar{3}\bar{1}\bar{2}'; 1'2'3'), \\ [\mathcal{K}_{32}[Y]\chi](312; 1'2'3') &= G(3\bar{3})G(2\bar{2})Y(\bar{3}\bar{2}; \bar{2}'\bar{3}') \\ &\quad \times \chi(\bar{3}\bar{1}\bar{2}'; 1'2'3'). \end{aligned} \quad (16)$$

Eq. (15) is shown diagrammatically in Fig. 1.

Cutting the two lines of pair α separates the outgoing from the incoming legs of Y in $\mathcal{K}_\alpha[Y]$, *i.e.* all three pairs scatter in the pp channel of R . In Σ^c , Eq. (13), line 3 becomes the hole for $t_{1'} < t_1$ (block II) or the particle for $t_1 < t_{1'}$ (block I). The pair (12) then gives the ladder-type self-energy diagrams of the pp channel, and the pairs (31) and (32) give the ring-type diagrams of the eh and $\bar{\text{eh}}$ channels. From here on, we label the pairs by these channels, $\alpha \in \{\text{pp}, \text{eh}, \bar{\text{eh}}\}$ for (12), (31) and (32).

Eq. (15) is exact at first order, where all vertices are $\Gamma^{(1)}$. Beyond first order, iterating $\mathcal{K}_{\text{eh}}[\Gamma^{\text{pp}}]$ or $\mathcal{K}_{\bar{\text{eh}}}[\Gamma^{\text{pp}}]$ alone would count the rungs of the ladders in these pairs twice, since $\mathcal{G}$ is not antisymmetrized in the lines 3 and 1 (3 and 2). The pp channel of R in these pairs is the eh channel of F , whose ladder Φ^{eh} is already contained in Γ^{pp} .⁸ We therefore replace Γ^{pp} in $\mathcal{K}_\alpha$ by a pair vertex Γ_α , fixed such that the pair α alone reproduces its full two-body scattering F . For the pp pair, $\Gamma_{\text{pp}} = \Gamma^{\text{pp}}$, and at first order $\Gamma_\alpha = \Gamma^{(1)}$ for all pairs. Their explicit form is not needed, since only F appears in the Faddeev form derived next.

D. Faddeev form of Eq. (13): the ADC-parquet link

1. The Faddeev Series

Eq. (15) contains the pair-irreducible vertex Γ^{pp} in all three pairs. The aim of this subsection is to rewrite R in Eq. (13) such that only the full vertex F appears in all three pairs. This is the Faddeev form of the three-body problem.¹

Based on the Faddeev construction, we will then show that GF n is the truncation of the Faddeev series, at total order $n - 2$ in v . The ADC(n) matrices then follow from Schirmer's matching,¹¹ in which the resolvent form of ADC(n) is imposed and matched to the self-energy order by order. ADC(3) is worked out as an illustrative case in App. D.

With the pair vertices Γ_α in place of Γ^{pp} , Eq. (15) reads

$$R = \mathcal{G} + \sum_{\alpha} \mathcal{K}_\alpha[\Gamma_\alpha]R, \quad (17)$$

where $\mathcal{K}_\alpha[\Gamma_\alpha]$ acts only on the pair α , with the third line as a free spectator [Fig. 1(b), with the thick bars standing for Γ_α]. We collect the terms of R in which the last interaction, *i.e.* the one closest to the outgoing triple, acts on the pair α into $R_\alpha \equiv \mathcal{K}_\alpha[\Gamma_\alpha]R$. Solving $R_\alpha = \mathcal{K}_\alpha[\Gamma_\alpha](\mathcal{G} + R_\alpha + \sum_{\beta \neq \alpha} R_\beta)$ for R_α gives

$$\begin{aligned} R &= \mathcal{G} + R_{\text{pp}} + R_{\text{eh}} + R_{\bar{\text{eh}}}, \\ R_\alpha &= (\mathbf{1} - \mathcal{K}_\alpha[\Gamma_\alpha])^{-1} \mathcal{K}_\alpha[\Gamma_\alpha] \left(\mathcal{G} + \sum_{\beta \neq \alpha} R_\beta \right), \end{aligned} \quad (18)$$

where $(\mathbf{1} - \mathcal{K}_\alpha[\Gamma_\alpha])^{-1} \mathcal{K}_\alpha[\Gamma_\alpha]$ sums all consecutive interactions in the pair α .

Acting on $\mathcal{G}$, it gives all diagrams of R in which only the pair α interacts, while the third line remains a free spectator. By the definition of Γ_α , this is the full two-body scattering of the pair, $\mathcal{K}_\alpha[F] = \mathcal{K}_\alpha[\Gamma_\alpha] + \mathcal{K}_\alpha[\Gamma_\alpha]\mathcal{K}_\alpha[F]$, so that

$$(\mathbf{1} - \mathcal{K}_\alpha[\Gamma_\alpha])^{-1} \mathcal{K}_\alpha[\Gamma_\alpha] = \mathcal{K}_\alpha[F]. \quad (19)$$

Inserting Eq. (19) into Eq. (18) gives the Faddeev equations^{5,8}

$$R_\alpha = \mathcal{K}_\alpha[F] \left(\mathcal{G} + \sum_{\beta \neq \alpha} R_\beta \right). \quad (20)$$

Iterating Eq. (20), *i.e.* inserting the R_β on the right-hand side again and again, gives the Faddeev series^{1,5,12}

$$\begin{aligned} R &= \mathcal{G} + \sum_{\alpha} \mathcal{K}_\alpha[F]\mathcal{G} + \sum_{\alpha \neq \beta} \mathcal{K}_\alpha[F]\mathcal{K}_\beta[F]\mathcal{G} \\ &\quad + \sum_{\substack{\alpha \neq \beta \\ \beta \neq \gamma}} \mathcal{K}_\alpha[F]\mathcal{K}_\beta[F]\mathcal{K}_\gamma[F]\mathcal{G} + \dots, \end{aligned} \quad (21)$$

in which consecutive pairs are always different (Fig. 2). Eq. (21) is the starting point for two approximations. Truncated at fixed order in v , it gives GF n and, with Schirmer's matching, ADC(n). Keeping only the own-channel ladders without overlaps, but to all orders, gives Eq. (26), from which an effective Hamiltonian is derived in Sec. III.




FIG. 1. (a) The Faddeev kernel, Eqs. (15) and (16). Lines run from the second triple on the right to the first triple on the left. The thick bar is the two-particle vertex Γ^{pp} : it connects the two lines of one pair, while the third line (gray) is a spectator that passes the vertex without interacting. (b) The Dyson equation for R , Eq. (15), sorted by the interacting pair. The exchange of the lines 1 and 2 in $\mathcal{G}$ is implied.




FIG. 2. Regrouping of the iterated Eq. (17) into the Faddeev series. Left: one term of the iteration, a sequence of pair vertices $\mathcal{K}_\alpha[\Gamma_\alpha]$ (thick bars). Right: each sequence of consecutive vertices in the same pair is part of the full two-body scattering $\mathcal{K}_\alpha[F]$ of that pair (shaded box), so that consecutive boxes always belong to different pairs.

2. Truncation at fixed order: ADC

To relate Eq. (21) to ADC, we expand F in powers of v . By Eqs. (A3) and (A4), Λ has no second-order term, so that

$$\begin{aligned} F^{(1)} &= \Gamma^{(1)}, \\ F^{(2)} &= \Phi^{\text{pp},(2)} + \Phi^{\text{eh},(2)} + \Phi^{\bar{\text{eh}},(2)}. \end{aligned} \quad (22)$$

The self-energy, Eq. (13), contains two explicit interactions v . A term of order m in v of R therefore contributes to Σ at order $m+2$. ADC(n) is complete through order n , so that R is needed through order $n-2$ in v . Collecting the terms of Eq. (21) up to this order,

$$\begin{aligned} R_{\text{ADC}(2)} &= \mathcal{G}, \\ R_{\text{ADC}(3)} &= \mathcal{G} + \sum_{\alpha} \mathcal{K}_{\alpha}[F^{(1)}] \mathcal{G}_{\text{HF}}, \\ R_{\text{ADC}(4)} &= \mathcal{G} + \sum_{\alpha} \mathcal{K}_{\alpha}[F^{(1)} + F^{(2)}] \mathcal{G}_{\text{HF}} \\ &\quad + \sum_{\alpha \neq \beta} \mathcal{K}_{\alpha}[F^{(1)}] \mathcal{K}_{\beta}[F^{(1)}] \mathcal{G}_{\text{HF}}. \end{aligned} \quad (23)$$

Here, $\mathcal{G}$ in the first term, $G(\bar{3}'\bar{4})$ and the static part Σ_{HF} are evaluated with G expanded to the corresponding order, while in all terms with at least one F , all lines can be taken at the HF level; $\mathcal{G}_{\text{HF}}$ denotes Eq. (14) with G_{HF} lines.

3. Own-Channel and No-Overlap Approximations

We now introduce the own-channel approximation to Eq. (21). For this, we approximate the BSEs Eq. (A7) in the own-channel approximation, for the special case of the first-order Faddeev kernel, so that $\Gamma^{\lambda} \rightarrow \Gamma^{(1)}$. The two BSEs then define two separate vertices

$$\begin{aligned} F_{\text{oc}}^{\text{pp}} &= \Gamma^{(1)} - \frac{1}{2} \Gamma^{(1)} \circ_{\text{pp}} F_{\text{oc}}^{\text{pp}}, \\ F_{\text{oc}}^{\text{eh}} &= \Gamma^{(1)} + \Gamma^{(1)} \circ_{\text{eh}} F_{\text{oc}}^{\text{eh}}. \end{aligned} \quad (24)$$

This is also done in the FLEX approximation.^{13,14}

In the own-channel approximation to Eq. (21), each pair scatters only through the ladder of its own channel with the bare kernel. The own-channel approximation amounts to neglecting the channel crossings of parquet (and of coupled cluster) inside a pair.⁵ Barbieri and Dickhoff⁵ combined this approximation additionally with a no-overlap approximation that we denote here by $[\cdots]_{2\text{p1h}}$. It neglects the time orderings in which the ladders of different pairs overlap in time, *i.e.* consecutive pairs are separated by a 2p1h cut, a subinterval spanned only by the three lines of a 2p1h state (Sec. III A). Due to this restriction, at first-order Faddeev the effective Hamiltonian only contains 2p1h (2h1p) configurations. At fourth order, the neglected terms are added perturbatively together with the 3p2h (3h2p) space. Each pair scatters with the own-channel vertex of its channel,

$$F_{\text{pp}} = F_{\text{oc}}^{\text{pp}}, \quad F_{\text{eh}} = F_{\text{eh}}^{\text{eh}} = F_{\text{oc}}^{\text{eh}}. \quad (25)$$

With these approximations, Eq. (21) becomes

$$R_{\text{oc}} = \mathcal{G} + \sum_{\alpha} \mathcal{K}_{\alpha}[F_{\alpha}] \mathcal{G} + \sum_{\alpha \neq \beta} [\mathcal{K}_{\alpha}[F_{\alpha}] \mathcal{K}_{\beta}[F_{\beta}] \mathcal{G}]_{2\text{p1h}} + \sum_{\substack{\alpha \neq \beta \\ \beta \neq \gamma}} [\mathcal{K}_{\alpha}[F_{\alpha}] \mathcal{K}_{\beta}[F_{\beta}] \mathcal{K}_{\gamma}[F_{\gamma}] \mathcal{G}]_{2\text{p1h}} + \dots \quad (26)$$

Expanded in v , Eq. (26) reproduces Eq. (23) through $R_{\text{ADC}(3)}$. At the next order, it misses the crossed-channel parts of $F^{(2)}$ in each pair, Eq. (22), and the overlapping orderings of $\mathcal{K}_{\alpha}[F^{(1)}] \mathcal{K}_{\beta}[F^{(1)}] \mathcal{G}$.

III. PARQUET IN EFFECTIVE-HAMILTONIAN FORM

In this section, we insert the own-channel and no-overlap approximation, Eq. (26), into Eq. (13),

$$\Sigma^c(11') = v(1\bar{2}; \bar{3}'\bar{2}') G(\bar{3}'\bar{4}) v(\bar{4}\bar{5}; \bar{6}'\bar{7}') \times R_{\text{oc}}(\bar{2}'\bar{6}'\bar{7}'; \bar{3}\bar{2}\bar{5}^+) G^{-1}(\bar{3}1') \quad (27)$$

We then show that, with G_{HF} lines, the self-energy becomes

$$\Sigma(\omega) = \Sigma(\infty) + \mathcal{U}^{\text{II}}(\omega \mathcal{N}^{\text{II}} - \mathcal{H}^{\text{II}})^{-1} \mathcal{U}^{\text{II}\dagger} + \mathcal{U}^{\text{I}}(\omega \mathcal{N}^{\text{I}} + \mathcal{H}^{\text{I}})^{-1} \mathcal{U}^{\text{I}\dagger}, \quad (28)$$

with Hermitian $\mathcal{H}$ and $\mathcal{N}$, and the static self-energy $\Sigma(\infty) = \Sigma_{\text{HF}}[G]$, Eq. (12). At the level of the first-order Faddeev kernel, this is essentially the Faddeev random-phase approximation (FRPA) of Barbieri and co-workers.⁵⁻⁸ The difference is that the effective Hamiltonian derived here remains Hermitian as long as its metric is positive definite, whereas the Faddeev eigenvalue problem of FRPA is non-Hermitian and contains spurious solutions.⁸

A. Resummation of the no-overlap own-channel Faddeev series

Our first goal is to evaluate the terms of Eq. (26) up to first order in $\mathcal{K}_{\alpha}$. It is convenient to rewrite these contributions as

$$\mathcal{G} + \sum_{\alpha} \mathcal{K}_{\alpha}[F_{\alpha}] \mathcal{G} = \sum_{\alpha} (\mathcal{G} + \mathcal{K}_{\alpha}[F_{\alpha}] \mathcal{G}) - 2\mathcal{G}, \quad (29)$$

which can then be evaluated channel by channel. The terms inside the sum over α are interacting single-channel three-particle propagators, for which we introduce the shorthand Q_{α} ,

$$Q_{\alpha} = \mathcal{G} + \mathcal{K}_{\alpha}[F_{\alpha}] \mathcal{G}, \quad (30)$$

so that

$$\mathcal{G} + \sum_{\alpha} \mathcal{K}_{\alpha}[F_{\alpha}] \mathcal{G} = \sum_{\alpha} Q_{\alpha} - 2\mathcal{G}. \quad (31)$$

Truncating the Faddeev series, Eq. (26), after these terms is the one-shot FLEX approximation^{13,14} (Sec. IV D). We also

require that these propagators obey a three-particle Dyson-type equation,

$$Q_{\alpha} = \mathcal{G} + \mathcal{G} C_{\alpha} Q_{\alpha}, \quad (32)$$

or

$$Q_{\alpha}^{-1} = \mathcal{G}^{-1} - C_{\alpha}(\omega).$$

This serves as a definition for C_{α} , as soon as we have specified the explicit form of Q_{α} .

Once we have these propagators at hand, we must evaluate the remaining terms in Eq. (27), i.e. the terms of the form

$$\mathcal{K}_{\alpha}[F_{\alpha}] \mathcal{K}_{\beta}[F_{\beta}] \mathcal{G}, \quad \alpha \neq \beta, \quad (33)$$

and so on. We can rewrite these terms using the identity and Eq. (30)

$$\begin{aligned} \mathcal{K}_{\alpha}[F_{\alpha}] \mathcal{K}_{\beta}[F_{\beta}] \mathcal{G} &= \mathcal{K}_{\alpha}[F_{\alpha}] \mathcal{G} \mathcal{G}^{-1} \mathcal{K}_{\beta}[F_{\beta}] \mathcal{G} \\ &= (Q_{\alpha} - \mathcal{G}) \mathcal{G}^{-1} (Q_{\beta} - \mathcal{G}), \end{aligned} \quad (34)$$

and therefore, Eq. (26) becomes

$$\begin{aligned} R_{\text{oc}}(\omega) &= \mathcal{G} + \sum_{\alpha} (Q_{\alpha} - \mathcal{G}) + \sum_{\alpha \neq \beta} (Q_{\alpha} - \mathcal{G}) \mathcal{G}^{-1} (Q_{\beta} - \mathcal{G}) \\ &\quad + \sum_{\substack{\alpha \neq \beta \\ \beta \neq \gamma}} (Q_{\alpha} - \mathcal{G}) \mathcal{G}^{-1} (Q_{\beta} - \mathcal{G}) \mathcal{G}^{-1} (Q_{\gamma} - \mathcal{G}) + \dots \end{aligned} \quad (35)$$

Notice that this step strictly requires the no-overlap approximation, otherwise we could not just insert $\mathcal{G} \mathcal{G}^{-1}$ in each term.

To resum the series, we collect all terms whose leftmost interacting pair is α into a Faddeev component R_{α} . Thus,

$$R_{\text{oc}} = \mathcal{G} + \sum_{\alpha} R_{\alpha}. \quad (36)$$

The constraint that two consecutive factors $Q_{\alpha} - \mathcal{G}$ must belong to different pairs implies

$$\begin{aligned} R_{\alpha} &= (Q_{\alpha} - \mathcal{G}) \mathcal{G}^{-1} \left(\mathcal{G} + \sum_{\beta \neq \alpha} R_{\beta} \right) \\ &= (Q_{\alpha} - \mathcal{G}) \mathcal{G}^{-1} (R_{\text{oc}} - R_{\alpha}). \end{aligned} \quad (37)$$

By the Dyson equation Eq. (33) and after some algebra, we obtain

$$(1 + Q_{\alpha} C_{\alpha}) R_{\alpha} = Q_{\alpha} C_{\alpha} R_{\text{oc}}. \quad (38)$$

Since

$$1 + Q_{\alpha} C_{\alpha} = Q_{\alpha} (Q_{\alpha}^{-1} + C_{\alpha}) = Q_{\alpha} \mathcal{G}^{-1}, \quad (39)$$

it follows that

$$R_{\alpha} = \mathcal{G} C_{\alpha} R_{\text{oc}}. \quad (40)$$

Summing Eq. (40) over the three pair channels and using Eq. (36) gives

$$R_{\text{oc}} = \mathcal{G} + \mathcal{G} (C_{\text{pp}} + C_{\text{ch}} + C_{\text{eh}}) R_{\text{oc}}. \quad (41)$$

Hence,

$$R_{\text{oc}}(\omega) = (\mathcal{G}^{-1} - C_{\text{pp}} - C_{\text{eh}} - C_{\text{eh}}^{-})^{-1}, \quad (42)$$

or, equivalently,

$$C_{\alpha} = \mathcal{G}^{-1} - Q_{\alpha}^{-1}, \quad (43)$$

$$R_{\text{oc}}^{-1} = Q_{\text{pp}}^{-1} + Q_{\text{eh}}^{-1} + Q_{\text{eh}}^{-1} - 2\mathcal{G}^{-1}. \quad (44)$$

Eq. (44) holds in the space $\mathcal{P}$ of the 2p1h and 2h1p configurations, where $\mathcal{G}$ lives. To insert it into Eq. (27), we still need to know how the two bare interactions of Eq. (27), at t_1 and $t_{1'}$ (the time arguments of the self-energy), connect the orbitals p and q of Σ_{pq}^c to the three-line configurations of R_{oc} .

In the orbital basis, each line of Eq. (27) is a particle or a hole, depending on its time direction. For $R = \mathcal{G}$, all lines run between $t_{1'}$ and t_1 , and only the configurations in $\mathcal{P}$ contribute. In this case, the free propagation between two equal-time three-line states is

$$\mathcal{G}(\omega) = (\omega - K)^{-1}, \quad (45)$$

with $\omega + i0^+$ in the 2p1h and $\omega - i0^+$ in the 2h1p block, and

$$K = K^{\text{II}} \oplus (-K^{\text{I}}),$$

$$K_{abi,cdj}^{\text{II}} = \Delta_{abi} \delta_{ac} \delta_{bd} \delta_{ij},$$

$$K_{ija,klb}^{\text{I}} = -\Delta_{ija} \delta_{ik} \delta_{jl} \delta_{ab},$$

with the shorthand

$$\Delta_{xyz} = \epsilon_x + \epsilon_y - \epsilon_z. \quad (46)$$

The first-order couplings of the one-particle states to the three-line states are

$$U^{(1)} = (U^{\text{II},(1)}, U^{\text{I},(1)}), \quad (47)$$

with

$$\begin{aligned} U_{p,abi}^{\text{II},(1)} &= \langle pi||ab \rangle, \\ U_{p,ija}^{\text{I},(1)} &= \langle pa||ij \rangle. \end{aligned} \quad (48)$$

In these expressions, $a < b$ and $i < j$.

These are not all configurations that the two bare interactions of Eq. (27) can reach. If the last interaction of a pair α happens after t_1 , or its first one before $t_{1'}$, the two lines of the pair arrive at the vertex in reversed time direction. For the pp pair, for instance, the vertex at t_1 then meets the three holes ($kl; i$) instead of ($ab; i$). Fig. 3 shows the lowest-order example. In (a), the pp rung lies between the two vertices, and all intermediate states are 2p1h configurations. In (b), the rung lies after t_1 . At t_1 , the bare interaction $\langle pi||kl \rangle$ removes the electrons k and l , i.e. it creates the holes k and l , which end at the later rung together with particles a and b . The pair attached at t_1 is thus the hole pair (kl), and the rung enters with the static denominator $\epsilon_k + \epsilon_l - \epsilon_a - \epsilon_b$. This is the coupling $U^{(2)}$ of ADC(3).




FIG. 3. Time-ordered third-order self-energy diagrams with one pp rung (block II, time upward, dashed lines: bare interactions, external legs drawn horizontally, so that each horizontal cut between two interactions crosses only the lines of one intermediate state). (a) Rung at $t_{1'} < \tau < t_1$: the pair propagates in $\mathcal{P}$. (b) Rung at $\tau > t_1$: at t_1 , the pair slot holds the holes k and l , the reversed configuration ($kl; i$) in $\bar{\mathcal{P}}_{\text{pp}}$, and the rung gives the first-order amplitude t_{kl}^{ab} of $U^{(2)}$.

The diagram in Fig. 3(b) and its mirror image, with the rung before $t_{1'}$, are the GF3 terms of Eqs. (D21) and (D22); the eh analogues are Eqs. (D27) and (D28). They arise from the time orderings of the three-particle propagator in which the pair lines run outside $[t_{1'}, t_1]$. When the pair interactions are resummed, reversed configurations appear in two places:

- At the two outer vertices, when the ladder or ring of a pair continues beyond t_1 or before $t_{1'}$. These configurations are folded into the couplings, $\mathcal{U} = U^{(1)} + \sum_{\alpha} \bar{U}_{\alpha}^{(1)} T_{\alpha}$, and enter Σ from third order on.
- Inside $[t_{1'}, t_1]$, within one pair, when the pair turns backward in time and forward again. These orderings give the terms $B_{\alpha} T_{\alpha}$ in $\mathcal{H}$ and the metric $\mathcal{N} = \mathbf{1} - \sum_{\alpha} T_{\alpha}^{\dagger} T_{\alpha}$, and enter Σ from fourth order on.

Between two different pairs, reversed configurations do not occur, since the no-overlap approximation separates consecutive pairs by a 2p1h cut, the factor $\mathcal{G}^{-1}$ in Eq. (35).

We call $\mathcal{P}$ the 2p1h and 2h1p configurations of Eq. (45), and $\bar{\mathcal{P}}_{\alpha}$ the reversed configurations of the pair α . Then:

- $\mathcal{G}$ lives on $\mathcal{P}$ only.
- Q_{α} lives on $\mathcal{P} \oplus \bar{\mathcal{P}}_{\alpha}$. We keep the symbol Q_{α} for its block on $\mathcal{P}$ and write $[Q_{\alpha}]_{\bar{\mathcal{P}}_{\alpha}, \mathcal{P}}$ for its block from $\mathcal{P}$ to $\bar{\mathcal{P}}_{\alpha}$.
- The bare interaction couples the orbital p to $\mathcal{P}$ through $U^{(1)}$, Eq. (47), and to $\bar{\mathcal{P}}_{\alpha}$ through antisymmetrized integrals $\bar{U}_{\alpha}^{(1)}$.

Eq. (27) is therefore a sum over the blocks of R ,

$$\Sigma^c(\omega) = \sum_{X,Y} V_X [R(\omega)]_{XY} V_Y^{\dagger}, \quad (49)$$

with $X, Y \in \{\mathcal{P}, \bar{\mathcal{P}}_{\text{pp}}, \bar{\mathcal{P}}_{\text{eh}}, \bar{\mathcal{P}}_{\text{eh}}^{-}\}$, $V_{\mathcal{P}} = U^{(1)}$ and $V_{\bar{\mathcal{P}}_{\alpha}} = \bar{U}_{\alpha}^{(1)}$. For $R = \mathcal{G}$, only $X = Y = \mathcal{P}$ contributes, and $\Sigma^c = U^{(1)} \mathcal{G} U^{(1)\dagger} = \Sigma^{(2)}$.

The block on $\mathcal{P}$ is R_{oc} , Eq. (44). The other blocks follow in three steps.

First, in Eq. (35), consecutive factors are joined by $\mathcal{G}^{-1}$, which lives on $\mathcal{P}$. Only the first factor of a term can reach $\tilde{\mathcal{P}}_\alpha$, and only if it is $\mathcal{Q}_\alpha - \mathcal{G}$. These terms make up R_α , Eq. (37), so that

$$[R]_{\tilde{\mathcal{P}}_\alpha \mathcal{P}} = [\mathcal{Q}_\alpha]_{\tilde{\mathcal{P}}_\alpha \mathcal{P}} \mathcal{G}^{-1} (R_{\text{oc}} - R_\alpha) . \quad (50)$$

Second, we eliminate R_α . Adding $R_{\text{oc}} - R_\alpha$ to both sides of Eq. (37) gives

$$R_{\text{oc}} = \mathcal{Q}_\alpha \mathcal{G}^{-1} (R_{\text{oc}} - R_\alpha) ,$$

i.e.

$$\mathcal{G}^{-1} (R_{\text{oc}} - R_\alpha) = \mathcal{Q}_\alpha^{-1} R_{\text{oc}} .$$

Inserted into Eq. (50),

$$[R]_{\tilde{\mathcal{P}}_\alpha \mathcal{P}} = [\mathcal{Q}_\alpha]_{\tilde{\mathcal{P}}_\alpha \mathcal{P}} \mathcal{Q}_\alpha^{-1} R_{\text{oc}} .$$

Third, the same argument for the last factor of a term gives the columns, and both together the corners,

$$\begin{aligned} [R]_{\mathcal{P} \tilde{\mathcal{P}}_\beta} &= R_{\text{oc}} \mathcal{Q}_\beta^{-1} [\mathcal{Q}_\beta]_{\mathcal{P} \tilde{\mathcal{P}}_\beta} , \\ [R]_{\tilde{\mathcal{P}}_\alpha \tilde{\mathcal{P}}_\beta} &= [\mathcal{Q}_\alpha]_{\tilde{\mathcal{P}}_\alpha \mathcal{P}} \mathcal{Q}_\alpha^{-1} R_{\text{oc}} \mathcal{Q}_\beta^{-1} [\mathcal{Q}_\beta]_{\mathcal{P} \tilde{\mathcal{P}}_\beta} . \end{aligned}$$

All blocks of R are thus R_{oc} , multiplied from the left by $[\mathcal{Q}_\alpha]_{\tilde{\mathcal{P}}_\alpha \mathcal{P}} \mathcal{Q}_\alpha^{-1}$ for a row in $\tilde{\mathcal{P}}_\alpha$ and from the right by its adjoint for a column in $\tilde{\mathcal{P}}_\beta$. Inserted into Eq. (49), the sum over the blocks factorizes,

$$\Sigma^c(\omega) = \mathcal{U} R_{\text{oc}}(\omega) \mathcal{U}^\dagger ,$$

with the couplings

$$\mathcal{U} = U^{(1)} + \sum_\alpha \bar{U}_\alpha^{(1)} [\mathcal{Q}_\alpha]_{\tilde{\mathcal{P}}_\alpha \mathcal{P}} \mathcal{Q}_\alpha^{-1} . \quad (51)$$

The reversed configurations are thus folded into the couplings, and R_{oc} is needed only on $\mathcal{P}$. It remains to specify the three-particle propagators in the pp and eh channels, which we do in the next section.

B. Pair and three-particle propagators

The Faddeev series correlates two particles at a time, while the third one is a spectator. The three-particle propagators $\mathcal{Q}_\alpha$ of Eq. (44) therefore follow from two-body pair problems and the free propagation of the spectator. The two-body problems are well known for the herein used first-order Faddeev kernel.¹⁵ The eh propagator is obtained by solving the TDHF equations Eq. (B6), and the eh propagator follows by crossing. Analogously, the pp propagator follows from the solution of the ppRPA, Eq. (B7). Coupled to two external orbitals, they give the two-body pair vertices $M_{pq,n}^{\text{eh}}$ and $M_{pq,n}^{\text{pp}}$ of the reducible vertices, Eqs. (B8) and (B9).

With one interacting pair, the three-particle propagator is the pair propagator times the free propagation of the spectator. For the pp pair with spectator r ,

$$\begin{aligned} Q_{\text{pp};xyr,uwr}(\omega) &= n_r \sum_{n \in N+2} \frac{Z_{xy}^n Z_{uw}^n}{\omega + \epsilon_r - \Omega_n^+ + i0^+} \\ &+ \bar{n}_r \sum_{m \in N-2} \frac{Z_{xy}^m Z_{uw}^m}{\omega + \epsilon_r - \Omega_m^- - i0^+} , \end{aligned} \quad (52)$$

and for the eh pair with spectator s ,

$$\begin{aligned} Q_{\text{eh};xrs,yus}(\omega) &= \bar{n}_s \sum_{\Omega_n > 0} \frac{Z_{xr}^n Z_{yu}^n}{\omega - \epsilon_s - \Omega_n + i0^+} \\ &+ n_s \sum_{\Omega_n > 0} \frac{\tilde{Z}_{xr}^n \tilde{Z}_{yu}^n}{\omega - \epsilon_s + \Omega_n - i0^+} . \end{aligned} \quad (53)$$

Here, Z_{xy}^n are the components of the two-body eigenvectors of Eq. (B5) on the pair indices (xy) , forward (X) or backward (Y). Q_{eh} follows from Q_{eh} with the lines 1 and 2 exchanged. For a particle spectator only the forward eh excitations Z_n survive, for a hole spectator only the backward ones, $\tilde{Z}_n$.

Both propagators are block diagonal in $2p1h \oplus 2h1p$. The occupation factors select the first terms of Eqs. (52) and (53) in block II, where the spectator of the pp pair is the hole i and the spectators of the eh and eh pairs are the particles b and a . The second term is selected in block I, where the spectator of the pp pair is the particle a and the spectators of the eh and eh pairs are the holes j and i . Between the configurations of $2p1h \oplus 2h1p$, Eqs. (52) and (53) read, in matrix notation,

$$Q_{\text{pp}}(\omega) = Z_{\text{pp}}^{\text{II}}(\omega + \epsilon_i - \Omega^+)^{-1} Z_{\text{pp}}^{\text{II}\dagger} \oplus Z_{\text{pp}}^{\text{I}}(\omega + \epsilon_a - \Omega^-)^{-1} Z_{\text{pp}}^{\text{I}\dagger} , \quad (54)$$

with the $N + 2$ solutions in block II and the $N - 2$ solutions in block I, and

$$\begin{aligned} Q_{\text{eh}}(\omega) &= Z_{\text{eh}}(\omega - \epsilon_b - \Omega)^{-1} Z_{\text{eh}}^\dagger \oplus Z_{\text{eh}}(\omega - \epsilon_j + \Omega)^{-1} Z_{\text{eh}}^\dagger , \\ Q_{\text{eh}}(\omega) &= Z_{\text{eh}}(\omega - \epsilon_a - \Omega)^{-1} Z_{\text{eh}}^\dagger \oplus Z_{\text{eh}}(\omega - \epsilon_i + \Omega)^{-1} Z_{\text{eh}}^\dagger . \end{aligned} \quad (55)$$

Here, Z_α contains the components of the pair eigenvectors on the configurations of the block, $(ab; i)$, $(ia; b)$ in block II and $(ij; a)$, $(ia; j)$ in block I. The components on the remaining configurations of the pair, $\tilde{Z}_\alpha$, enter only at the external vertices, Sec. III C.

1. Amplitude Form of Propagators

It is useful to replace the propagators above by a form that uses CC-amplitudes. This is possible as the RPA eigenvalue problem in each channel can equivalently be expressed as the Riccati equation (see Appendix C, and Refs. 9,16,17)

$$B^\dagger + DT + TA + TBT = 0 . \quad (56)$$

The amplitudes map the forward components on the backward components,

$$Y = TX . \quad (57)$$

Eq. (56) is the ring-CCD (eh) and ladder-CCD (pp) amplitude equation of the channel.⁹ The relations between GW , the RPA and (equation-of-motion) coupled cluster theory are analyzed in Refs. 16,18–20. Each channel introduces its own set of amplitudes, but they remain decoupled. Full CCD couples them through the crossed-ring and mosaic terms of the amplitude equation.¹⁷ These couplings play the role of the crossed channels that the own-channel approximation neglects.

One can then show the relations

$$X\Omega X^{-1} = A + BT. \quad (58)$$

and

$$XX^\dagger = (\mathbf{1} - T^\dagger T)^{-1}. \quad (59)$$

These relations are readily upgraded to the three-particle space in each channel,

$$\begin{aligned} \tilde{Z}_\alpha &= T_\alpha Z_\alpha, \\ Z_\alpha \Omega_\alpha Z_\alpha^{-1} &= A_\alpha + B_\alpha T_\alpha, \\ Z_\alpha Z_\alpha^\dagger &= (\mathbf{1} - T_\alpha^\dagger T_\alpha)^{-1}, \end{aligned} \quad (60)$$

with $\Omega_{pp} = \Omega^+$ and $\Omega_{eh} = \Omega_{eh}^- = \Omega$.

For block II (block I follows equivalently), Eqs. (54) and (55) are $Q_\alpha = Z_\alpha(\omega - s_\alpha - \Omega_\alpha)^{-1} Z_\alpha^\dagger$, with the spectator energies

$$s_{pp} = -\epsilon_i, \quad s_{eh} = \epsilon_b, \quad s_{eh}^- = \epsilon_a. \quad (61)$$

With Eq. (60), it can be written as

$$\begin{aligned} Q_\alpha(\omega) &= Z_\alpha(\omega - s_\alpha - \Omega_\alpha)^{-1} Z_\alpha^{-1} Z_\alpha Z_\alpha^\dagger \\ &= [Z_\alpha(\omega - s_\alpha - \Omega_\alpha)^{-1}]^{-1} Z_\alpha Z_\alpha^\dagger \\ &= (\omega - s_\alpha - Z_\alpha \Omega_\alpha Z_\alpha^{-1})^{-1} Z_\alpha Z_\alpha^\dagger \\ &= (\omega - s_\alpha - A_\alpha - B_\alpha T_\alpha)^{-1} (\mathbf{1} - T_\alpha^\dagger T_\alpha)^{-1}. \end{aligned} \quad (62)$$

Next, the spectator energy and the orbital energies in A_α add up to K , and the interaction part of A_α is the pair interaction C_α^{II} . For the pp pair in the configuration $(ab; i)$ and the eh pair in $(ia; b)$,

$$\begin{aligned} s_{pp} + A_{ab,cd}^{\text{pp}} &= (\epsilon_a + \epsilon_b - \epsilon_i) \delta_{ac} \delta_{bd} + \langle ab || cd \rangle, \\ s_{eh} + A_{ia,jc}^{\text{eh}} &= (\epsilon_a + \epsilon_b - \epsilon_i) \delta_{ac} \delta_{ij} + \langle a j || i c \rangle, \end{aligned}$$

i.e. $s_\alpha + A_\alpha = K + C_\alpha^{\text{II}}$, and

$$Q_\alpha(\omega) = (\omega - K - C_\alpha^{\text{II}} - B_\alpha T_\alpha)^{-1} (\mathbf{1} - T_\alpha^\dagger T_\alpha)^{-1}.$$

After a similar treatment for block I, and with

$$\begin{aligned} C_\alpha &= C_\alpha^{\text{II}} \oplus (-C_\alpha^{\text{I}}), \\ B_\alpha T_\alpha &= B_\alpha^{\text{II}} T_\alpha^{\text{II}} \oplus (-B_\alpha^{\text{I}} T_\alpha^{\text{I}}), \end{aligned} \quad (63)$$

both blocks combine into

$$Q_\alpha(\omega) = (\omega - K - C_\alpha - B_\alpha T_\alpha)^{-1} (\mathbf{1} - T_\alpha^\dagger T_\alpha)^{-1}. \quad (64)$$

In the Tamm–Dancoff limit, $B_\alpha = 0$ and $T_\alpha = 0$, and

$$Q_\alpha(\omega) = (\omega - K - C_\alpha)^{-1}, \quad (65)$$

where $K + C_\alpha$ is the Tamm–Dancoff Hamiltonian of the pair, shifted by the spectator energy.

C. External vertices and effective Hamiltonian

We are now in a position to state the central result of these derivations. Inserting the amplitude form of Q_α defined by Eq. (64) into Eq. (44), we obtain

$$\begin{aligned} R_{\text{oc}}^{-1}(\omega) &= \omega \mathcal{N} - \mathcal{H}, \\ \mathcal{H} &= K + C + \sum_\alpha B_\alpha T_\alpha \\ &\quad - \sum_\alpha T_\alpha^\dagger T_\alpha (K + C_\alpha + B_\alpha T_\alpha), \\ \mathcal{N} &= \mathbf{1} - \sum_\alpha T_\alpha^\dagger T_\alpha, \end{aligned} \quad (66)$$

in both blocks. $\mathcal{H}$ and the metric $\mathcal{N}$ are Hermitian and block diagonal,

$$\mathcal{N} = \mathcal{N}^{\text{II}} \oplus \mathcal{N}^{\text{I}}, \quad \mathcal{H} = \mathcal{H}^{\text{II}} \oplus (-\mathcal{H}^{\text{I}}).$$

For the couplings, Eq. (51), we need the blocks of the three-particle propagators between $\tilde{\mathcal{P}}_\alpha$ and $\mathcal{P}$. With all rows of the pair eigenvectors, Eqs. (52) and (53) have the block $\tilde{Z}_\alpha(\omega - s_\alpha - \Omega_\alpha)^{-1} Z_\alpha^\dagger$ between $\tilde{\mathcal{P}}_\alpha$ and $\mathcal{P}$. With $\tilde{Z}_\alpha = T_\alpha Z_\alpha$, Eq. (60),

$$[Q_\alpha]_{\tilde{\mathcal{P}}_\alpha \mathcal{P}} = T_\alpha Z_\alpha(\omega - s_\alpha - \Omega_\alpha)^{-1} Z_\alpha^\dagger = T_\alpha Q_\alpha,$$

so that Eq. (51) becomes $\mathcal{U} = U^{(1)} + \sum_\alpha \tilde{U}_\alpha^{(1)} T_\alpha$. With all indices given explicitly, the couplings become

$$\begin{aligned} \mathcal{U} &= (\mathcal{U}^{\text{II}}, \mathcal{U}^{\text{I}}) = U^{(1)} + \sum_\alpha \tilde{U}_\alpha^{(1)} T_\alpha, \\ \mathcal{U}_{p;abi}^{\text{II}} &= \langle pi || ab \rangle + \hat{P}_{ab}^- \sum_{cj} \langle pc || jb \rangle T_{jc,ia}^{\text{eh}} + \sum_{k<l} \langle pi || kl \rangle T_{kl,ab}^{\text{pp}}, \\ \mathcal{U}_{p;ija}^{\text{I}} &= \langle pa || ij \rangle - \hat{P}_{ij}^- \sum_{kb} \langle pk || bj \rangle T_{ib,ka}^{\text{eh}} + \sum_{b<c} \langle pa || bc \rangle T_{ij,bc}^{\text{pp}}. \end{aligned} \quad (67)$$

Here, $\tilde{U}_\alpha^{(1)}$ couples to the backward configurations of the pair α (all but the first terms). In contrast to ADC(3), the couplings contain the channel-restricted CCD amplitudes instead of the first-order doubles ones. With the substitutions

$$T_{jc,ia}^{\text{eh}} \rightarrow t_{ij}^{ac} \quad (68)$$

and

$$T_{kl,ab}^{\text{pp}} \rightarrow t_{kl}^{ab}, \quad (69)$$

these are the ADC(3) couplings, that describe the first-order coupling of the backward amplitudes to single-particle states.

Projected onto $a < b$ and $i < j$, and with

$$\omega \mathcal{N} - \mathcal{H} = \mathcal{N}^{1/2} (\omega - H_{\text{eff}}) \mathcal{N}^{1/2}$$

for positive definite $\mathcal{N}$, we obtain the self-energy

$$\begin{aligned} \Sigma^c(\omega) &= \mathcal{U} R_{\text{oc}}(\omega) \mathcal{U}^\dagger = \mathcal{U} (\omega \mathcal{N} - \mathcal{H})^{-1} \mathcal{U}^\dagger \\ &= (\mathcal{U} \mathcal{N}^{-1/2}) (\omega - H_{\text{eff}})^{-1} (\mathcal{U} \mathcal{N}^{-1/2})^\dagger, \\ H_{\text{eff}} &= \mathcal{N}^{-1/2} \mathcal{H} \mathcal{N}^{-1/2}, \end{aligned} \quad (70)$$

which, written out block by block, is Eq. (28). The quasiparticle energies are the eigenvalues of the corresponding Hermitian Dyson supermatrix

$$\begin{pmatrix} h_0 + \Sigma(\infty) & \mathcal{U} \mathcal{N}^{-1/2} \\ \mathcal{N}^{-1/2} \mathcal{U}^\dagger & H_{\text{eff}} \end{pmatrix},$$

with h_0 the one-body Hamiltonian of Eq. (2). In block II, the matrix elements of Eq. (66) are

$$\begin{aligned} \mathcal{H}_{abi,cdj} &= K_{abi,cdj}^{\text{II}} + C_{abi,cdj}^{\text{II}} + \delta_{ij} h_{ab,cd}^{\text{pp}} \\ &\quad + \hat{P}_{ab}^- \hat{P}_{cd}^- \delta_{bd} h_{ia,jc}^{\text{eh}}, \\ \mathcal{N}_{abi,cdj} &= \delta_{ac} \delta_{bd} \delta_{ij} - \delta_{ij} \sum_{k<l} T_{kl,ab}^{\text{pp}} T_{kl,cd}^{\text{pp}} \\ &\quad - \hat{P}_{ab}^- \hat{P}_{cd}^- \delta_{bd} \sum_{ke} T_{ke,ia}^{\text{eh}} T_{ke,jc}^{\text{eh}}, \end{aligned} \quad (71)$$

with

$$\begin{aligned} h_{ab,cd}^{\text{pp}} &= \sum_{k<l} \left[\langle ab||kl \rangle T_{kl,cd}^{\text{pp}} + T_{kl,ab}^{\text{pp}} \langle kl||cd \rangle \right] \\ &\quad + \sum_{k<l} \sum_{m<n} T_{kl,ab}^{\text{pp}} (D_{kl,mn}^{\text{pp}} + \epsilon_i \delta_{km} \delta_{ln}) T_{mn,cd}^{\text{pp}}, \\ h_{ia,jc}^{\text{eh}} &= \sum_{ke} \left[\langle ae||ik \rangle T_{ke,jc}^{\text{eh}} + T_{ke,ia}^{\text{eh}} \langle ce||jk \rangle \right] \\ &\quad + \sum_{ke} \sum_{lf} T_{ke,ia}^{\text{eh}} (A_{ke,lf}^{\text{eh}} - \epsilon_b \delta_{kl} \delta_{ef}) T_{lf,jc}^{\text{eh}}. \end{aligned} \quad (72)$$

D. Phonon form

It is instructive to write the effective Hamiltonian in terms of the pair excitations (phonons). We do this for block II; block I follows in the same way with the block-I parts of Eqs. (54) and (55). By Eqs. (44) and (66),

$$\begin{aligned} \omega \mathcal{N} - \mathcal{H} &= \sum_{\alpha} \mathcal{Q}_{\alpha}^{-1} - 2\mathcal{G}^{-1}, \\ \mathcal{Q}_{\alpha}^{-1} &= Z_{\alpha}^{-\dagger} (\omega - s_{\alpha} - \mathbf{\Omega}_{\alpha}) Z_{\alpha}^{-1}. \end{aligned} \quad (73)$$

Here, $\mathbf{\Omega}_{\alpha}$ is the diagonal matrix of the pair energies, $\mathbf{\Omega}^+$ for the pp pair and $\mathbf{\Omega}$ for the eh and eh pairs, and s_{α} is the spectator energy, Eq. (61). Its ω -linear part gives $\mathcal{N}$, its constant part $\mathcal{H}$. With Eqs. (55) and (54), as well as the normalization in Eq. (64), the quantities appearing in the self-energy defined by Eq. (70) are

$$\begin{aligned} \mathcal{N} &= \mathbf{1} + \sum_{\alpha} \left[(Z_{\alpha} Z_{\alpha}^{\dagger})^{-1} - \mathbf{1} \right], \\ \mathcal{H} &= K^{\text{II}} + C^{\text{II}} + \sum_{\alpha} \left[Z_{\alpha}^{-\dagger} (\mathbf{\Omega}_{\alpha} + s_{\alpha}) Z_{\alpha}^{-1} - K^{\text{II}} - C_{\alpha} \right], \\ \mathcal{U} &= U^{(1)} + \sum_{\alpha} \left[M_{\alpha} Z_{\alpha}^{-1} - U^{(1)} \right], \end{aligned} \quad (74)$$

with the three-body couplings of the pairs $M_{\alpha} = U^{(1)} Z_{\alpha} + \bar{U}_{\alpha}^{(1)} \bar{Z}_{\alpha}$. Their entries are the two-body vertices of Eqs. (B8) and (B9), with the spectator as second orbital index,

$$\begin{aligned} M_{\text{pp},p;ni} &= M_{pi,n}^{\text{pp}}, \\ M_{\text{eh},p;nb} &= -M_{bp,n}^{\text{eh}}. \end{aligned}$$

IV. RELATIONS TO VARIOUS SCHEMES TO CONSTRUCT ELECTRONIC SELF-ENERGIES

A. Tamm–Dancoff limit and ADC(3)

In the Tamm–Dancoff limit, $B_{\alpha} = 0$, Eq. (56) gives $T_{\alpha} = 0$, and the three-particle propagators reduce to Eq. (65). Eq. (44) then gives

$$R_{\text{oc}}(\omega) = (\omega - K - C)^{-1} = (\omega - K^{\text{II}} - C^{\text{II}})^{-1} \oplus (\omega + K^{\text{I}} + C^{\text{I}})^{-1}, \quad (75)$$

with $C = C_{\text{pp}} + C_{\text{eh}} + C_{\text{eh}}^{-}$, *i.e.* $\mathcal{N} = \mathbf{1}$ and $\mathcal{H} = K + C$. Since $T_{\alpha} = 0$, the couplings reduce to $\mathcal{U} = U^{(1)}$, and the self-energy is that of ADC(2)-X,²¹ also known as the 2ph-TDA,^{22–24} or multi-channel Dyson equation (MCDE).^{25–27}

ADC(3) is not the strict Tamm–Dancoff limit. Instead, the amplitudes are restricted to second order,

$$\mathcal{U} = U^{(1)} + U^{(2)} + O(v^3).$$

The TDA-limit [ADC(2)-X] and ADC(3) differ, in principle, only in the couplings. However, since ADC(3) is usually constructed to be third order complete, $\Sigma(\infty)$ is chosen to be the third-order static self-energy, while ADC(2)-X keeps it at the Hartree–Fock level. For molecules, ADC(2)-X is a very crude approximation anyway, with large errors for charged excitations, while offering no tangible numerical advantage over ADC(3). Changing the static part has a non-negligible effect, but its accuracy remains poor either way.²⁸

Recent work has used CC-amplitudes in ADC(3). The CCD-ADC(3) from Ref. 29 uses exactly Eq. (67), but solves the full CCD equations to couple all channel amplitudes. However, the effective Hamiltonian is kept at the TDA level. Their EN-ADC(3) is the linearized, diagonal version of the ladder-CCD couplings.

Superficially, this scheme seems related to recent work by Coveney³⁰, since also Coveney³⁰ uses the full CCD amplitudes. However, his construction differs in key points. His Hamiltonian is constructed in the CC similarity transform structure. It does not contain any metric and is non-Hermitian due to the similarity transformation. It also contains a line dressing that ADC(4) contains only at 4th order.

B. Connections to (vertex-corrected) GW

The GW approximation and other schemes derived in the framework of Hedin’s equations³¹ are formulated in the eh channel only. It is therefore instructive to retain only the eh pair in Eq. (74). The quantities entering the effective Hamiltonian

then simplify to

$$\begin{aligned}
\mathcal{N}_{iab,jcd}^{\text{II}} &= [(Z_{\text{eh}} Z_{\text{eh}}^\dagger)^{-1}]_{ia,jc} \delta_{bd}, \\
\mathcal{H}_{iab,jcd}^{\text{II}} &= [Z_{\text{eh}}^{-\dagger} (\Omega_{\text{eh}} + \epsilon_b) Z_{\text{eh}}^{-1}]_{ia,jc} \delta_{bd}, \\
\mathcal{U}_{p;iab}^{\text{II}} &= \sum_n M_{\text{eh},p;nb}^{\text{II}} [Z_{\text{eh}}^{-1}]_{n,ia}, \\
\mathcal{N}_{iaj,kbl}^{\text{I}} &= [(Z_{\text{eh}} Z_{\text{eh}}^\dagger)^{-1}]_{ia,kb} \delta_{jl}, \\
\mathcal{H}_{iaj,kbl}^{\text{I}} &= [Z_{\text{eh}}^{-\dagger} (\Omega_{\text{eh}} - \epsilon_j) Z_{\text{eh}}^{-1}]_{ia,kb} \delta_{jl}, \\
\mathcal{U}_{p;iaj}^{\text{I}} &= \sum_n M_{\text{eh},p;nj}^{\text{I}} [Z_{\text{eh}}^{-1}]_{n,ia},
\end{aligned} \tag{76}$$

where we have specified the indices explicitly for later reference. In block II, the pair (ia) has the spectator particle b , in block I the spectator hole j , and the pair and its spectator are independent. With Eq. (28),

$$\begin{aligned}
\omega \mathcal{N}^{\text{II}} - \mathcal{H}^{\text{II}} &= Z_{\text{eh}}^{-\dagger} (\omega - \epsilon_b - \Omega_{\text{eh}}) Z_{\text{eh}}^{-1}, \\
\omega \mathcal{N}^{\text{I}} + \mathcal{H}^{\text{I}} &= Z_{\text{eh}}^{-\dagger} (\omega - \epsilon_j + \Omega_{\text{eh}}) Z_{\text{eh}}^{-1},
\end{aligned}$$

and the self-energy becomes

$$\Sigma_{\text{eh},pq}(\omega) = \sum_{nb} \frac{M_{\text{eh},p;nb}^{\text{II}} M_{\text{eh},q;nb}^{\text{II}*}}{\omega - \epsilon_b - \Omega_n} + \sum_{nj} \frac{M_{\text{eh},p;nj}^{\text{I}} M_{\text{eh},q;nj}^{\text{I}*}}{\omega - \epsilon_j + \Omega_n}. \tag{77}$$

The eh-first-order reducible vertices have been defined in Eq. (B8), with $M_{\text{eh},p;nb} = -M_{bp,n}^{\text{eh}}$. The self-energy Eq. (77) is completely equivalent to the PSD completion of the Σ^{TDHF} self-energy³² from Ref. 33, therein denoted as Σ^{PSDII} . The only difference is that they construct Eq. (B8) with a statically screened TDHF kernel. This also illustrates the interpretation of the PSD self-energy of Ref. 33 as a “single-channel parquet”-type approximation.

1. Double counting

It has been remarked in Ref. 33 that Σ^{PSDII} double counts the $\Sigma^{(2)}$ contribution. This raises the natural question of how this can be the case, since the Faddeev-ADC(3) construction performed here is double-counting-free by construction.

To see how the double counting of $\Sigma^{(2)}$ arises, we take $v \rightarrow 0$. Then $Z_{\text{eh}} \rightarrow \mathbf{1}$, $\bar{Z}_{\text{eh}} \rightarrow 0$, $\Omega_n \rightarrow \epsilon_a - \epsilon_i$, $\mathcal{N} \rightarrow \mathbf{1}$, $\mathcal{H}^{\text{II}} \rightarrow \Delta_{abi}$, $\mathcal{H}^{\text{I}} \rightarrow -\Delta_{ija}$, and the couplings become those of Eq. (48), $\mathcal{U}_{p;iab}^{\text{II}} \rightarrow \langle pi||ab \rangle$ and $\mathcal{U}_{p;iaj}^{\text{I}} \rightarrow \langle pa||ij \rangle$. Eq. (77) gives

$$\begin{aligned}
\Sigma_{\text{eh},pq} &\rightarrow \sum_{abi} \frac{\langle pi||ab \rangle \langle ab||qi \rangle}{\omega + \epsilon_i - \epsilon_a - \epsilon_b} + \sum_{ija} \frac{\langle pa||ij \rangle \langle ij||qa \rangle}{\omega + \epsilon_a - \epsilon_i - \epsilon_j} \\
&= 2 \Sigma_{pq}^{(2)}.
\end{aligned} \tag{78}$$

Every 2p1h and 2h1p configuration appears twice: first, as the pair (ia) with spectator b and then as the pair (ib) with spectator a . Of course, it would be possible to subtract this contribution

$$\Sigma^c = \Sigma_{\text{eh}} - \Sigma^{(2)}, \tag{79}$$

as is done in the FLEX approximation^{13,14,34,35} and in the several flavors of combinations of the $GW + T$ -matrix approximations.^{36,37} This would however defeat the purpose of the PSD-construction, as the resulting self-energy is then not a sum of squares anymore and hence not PSD.

2. GW

Constructing Eq. (B8) with the standard (direct electron–hole) RPA, and keeping only the direct vertex, gives the GW approximation,

$$\Sigma_{GW}(\omega) = M_{\text{eh}}^{\text{D}} [\omega - s_{\text{eh}} - \Omega_{\text{eh}}]^{-1} M^{\text{D}\dagger}. \tag{80}$$

Written as an eigenvalue problem in the 2p1h and 2h1p spaces, this is the upfolded form of GW .^{38,39} The functional form of the GW self-energy is retained independent of the linear-response problem that is solved to construct the M_{eh}^{D} and to obtain Ω_{eh} . Usually, the direct RPA is used, which is the TDHF form of the RPA without the direct electron–hole interaction. In Hedin’s framework, the GW approximation naturally goes together with the RPA, since both arise from the lowest-order vertex, $\Gamma = 1$, when the non-interacting propagator is calculated at the Hartree level. In practice, the non-interacting Green’s function is almost always chosen to be a (generalized) Kohn–Sham one.⁴⁰ This, together with the availability of TDDFT codes, has motivated the use of TDDFT to construct Eq. (80) in GW calculations for solids,^{41–50} but also in molecules,^{49,51–54} and it is to be noted that local vertices from the same kernels have also been placed in the self-energy.^{42–45,48–51,53–65} More recently, the solid state community has used statically screened TDHF, i.e. the usual BSE. This approach is quite successful as it lowers the overestimated band gaps of quasiparticle self-consistent GW in solids.^{66–69} For molecules, this approach significantly worsens ionization potentials⁷⁰ compared to standard qs GW . For molecules, TDHF screening has been tested,^{71,72} and even polarizabilities from EOM-CCSD.⁷³ Combined with HF Green’s functions, they somewhat improve over the RPA. However, they are unbalanced and miss the corresponding vertex in the self-energy.³²

3. Σ^{TDHF}

This naturally motivates the choice of the Σ^{TDHF} self-energy that has been introduced by Maggio and Kresse⁷⁴ and investigated later by several authors.^{70,72,75–78} A first-order vertex constructed to satisfy the Ward identity had been used before in self-consistent calculations for atoms and small molecules.^{79,80} It operates in Hedin’s framework. It solves the BSE for the eh channel with the complete first-order vertex (TDHF), and uses the same vertex in the self-energy. This self-energy is the natural choice if based on a HF non-interacting reference, since the TDHF vertex is the functional derivative of the HF self-energy with respect to G .

Hedin’s formalism operates with a self-energy with two vertices, but one of them stays bare at first order. The resulting

self-energies cannot be written in a Hermitian (PSD) resolvent form, which has been the motivation to introduce the PSD construction from above³³ in the first place. Of course, the idea of the PSD completion of the self-energy is much older.⁸¹ Either way, the Σ^{TDHF} self-energy can be obtained by replacing one of the two vertices of Eq. (77) by its direct part M_{eh}^{D} ,

$$\Sigma_{\text{GWT}}(\omega) = M_{\text{eh}}^{\text{D}}[\omega - s_{\text{eh}} - \mathbf{\Omega}_{\text{eh}}]^{-1} M_{\text{eh}}^{\dagger}. \quad (81)$$

This clearly looks like a hybrid between the GW approximation Eq. (80) and the PSDII self-energy Eq. (77). From the perspective of the present review, such a hybrid approximation seems to make very little sense. We however emphasize that, unlike the PSDII self-energy, it is derived rigorously in Hedin's framework. It of course comes with the downside that it stops being PSD. It also eliminates the contribution from the eh-triplet channel. The eh-triplet channel turns out to be absolutely devastating for numerical accuracy. In fact, in Ref. 33, the triplet channel has been eliminated completely, and the resulting self-energy, coined PSD1, was found to give excellent agreement with experiment. This is of course not a very satisfactory ad hoc approximation, albeit a physically well-motivated one.

A significant upside of the Σ^{TDHF} self-energy is that it is double-counting-free, by virtue of the fact that Hedin's equations only create a single diagram once. However, at third order in perturbation theory, it only contains half of the eh contribution. In the correlation channel language adopted here, it contains the contribution from C_{eh} (through $U^{\text{D}} \mathcal{G} C_{\text{eh}} \mathcal{G} U^{(1)\dagger}$), but not from C_{eh}^{-} (which is also the reason why it is double-counting-free at second order).

In Hedin's scheme, these missing diagrams would be added to the self-energy through the second-order vertex. In addition to the missing half of the eh channel, the second-order vertex also adds the missing pp channel to the self-energy; the T matrix is fully contained in Hedin's equations.⁸² It is a feature of Hedin's equations that they never add diagrammatically complete sets of diagrams at an iteration order. This can be understood from the structure of the equations and has for instance been pointed out by Schindlmayr and Godby⁸³. Being diagrammatically complete at a certain order, a quality closely linked but not equivalent to being crossing symmetric, is a feature we consider an important property of any theory of the electronic self-energy.

4. $G3W2$ -type vertex corrections

Within Hedin's equations, some of the simplest possible vertex corrections arise from a first-order (3-point) vertex of the form^{78,84}

$$\Gamma(123) = \delta(12)\delta(13) + iW(21)G(13)G(32), \quad (82)$$

that, inserted into Hedin's expression for Σ , results in the family of the so-called $G3W2$ -type self-energies,

$$\Sigma_{\text{xc}}(11') = \Sigma^{\text{GW}} - G(14)W(1'4)G(42)G(21')W(21) \quad (83)$$

that have been investigated first for solids^{85–89} and later for molecules.^{78,90} By splitting the screened Coulomb interaction into the bare interaction and its polarizable part $W = v + W^c$, the $G3W2$ term splits into 4 contributions, schematically

$$GWGWG = GvGvG + GW^cGvG + GvGW^cG + GW^cGW^cG. \quad (84)$$

The sum of the first two terms is known as the SOSEX-approximation.^{91,92} Bruneval and Förster⁷⁸ have coined the sum of the first three terms the 2SOSEX-approximation, pointing out that it constitutes a more systematic approximation to the full $G3W2$. We note that also a statically screened $G3W2$ approximation has been investigated both for solids⁹³ and molecules,⁹⁴ and more recently all these approximations to $G3W2$ have been used in a fully self-consistent scheme by Pokhilko *et al.*⁹⁵ and assessed for molecules⁹⁶ (notice that the authors of this method denote the static approximation to $G3W2$ as $G3W2$). Resumming the statically screened $G3W2$ vertex to all orders as an eh ladder gives the screened eh T matrix, recast as a vertex.⁸⁴ With W in the static limit, as used in practice by Romaniello, Bechstedt, and Reining⁸⁴, this is the vertex of Σ^{BSE} of Ref. 32, but in the self-energy only, while the eh propagator is kept at the level of the direct RPA.

Without any vertex corrections to the inner vertex that is used to generate W , the performance of all of these approximations is exceptionally poor. The performance of the statically screened $G3W2$ self-energy is satisfactory, but only because the magnitude of its effect in QP energies is very small.⁹⁴ However, when paired with vertex corrections for W , $G3W2$ -type approximations perform well.^{32,93}

TABLE II. Approximations to Eq. (70) built from the pairs of Eq. (35): exact at second order, eh and pp parts of ADC(3) at third order, and positive semidefinite self-energy (Faddeev: for positive definite $\mathcal{N}$). Faddeev, eh only: Eq. (44) with $Q_{\text{pp}} \rightarrow \mathcal{G}$.

	$\Sigma^{(2)}$	eh	pp	PSD	Ref.
Σ_{eh} (PSD)	$2\Sigma^{(2)}$	$C_{\text{eh}} + C_{\text{eh}}^{-}$	–	yes	33
$\frac{1}{2}\Sigma_{\text{eh}}$	exact	$\frac{1}{2}(C_{\text{eh}} + C_{\text{eh}}^{-})$	–	yes	33
$\Sigma_{\text{eh}} - \Sigma^{(2)}$	exact	$C_{\text{eh}} + C_{\text{eh}}^{-}$	–	no	this work
Σ_{GWT}	exact	C_{eh}	–	no	32,74
Σ_{FLEX}	exact	$C_{\text{eh}} + C_{\text{eh}}^{-}$	C_{pp}	no	13,14
Faddeev-ADC(3)	exact	$C_{\text{eh}} + C_{\text{eh}}^{-}$	C_{pp}	yes	5–8

5. PSD-version of $G3W2$ -type self-energies

a. PSD-2SOSEX. Like all vertex-corrected self-energies derived in Hedin's formalism, also the ones of $G3W2$ -type cannot be written in resolvent form. Instead, their pole structures are rather complicated. Recently, the PSD-completion procedure of Stefanucci *et al.*⁸¹ has been applied to derive a PSD-version of the $GW + 2\text{SOSEX}$ self-energy,¹²⁵ and it is interesting to investigate how it fits into the effective Hamiltonian framework of Faddeev-ADC(3). Like GW , the $GW + 2\text{SOSEX}$ self-energy is of the form of Eq. (77) and can be written in the

TABLE III. Treatment and combination of correlation channels in self-energy approximations: second-order exactness in the bare interaction and positive semidefiniteness (PSD). Second order of the PSD completions in finite systems: Ref. 33 and this work.

Class	Type	Approximation	$\Sigma^{(2)}$ exact	PSD	Ref.
pp pair	T matrix	G_0T_0 @ppRPA (one-shot)	yes	yes	15,97
	T matrix	$G_{RS}T_{RS}$ (renormalized singles)	yes	yes	98
	T matrix	first order of the pp Hedin equations	yes	yes	99
	T matrix	Kadanoff–Baym T matrix (self-consistent)	yes	yes	100–102
	T matrix	pp and eh T matrix in the G1–G2 scheme (GKBA)	yes	not discussed	103–106
	T matrix	generalized T matrix (self-consistent, Gorkov)	yes	yes	107,108
pp pair, screened kernel	screened T matrix	screened ladder (W in the rungs)	–	in general no	109–111
	screened T matrix	pp Hedin vertex corrections (perturbative in T)	yes	not discussed	99
eh pair	GW	GW	no ($\Sigma_D^{(2)}$ only)	yes	31
	eh T matrix	eh T matrix	yes	no	15
	$\Sigma^{\text{TDHF}} + \text{PSD}$	PSDI	no	yes	33
	$\Sigma^{\text{TDHF}} + \text{PSD}$	PSDII (Σ_{eh})	no	yes	33
	vertex in W only	GW with beyond-RPA screening	no ($\Sigma_D^{(2)}$ only)	yes	46,66,67,69,71,73
	vertex in W only	GW with beyond-RPA screening	no ($\Sigma_D^{(2)}$ only)	yes	46,66,67,69,71,73
channel sums	$GW + T$ matrix	FLEX	yes	no	13,14,34,35
	$GW + T$ matrix	$GW + T$	yes	no	36,37
	$GW + T$ matrix	statically screened FLEX (Hedin form)	yes	no	112
	charge + spin ladders	ladder DGA, TPSC (lattice models)	–	–	113–115
	$GW + \text{spin-flip ladder}$	$GW\Gamma$ (magnons)	–	–	116,117
channels coupled	all channels	dynamically screened ladder ($g_{123} \rightarrow 0$)	yes	no guarantee	118,119
	all channels	parquet (static reducible vertices, TDA)	yes	no guarantee	35
	all channels	parquet approximation (lattice models)	yes	no guarantee	120,121
three-particle	Faddeev	real-time Faddeev (NEGF, bare v)	yes	not discussed	122,123
	Faddeev	Faddeev-ADC(3) (RPA pairs, metric N)	yes	yes	this work
PSD completions	vertex-corrected GW	half-diagram squares ($GW + \text{vertex}$)	no (finite systems)	yes	81,89,124
	vertex-corrected GW	$GW+2\text{SOSEX-psd}$	no	yes	125
	vertex-corrected GW	ADC-G3W2	no	yes	126
non-PSD vertex	$G3W2$ -type	SOSEX	yes	no	91
	$G3W2$ -type	2SOSEX	yes	no	78
	$G3W2$ -type	$G3W2$	yes	no	78,87,95,96
	$G3W2$ -type	statically screened $G3W2$	yes	no	93,94
	$G3W2$ -type	screened T matrix as vertex	yes	no	84
	$GW\Gamma$	$GW\Gamma$ (TDHF vertex, Σ^{TDHF})	yes	no	32,74,77,79,80
	$GW\Gamma$	$GW\Gamma$ (BSE vertex, Σ^{BSE})	yes	no	32,70
	$GW\Gamma$	$GW\Gamma$ (local TDDFT vertex)	no	no	44,48,59
extended CC	vertex-corrected GW	ECC exchange corrections	no ($\Sigma_D^{(2)}$ only)	not discussed	16

form of Eq. (80),

$$\Sigma_{pq}^{\text{PSD},p}(\omega) = \sum_{c,bi} \frac{(M_{pc,bi}^D + \tilde{M}_{pc,bi}^D)(M_{qc,bi}^D + \tilde{M}_{qc,bi}^D)^*}{\omega - \epsilon_c - \Delta_{bi}}. \quad (85)$$

where

$$\tilde{M}_{pc,n}^D = \sum_{aj} M_{aj,n}^D \left[\frac{\langle pa|jc \rangle}{\epsilon_a - \epsilon_j - \Omega_n} + \frac{\langle pj|ac \rangle}{\epsilon_a - \epsilon_j + \Omega_n} \right]. \quad (86)$$

Clearly, in their first-order limit, $\Omega_n \rightarrow \epsilon_b - \epsilon_i$ and $M_{aj,n}^D \rightarrow \langle ab|ij \rangle$, Eq. (86) becomes

$$\tilde{M}_{pc,bi}^{D,(1)} = \sum_{aj} \langle ab|ji \rangle \left[\frac{\langle pa|jc \rangle}{\epsilon_a + \epsilon_i - \epsilon_b - \epsilon_j} + \frac{\langle pj|ac \rangle}{\epsilon_a + \epsilon_b - \epsilon_i - \epsilon_j} \right]. \quad (87)$$

The second term in this expression is a non-antisymmetrized second-order doubles amplitude, that is contained in ADC(3), and its renormalized version Eq. (86) is completely contained

in Faddeev-ADC(3) as part of the ring-CCD amplitudes. The first term contains instead the transition-energy difference

$$\Delta_{aj} - \Delta_{bi} = (\epsilon_a - \epsilon_j) - (\epsilon_b - \epsilon_i). \quad (88)$$

This is the denominator for first-order mixing of the two forward particle–hole transitions (bi) and (aj), and it is part neither of the pp nor the eh-type couplings in Eq. (67).

To see how it fits into Faddeev-ADC(3), define the direct electron–hole scattering matrix between the $2p1h$ configurations ($c; bi$) and ($c; aj$) by

$$C_{c,bi;c,aj}^{\text{eh},D,(1)} \equiv -\langle ab|ji \rangle. \quad (89)$$

where c is a fixed spectator. Their unperturbed energies are $E_{c,bi} = \epsilon_c + \Delta_{bi}$ and $E_{c,aj} = \epsilon_c + \Delta_{aj}$. They enter the Faddeev-ADC(3) self-energy, Eq. (70), via the expansion of the resol-

vent $(\omega N - \mathcal{H})^{-1}$ via Eq. (66) through the contribution

$$\begin{aligned}\Sigma_{\text{fwd,pq}}^{(3)}(\omega) &= \left[U^{(1)} g(\omega) C^{\text{eh,(1)}} g(\omega) U^{(1)\dagger} \right]_{pq} \\ &= \sum_{c,bi,aj} \frac{m_{p;c,bi} C_{c,bi;c,aj}^{\text{eh,(1)}} m_{q;c,aj}^*}{(\omega - E_{c,bi})(\omega - E_{c,aj})} \\ &= - \sum_{c,bi,aj} \frac{\langle pb|ci \rangle \langle ab|ji \rangle \langle qa|cj \rangle}{(\omega - E_{c,bi})(\omega - E_{c,aj})}.\end{aligned}\quad (90)$$

The denominator of this term can be expanded through a partial fraction decomposition as

$$\frac{1}{(\omega - E_{c,bi})(\omega - E_{c,aj})} = - \frac{1}{\Delta_{aj} - \Delta_{bi}} \left[\frac{1}{\omega - E_{c,bi}} - \frac{1}{\omega - E_{c,aj}} \right], \quad (91)$$

which shows that it contains the denominator of the first term in Eq. (87). The numerators are the same as the ones of the third-order terms of Eq. (85),

$$\Sigma_{pq}^{(3),\text{PSD},p}(\omega) = \sum_{c,bi} \frac{\tilde{M}_{pc,bi}^{D,(1)} M_{qc,bi}^{D,(0)*} + M_{pc,bi}^{D,(0)} \tilde{M}_{qc,bi}^{D,(1)*}}{\omega - \epsilon_c - \Delta_{bi}}. \quad (92)$$

Of course, by expanding the resolvent, one can match the terms in the PSD-2SOSEX and Eq. (70) order by order.

Like its PSDII sibling, PSD-2SOSEX double counts part of $\Sigma^{(2)}$. The origin is the first term of Eq. (86). For the phonon n that reduces to the transition (aj) for $v \rightarrow 0$, its denominator is the RPA shift of that transition,

$$\epsilon_a - \epsilon_j - \Omega_n = O(v),$$

and the term is of first order, not of second. In Eq. (87) it is the diagonal term $(aj) = (bi)$, where $\Delta_{aj} - \Delta_{bi} = 0$. With the singlet phonons and spin-adapted vertices of Refs. 125,126, $M_{aj,n}^D/(\epsilon_a - \epsilon_j - \Omega_n) \rightarrow -1/\sqrt{2}$, and the vertex of Eq. (85) becomes, to first order,

$$M_{pc,aj}^D + \tilde{M}_{pc,aj}^D \rightarrow \frac{1}{\sqrt{2}} (2 \langle pa|cj \rangle - \langle pa|jc \rangle),$$

the singlet antisymmetrized vertex of PSDI. Summed over a, c and j ,

$$\Sigma^{\text{PSD}} = \frac{5}{4} \Sigma_D^{(2)} + 2 \Sigma_X^{(2)} + O(v^3),$$

the same second order as PSDI. The cross term $2 \Sigma_X^{(2)}$ is the second-order exchange that $GW+2\text{SOSEX}$ -aug already counts twice;¹²⁵ completing the square adds $\frac{1}{4} \Sigma_D^{(2)}$.

In summary, double counting aside, the PSD-2SOSEX self-energy is a (severely) channel-restricted subset of the Faddeev-ADC(3) self-energy, but not through a one-to-one correspondence of individual pieces. As it does not assume the TDA, it contains terms that are not contained in ADC(3).

b. PSD-G3W2. Marie, Tölle, and Loos¹²⁶ applied the ADC construction to the expansion of the G3W2 self-energy in W , with the goal to complete it into a PSD expression. The configurations are those of Eq. (76), with direct-RPA phonons,

$K = \epsilon_a + \Omega_v (\epsilon_i - \Omega_v)$. Their PSD-G3W2 effective Hamiltonian goes beyond the Faddeev-ADC(3) configuration space and adds 3p2h configurations that would show up in ADC(4) and ADC(5). It also correlates the 5-particle block, terms that would otherwise only show up in ADC(5). Of course, since they solve the RPA problem, their scheme goes beyond ADC, but it is reasonable to assume that their self-energy is obtained in Faddeev-ADC(5). In the bare interaction PSD-G3W2 contains PSD-2SOSEX. A doubly resonant term of $U^{(3)}$ adds a further first-order exchange vertex, so that ADC-G3W2 gives $\frac{17}{16} \Sigma_D^{(2)} + \Sigma_X^{(2)}$ in total.

All constructions that make self-energies derived in Hedins method PSD seem to share this shortcoming, and it is unclear how it can be overcome. It is also reasonable to assume that the PSD completions of higher-order self-energy derived in Hedins equations would also double count contributions at higher orders. As already pointed out, the double counted contributions can of course be subtracted, but this would make the self-energies non-PSD. For this reason, the current PSD-completions do not seem to constitute a path toward systematically improvable self-energy approximations

C. The pp channel

Very similar to working in the eh channel only, it is of course also possible to restrict Eq. (74) to the pp channel only. The canonical self-energy approximation that operates in this channel is the T -matrix approximation.^{15,99} Unlike in the eh channel, there is no double counting ambiguity in the pp channel, simply due to the fact that it is not related by a simple symmetry to another channel in the same way as the eh channel. In the same way as the PSDII self-energy, the resulting self-energy is complete to third order in the pp channel, and of course, by construction, it is also PSD.

T -matrix self-energies have been used and extended in several communities. In molecules, the G_0T_0 approximation is built on the ppRPA,⁹⁷ on a renormalized-singles reference,⁹⁸ compared to GW and the eh T matrix,¹⁵ used in the Bethe–Salpeter equation,¹²⁷ and obtained as the first order of the pp Hedin equations.⁹⁹ Self-consistent T matrices are used in Kadanoff–Baym dynamics,^{100–102,104,105} and, with anomalous propagators, in nuclear matter.^{107,108} Screening enters the pp kernel in the screened ladder approximation of plasma physics,^{109–111} through the Gorkov–Melik-Barkhudarov correction,¹²⁸ and through the vertex corrections of the pp Hedin equations.⁹⁹

D. FLEX and other combinations of eh and pp channels

Truncating the Faddeev series, Eq. (35), after the terms with a single interacting pair, Eq. (31), gives the one-shot FLEX approximation,^{13,14}

$$R_{\text{FLEX}} = \mathcal{G} + \sum_{\alpha} (Q_{\alpha} - \mathcal{G}) = Q_{\text{pp}} + Q_{\text{eh}} + Q_{\text{eh}} - 2 \mathcal{G}.$$

Each pair couples to the one-particle states through its own couplings, $\mathcal{U}_\alpha = M_\alpha Z_\alpha^{-1}$, and the free propagation through $U^{(1)}$,

$$\Sigma_{\text{FLEX}} = \sum_\alpha \mathcal{U}_\alpha \mathcal{Q}_\alpha \mathcal{U}_\alpha^\dagger - 2 U^{(1)} \mathcal{G} U^{(1)\dagger}. \quad (93)$$

Each pair term contains $\Sigma^{(2)}$ once, and $-2\Sigma^{(2)}$ removes the excess. Faddeev-ADC(3) and FLEX agree through first order in C ; both are exact through third order. FLEX however misses the sequences of different pairs in Eq. (35), which enter at fourth order. Since the subtraction is outside the resolvent, Σ_{FLEX} has negative residues and no Hermitian effective Hamiltonian. In the Tamm–Dancoff limit,

$$R_{\text{FLEX}} = \sum_\alpha (\omega - K - C_\alpha)^{-1} - 2(\omega - K)^{-1}.$$

FLEX is usually motivated as a drastic approximation to the parquet approximation,^{14,35} but closely related approximations can also be motivated as a rather heuristic combination of correlation channels. For instance, to place eh and pp on the same footing, GW and the T matrix have been summed with a subtraction of the common second-order terms in $GW + T$.^{36,37} Starting from Hedin’s equations, also Cunningham¹¹² has rederived FLEX, but he suggested screening all pair channels statically in the TDA. In the eh channel he suggested to use the kernel $v - W$, and in the pp channel he screened all interactions ($W - W$, schematically). The same has been done for instance also by Romaniello and Berger¹²⁹. Instead of being summed, the channels are coupled within one two-particle correlation function in the dynamically screened ladder approximation^{118,119} and in the parquet approximation.^{35,120,121}

The static screening approximation is of course motivated by the usual GW -BSE approximation for neutral excitations, and many authors have solved the eh channel RPA with a statically screened HF problem.^{33,67,69,70,72,78} This is well motivated for solids, where TDHF is problematic and will even diverge for metals. For molecules, this replacement is motivated by the superior performance of GW -BSE over TDHF. However, due to the fact that static screening effects are usually rather weak in molecules, the effects on ionization potentials are minor.^{28,32}

V. NUMERICAL RESULTS

Most of these approximations have never been evaluated for the same set of molecules in a reasonable basis set. Here, we therefore evaluate many of them for the set of 58 IPs of 23 small molecules for which Marie and Loos¹³⁰ have recently calculated reference values of near FCI quality in the aug-cc-pVQZ basis. We do not consider statically screened interactions to calculate the ph/pp RPA phonons. For once, these are heuristic adjustments, and, if not applied carefully, they can lead to double counting issues. Moreover, they can only be expected to have a minor influence on the final results. A method that does not perform well with bare interactions will not perform well with statically screened interaction either.

Table IV summarizes the errors, and all raw numerical data is shown in Table V. All calculations use RHF references, are

TABLE IV. Errors (eV) with respect to sCI over the 48 states of Table V with a value for every method, and MAD over their 20 first IPs. FLEX CCD: running, 44 states so far.

	N	MAD	MSE	max	MAD first IP
ADC(3)	48	0.30	+0.30	1.12	0.26
CCD-ADC(3)	48	0.27	+0.26	0.71	0.23
EN-ADC(3)	48	0.07	−0.01	0.25	0.05
Faddeev-ADC(3) RPA ^a	48	2.05	+2.03	21.03	1.54
Faddeev-ADC(3) T1	48	0.33	+0.31	1.39	0.29
Faddeev-ADC(3) T0	48	0.51	+0.45	3.47	0.28
CCD-Faddeev-ADC(3)	48	0.19	+0.11	1.08	0.14
Faddeev-ADC(3) dRPA	48	0.67	+0.60	3.86	0.40
FLEX RPA	48	1.42	−0.33	5.87	1.24
FLEX T1	48	1.09	−1.01	3.96	0.85
FLEX T0	48	0.49	+0.44	3.24	0.27
FLEX CCD	44	1.24	−1.16	4.09	1.07
PSD1	48	0.51	−0.17	2.88	0.35
PSD1 $-D/4 - X$	48	0.29	−0.08	1.64	0.19
PSD2	48	2.16	−1.45	5.84	1.83
PSD2 $-GF2$	48	1.60	−0.59	5.39	1.40
PSD2 T1 $-GF2$	48	1.38	−1.29	4.44	1.14
CCSD	48	0.16	+0.10	0.92	0.08
EOM-CCD	48	0.24	+0.12	1.02	0.19
GW	48	0.52	+0.50	2.14	0.44
$GF2$	48	0.78	−0.66	1.94	0.78
All 56 states with an sCI value (23 first IPs), C_2 , BN and F_2 included					
ADC(3)	56	0.33	+0.31	1.12	0.28
CCD-ADC(3)	56	0.29	+0.29	0.77	0.24
EN-ADC(3)	56	0.09	−0.01	0.64	0.06
CCD-Faddeev-ADC(3)	56	0.23	+0.14	1.08	0.20
Faddeev-ADC(3) dRPA	56	0.76	+0.63	3.86	0.45
FLEX CCD (51 so far)	51	1.38	−1.12	4.57	1.25
CCSD	56	0.18	+0.12	1.08	0.10
EOM-CCD	56	0.25	+0.13	1.15	0.21
GW	56	0.53	+0.49	2.14	0.45

^a Faddeev-ADC(3) RPA for BF and CH₂O. The TDHF triplet channel is close to unstable (lowest triplet excitation 1.4 and 2.1 eV). The Davidson root of the HOMO (BF) and of 1b₁ (CH₂O) does not converge, so the entries are the remaining converged roots in energy order, not assigned states.

performed in an all-electron framework, make use of density fitting with the aug-cc-pVQZ-RI basis, and, except GW , add the MP2-relaxed static self-energy. Faddeev-ADC(3) and the ADC(3) variants are solved as Dyson supermatrices; FLEX, PSD and GF2 in the diagonal quasiparticle approximation. T1 and T0 denote the triplet eh channel at first order and switched off, respectively. C_2 and BN are excluded, since their RHF references are unstable in the TDHF singlet (and triplet) channel, and F_2 , since its TDHF triplet is unstable. The statistics run over the 48 states with a value for every method. FLEX finds no quasiparticle root for NH₃ (3) (RPA) and Ne (2) (T0).

From a formal perspective, Faddeev-ADC(3) has many desirable properties. It is exact to third order, double-counting-free, PSD, Hermitian, and includes renormalized amplitudes via the solution of the ring- and ladder-CCD equations. However, its numerical performance is abysmal, coming with a MAD of 2.05 eV. FLEX performs nearly equally badly, and the only method that performs even worse than Faddeev-ADC(3) is the PSDII approximation of Ref. 33. What is the reason for this bad performance of these methods? All of them have

TABLE V. IPs (eV) of the set of Marie, Romaniello, and Loos⁹⁹ in the aug-cc-pVQZ basis. ADC(3), CCD-ADC(3) and spin-summed doubles-only EN-ADC(3) from Ref. 29; all ADC variants and FLEX with the MP2-relaxed static self-energy. Faddeev-ADC(3) and FLEX with RPA pair channels, with the triplet particle-hole channel at first order (T1) or switched off (T0); dRPA: direct-RPA eh channels; CCD: one CCD T₂ in every pair channel. PSD1 and PSD2 on bare TDHF phonons; *D* and *X* are the direct and second-order-exchange parts of GF2. CCSD: EOM-IP-CCSD, this work. *GW* (*G*₀*W*₀@HF, without the MP2-relaxed static self-energy) from the SupInf reference data; EOM-CCD on the CCD of CCD-Faddeev-ADC(3). – : no sCI value, method not defined (unstable TDHF channel), or no quasiparticle. MAD and MSE are over the 48 states with a value for every method (C₂, BN and F₂ excluded). All calculations have been performed in the diagonal one-shot approximation. FLEX CCD: running, 44 states so far.

IP	sCI	ADC(3)	CCD-ADC(3)	EN-ADC(3)	Faddeev-ADC(3)					FLEX				PSD1		PSD2		CCSD	EOM-CCD	GW	GF2	
					RPA	T1	T0	CCD	dRPA	RPA	T1	T0	CCD	bare	$-D/4-X$	bare	-GF2					T1-GF2
Ar (1)	15.74	15.84	15.86	15.71	16.55	16.02	15.77	15.86	15.79	16.18	15.66	15.83	15.58	15.67	15.73	15.63	16.08	15.57	15.80	15.84	15.93	15.28
Ar (2)	29.18	29.76	29.78	29.26	29.25	29.29	32.65	29.23	33.05	25.12	25.23	32.38	25.10	32.02	30.82	23.34	24.66	24.74	29.96	30.00	31.22	30.26
BeO (1)	10.02	10.72	10.40	9.91	13.47	10.62	10.66	10.14	10.74	7.10	7.41	10.17	7.08	9.17	9.29	6.49	6.92	7.05	9.96	9.66	9.91	8.70
BeO (2)	11.12	12.23	11.82	11.12	13.79	11.80	12.03	11.46	12.27	8.10	8.15	11.52	7.95	9.77	10.11	6.52	7.62	7.47	11.10	10.78	11.20	9.17
BF (1)	11.18	11.21	11.32	11.14	18.69 ^a	11.67	10.88	11.63	10.89	17.05	11.48	11.09	12.02	11.05	11.20	15.13	16.57	11.55	11.29	11.38	11.42	10.75
BF (2)	18.66	19.27	19.13	18.79	21.54 ^a	19.14	19.38	18.91	19.71	17.12	16.99	19.39	16.88	18.27	18.41	17.05	17.10	16.44	18.30	18.26	18.95	17.58
BF (3)	20.97	21.62	21.51	21.19	42.00 ^a	22.36	22.26	22.05	22.66	22.20	19.82	22.16	19.68	20.73	21.05	17.25	22.05	19.15	21.36	21.27	21.78	20.30
BH ₃ (1)	13.33	13.36	13.40	13.28	14.22	13.55	13.30	13.45	13.44	13.88	13.27	13.31	13.31	13.05	13.28	13.62	13.83	13.24	13.39	13.44	13.78	13.10
BH ₃ (2)	18.33	18.35	18.41	18.23	19.10	18.50	18.67	18.41	18.83	18.24	17.76	18.69	17.78	18.56	18.48	17.69	18.11	17.63	18.42	18.48	18.78	18.28
BN (1)	12.02	12.54	12.42	11.88	-	-	-	-	12.92	12.22	-	-	-	9.40	-	-	-	-	12.04	11.90	11.91	-
BN (2)	13.71	13.89	14.00	13.06	-	-	-	-	13.91	12.96	-	-	-	15.23	-	-	-	-	13.72	13.34	13.59	-
C ₂ (1)	12.50	12.24	12.57	12.66	-	-	-	-	13.22	11.57	-	-	-	14.44	-	-	-	-	13.06	13.33	13.08	-
CH ₂ O (1)	10.95	11.29	11.24	10.88	11.46 ^a	11.13	11.23	10.96	11.50	9.43	9.44	11.15	9.36	10.14	10.49	8.01	9.05	9.01	10.94	10.93	11.56	9.87
CH ₂ O (2)	14.62	14.72	14.74	14.50	16.75 ^a	15.01	14.54	14.67	14.57	14.04	13.71	14.52	13.36	14.23	14.42	12.75	14.03	13.68	14.69	14.63	14.76	13.92
CH ₂ O (3)	16.16	16.70	16.63	16.16	18.21 ^a	16.47	16.74	16.29	17.04	14.55	14.36	16.67	14.22	15.44	15.82	13.06	14.08	14.04	16.20	16.17	16.79	15.02
CH ₂ O (4)	17.16	17.43	17.42	17.18	22.88 ^a	17.20	17.85	17.59	18.21	14.85	14.57	16.79	14.42	16.10	16.29	14.00	14.70	14.09	17.57	17.68	18.16	16.46
CH ₄ (1)	14.41	14.52	14.55	14.35	15.24	14.62	14.38	14.52	14.53	14.72	14.16	14.39	-	13.96	14.23	14.08	14.58	14.03	14.44	14.52	14.87	13.87
CH ₄ (2)	23.15	23.42	23.48	23.13	23.65	23.34	24.36	23.28	24.64	21.79	21.60	24.36	-	24.04	23.60	20.30	21.40	21.20	23.41	23.52	23.99	23.15
CO (1)	13.97	14.15	14.23	14.09	15.89	14.26	14.14	14.03	14.24	14.83	13.47	14.30	13.47	13.92	14.06	14.10	14.66	13.32	14.27	14.50	14.91	13.41
CO (2)	17.02	17.19	17.13	16.97	18.79	17.41	17.20	17.02	17.19	16.46	15.98	17.14	15.61	16.79	16.90	15.26	16.36	15.86	17.12	17.07	17.26	16.36
CO (3)	19.70	20.39	20.23	19.77	22.11	19.99	20.80	19.64	21.15	17.32	17.39	20.70	17.22	19.29	19.52	16.03	16.77	16.77	19.88	19.82	20.48	18.58
CO ₂ (1)	13.82	14.02	13.98	13.82	14.93	14.03	14.06	13.69	14.16	13.18	12.70	14.13	12.39	13.45	13.66	12.14	12.92	12.43	13.90	13.97	14.43	12.86
CO ₂ (2)	17.62	18.16	18.16	17.87	18.24	17.97	18.38	17.57	18.59	15.30	15.55	18.11	15.23	17.41	17.50	14.34	15.08	15.27	18.11	18.21	18.70	16.83
CO ₂ (3)	-	18.80	18.69	18.33	18.76	18.47	18.85	18.23	19.15	16.64	16.55	18.94	16.34	17.59	17.94	14.37	16.04	15.92	18.28	18.36	18.99	16.91
CO ₂ (4)	-	20.19	20.06	19.65	20.04	19.76	20.31	19.55	20.68	17.49	17.40	20.28	17.24	18.86	19.14	15.09	16.86	16.73	19.61	19.67	20.33	18.09
CS (1)	11.36	11.59	11.62	11.38	16.11	11.53	11.57	11.14	11.61	11.52	9.99	11.65	9.84	11.00	11.27	11.03	11.41	9.83	11.62	11.92	13.06	10.43
CS (2)	12.94	12.85	12.89	12.87	18.38	13.44	12.72	13.03	12.47	17.55	13.54	12.84	13.18	13.07	13.15	16.74	17.47	13.74	13.06	13.09	12.52	12.64
CS (3)	17.96	18.16	18.06	17.79	20.05	17.83	18.03	17.57	18.14	19.20	18.87	18.63	18.57	18.21	18.17	18.82	19.16	18.65	17.33	17.41	18.12	16.66
F ₂ (1)	15.81	16.33	16.22	15.69	-	-	-	-	15.75	17.07	-	13.42	16.39	13.34	-	-	-	-	15.75	15.73	16.56	-
F ₂ (2)	18.94	19.51	19.37	18.76	-	-	-	-	18.21	20.69	-	14.49	18.92	14.38	-	-	-	-	19.08	19.04	20.10	-
F ₂ (3)	21.15	21.36	21.45	21.14	-	-	-	-	22.03	20.87	-	21.92	20.36	22.39	-	-	-	-	21.26	21.13	21.03	-
H ₂ O (1)	12.74	13.15	13.07	12.72	13.30	12.99	13.14	12.81	13.37	11.27	11.25	13.18	11.10	12.22	12.35	9.41	10.73	10.68	12.70	12.68	13.08	11.54
H ₂ O (2)	14.96	15.40	15.32	14.95	15.73	15.27	15.33	15.09	15.54	13.97	13.77	15.39	13.66	14.45	14.66	12.27	13.49	13.27	14.91	14.88	15.29	13.88
H ₂ O (3)	19.06	19.43	19.38	19.05	20.09	19.43	19.22	19.28	19.41	18.78	18.37	19.16	18.31	18.39	18.73	17.62	18.45	18.04	19.04	19.01	19.29	18.24
H ₂ S (1)	10.46	10.48	10.51	10.38	11.62	10.71	10.32	10.54	10.29	11.20	10.39	10.41	10.35	10.34	10.42	10.99	11.18	10.40	10.49	10.55	10.66	9.97
H ₂ S (2)	13.46	13.52	13.55	13.38	14.80	13.72	13.38	13.58	13.41	14.28	13.32	13.44	13.32	13.23	13.39	13.91	14.21	13.29	13.50	13.56	13.76	12.97
H ₂ S (3)	15.67	15.73	15.76	15.60	16.60	15.90	15.68	15.77	15.74	16.52	15.33	15.68	15.26	15.41	15.61	16.27	16.48	15.29	15.73	15.80	16.06	15.27
HCl (1)	12.77	12.84	12.87	12.71	13.74	13.04	12.73	12.87	12.73	13.29	12.64	12.80	12.57	12.64	12.72	12.86	13.21	12.58	12.81	12.86	12.97	12.26
HCl (2)	16.73	16.82	16.84	16.68	17.84	17.01	16.72	16.85	16.75	17.45	16.64	16.77	16.61	16.55	16.70	17.00	17.37	16.59	16.78	16.83	17.03	16.28
HF (1)	16.21	16.79	16.66	16.23	16.74	16.56	16.85	16.41	17.17	14.62	14.68	16.89	14.56	15.61	15.80	12.01	13.86	13.89	16.14	16.04	16.45	14.91
HF (2)	20.10	20.57	20.47	20.11	20.99	20.50	20.49	20.32	20.72	19.36	19.14	20.53	19.04	19.60	19.84	17.62	18.86	18.63	20.03	19.93	20.26	19.15
LiCl (1)	10.01	10.16	10.17	9.97	10.92	10.33	10.11	10.14	10.12	10.06	9.61	10.15	9.50	9.89	9.96	9.49	9.91	9.48	10.07	10.09	10.18	9.49
LiCl (2)	10.70	10.87	10.87	10.66	11.63	11.03	10.81	10.84	10.84	10.80	10.32	10.83	10.21	10.51	10.61	10.14	10.63	10.16	10.76	10.78	10.90	10.16
LiF (1)	11.54	12.35	12.13	11.64	12.29	12.09	12.41	11.89	12.70	9.42	9.67	12.42	9.51	10.99	11.13	6.98	8.67	8.85	11.51	11.31	11.59	10.22
LiF (2)	12.02	12.96	12.72	12.16	12.83	12.65	12.98	12.47	13.33	10.05	10.21	12.90	10.09	11.24	11.53	7.28	9.19	9.30	12.00	11.78	12.14	10.63
N ₂ (1)	15.54	15.85	15.82	15.61	15.88	15.67	15.88	15.42	16.06	14.28	14.18	16.04	13.96	15.18	15.40	12.77	13.90	13.77	15.74	15.95	16.52	14.46
N ₂ (2)	16.98	16.95	17.11	17.03	17.71	17.70	16.71	17.29	16.45	15.61	16.04	16.81	15.84	16.99	17.16	14.16	15.26	15.59	17.32	17.43	17.26	16.67
N ₂																						




FIG. 4. Errors of the IPs with respect to sCI (aug-cc-pVQZ) for ADC(3), CCD-ADC(3), EN-ADC(3), CCD-Faddeev-ADC(3) and CCSD, over the 48 states of the first block of Table IV.

This improves the MAD from more than 2.0 eV to about 0.5 eV, which is not great, but somewhat reasonable, in the same ballpark as G_0W_0 . By making it double-counting-free, its MAD nearly halves, to 0.29 eV, on par with ADC(3).

This motivates evaluating Faddeev-ADC(3) and FLEX without triplet excitations. When the triplet channel is omitted altogether (T0), both methods significantly improve, and the MADs become 0.51 and 0.49 eV. As a less drastic approximation, one can replace the triplet eh amplitudes by their first-order doubles counterparts that are also used in ADC(3). For Faddeev-ADC(3), this reduces the error further to 0.33 eV. For FLEX it does not help; the error increases again to more than 1.0 eV. Another way to avoid triplet instabilities is to use direct RPA instead of TDHF in the eh channel. This can be done both for Faddeev-ADC(3) and FLEX, but we have only numerically evaluated it for Faddeev-ADC(3). For the subset of 48 states for which Faddeev-ADC(3) is defined this gives a MAD of 0.67 eV, which is somewhat more reasonable than the 2.05 eV of Faddeev-ADC(3) with the RPA channels, but still far from good.

The results clearly show that channel coupling is important. The same has been found for correlation energies in coupled-cluster theory, where each single channel of CCD, rings, ladders or crossed rings, individually, diverges for the electron gas. Only the terms that couple the channels (the mosaic terms) remove the divergence.^{9,17} The reason is that the triplet eh channel resummed with the bare interaction is close to unstable, and sometimes even completely unstable, as for instance seen for F_2 . As shown in Fig. 5, the error of Faddeev-ADC(3) grows as the lowest TDHF triplet approaches zero.

The observation that keeping it at first order (T1) or switching it off (T0) improves the results parallels the instabilities of RPA with exact exchange, which are removed by keeping exchange exact only at first order^{131,132} and the improved stability of singlet-paired coupled cluster.^{133,134}

This motivates the replacement of the uncoupled doubles amplitudes in the Faddeev-ADC(3) self-energy by CCD ones, which contain the couplings between the channels. The resulting method, to which we refer as CCD-Faddeev-ADC(3), is




FIG. 5. Mean absolute error per molecule of Faddeev-ADC(3) (RPA pair channels) against sCI, versus the lowest TDHF triplet excitation energy of the RHF reference (aug-cc-pVDZ). For unstable references, where Faddeev-ADC(3) has no solution (open circles), the most negative eigenvalue of $A \pm B$ is shown; (S): also unstable in the singlet channel.

the same substitution that turns ADC(3) into CCD-ADC(3).²⁹ Since no RPA problem has to be solved, it is also defined in case of a triplet instability. The triplet problem disappears. The MAD over all 56 states with an sCI reference value is 0.23 eV, against 0.33 eV for ADC(3), 0.29 eV for CCD-ADC(3) and 0.18 eV for CCSD. The error distributions are shown in Fig. 4. For the subset of 48 states for which Faddeev-ADC(3) is actually defined, CCD-Faddeev-ADC(3) is a substantial improvement over CCD-ADC(3), with a MAD of 0.19 eV. This is very close to CCSD that comes with a MAD of 0.16 eV.

Maybe a bit surprisingly, the CCD amplitudes do not rescue FLEX. With an MAD of 1.24 eV, it is almost equally bad FLEX with RPA channels (MAD of 1.42 eV). The CCD channels lower the IPs, which FLEX already underestimates. The deficiency of FLEX is not the quality of its pair channels but the way it combines them.

VI. CONCLUSIONS

Through the response function R , Eqs. (13) and (15), the self-energy is expressed in terms of the three-particle kernel. In the Faddeev form, Eq. (20), the only two-body input is the parquet vertex F . ADC(n) is the Schirmer matching of the Faddeev series at total order $n - 2$ in v . The orders through which the various approximations are exact are collected in Table I.

Mapping the Faddeev form of parquet onto an effective Hamiltonian of the 2p1h (2h1p) space involves a few approximations. First, the parquet approximation, $\Lambda = \Lambda^{(1)}$, is combined with a static reference, $G = G_{\text{HF}}$ in all lines of the vertex. Every building block of the vertex is then an instantaneous interaction $\bar{v}$ or a G_{HF} line, and the frequency integrals of the

pair propagators turn into resolvents of configuration spaces (Sec. III B). Next, an own-channel approximation is invoked that consists in letting each pair scatter only through the ladder of its own channel with the bare kernel. Moreover, the ladders of different pairs do not overlap in time (Sec. III A).

In the own-channel and no-overlap approximation, Hermitian effective Hamiltonians (for a positive definite metric) are obtained that go beyond ADC by going beyond the TDA. At the ADC(3) level, this amounts to an effective Hamiltonian that replaces the first-order double amplitudes in the couplings with the ring-CCD (for the eh channel) and the ladder-CCD amplitudes (for the pp channel), and adds the metric $\mathcal{N}$ and the terms $B_\alpha T_\alpha$ to the matrix. Channel crossings that are not already part of the Faddeev kernel itself, as in the full CCD amplitude equations, are not considered. At first order, the Faddeev kernel does not contain any channel crossings.

Starting from the second-order Faddeev kernel, crossed channels already enter before any resummation, by Eq. (A8) and Eq. (23). These crossed ladders, together with the overlapping orderings of two first-order kernels and the line dressing, lead to the presence of 3p2h configurations. Obtaining an effective Hamiltonian requires including this configuration space, in the same way as in the ADC(4) approximation.

An advantage of our Faddeev construction is that it preserves the resolvent structure of the self-energy through a Hermitian effective Hamiltonian formalism, in the same way as in ADC. Unlike ADC, it also introduces a significantly larger amount of ground-state correlation through an infinite-order amplitude renormalization. The derivations shown here allow us to make precise connections to recent PSD completions of vertex-corrected self-energies.^{33,125,126} The formal properties of the self-energies discussed here are summarized in Table VI.

TABLE VI. Formal properties of self-energy approximations. DC free: no diagram counted twice. 2nd, 3rd: exact through second and third order in v . Hermitian: $\Sigma(\omega) = \Sigma^\dagger(\omega)$ for real ω . Resolvent: $\Sigma(\omega) = \Sigma^\infty + \mathcal{U}(\omega - \mathcal{H})^{-1}\mathcal{V}$ with ω -independent matrices of finite dimension. Coupled: the amplitudes of different channels are coupled.

	DC free	2nd	3rd	Hermitian	Resolvent	Coupled
Faddeev-ADC(3)	✓	✓	✓	✓	✓	×
CCD-Faddeev-ADC(3)	✓	✓	✓	✓	✓	✓
CCSD	✓	✓	✓	×	✓	✓
GW	✓	×	×	✓	✓	×
PSDII	×	×	×	✓	✓	×
Σ^{TDHF}	✓	✓	×	×	✓ ^a	×
Hedin, general	✓	✓ ^b	✓ ^b	×	×	✓ ^b

^aEq. (81), with $\mathcal{V} \neq \mathcal{U}$.

^bIf the vertex is iterated to the corresponding order.

^cNot for a dynamical vertex in general.

However, the numerical results of Faddeev-ADC(3) are very poor, which can be linked to the triplet channel. The same goes for the FLEX approximation and the PSDII approximation of Pavlyukh, Bruneval, and Förster³³. Switching off the triplet channel entirely or leaving the triplet amplitudes at first order, like in ADC(3), significantly improves performance. Nevertheless, these schemes are clearly not competitive with methods based on the CC scheme, for instance diagonalization of the

CCSD Hamiltonian. Clearly, coupling of the different correlation channels is mandatory whenever single-particle states are coupled to triplets beyond the TDA. To support this point, the replacement of the uncoupled amplitudes in Faddeev-ADC(3) with CCD ones results in improved results, but ADC(3) is only outperformed slightly. The performance is almost identical to CCD-ADC(3), and does not reach the performance of CCSD.

Appendix A: Parquet vertices

The Faddeev kernel is expanded order by order in $\bar{v}$,

$$\mathcal{K} = \mathcal{K}^{(1)} + \mathcal{K}^{(2)} + \mathcal{K}^{(3)} + O(v^4). \quad (\text{A1})$$

At first order, all channels contain only the bare antisymmetrized interaction,

$$\begin{aligned} \Gamma^{\text{pp},(1)} &= \Gamma^{\text{eh},(1)} = \Gamma^{\bar{\text{eh}},(1)} = \Lambda^{(1)} \equiv \Gamma^{(1)}, \\ \Gamma^{(1)}(12; 1'2') &= -i \bar{v}(12; 1'2'), \\ \bar{v}(12; 1'2') &\equiv v(12; 1'2') - v(12; 2'1'). \end{aligned} \quad (\text{A2})$$

To all orders,

$$\begin{aligned} \Gamma^{\text{eh}} &= \Lambda + \Phi^{\bar{\text{eh}}} + \Phi^{\text{pp}}, \\ \Gamma^{\bar{\text{eh}}} &= \Lambda + \Phi^{\text{eh}} + \Phi^{\text{pp}}, \\ \Gamma^{\text{pp}} &= \Lambda + \Phi^{\text{eh}} + \Phi^{\bar{\text{eh}}}, \\ F &= \Gamma^\lambda + \Phi^\lambda = \Lambda + \Phi^{\text{pp}} + \Phi^{\text{eh}} + \Phi^{\bar{\text{eh}}}. \end{aligned} \quad (\text{A3})$$

for each channel $\lambda \in \{\text{pp}, \text{eh}, \bar{\text{eh}}\}$. Note that³⁵

$$\Lambda = \Lambda^{(1)} + O(v^4). \quad (\text{A4})$$

Defining

$$(A \circ_{\text{pp}} B)(12; 1'2') \equiv A(12; \bar{1}\bar{2}) G(\bar{1}\bar{1}') G(\bar{2}\bar{2}') \times B(\bar{1}'\bar{2}'; 1'2'), \quad (\text{A5})$$

$$(A \circ_{\text{eh}} B)(12; 1'2') \equiv A(1\bar{4}; 1'\bar{3}) G(\bar{3}\bar{3}') G(\bar{4}'\bar{4}) \times B(\bar{3}'\bar{2}; 4'2'), \quad (\text{A6})$$

the pp and eh BSEs are⁴

$$\begin{aligned} F &= \Gamma^{\text{pp}} - \frac{1}{2} \Gamma^{\text{pp}} \circ_{\text{pp}} F, \\ F &= \Gamma^{\text{eh}} + \Gamma^{\text{eh}} \circ_{\text{eh}} F. \end{aligned} \quad (\text{A7})$$

At second order, the reducible vertices are

$$\begin{aligned} \Phi^{\text{eh},(2)} &= \Gamma^{(1)} \circ_{\text{eh}} \Gamma^{(1)}, \\ \Phi^{\text{pp},(2)} &= -\frac{1}{2} \Gamma^{(1)} \circ_{\text{pp}} \Gamma^{(1)}, \end{aligned} \quad (\text{A8})$$

and at third order

$$\begin{aligned} \Phi^{\text{eh},(3)} &= \Gamma^{\text{eh},(2)} \circ_{\text{eh}} \Gamma^{(1)} + \Gamma^{(1)} \circ_{\text{eh}} \Gamma^{\text{eh},(2)} \\ &\quad + \Gamma^{(1)} \circ_{\text{eh}} \Gamma^{(1)} \circ_{\text{eh}} \Gamma^{(1)}, \\ \Phi^{\text{pp},(3)} &= -\frac{1}{2} (\Gamma^{\text{pp},(2)} \circ_{\text{pp}} \Gamma^{(1)} + \Gamma^{(1)} \circ_{\text{pp}} \Gamma^{\text{pp},(2)}) \\ &\quad + \frac{1}{4} \Gamma^{(1)} \circ_{\text{pp}} \Gamma^{(1)} \circ_{\text{pp}} \Gamma^{(1)}. \end{aligned} \quad (\text{A9})$$

Together with Eqs. (A3) and (A4) and fermionic crossing,

$$\Phi^{\text{eh}}(12; 1'2') = -\Phi^{\text{eh}}(21; 1'2'), \quad (\text{A10})$$

this gives $\Gamma^{\lambda, (2)}$ and $\Gamma^{\lambda, (3)}$.

Appendix B: Pair and three-particle propagators

Here we derive the propagators stated in Sec. III B.

1. Notation

We use real canonical spin-orbitals and map numbers to orbitals with $1, 2, 1', 2' \rightarrow p, q, r, s$. Since $\bar{v}$, Eq. (A2), is instantaneous,

$$\bar{v}(12; 1'2') \rightarrow \langle pq||rs \rangle \delta(t_1 - t_2) \delta(t_1 - t_{1'}) \delta(t_2 - t_{2'}), \quad (\text{B1})$$

with

$$\langle pq||rs \rangle = \langle pq|rs \rangle - \langle pq|sr \rangle. \quad (\text{B2})$$

The antisymmetrizer of a pair of indices is

$$\hat{P}_{ab}^- = 1 - \hat{P}_{ab}, \quad (\text{B3})$$

where $\hat{P}_{ab}$ exchanges a and b . In canonical spin-orbitals p with energies ϵ_p , occupations n_p ($n_i = 1$, $n_a = 0$, $\bar{n}_p = 1 - n_p$), and $\sigma_p = \bar{n}_p - n_p$, the HF Green's function is

$$G_{\text{HF}, p}(\omega) = \frac{\bar{n}_p}{\omega - \epsilon_p + i0^+} + \frac{n_p}{\omega - \epsilon_p - i0^+}. \quad (\text{B4})$$

2. Pair problems and reducible vertices

In the own-channel approximation, each pair scatters with the bare kernel. In the eh channel this is the TDHF problem, in the pp channel the ppRPA problem,

$$\eta^\lambda \mathbf{H}^\lambda Z_n = \Omega_n Z_n, \quad \eta^\lambda = \text{diag}(\mathbf{1}, -\mathbf{1}), \quad (\text{B5})$$

with

$$\mathbf{H}^{\text{eh}} = \begin{pmatrix} A^{\text{eh}} & B^{\text{eh}} \\ B^{\text{eh}} & A^{\text{eh}} \end{pmatrix}, \quad (\text{B6})$$

$$A_{ia,jb}^{\text{eh}} = (\epsilon_a - \epsilon_i) \delta_{ab} \delta_{ij} + \langle aj||ib \rangle,$$

$$B_{ia,jb}^{\text{eh}} = \langle ab||ij \rangle,$$

and

$$\mathbf{H}^{\text{pp}} = \begin{pmatrix} A^{\text{pp}} & B^{\text{pp}} \\ B^{\text{pp}\dagger} & D^{\text{pp}} \end{pmatrix},$$

$$A_{cd,ef}^{\text{pp}} = (\epsilon_c + \epsilon_d) \delta_{ce} \delta_{df} + \langle cd||ef \rangle,$$

$$B_{cd,kl}^{\text{pp}} = \langle cd||kl \rangle,$$

$$D_{kl,mn}^{\text{pp}} = -(\epsilon_k + \epsilon_l) \delta_{km} \delta_{ln} + \langle kl||mn \rangle, \quad (\text{B7})$$

with $c < d$, $k < l$. The eh eigenvalues come in pairs $\pm \Omega_n$, with eigenvectors $Z_n = (X_n, Y_n)$ and $\bar{Z}_n = (Y_n, X_n)$. The pp spectrum is not symmetric: the positive-norm eigenvectors are the $N + 2$ states (Ω_n^+), the negative-norm ones the $N - 2$ states (Ω_n^-).

The pair propagators give the reducible vertices of the own-channel approximation, the resummed counterparts of $\Phi^{\lambda, (2)}$ in Eq. (A8), as sums over pair excitations with the two-body pair vertices M^{eh} and M^{pp} ,

$$i\Phi_{pqrs}^{\text{eh}}(\Omega) = \sum_{\Omega_n > 0} \left[\frac{M_{pr,n}^{\text{eh}} M_{qs,n}^{\text{eh}}}{\Omega - \Omega_n + i0^+} - \frac{M_{rp,n}^{\text{eh}} M_{sq,n}^{\text{eh}}}{\Omega + \Omega_n - i0^+} \right],$$

$$M_{pr,n}^{\text{eh}} = \sum_{ia} \left[\langle pa||ri \rangle X_{ia}^n + \langle pi||ra \rangle Y_{ia}^n \right], \quad (\text{B8})$$

and

$$i\Phi_{pqrs}^{\text{pp}}(\Omega) = \sum_{n \in N+2} \frac{M_{pq,n}^{\text{pp}} M_{rs,n}^{\text{pp}}}{\Omega - \Omega_n^+ + i0^+} - \sum_{m \in N-2} \frac{M_{pq,m}^{\text{pp}} M_{rs,m}^{\text{pp}}}{\Omega - \Omega_m^- - i0^+},$$

$$M_{pq,n}^{\text{pp}} = \sum_{c < d} \langle pq||cd \rangle X_{cd}^n + \sum_{k < l} \langle pq||kl \rangle Y_{kl}^n. \quad (\text{B9})$$

For real orbitals the backward eh vertex is the forward one with exchanged orbital indices, $M_{rp,n}^{\text{eh}}$. The same first-order vertex appears in the vertex-corrected GW self-energies of Refs. 32,74,77. In the eh variables, Φ^{eh} depends only on the bosonic transfer frequency Ω ; the crossed channel, eh, follows by crossing, Eq. (A10).

3. Pair propagators

The non-interacting particle-particle propagator is

$$L_0^{\text{pp}}(12; 1'2') = G(11')G(22') - G(12')G(21'), \quad (\text{B10})$$

and performing the frequency integrations,

$$L_{0,xy}^{\text{pp}}(\Omega) = \int \frac{d\nu}{2\pi} G_x(\nu) G_y(\Omega - \nu)$$

$$= \frac{-i\sigma_x \delta_{\sigma_x \sigma_y}}{\Omega - (\epsilon_x + \epsilon_y) + i\sigma_x 0^+}, \quad (\text{B11})$$

with its indices restricted to $(xy) \in \{(cd), c < d\} \cup \{(kl), k < l\}$. The particle-hole propagator is

$$L_0^{\text{eh}}(31; 1'3') = G(33')G(11'). \quad (\text{B12})$$

After performing the frequency integrations, we obtain

$$L_{0,xr}^{\text{eh}}(\Omega) = \int \frac{d\nu}{2\pi} G_x(\nu) G_r(\nu + \Omega)$$

$$= \frac{i\sigma_r \delta_{\sigma_x, -\sigma_r}}{\Omega - (\epsilon_r - \epsilon_x) + i\sigma_r 0^+}, \quad (\text{B13})$$

with its indices restricted to $(xr) \in \{(ia)\} \cup \{(ai)\}$.

The interacting propagators follow from the own-channel BSEs, Eq. (24). Restricting the pp pairs to $c < d$, $k < l$ absorbs

the $\frac{1}{2}$ of the pp BSE, and $i\Gamma_{pqrs}^{(1)} = \langle pq||rs \rangle$ by Eqs. (A2) and (B1). With the pair Hamiltonians of Eqs. (B6) and (B7),

$$L^{\text{eh}}(\Omega) = i(\Omega\eta^{\text{eh}} - \mathbf{H}^{\text{eh}})^{-1},$$

$$L^{\text{pp}}(\Omega) = -i(\Omega\eta^{\text{pp}} - \mathbf{H}^{\text{pp}})^{-1},$$

with the spectral representations

$$L^{\text{eh}}(\Omega) = i \sum_{\Omega_n > 0} \left[\frac{Z_n Z_n^\dagger}{\Omega - \Omega_n + i0^+} - \frac{\tilde{Z}_n \tilde{Z}_n^\dagger}{\Omega + \Omega_n - i0^+} \right] \quad (\text{B14})$$

and

$$L^{\text{pp}}(\Omega) = -i \sum_{n \in N+2} \frac{Z_n Z_n^\dagger}{\Omega - \Omega_n^+ + i0^+} + i \sum_{m \in N-2} \frac{Z_m Z_m^\dagger}{\Omega - \Omega_m^- - i0^+}. \quad (\text{B15})$$

Inserting them into the own-channel reducible vertices gives Eqs. (B8) and (B9).

4. Three-particle propagators

For the pp pair, Eqs. (14), (16) and (B10) give

$$\begin{aligned} \mathcal{G} + \mathcal{K}_{\text{pp}}[F_{\text{oc}}^{\text{pp}}]\mathcal{G} &= G(33') \left[L_0^{\text{pp}} - \frac{1}{4} L_0^{\text{pp}} F_{\text{oc}}^{\text{pp}} L_0^{\text{pp}} \right] (12; 1'2') \\ &= G(33') L^{\text{pp}}(12; 1'2'), \end{aligned} \quad (\text{B16})$$

Working out the frequency integral with Eqs. (B4) and (B15), with r the spectator,

$$\begin{aligned} Q_{\text{pp};xyr,uwr}(\omega) &= \int \frac{d\Omega}{2\pi} L_{xy,uw}^{\text{pp}}(\Omega) G_r(\Omega - \omega) \\ &= n_r \sum_{n \in N+2} \frac{Z_{xy}^n Z_{uw}^n}{\omega + \epsilon_r - \Omega_n^+ + i0^+} \\ &\quad + \bar{n}_r \sum_{m \in N-2} \frac{Z_{xy}^m Z_{uw}^m}{\omega + \epsilon_r - \Omega_m^- - i0^+}. \end{aligned}$$

which is Eq. (52).

For the eh pair, Eqs. (14), (16) and (B12) give

$$\begin{aligned} \mathcal{G} + \mathcal{K}_{\text{eh}}[F_{\text{oc}}^{\text{eh}}]\mathcal{G} &= G(22') \left[L_0^{\text{eh}} + L_0^{\text{eh}} F_{\text{oc}}^{\text{eh}} L_0^{\text{eh}} \right] (31; 1'3') \\ &\quad - G(21') \left[L_0^{\text{eh}} + L_0^{\text{eh}} F_{\text{oc}}^{\text{eh}} L_0^{\text{eh}} \right] (31; 2'3') \\ &= G(22') L^{\text{eh}}(31; 1'3') - G(21') L^{\text{eh}}(31; 2'3'), \end{aligned} \quad (\text{B17})$$

The propagator of the eh pair follows with $1 \leftrightarrow 2$. Unlike the pp pair, the eh pair is not antisymmetrized in $\mathcal{G}$, so that the exchange of the lines 1 and 2 appears as the second term. Working out the frequency integral with Eqs. (B4) and (B14),

$$\begin{aligned} Q_{\text{eh};xrs,yus}(\omega) &= \int \frac{d\Omega}{2\pi} L_{xr,yu}^{\text{eh}}(\Omega) G_s(\omega - \Omega) \\ &= \bar{n}_s \sum_{\Omega_n > 0} \frac{Z_{xr}^n Z_{yu}^n}{\omega - \epsilon_s - \Omega_n + i0^+} \\ &\quad + n_s \sum_{\Omega_n > 0} \frac{\tilde{Z}_{xr}^n \tilde{Z}_{yu}^n}{\omega - \epsilon_s + \Omega_n - i0^+}. \end{aligned}$$

which is Eq. (53). With L_0^{pp} in place of L^{pp} , Eq. (52) gives the free propagation $\mathcal{G}$, Eq. (45).

Appendix C: Riccati equations and decoupling conditions

Consider a space $\mathcal{P} \oplus \mathcal{Q}$, where $\mathcal{P}$ is some active space. Let $x \in \mathcal{P} \oplus \mathcal{Q}$ be an eigenvector of H and write

$$x = \begin{pmatrix} X \\ Y \end{pmatrix} = \begin{pmatrix} X \\ TX \end{pmatrix} \quad (\text{C1})$$

which implies $Y = TX$. We then define the graph of T as

$$\mathcal{P}_T = \text{graph}(T) = \left\{ \begin{pmatrix} X \\ TX \end{pmatrix}, X \in \mathcal{P} \right\} \quad (\text{C2})$$

and require it to be invariant under the action of the partitioned Hamiltonian

$$\begin{pmatrix} H_{PP} & H_{PQ} \\ H_{QP} & H_{QQ} \end{pmatrix} \begin{pmatrix} X \\ Y \end{pmatrix} = \begin{pmatrix} H_{PP} & H_{PQ} \\ H_{QP} & H_{QQ} \end{pmatrix} \begin{pmatrix} X \\ TX \end{pmatrix} = \begin{pmatrix} (H_{PP} + H_{PQ}T)X \\ (H_{QP} + H_{QQ}T)X \end{pmatrix}. \quad (\text{C3})$$

Invariance under the action of H then gives the condition

$$(H_{QP} + H_{QQ}T)X = T(H_{PP} + H_{PQ}T)X, \quad (\text{C4})$$

and as this must be valid for all $X \in \mathcal{P}$, this yields the Riccati equation

$$H_{QP} + H_{QQ}T = TH_{PP} + TH_{PQ}T. \quad (\text{C5})$$

The rewriting of the pp and eh-RPA equations, Eqs. (B7) and (B6), as the Riccati equations, Eq. (56), is easily seen to be a special case of this general transformation. Equivalently, with

$$\Theta = \begin{pmatrix} \mathbf{1} & 0 \\ T & \mathbf{1} \end{pmatrix}, \quad \Theta^{-1} = \begin{pmatrix} \mathbf{1} & 0 \\ -T & \mathbf{1} \end{pmatrix}, \quad (\text{C6})$$

whose first block column spans $\mathcal{P}_T$,

$$\Theta^{-1}H\Theta = \begin{pmatrix} H_{PP} + H_{PQ}T & H_{PQ} \\ H_{QP} + H_{QQ}T - TH_{PP} - TH_{PQ}T & H_{QQ} - TH_{PQ} \end{pmatrix}. \quad (\text{C7})$$

The lower-left block is the Riccati equation, and the invariance of $\mathcal{P}_T$ is the decoupling condition

$$Q\Theta^{-1}H\Theta P = 0, \quad (\text{C8})$$

with P and Q the projectors onto $\mathcal{P}$ and $\mathcal{Q}$. $\Theta^{-1}H\Theta$ is then block upper triangular, and $H_{PP} + H_{PQ}T$ carries the eigenvalues of H on $\mathcal{P}_T$. For $H = \eta^l \mathbf{H}^l$ of Eqs. (B6) and (B7), i.e. $H_{PP} = A_\alpha$, $H_{PQ} = B_\alpha$, $H_{QP} = -B_\alpha^\dagger$ and $H_{QQ} = -D_\alpha$, the lower-left block is minus the left-hand side of Eq. (56).

As an example, consider the wave operator

$$\mathcal{W} = \Theta P = P + T, \quad P\mathcal{W} = P, \quad (\text{C9})$$

which maps $\mathcal{P}$ onto $\mathcal{P}_T$. The invariance of $\mathcal{P}_T$ is then the Bloch equation^{135–137}

$$H\mathcal{W} = \mathcal{W}H\mathcal{W}, \quad (\text{C10})$$

whose QP block is the Riccati equation and whose PP block is an identity. The effective Hamiltonian is the upper-left block,

$$H_T = P\mathcal{W}H\mathcal{W} = P\Theta^{-1}H\Theta P = H_{PP} + H_{PQ}T. \quad (\text{C11})$$

For $H = H_0 + V$ with $[H_0, P] = 0$, Eq. (C10) becomes the generalized Bloch equation

$$[\mathcal{W}, H_0] = V\mathcal{W} - \mathcal{W}PV\mathcal{W}, \quad (\text{C12})$$

the starting point of Rayleigh–Schrödinger perturbation theory for $\mathcal{W}$. For a one-dimensional $\mathcal{P}$ spanned by Φ , T contains the CI coefficients of $\Psi = \mathcal{W}\Phi$ in intermediate normalization, $H_T = E$, and Eq. (C10) is the Schrödinger equation projected onto $\mathcal{Q}$.

1. Comparison to Löwdin downfolding

The Bloch equation should be compared with Löwdin downfolding, which generates the frequency-dependent effective Hamiltonian acting in $\mathcal{P}$ -space only,

$$H_L(\omega) = H_{PP} + H_{PQ}(\omega - H_{QQ})^{-1}H_{QP}, \quad (\text{C13})$$

Comparison with Eq. (C11) then shows that

$$T(\omega) = (\omega - H_{QQ})^{-1}H_{QP}. \quad (\text{C14})$$

$H_L(\omega)$ is Hermitian for Hermitian H and has the form of a self-energy, and $\det(\omega - H_L(\omega)) = 0$ yields all eigenvalues of H with a non-vanishing $\mathcal{P}$ component, but it is energy-dependent and has to be solved root by root. The Bloch amplitude is instead a single, energy-independent map for the $\dim \mathcal{P}$ selected roots,

$$T = (T(\omega_1)x_1, \dots, T(\omega_{\dim \mathcal{P}})x_{\dim \mathcal{P}})X^{-1}, \quad (\text{C15})$$

at the price of a nonlinear equation, a non-Hermitian H_T , and one solution for each choice of roots. Expanded in V , Eq. (C13) gives Brillouin–Wigner and Eq. (C10) Rayleigh–Schrödinger perturbation theory. The two are related by a Taylor expansion of $T(\omega_k)$ around a reference energy E_0 : with $g_0 = (E_0 - H_{QQ})^{-1}$ and $\omega_k x_k = H_T x_k$,

$$T = \sum_{n \geq 0} g_0^{n+1} H_{QP} (E_0 - H_T)^n, \quad (\text{C16})$$

i.e. the energy dependence of $T(\omega)$ becomes the dependence on H_T , which is the same for all selected roots.

Appendix D: The ADC hierarchy

In this appendix, we work out ADC(3) explicitly, as an illustrative case of Sec. II D. ADC(1), *i.e.* HF, needs no three-particle propagator at all, and ADC(2), *i.e.* GF2, only the 2p1h propagator $\mathcal{G}$.

1. ADC(3) from the first-order Faddeev kernel

We here derive the ADC(3) equation starting from Eq. (13) with $R_{\text{ADC}(3)}$ of Eq. (23), *i.e.* with the first-order Faddeev kernel acting once on $\mathcal{G}$. We then collect all diagrams by the

pair on which $\mathcal{K}^{(1)}$ acts, *i.e.* in the pp and eh channels. This will give us the GF3 self-energy. Then we use Schirmer’s approach of imposing the resolvent form of the self-energy and diagrammatic matching to get ADC(3).

The self-energy to third order is

$$\Sigma - \Sigma_{\text{HF}}[G_{\text{HF}}] = \Sigma^{(2)} + \Sigma^{(3)} + O(v^4). \quad (\text{D1})$$

The Green’s function is expanded to second order, which will later give the static self-energy to third order (notice that the $G_{\text{HF}}\Sigma_{\text{HF}}G_{\text{HF}}\Sigma_{\text{HF}}G_{\text{HF}}$ term has been absorbed in the mean field, $G^{-1} = G_{\text{HF}}^{-1} + O(v^2)$),

$$G = G_{\text{HF}} + G_{\text{HF}}\Sigma^{(2)}G_{\text{HF}} + O(v^3), \quad (\text{D2})$$

$\mathcal{G}$ is built from G_{HF} ,

$$\mathcal{G} = \mathcal{G}_{\text{HF}} + O(v^2). \quad (\text{D3})$$

For ADC(3), we must consider the first-order Faddeev kernel,

$$\mathcal{K} = \mathcal{K}^{(1)} + O(v^2), \quad (\text{D4})$$

that only contains the bare antisymmetrized interactions in all channels, Eq. (A2). Recall that ADC(3) follows from the second line of Eq. (23), where

$$\sum_{\alpha} \mathcal{K}_{\alpha}[F^{(1)}] = \mathcal{K}^{(1)}.$$

The third-order self-energy splits into static and dynamical contributions,

$$\Sigma^{(3)}(11') = \Sigma_{\infty}^{(3)}(11') + \Sigma_{\mathcal{K}}^{(3)}(11'), \quad (\text{D5})$$

$$\Sigma_{\infty}^{(3)}(11') = -i[v(1\bar{2}; 1'\bar{2}') - v(1\bar{2}; \bar{2}'1')] \times [G_{\text{HF}}\Sigma^{(2)}G_{\text{HF}}](\bar{2}'\bar{2}^+), \quad (\text{D6})$$

$$\Sigma_{\mathcal{K}}^{(3)}(11') = v(1\bar{2}; \bar{3}'\bar{2}')G_{\text{HF}}(\bar{3}\bar{4})v(\bar{4}\bar{5}; \bar{6}'\bar{7}') \times [\mathcal{K}^{(1)}\mathcal{G}_{\text{HF}}](\bar{2}'\bar{6}'\bar{7}'; \bar{3}\bar{2}\bar{5}^+) \times G_{\text{HF}}^{-1}(\bar{3}1'). \quad (\text{D7})$$

Eq. (D6) is the HF self-energy evaluated with the second-order density matrix,

$$\Sigma_{\infty, pq}^{(3)} = \sum_{rs} \langle pr || qs \rangle \gamma_{sr}^{(2)}. \quad (\text{D8})$$

Defining the first-order doubles

$$t_{ij}^{ab} = \frac{\langle ab || ij \rangle}{\epsilon_i + \epsilon_j - \epsilon_a - \epsilon_b}, \quad (\text{D9})$$

and second-order singles amplitudes

$$t_i^{a(2)} = \frac{1}{\epsilon_i - \epsilon_a} \left[\frac{1}{2} \sum_{kbc} t_{ik}^{bc} \langle bc || ak \rangle - \frac{1}{2} \sum_{jkc} t_{jk}^{ac} \langle ic || jk \rangle \right], \quad (\text{D10})$$

the blocks of $\gamma^{(2)}$ are²⁸

$$\begin{aligned} \gamma_{ij}^{(2)} &= -\frac{1}{2} \sum_{kab} t_{ik}^{ab} t_{jk}^{ab}, \\ \gamma_{ab}^{(2)} &= \frac{1}{2} \sum_{ijc} t_{ij}^{ac} t_{ij}^{bc}, \\ \gamma_{ia}^{(2)} &= \gamma_{ai}^{(2)} = t_i^{a(2)}. \end{aligned} \quad (\text{D11})$$

Using Eqs. (15) and (16) in Eq. (D7) with $\chi = \mathcal{G}$, we obtain the contributions of the three channels to $\Sigma_{\mathcal{K}}^{(3)}$. Here and below, $G \equiv G_{\text{HF}}$, and all interactions appear antisymmetrized, Eq. (A2). For the pp channel, we get

$$\begin{aligned} \Sigma_{\mathcal{K}}^{\text{pp}}(11') &= \frac{i}{4} \bar{v}(1\bar{2}; \bar{3}'\bar{2}') G(\bar{3}'\bar{4}) G(\bar{2}'\bar{5}^+) \\ &\quad \times \bar{v}(\bar{4}\bar{5}; \bar{6}'\bar{7}') G(\bar{6}'\bar{8}) G(\bar{7}'\bar{9}) \\ &\quad \times \bar{v}(\bar{8}\bar{9}; 1'\bar{9}') G(\bar{9}'\bar{2}) . \end{aligned} \quad (\text{D12})$$

For the eh, $\bar{\text{eh}}$ channels, we obtain

$$\begin{aligned} \Sigma_{\mathcal{K}}^{\text{eh}}(11') &= -\frac{i}{2} \bar{v}(1\bar{2}; \bar{3}'\bar{2}') G(\bar{3}'\bar{4}) \bar{v}(\bar{4}\bar{5}; \bar{6}'\bar{7}') \\ &\quad \times G(\bar{2}'\bar{8}) G(\bar{6}'\bar{9}) \bar{v}(\bar{8}\bar{9}; 1'\bar{8}') \\ &\quad \times G(\bar{8}'\bar{5}^+) G(\bar{7}'\bar{2}) = \Sigma_{\mathcal{K}}^{\bar{\text{eh}}}(11') . \end{aligned} \quad (\text{D13})$$

In the time domain, the non-interacting pair propagators follow from $G_{\text{HF},p}(t) = -i\sigma_p \theta(\sigma_p t) e^{-i\epsilon_p t}$,

$$L_{0,xy}^{\text{pp}}(t) = G_x(t) G_y(t) = -\delta_{\sigma_x \sigma_y} \theta(\sigma_x t) e^{-i(\epsilon_x + \epsilon_y)t} , \quad (\text{D14})$$

and

$$L_{0,xr}^{\text{eh}}(t) = G_x(-t) G_r(t) = \delta_{\sigma_x, -\sigma_r} \theta(\sigma_r t) e^{-i(\epsilon_r - \epsilon_x)t} . \quad (\text{D15})$$

Exploiting the temporal structure of $\bar{v}$ and the space-time to orbital index mapping of Eq. (B1), and with Eq. (D14) the pp term becomes

$$\begin{aligned} \Sigma_{\mathcal{K},pq}^{\text{pp}}(t_1, t_1') &= \frac{i}{4} \sum_{rxyuw} \langle pr||xy \rangle \langle xy||uw \rangle \langle uw||qr \rangle \\ &\quad \times \int d\tau L_{0,xy}^{\text{pp}}(t_1 - \tau) L_{0,uw}^{\text{pp}}(\tau - t_1') \\ &\quad \times G_r(t_1' - t_1) , \end{aligned} \quad (\text{D16})$$

in the orbital basis. Summing up the identical eh and $\bar{\text{eh}}$ channels and with Eq. (D15), the sum of Σ^{eh} and $\Sigma^{\bar{\text{eh}}}$ becomes

$$\begin{aligned} \Sigma_{\mathcal{K},pq}^{\text{eh}+\bar{\text{eh}}}(t_1, t_1') &= -i \sum_{rsxyu} \langle pr||xy \rangle \langle xs||ur \rangle \langle yu||qs \rangle \\ &\quad \times \int d\tau L_{0,rx}^{\text{eh}}(t_1 - \tau) L_{0,su}^{\text{eh}}(\tau - t_1') \\ &\quad \times G_y(t_1 - t_1') . \end{aligned} \quad (\text{D17})$$

Not coincidentally, one can identify the second-order pp and eh reducible vertices, Eq. (A8), of the parquet equations.

a. GF3 self-energy

We perform the integrals over τ by regarding each space-time term as a circuit and using energy conservation at the intermediate vertices, which leaves contour integrals over the channel frequency. For the pp channel, one then gets

$$\begin{aligned} \Sigma_{\mathcal{K},pq}^{\text{pp}}(\omega) &= \frac{i}{4} \sum_{rxyuw} \langle pr||xy \rangle \langle xy||uw \rangle \langle uw||qr \rangle \\ &\quad \times \int \frac{d\Omega}{2\pi} L_{0,xy}^{\text{pp}}(\Omega) L_{0,uw}^{\text{pp}}(\Omega) G_r(\Omega - \omega) , \end{aligned} \quad (\text{D18})$$

with Eq. (B11). In the eh + $\bar{\text{eh}}$ channel, Eq. (D17), we obtain

$$\begin{aligned} \Sigma_{\mathcal{K},pq}^{\text{eh}+\bar{\text{eh}}}(\omega) &= -i \sum_{rsxyu} \langle pr||xy \rangle \langle xs||ur \rangle \langle yu||qs \rangle \\ &\quad \times \int \frac{d\Omega}{2\pi} L_{0,rx}^{\text{eh}}(\Omega) L_{0,su}^{\text{eh}}(\Omega) G_y(\omega - \Omega) , \end{aligned} \quad (\text{D19})$$

with Eq. (B13). In both channels, the kernel factorizes into two one-frequency pair propagators in the channel transfer frequency Ω ; only the Ω integral couples them to the spectator line (r in pp, y in eh). What remains is to perform the contour integrals over Ω . For the pp channel, this gives

$$\Sigma_{\mathcal{K},pq}^{\text{pp}}(\omega) = \frac{1}{4} \sum_{abcdi} \frac{\langle pi||ab \rangle \langle ab||cd \rangle \langle cd||qi \rangle}{(\omega - \Delta_{abi})(\omega - \Delta_{cdi})} \quad (\text{D20})$$

$$+ \frac{1}{4} \sum_{abkli} \frac{\langle pi||ab \rangle t_{kl}^{ab} \langle kl||qi \rangle}{\omega - \Delta_{abi}} \quad (\text{D21})$$

$$+ \frac{1}{4} \sum_{cdkli} \frac{\langle pi||kl \rangle t_{kl}^{cd} \langle cd||qi \rangle}{\omega - \Delta_{cdi}} \quad (\text{D22})$$

$$- \frac{1}{4} \sum_{ijkla} \frac{\langle pa||ij \rangle \langle ij||kl \rangle \langle kl||qa \rangle}{(\omega - \Delta_{ija})(\omega - \Delta_{kla})} \quad (\text{D23})$$

$$+ \frac{1}{4} \sum_{bckla} \frac{\langle pa||bc \rangle t_{kl}^{bc} \langle kl||qa \rangle}{\omega - \Delta_{kla}} \quad (\text{D24})$$

$$+ \frac{1}{4} \sum_{ijbca} \frac{\langle pa||ij \rangle t_{ij}^{bc} \langle bc||qa \rangle}{\omega - \Delta_{ija}} . \quad (\text{D25})$$

For the eh + $\bar{\text{eh}}$ channel,

$$\Sigma_{\mathcal{K},pq}^{\text{eh}+\bar{\text{eh}}}(\omega) = \sum_{abcij} \frac{\langle pi||ab \rangle \langle aj||ci \rangle \langle bc||qj \rangle}{(\omega - \Delta_{abi})(\omega - \Delta_{bcj})} \quad (\text{D26})$$

$$- \sum_{abcik} \frac{\langle pi||ab \rangle t_{ik}^{ac} \langle bk||qc \rangle}{\omega - \Delta_{abi}} \quad (\text{D27})$$

$$+ \sum_{abcjk} \frac{\langle pc||ja \rangle t_{jk}^{bc} \langle ab||qk \rangle}{\omega - \Delta_{abk}} \quad (\text{D28})$$

$$- \sum_{ijkab} \frac{\langle pb||ij \rangle \langle ia||kb \rangle \langle jk||qa \rangle}{(\omega - \Delta_{ijb})(\omega - \Delta_{jka})} \quad (\text{D29})$$

$$- \sum_{ijkab} \frac{\langle pj||ai \rangle t_{jk}^{ab} \langle ik||qb \rangle}{\omega - \Delta_{ikb}} \quad (\text{D30})$$

$$+ \sum_{ijkab} \frac{\langle pb||ij \rangle t_{ik}^{ab} \langle ja||qk \rangle}{\omega - \Delta_{ijb}} . \quad (\text{D31})$$

We have used the shorthand of Eq. (46). These terms constitute the complete dynamic GF3 self-energy, which is for instance given in the appendix of Ref. 21.

b. From GF3 to ADC(3)

What remains is to match the GF3 terms with the ADC(3) ones, following the method introduced by Schirmer.¹¹ The

procedure is rather simple: we impose the resolvent structure on the self-energy

$$\Sigma_{\text{ADC}(3)}(\omega) = \Sigma(\infty) + U^{\text{II}}(\omega - K^{\text{II}} - C^{\text{II}})^{-1} U^{\text{II}\dagger} + U^{\text{I}}(\omega + K^{\text{I}} + C^{\text{I}})^{-1} U^{\text{I}\dagger}, \quad (\text{D32})$$

with the static self-energy $\Sigma(\infty) = \Sigma_{\text{HF}} + \Sigma_{\infty}^{(3)}$ and

$$U^{\text{II}} = U^{\text{II}(1)} + U^{\text{II}(2)}, \quad U^{\text{I}} = U^{\text{I}(1)} + U^{\text{I}(2)}. \quad (\text{D33})$$

All terms are then classified according to their number of dynamical energy denominators. The ones that only contain one go inside U , and the ones that contain two go inside C . Terms with the dynamical denominator $\omega - \Delta_{abi} (2p1h, N+1)$ belong to block II, terms with $\omega - \Delta_{ija} (2h1p, N-1)$ to block I. We use the zeroth-order matrices and first-order couplings of Eq. (48). The unrestricted sums of the GF3 terms are converted with

$$\begin{aligned} \sum_{a<b} \sum_{c<d} x_{ab} \hat{P}_{ab}^- \hat{P}_{cd}^- f_{ab,cd} y_{cd} &= \sum_{abcd} x_{ab} f_{ab,cd} y_{cd}, \\ \sum_{a<b} x_{ab} \hat{P}_{ab}^- g_{ab} &= \sum_{ab} x_{ab} g_{ab}, \end{aligned} \quad (\text{D34})$$

for x, y antisymmetric in their pair indices, with $\hat{P}_{ab}^-$ of Eq. (B3).

In block II, the two-denominator terms, Eqs. (D20) and (D26), combine into

$$\begin{aligned} \Sigma_C^{\text{II}}(\omega) &= U^{\text{II}(1)}(\omega - K^{\text{II}})^{-1} C^{\text{II}}(\omega - K^{\text{II}})^{-1} U^{\text{II}(1)\dagger}, \\ C_{abi,cdj}^{\text{II}} &= \langle ab||cd \rangle \delta_{ij} - \hat{P}_{ab}^- \hat{P}_{cd}^- \langle a||c \rangle \delta_{bd}, \end{aligned} \quad (\text{D35})$$

where we used $\langle bc||qj \rangle = -\langle qj||cb \rangle$. The one-denominator terms, Eqs. (D21), (D22), (D27) and (D28), give

$$\begin{aligned} \Sigma_U^{\text{II}}(\omega) &= U^{\text{II}(2)}(\omega - K^{\text{II}})^{-1} U^{\text{II}(1)\dagger} + U^{\text{II}(1)}(\omega - K^{\text{II}})^{-1} U^{\text{II}(2)\dagger}, \\ U_{p;abi}^{\text{II}(2)} &= \frac{1}{2} \sum_{jl} t_{jl}^{ab} \langle p||j \rangle - \hat{P}_{ab}^- \sum_{kc} t_{ki}^{ac} \langle p||k \rangle, \end{aligned} \quad (\text{D36})$$

where we used $t_{ik}^{ac} \langle bk||qc \rangle = t_{ki}^{ac} \langle qc||kb \rangle$.

In block I, the two-denominator terms, Eqs. (D23) and (D29), combine into

$$\begin{aligned} \Sigma_C^{\text{I}}(\omega) &= -U^{\text{I}(1)}(\omega + K^{\text{I}})^{-1} C^{\text{I}}(\omega + K^{\text{I}})^{-1} U^{\text{I}(1)\dagger}, \\ C_{ija,klb}^{\text{I}} &= \langle ij||kl \rangle \delta_{ab} - \hat{P}_{ij}^- \hat{P}_{kl}^- \langle ib||ka \rangle \delta_{jl}, \end{aligned} \quad (\text{D37})$$

where we used $\langle jk||qa \rangle = -\langle qa||kj \rangle$. The sign is such that Σ_C^{I} is the first-order term of $U^{\text{I}(1)}(\omega + K^{\text{I}} + C^{\text{I}})^{-1} U^{\text{I}(1)\dagger}$. The one-denominator terms, Eqs. (D24), (D25), (D30) and (D31), give

$$\begin{aligned} \Sigma_U^{\text{I}}(\omega) &= U^{\text{I}(2)}(\omega + K^{\text{I}})^{-1} U^{\text{I}(1)\dagger} + U^{\text{I}(1)}(\omega + K^{\text{I}})^{-1} U^{\text{I}(2)\dagger}, \\ U_{p;ija}^{\text{I}(2)} &= \frac{1}{2} \sum_{bc} t_{ij}^{bc} \langle pa||bc \rangle - \hat{P}_{ij}^- \sum_{kb} t_{ik}^{ba} \langle pk||bj \rangle. \end{aligned} \quad (\text{D38})$$

With Eqs. (D35)–(D38) and $\Sigma(\infty)$, all third-order terms are reproduced by Eq. (D32), which completes ADC(3).

- ¹L. D. Faddeev, Soviet Physics JETP **12**, 1014 (1961), russian original: Zh. Eksp. Teor. Fiz. **39**, 1459–1467 (1960). Reprinted in *Fifty Years of Mathematical Physics: Selected Works of Ludwig Faddeev*, World Scientific, 2016, pp. 37–42.
- ²J. Winter, Nuclear Physics A **194**, 535 (1972).
- ³S. Ethofer and P. Schuck, Z. Physik **228**, 264 (1969).
- ⁴F. B. Kugler, *Renormalization group approaches to strongly correlated electron systems*, Ph.D. thesis, Ludwig-Maximilians-Universität München (2018).
- ⁵C. Barbieri and W. H. Dickhoff, Physical Review C - Nuclear Physics **63**, 034313 (2001).
- ⁶C. Barbieri and W. Dickhoff, Physical Review C **68**, 014311 (2003).
- ⁷C. Barbieri, D. Van Neck, and W. Dickhoff, Physical Review A—Atomic, Molecular, and Optical Physics **76**, 052503 (2007).
- ⁸M. Degroote, D. V. Neck, and C. Barbieri, Physical Review A **83**, 042517 (2011).
- ⁹G. E. Scuseria, T. M. Henderson, and I. W. Bulik, The Journal of Chemical Physics **139**, 104113 (2013), arXiv:1306.6360.
- ¹⁰P. C. Martin and J. Schwinger, Physical Review **115**, 1342 (1959).
- ¹¹J. Schirmer, L. S. Cederbaum, and O. Walter, Physical Review A **28**, 1237 (1983).
- ¹²K. M. Watson, Physical Review **89**, 575 (1953).
- ¹³N. E. Bickers and D. J. Scalapino, Annals of Physics **193**, 206 (1989).
- ¹⁴N. E. Bickers, “Self-consistent many-body theory for condensed matter systems,” in *Theoretical Methods for Strongly Correlated Electrons*, edited by D. Sénéchal, A.-M. Tremblay, and C. Bourbonnais (Springer, 2004) pp. 237–296.
- ¹⁵R. Orlando, P. Romaniello, and P.-F. Loos, The Journal of Chemical Physics **159**, 184113 (2023).
- ¹⁶J. Tölle, M.-P. Kitsaras, A. Irmeler, A. Grüneis, and P.-F. Loos, Journal of Chemical Theory and Computation **22**, 6471 (2026).
- ¹⁷J. J. Shepherd, T. M. Henderson, and G. E. Scuseria, The Journal of Chemical Physics **140**, 124102 (2014), arXiv:1310.6806.
- ¹⁸M. F. Lange and T. C. Berkelbach, Journal of Chemical Theory and Computation **14**, 4224 (2018).
- ¹⁹R. Quintero-Monsebaiz, E. Monino, A. Marie, and P.-F. Loos, The Journal of Chemical Physics **157**, 231102 (2022).
- ²⁰J. Tölle and G. K.-L. Chan, The Journal of Chemical Physics **158**, 124123 (2023).
- ²¹W. von Niessen, J. Schirmer, and L. S. Cederbaum, Computer Physics Reports **1**, 57 (1984).
- ²²L. S. Cederbaum, J. Schirmer, W. Domcke, and W. V. Niessen, Journal of Physics B: Atomic and Molecular Physics **10**, L549 (1977).
- ²³L. Cederbaum and W. Domcke, Advances in chemical physics **36**, 205 (1977).
- ²⁴J. Schirmer and L. Cederbaum, Journal of Physics B: Atomic and Molecular Physics **11**, 1889 (1978).
- ²⁵G. Riva, P. Romaniello, and J. A. Berger, Physical Review Letters **131**, 216401 (2023).
- ²⁶G. Riva, P. Romaniello, and J. A. Berger, Physical Review B **110**, 115140 (2024).
- ²⁷G. Riva, T. Fischer, S. Paggi, J. A. Berger, and P. Romaniello, Physical Review B **111**, 195133 (2025).
- ²⁸M. Keizer, S. Paggi, J. A. Berger, P. Romaniello, and A. Förster, arXiv:2608.25669, 1 (2026).
- ²⁹A. Förster, arXiv preprint arXiv:2608.28424 (2026).
- ³⁰C. J. Coveney, Journal of Physical Chemistry A **129**, 8689 (2025).
- ³¹L. Hedin, Physical Review **139**, A796 (1965).
- ³²A. Förster and F. Bruneval, The Journal of Physical Chemistry Letters **15**, 12526 (2024).
- ³³Y. Pavlyukh, F. Bruneval, and A. Förster, arXiv:2607.29359, 1 (2026).
- ³⁴N. E. Bickers, D. J. Scalapino, and S. R. White, Physical Review Letters **62**, 961 (1989).
- ³⁵A. Marie and P.-F. Loos, J. Chem. Phys. **163**, 194115 (2025).
- ³⁶M. Springer, F. Aryasetiawan, and K. Karlsson, Physical Review Letters **80**, 2389 (1998).
- ³⁷V. P. Zhukov, E. V. Chulkov, and P. M. Echenique, Physical Review Letters **93**, 096401 (2004).
- ³⁸S. J. Bintrim and T. C. Berkelbach, The Journal of Chemical Physics **154**, 041101 (2021).

- ³⁹E. Monino and P.-F. Loos, *The Journal of Chemical Physics* **159**, 034105 (2023).
- ⁴⁰M. S. Hybertsen and S. G. Louie, *Physical Review B* **34**, 5390 (1986).
- ⁴¹M. S. Hybertsen and S. G. Louie, *Physical Review B* **35**, 5585 (1987).
- ⁴²J. E. Northrup, M. S. Hybertsen, and S. G. Louie, *Physical Review Letters* **59**, 819 (1987).
- ⁴³J. E. Northrup, M. S. Hybertsen, and S. G. Louie, *Physical Review B* **39**, 8198 (1989).
- ⁴⁴R. D. Sole, L. Reining, and R. W. Godby, *Physical Review B* **49**, 8024 (1994).
- ⁴⁵A. J. Morris, M. Stankovski, K. T. Delaney, P. Rinke, P. García-González, and R. W. Godby, *Physical Review B* **76**, 155106 (2007).
- ⁴⁶W. Chen and A. Pasquarello, *Physical Review B* **92**, 041115 (2015).
- ⁴⁷S. Sharma, J. K. Dewhurst, A. Sanna, and E. K. U. Gross, *Physical Review Letters* **107**, 186401 (2011).
- ⁴⁸P. S. Schmidt, C. E. Patrick, and K. S. Thygesen, *Physical Review B* **96**, 205206 (2017).
- ⁴⁹H. Ma, M. Govoni, F. Gygi, and G. Galli, *Journal of Chemical Theory and Computation* **15**, 154 (2019).
- ⁵⁰T. Olsen, C. E. Patrick, J. E. Bates, A. Ruzsinszky, and K. S. Thygesen, *npj Computational Materials* **5**, 106 (2019).
- ⁵¹M. L. Tiago and J. R. Chelikowsky, *Physical Review B* **73**, 205334 (2006).
- ⁵²M. J. van Setten, F. Weigend, and F. Evers, *Journal of Chemical Theory and Computation* **9**, 232 (2013).
- ⁵³L. Hung, F. H. da Jornada, J. Souto-Casares, J. R. Chelikowsky, S. G. Louie, and S. Ögüt, *Physical Review B* **94**, 085125 (2016).
- ⁵⁴L. Hung, F. Bruneval, K. Baishya, and S. Ögüt, *Journal of Chemical Theory and Computation* **13**, 2135 (2017).
- ⁵⁵G. D. Mahan and B. E. Sernelius, *Physical Review Letters* **62**, 2718 (1989).
- ⁵⁶E. L. Shirley, *Physical Review B* **54**, 7758 (1996).
- ⁵⁷M. Hindgren and C.-O. Almbladh, *Physical Review B* **56**, 12832 (1997).
- ⁵⁸Y. Takada, *Physical Review Letters* **87**, 226402 (2001).
- ⁵⁹F. Bruneval, F. Sottile, V. Olevano, R. D. Sole, and L. Reining, *Physical Review Letters* **94**, 186402 (2005).
- ⁶⁰P. Romaniello, S. Guyot, and L. Reining, *The Journal of Chemical Physics* **131**, 154111 (2009).
- ⁶¹M. Hellgren, N. Colonna, and S. de Gironcoli, *Physical Review B* **98**, 045117 (2018).
- ⁶²A. Tal, W. Chen, and A. Pasquarello, *Physical Review B* **103**, L161104 (2021).
- ⁶³A. L. Kutepov, *Physical Review B* **105**, 045124 (2022).
- ⁶⁴A. Lorin, T. Bischoff, A. Tal, and A. Pasquarello, *Physical Review B* **108**, 245303 (2023).
- ⁶⁵A. Tal, T. Bischoff, and A. Pasquarello, *Proceedings of the National Academy of Sciences* **121**, e2311472121 (2024).
- ⁶⁶M. Shishkin, M. Marsman, and G. Kresse, *Physical Review Letters* **99**, 246403 (2007).
- ⁶⁷B. Cunningham, M. Grüning, P. Azarhoosh, D. Pashov, and M. van Schilf-gaarde, *Physical Review Materials* **2**, 034603 (2018).
- ⁶⁸H. Sakakibara, T. Kotani, M. Obata, and T. Oda, *Physical Review B* **101**, 205120 (2020).
- ⁶⁹B. Cunningham, M. Grüning, D. Pashov, and M. van Schilf-gaarde, *Physical Review B* **108** (2023), 10.1103/physrevb.108.165104.
- ⁷⁰A. Förster, *Journal of Chemical Theory and Computation* **21**, 1709-1721 (2025).
- ⁷¹F. Bruneval, N. Dattani, and M. J. van Setten, *Frontiers in Chemistry* **9**, 749779 (2021).
- ⁷²C. H. Patterson, *Journal of Chemical Theory and Computation* **20**, 7479 (2024).
- ⁷³A. M. Lewis and T. C. Berkelbach, *Journal of Chemical Theory and Computation* **15**, 2925 (2019).
- ⁷⁴E. Maggio and G. Kresse, *Journal of Chemical Theory and Computation* **13**, 4765 (2017).
- ⁷⁵V. Vlček, *Journal of Chemical Theory and Computation* **15**, 6254 (2019).
- ⁷⁶C. Mejuto-Zaera, G. Weng, M. Romanova, S. J. Cotton, K. B. Whaley, N. M. Tubman, and V. Vlček, *The Journal of Chemical Physics* **154**, 121101 (2021).
- ⁷⁷S. Vacondio, D. Varsano, A. Ruini, and A. Ferretti, *Journal of Chemical Theory and Computation* **20**, 4718 (2024).
- ⁷⁸F. Bruneval and A. Förster, *Journal of Chemical Theory and Computation* **20**, 3218-3230 (2024).
- ⁷⁹R. Kuwahara and K. Ohno, *Physical Review A* **90**, 032506 (2014).
- ⁸⁰R. Kuwahara, Y. Noguchi, and K. Ohno, *Physical Review B* **94**, 121116 (2016).
- ⁸¹G. Stefanucci, Y. Pavlyukh, A.-M. Uimonen, and R. van Leeuwen, *Physical Review B* **90**, 115134 (2014).
- ⁸²C. Mejuto-Zaera and V. Vlček, *Physical Review B* **106**, 165129 (2022).
- ⁸³A. Schindlmayr and R. W. Godby, *Phys. Rev. Lett.* **80**, 1702 (1998).
- ⁸⁴P. Romaniello, F. Bechstedt, and L. Reining, *Physical Review B* **85**, 155131 (2012).
- ⁸⁵P. A. Bobbert and W. van Haeringen, *Physical Review B* **49**, 10326 (1994).
- ⁸⁶R. T. M. Ummels, P. A. Bobbert, and W. van Haeringen, *Physical Review B* **57**, 11962 (1998).
- ⁸⁷A. L. Kutepov, *Physical Review B* **94**, 155101 (2016).
- ⁸⁸A. L. Kutepov, *Physical Review B* **95**, 195120 (2017).
- ⁸⁹Y. Pavlyukh, G. Stefanucci, and R. van Leeuwen, *Physical Review B* **102**, 045121 (2020).
- ⁹⁰K. Yoneyama, Y. Noguchi, and K. Ohno, *The Journal of Chemical Physics* **161**, 154102 (2024).
- ⁹¹X. Ren, N. Marom, F. Caruso, M. Scheffler, and P. Rinke, *Physical Review B* **92**, 081104(R) (2015).
- ⁹²S. Vacondio, D. Varsano, A. Ruini, and A. Ferretti, *Journal of Chemical Theory and Computation* **18**, 3703 (2022).
- ⁹³A. Grüneis, G. Kresse, Y. Hinuma, and F. Oba, *Physical Review Letters* **112**, 096401 (2014).
- ⁹⁴A. Förster and L. Visscher, *Physical Review B* **105**, 125121 (2022).
- ⁹⁵P. Pokhilko, C. N. Yeh, M. A. Morales, and D. Zgid, *Journal of Chemical Physics* **162**, 244110 (2025).
- ⁹⁶M. Wang, M. Wen, P. Pokhilko, C.-N. Yeh, M. A. Morales, and D. Zgid, *The Journal of Chemical Physics* **165**, 084107 (2026).
- ⁹⁷D. Zhang, N. Q. Su, and W. Yang, *The Journal of Physical Chemistry Letters* **8**, 3223 (2017).
- ⁹⁸J. Li, Z. Chen, and W. Yang, *The Journal of Physical Chemistry Letters* **12**, 6203 (2021).
- ⁹⁹A. Marie, P. Romaniello, and P.-F. Loos, *Physical Review B* **110**, 115155 (2024).
- ¹⁰⁰M. P. von Friesen, C. Verdozzi, and C.-O. Almbladh, *Physical Review Letters* **103**, 176404 (2009).
- ¹⁰¹M. Puig von Friesen, C. Verdozzi, and C.-O. Almbladh, *Physical Review B* **82**, 155108 (2010).
- ¹⁰²H. Köhler, N. Kwong, and H. A. Yousif, *Computer Physics Communications* **123**, 123 (1999).
- ¹⁰³N. Schlünzen, J.-P. Joost, F. Heidrich-Meisner, and M. Bonitz, *Physical Review B* **95**, 165139 (2017).
- ¹⁰⁴N. Schlünzen, J.-P. Joost, and M. Bonitz, *Physical Review Letters* **124**, 076601 (2020).
- ¹⁰⁵J.-P. Joost, N. Schlünzen, and M. Bonitz, *Physical Review B* **101**, 245101 (2020).
- ¹⁰⁶N. Schlünzen, S. Hermanns, M. Scharnke, and M. Bonitz, *Journal of Physics: Condensed Matter* **32**, 103001 (2020).
- ¹⁰⁷P. Božek, *Physical Review C* **65**, 034327 (2002).
- ¹⁰⁸P. Božek, *Physics Letters B* **551**, 93 (2003).
- ¹⁰⁹W.-D. Kraeft, D. Kremp, W. Ebeling, and G. Röpke, *Quantum Statistics of Charged Particle Systems* (Akademie-Verlag (De Gruyter); Plenum, 1986).
- ¹¹⁰D. Kremp, M. Schlangles, and W.-D. Kraeft, *Quantum Statistics of Nonideal Plasmas*, Springer Series on Atomic, Optical, and Plasma Physics, Vol. 25 (Springer, Berlin, 2005).
- ¹¹¹M. Bonitz, *Quantum Kinetic Theory* (Springer International Publishing, 2016).
- ¹¹²B. Cunningham, *Physical Review Research* **6**, 043277 (2024).
- ¹¹³A. Toschi, A. A. Katanin, and K. Held, *Physical Review B* **75**, 045118 (2007).
- ¹¹⁴A. A. Katanin, A. Toschi, and K. Held, *Physical Review B* **80**, 075104 (2009).
- ¹¹⁵Y. M. Vil'k and A.-M. Tremblay, *Journal de Physique I* **7**, 1309 (1997).
- ¹¹⁶M. C. T. D. Müller, S. Blügel, and C. Friedrich, *Physical Review B* **100**, 045130 (2019).
- ¹¹⁷D. Nabok, S. Blügel, and C. Friedrich, *npj Computational Materials* **7**, 178 (2021).
- ¹¹⁸J.-P. Joost, N. Schlünzen, H. Ohldag, M. Bonitz, F. Lackner, and I. Březina

- nová, *Physical Review B* **105**, 165155 (2022).
- ¹¹⁹M. Bonitz, J. Joost, C. Makait, E. Schroedter, T. Kalsberger, and K. Balzer, *physica status solidi (b)* **261**, 2300578 (2024).
- ¹²⁰N. E. Bickers and S. R. White, *Physical Review B* **43**, 8044 (1991).
- ¹²¹S. X. Yang, H. Fotso, J. Liu, T. A. Maier, K. Tomko, E. F. D’Azevedo, R. T. Scalettar, T. Pruschke, and M. Jarrell, *Physical Review E* **80**, 046706 (2009).
- ¹²²Y. Pavlyukh, E. Perfetto, and G. Stefanucci, *Physical Review B* **104**, 035124 (2021).
- ¹²³Y. Pavlyukh, E. Perfetto, D. Karlsson, R. van Leeuwen, and G. Stefanucci, *Physical Review B* **105**, 125134 (2022).
- ¹²⁴Y. Pavlyukh, A.-M. Uimonen, G. Stefanucci, and R. van Leeuwen, *Physical Review Letters* **117**, 206402 (2016).
- ¹²⁵F. Bruneval, A. Förster, and Y. Pavlyukh, *Journal of Chemical Theory and Computation* **21**, 10223 (2025).
- ¹²⁶A. Marie, J. Tölle, and P.-F. Loos, *The Journal of Physical Chemistry Letters* **17**, 8541 (2026).
- ¹²⁷P.-F. Loos and P. Romaniello, *The Journal of Chemical Physics* **156**, 164101 (2022).
- ¹²⁸L. Pisani, A. Perali, P. Pieri, and G. C. Strinati, *Physical Review B* **97**, 014528 (2018).
- ¹²⁹P. Romaniello and J. A. Berger, arXiv preprint arXiv:2603.27329 (2026).
- ¹³⁰A. Marie and P. F. Loos, *Journal of Chemical Theory and Computation* **20**, 4751–4777 (2024).
- ¹³¹N. Colonna, M. Hellgren, and S. de Gironcoli, *Physical Review B* **90**, 125150 (2014).
- ¹³²A. Heßelmann and A. Görling, *Physical Review Letters* **106**, 093001 (2011).
- ¹³³I. W. Bulik, T. M. Henderson, and G. E. Scuseria, *Journal of Chemical Theory and Computation* **11**, 3171 (2015).
- ¹³⁴J. A. Gomez, T. M. Henderson, and G. E. Scuseria, *The Journal of Chemical Physics* **144**, 244117 (2016).
- ¹³⁵C. Bloch, *Nuclear Physics* **6**, 329 (1958).
- ¹³⁶B. H. Brandow, *Reviews of Modern Physics* **39**, 771 (1967).
- ¹³⁷I. Shavitt and R. J. Bartlett, *Many-Body Methods in Chemistry and Physics: MBPT and Coupled-Cluster Theory* (Cambridge University Press, Cambridge, 2009).