

Simple Example

All records for students more than 18 years old

$$\{ S \mid S \in \text{Students} \wedge S.\text{age} > 18 \}$$

The set of tuples S such that S is in the table “Students” and has component “age” least 18.

This is like **list comprehension** in Haskell

```
[ s | s <- students, age s > 18 ]
```

and similar constructions in other languages.

All are based on “comprehensions” in set theory

Tuple Relational Calculus Basics

Queries in TRC have the general form

$$\{ T \mid P(T) \}$$

where T is a *tuple variable* and $P(T)$ is a logical formula.

Every tuple variable such as T has a *schema*, like rows in a relational table, with fields and their domains. In practice, the details of the schema are usually inferred from the way T appears in $P(T)$.

A *tuple variable* ranges over all possible *tuple values* matching its schema.

The result of the query

$$\{ T \mid P(T) \}$$

is then the set of all possible tuple values for T such that $P(T)$ is true.

Another Example

Names and ages of all students over 20

$$\{ T \mid \exists S . S \in \text{Students} \wedge S.\text{age} > 20 \\ \wedge T.\text{name} = S.\text{name} \wedge T.\text{age} = S.\text{age} \}$$

The set of tuples T such that there is an S in table “Students” with component “age” at least 20 and where S and T have the same values for “name” and “age”.

- Tuple variable S has schema matching the table “Students”.
- Tuple variable T has (only) fields “name” and “age”, with domains to match those of S .
- Even if S has other fields, they do not appear in T or the overall result.

Formula Syntax

Inside TRC expression $\{T \mid P(T)\}$ the logical formula $P(T)$ may be quite long, but is built up from standard logical components.

- Simple assertions: $(T \in \text{Table})$, $(T.\text{age} > 65)$, $(S.\text{name} = T.\text{name})$, ...
- Logical combinations: $(P \vee Q)$, $(P \wedge Q \wedge \neg Q')$, ...
- Quantification:


$\exists S . P(S)$ There exists a tuple S such that $P(S)$

$\forall T . Q(T)$ For all tuples T it is true that $Q(T)$

For convenience, we require that for $\exists S . P(S)$ the variable S must actually appear in $P(S)$; and the same for $\forall T . Q(T)$. We also write:

$\exists S \in \text{Table} . P(S)$ to mean $\exists S . S \in \text{Table} \wedge P(S)$

Students and Courses (1/5)



Students and Courses (1/5)

Students taking Informatics 1

$$\{ R \mid \exists S \in \text{Students} . \exists T \in \text{Takes} . \exists C \in \text{Courses} . \\ C.\text{name} = \text{"Informatics 1"} \wedge C.\text{code} = T.\text{code} \\ \wedge T.\text{mn} = S.\text{mn} \wedge S.\text{name} = R.\text{name} \}$$

Schema for S, T and C match those of the tables from which they are drawn. The schema for result R is a single field “name” with string domain, because that’s all that appears here.

One way to compute this in relational algebra:


$$\pi_{\text{name}}((\text{Students} \bowtie \text{Takes}) \bowtie (\sigma_{\text{name}=\text{"Informatics 1"}}(\text{Courses})))$$

Relational Algebra

The relational algebra expression can be rearranged without changing its value, but possibly affecting the time and memory needed for computation:

$$\pi_{\text{name}}((\text{Students} \bowtie \text{Takes}) \bowtie (\sigma_{\text{name}=\text{"Informatics 1"}}(\text{Courses})))$$
$$\pi_{\text{name}}(\text{Students} \bowtie (\text{Takes} \bowtie (\sigma_{\text{name}=\text{"Informatics 1"}}(\text{Courses}))))$$
$$\pi_{\text{name}}(\text{Students} \bowtie ((\sigma_{\text{name}=\text{"Informatics 1"}}(\text{Courses})) \bowtie \text{Takes}))$$

We can also visualise this as rearrangements of a tree:



Students and Courses (2/5)

Courses taken by students called "Joe"

$$\begin{aligned} \{ R \mid \exists S \in \text{Students}, T \in \text{Takes}, C \in \text{Courses} . \\ S.\text{name} = \text{"Joe"} \wedge S.\text{mn} = T.\text{mn} \\ \wedge C.\text{code} = T.\text{code} \wedge C.\text{name} = R.\text{name} \} \end{aligned}$$

Note the slightly abbreviated syntax for multiple quantification: we use comma-separated $\exists .., .., ..$ instead of $\exists .. \exists .. \exists ..$

Computing this in relational algebra:

$$\pi_{\text{name}}((\text{Courses} \bowtie \text{Takes}) \bowtie (\sigma_{\text{name}=\text{"Joe"}}(\text{Students})))$$

Students and Courses (3/5)

Students taking Informatics 1 or Geology 1

$$\{ R \mid \exists S \in \text{Students}, T \in \text{Takes}, C \in \text{Courses} . \\ (C.\text{name} = \text{"Informatics 1"} \vee C.\text{name} = \text{"Geology 1"}) \\ \wedge C.\text{code} = T.\text{code} \wedge T.\text{mn} = S.\text{mn} \wedge S.\text{name} = R.\text{name} \}$$

Now the logical formula becomes a little more elaborate.

Computing this in relational algebra:

$$\begin{aligned} & \pi_{\text{name}}((\text{Students} \bowtie \text{Takes}) \bowtie (\sigma_{\text{name}=\text{"Informatics 1"}}(\text{Courses}))) \\ & \cup \pi_{\text{name}}((\text{Students} \bowtie \text{Takes}) \bowtie (\sigma_{\text{name}=\text{"Geology 1"}}(\text{Courses}))) \\ & \pi_{\text{name}}((\text{Students} \bowtie \text{Takes}) \bowtie (\sigma_{(\text{name}=\text{"Informatics 1"} \vee \text{name}=\text{"Geology 1"})}(\text{Courses}))) \end{aligned}$$

Students and Courses (4/5)

Students taking both Informatics 1 and Geology 1

$$\{ R \mid \exists S \in \text{Students}, T, T' \in \text{Takes}, C, C' \in \text{Courses} .$$
$$C.\text{name} = \text{"Informatics 1"} \wedge C.\text{code} = T.\text{code} \wedge T.\text{mn} = S.\text{mn}$$
$$C'.\text{name} = \text{"Geology 1"} \wedge C'.\text{code} = T'.\text{code} \wedge T'.\text{mn} = S.\text{mn}$$
$$\wedge S.\text{name} = R.\text{name} \}$$

Computing this in relational algebra:

$$\pi_{\text{name}}((\text{Students} \bowtie \text{Takes}) \bowtie (\sigma_{\text{name}=\text{"Informatics 1"}}(\text{Courses})))$$
$$\cap \pi_{\text{name}}((\text{Students} \bowtie \text{Takes}) \bowtie (\sigma_{\text{name}=\text{"Geology 1"}}(\text{Courses})))$$

Students and Courses (5/5)

Students taking no courses

$$\{ R \mid \exists S \in \text{Students} . S.\text{name} = R.\text{name} \wedge \forall T \in \text{Takes} . T.\text{mn} \neq S.\text{mn} \}$$

Computing this in relational algebra:

$$\pi_{\text{name}}(\text{Students} - \pi_{\text{name}, \text{mn}}(\text{Students} \bowtie \text{Takes}))$$

★ Challenge: why not one of these instead?

$$\pi_{\text{name}}(\text{Students} - (\text{Students} \bowtie \text{Takes}))$$

$$\pi_{\text{name}}(\text{Students}) - \pi_{\text{name}}(\text{Students} \bowtie \text{Takes})$$