

PAPER 5 (Astrophysical Fluid Dynamics) – ANSWERS 2012

- 1 (a) (UNSEEN) Given dispersion relation applies when $\mathbf{k} \parallel \mathbf{v}_A$

Mode: (A) if $\mathbf{v}_A \cdot \mathbf{u} = 0$, we have motion transverse to \mathbf{v}_A and we get a wave with:

$$\frac{\omega^2}{k^2} = v_A^2$$

i.e. a pure MHD wave

[2]

Mode (B): if $\mathbf{u} \parallel \mathbf{k}$ then we have

$$(k^2 v_A^2 - \omega^2) \mathbf{u} + \left(\frac{c_s^2}{v_A^2} - 1 \right) k^2 v_A^2 \mathbf{u} = 0$$

or

$$(k^2 c_s^2 - \omega^2) \mathbf{u} = 0$$

i.e. a normal compressible sound wave

[2]

- (b) (UNSEEN, BUT SIMILAR TO EXAMPLE SHEET QUESTION)

$$\text{Cooling rate } \dot{Q} = A \rho T^\alpha - H = \frac{\mu A \rho T^{\alpha-1}}{k_B} - H$$

[2]

therefore

$$\dot{Q} = \dot{Q}_{T_0} + \left(\frac{\partial \dot{Q}}{\partial T} \right)_p \Delta T$$

$\left(\frac{\partial \dot{Q}}{\partial T} \right)_p < 0$ implies instability and

$$\left(\frac{\partial \dot{Q}}{\partial T} \right)_p = (\alpha - 1) \frac{\mu A \rho}{k_B} T^{\alpha-1}$$

therefore unstable when $\alpha < 1$ i.e. Bremsstrahlung is unstable.

[2]

- (c) (UNSEEN, BUT SIMILAR TO EXAMPLE SHEET QUESTION)

Sound crossing time $t_c = R/c_s$

Estimate the free-fall collapse time:

Energy

$$\frac{GM^2}{R} \sim \frac{M\dot{R}}{2}$$

therefore

$$\dot{R} \sim \left(\frac{2GM}{R} \right)^{1/2} = \frac{R}{t_{ff}}$$

therefore

$$t_{ff} = \frac{R}{\dot{R}} \sim \left(\frac{R^3}{2GM} \right)^{1/2} \sim \left(\frac{3}{8\pi G \rho_0} \right)^{1/2}$$

or quote exact result:

$$t_{ff} = \left(\frac{3\pi}{32G\rho_0} \right)^{1/2}$$

equating

$$R^2 = c_s^2 \left(\frac{3}{8\pi G\rho_0} \right)$$

[3]

if $t_{ff} < t_c$ will collapse. Could also note result is approximately Jeans length.

[1]

2 Brief notes:

(a) stellar winds;

- Spherical symmetry and similarity to accretion (time reversed equations)
- Consider steady state
- Equations which govern the flow are:
conservation of mass $\dot{M} = 4\pi r^2 \rho u$
Momentum

$$u \frac{d}{dr} = -\frac{1}{\rho} \frac{dp}{dr} - \frac{GM}{r^2}$$

Continuity

$$\frac{d \ln \rho}{dr} = -\frac{2}{r} - \frac{d \ln u}{dr}$$

- Can derive key equation

$$(u^2 - c_s^2) \frac{d \ln u}{dr} = \frac{2c_s^2}{r} \left[1 - \frac{GM}{2c_s^2 r} \right]$$

This defines point where flow transitions between subsonic and supersonic flow $r_s = GM/2c_s^2$.

- Stellar winds are driven into ISM from stellar surface – provide energy input to the ISM
- Able to blow bubbles in the ISM – could give examples or more astrophysical insights here
- Proceeding with analysis need to consider equation of state; isothermal or adiabatic are most obvious
- Need a solution with speed going to zero as we get to infinity – could discuss general solution and select appropriate solution either mathematically or via a sketch of the solutions; either would get full credit.

(b) the Rayleigh-Taylor instability;

- Unstable stratified configuration of fluid under gravity
- Relationship to convective instability. Discussion of the effects of the entropy gradient and the requirements for how the entropy gradient stabilises or not the fluid.
- Might mention at this point astrophysical applications to the convective stability of stars
- Discuss general stratified fluid and how one would approach the analysis – no details are required for full credit just diagrams indicating the general considerations.
- RT instability is the static case – either analysis of physical insight will give full credit. Physical insight is that a heavier fluid sits above a lighter one in a static gravitational field – any instability in the layer moves heavy gas to

(TURN OVER)

lower potential and forces cannot act to reverse the process due to having an equilibrium situation in which the stratified fluid is maintained.

- discuss astrophysical examples – e.g. filament formation in swept up gas behind a shock etc.

(c) Bernoulli's equation.

- Outline derivation of Bernoulli from time-dependent Euler equation
- Noting that

$$\mathbf{u} \cdot \nabla \mathbf{u} = \nabla \frac{1}{2} u^2 - \mathbf{u} \times (\nabla \times \mathbf{u})$$

and that the second term is the curl of the vorticity.

- We get to the critical equation

$$\frac{\partial \mathbf{u}}{\partial t} - \mathbf{u} \times (\nabla \times \mathbf{u}) = -\nabla \left(\int \frac{dp}{\rho} + \frac{1}{2} u^2 + \phi \right)$$

- The term in the parentheses

$$H = \int \frac{dp}{\rho} + \frac{1}{2} u^2 + \phi$$

is a constant if:

- We consider flow along a streamline which is steady - dotting with the velocity loses the curl term. H is then constant along a streamline
- We have vorticity free steady flow everywhere in which case H is the same constant throughout the fluid
- Examples of its use include: aircraft wings; motion of a shower curtain when you turn on the tap or any other relevant examples
- In the notes under this heading is the discussion of the conservation of vorticity in a fluid. Credit will be given if this is reproduced.

3 (BOOKWORK) Briefly describe how shock waves arise in interstellar space.

Answer: Arises from supersonic flow which is intrinsically unstable – gas moving faster than the sound speed means pressure waves cannot stabilise (also faster than molecular mean speed). result is a discontinuous change in which the kinetic energy of the gas is thermalised in a shock with the post shock gas moving subsonically (speed goes down and sound speed increases substantially).

[3]

Rankine-Hugoniot conditions at a normal adiabatic shock:

Conservation of mass

$$\rho_1 u_1 = \rho_2 u_2$$

or integrate the continuity equation across the shock front under stationary conditions.

[1]

Momentum Integrate Euler

$$\frac{\partial \rho u}{\partial t} = -\frac{\partial}{\partial x}(\rho u_x u_x + p)$$

again stationary condition across shock surface or use physical argument to give pressure balance directly.

[2]

Energy:

Conserve flow of total energy allowing for the internal energy and pdV work

$$\rho_1 u_1 \left(\frac{1}{2} u_1^2 e_1 + p_1 / \rho_1 \right) = \rho_2 u_2 \left(\frac{1}{2} u_2^2 e_2 + p_2 / \rho_2 \right)$$

but $e = 1/(1 - \gamma)p/\rho$ and use mass conservation to give

$$\frac{\gamma}{\gamma - 1} \frac{p_2}{\rho_2} + \frac{1}{2} u_2^2 = \frac{\gamma}{\gamma - 1} \frac{p_1}{\rho_1} + \frac{1}{2} u_1^2$$

Together we have

[3]

$$\begin{aligned} \rho_2 u_2 &= \rho_1 u_1 \\ p_2 + \rho_2 u_2^2 &= p_1 + \rho_1 u_1^2 \\ \frac{\gamma}{\gamma - 1} \frac{p_2}{\rho_2} + \frac{1}{2} u_2^2 &= \frac{\gamma}{\gamma - 1} \frac{p_1}{\rho_1} + \frac{1}{2} u_1^2 \end{aligned}$$

(UNSEEN, BUT SIMILAR TO EXAMPLE SHEET QUESTION)

Strong shock the upstream pressure can be neglected – terms in p_1 can be neglected. Hence we have to this approximation:

$$\rho_2 u_2 = \rho_1 u_1 \quad (1)$$

$$p_2 + \rho_2 u_2^2 \approx \rho_1 u_1^2 \quad (2)$$

$$\frac{2\gamma}{\gamma - 1} \frac{p_2}{\rho_2} + u_2^2 \approx u_1^2 \quad (3)$$

(TURN OVER)

From (2) $p_2 = \rho_1 u_1^2 - \rho_2 u_2^2 = \rho_2 u_2 (u_1 - u_2)$ therefore

$$\frac{2\gamma}{\gamma - 1} \frac{\rho_2 u_2 (u_1 - u_2)}{\rho_2} = u_1^2 - u_2^2 = (u_1 - u_2)(u_1 + u_2)$$

cancelling ρ_2 and $(u_1 - u_2)$ gives

$$\frac{\rho_2}{\rho_1} = \frac{u_1}{u_2} = \frac{\gamma + 1}{\gamma - 1}$$

[6]

(UNSEEN) Colliding clouds:

- Work in zero-momentum frame of colliding clouds – interface between clouds is stationary
- shock is driven back into each colliding cloud
- Shocked gas must have $u = 0$ since $u = 0$ at the interface between the clouds
- Unshocked gas must be moving at the original speed towards the interface – doesn't know the collision has occurred since no sound waves travel ahead of the shock

[3]

Speed of shock – transform from the frame considered in first part of question to the frame in which the shocked gas is stationary. Define shock speed at V_s

- V_s is the relative speed between the frames since shock was stationary in original frame
- Must have $u_2 = V_s$ to have shocked gas stationary
- Therefore $u_1 = V + V_s$

But

$$u_1 = \frac{\gamma + 1}{\gamma - 1} u_2 = \frac{\gamma + 1}{\gamma - 1} V_s$$

therefore

$$V = \left(\frac{\gamma + 1}{\gamma - 1} - 1 \right) V_s$$

Hence

$$V_s = \left(\frac{\gamma - 1}{2} \right) V$$

[4]

- As shock reaches back of cloud need to consider boundary conditions
- Cloud must be embedded in an external medium, think of shock as a wave – it will propagate into external gas
- There must be some form of reflected discontinuity into the cloud
- In fact a shock propagates and a rarefaction wave is reflected into the cloud but not expecting this level of detail

[3]

4 (BOOKWORK) Lagrangian derivative of some quantity Q : Consider $Q = Q(\mathbf{r}, t)$ then

$$\Delta Q = \frac{\partial Q}{\partial t} \Delta t + \Delta \mathbf{r} \cdot \nabla Q$$

Hence

$$\frac{\Delta Q}{\Delta t} = \frac{DQ}{Dt} = \frac{\partial Q}{\partial t} + \mathbf{u} \cdot \nabla Q$$

where $\mathbf{u} = \Delta \mathbf{r} / \Delta t$ is the fluid velocity.

[3]

Euler's equation for fluid flow: Consider a small volume $\delta\tau$ then

$$\rho \delta\tau \frac{D\mathbf{u}}{Dt} = \text{forces}$$

Gravitational forces:

$$\mathbf{g} = -\nabla\phi$$

and

$$\mathbf{F}_g = -\rho \delta\tau \nabla\phi$$

Pressure:

$$\mathbf{F}_p = \oint_S p d\mathbf{S} = - \int_V \nabla p d\tau$$

therefore $-\nabla p d\tau$ is the force on $d\tau$. Hence

$$\frac{D\mathbf{u}}{Dt} = \frac{\partial \mathbf{u}}{\partial t} + \mathbf{u} \cdot \nabla \mathbf{u} = -\frac{1}{\rho} \nabla p - \nabla\phi$$

[4]

Hydrostatic equilibrium:

$$\rho \nabla\phi = -\nabla p$$

Spherical cloud

$$\phi = -\frac{GM_r}{r}$$

$$M_r = \int_0^r 4\pi r'^2 \rho dr'$$

Therefore

[3]

$$\rho \frac{GM_r}{r^2} = -\frac{dp}{dr} = -c_s^2 \frac{d\rho}{dr}$$

Trial solution $\rho = A/r^2$ find $M_r = 4\pi Ar$ and

$$\frac{d\rho}{dr} = -\frac{2A}{r^3}$$

Therefore

[3]

$$\frac{G4\pi Ar}{r^2} = -c_s^2 \frac{1}{\rho} \frac{d\rho}{dr} = c_s^2 \frac{r^2}{A} \frac{2A}{r^3}$$

(TURN OVER

works if

$$A = \frac{c_s^2}{2\pi G}$$

[2]

Cloud of mass $M = 4\pi A r_0$ from integral above hence:

$$M = \frac{2c_s^2 r_0}{G}$$

and must have $p_0 = p(r_0)$ as a boundary condition hence

$$p_0 = c_s^2 \frac{A}{r_0^2} = \frac{c_s^4}{2\pi G r_0^2}$$

[2]

(UNSEEN) Virial analysis:

Gravitational PE

$$\Omega = - \int \frac{GM_r}{r} \rho d\tau = -2c_s^2 \int \rho d\tau = -2c_s^2 M$$

Internal energy

$$T = \int \frac{1}{\gamma - 1} p d\tau = \frac{3}{2} c_s^2 \int \rho d\tau = \frac{3}{2} c_s^2 M$$

hence

$$2T + \Omega = 3c_s^2 M - 2c_s^2 M = c_s^2 M$$

Now

$$4\pi r_0^3 p_0 = 4\pi r_0^3 \frac{c_s^4}{2\pi G r_0^2} = \frac{2c_s^4 r_0}{G} = c_s^2 M$$

Hence shown modified form of virial.

[6]

The extra term is an Area, times a pressure times a distance - it is related to the work done in embedding the cloud in the external medium - i.e. it is work term which is the most important thing to note.

[2]

END OF PAPER