

Artificial Intelligence II (CS4442 & CS9542)

Classification: Generative Models

Boyu Wang
Department of Computer Science
University of Western Ontario

Discriminative model vs. generative model

- Recall: in logistic regression, we model directly $p(y|x)$:

$$p(y = 1|x; w) \triangleq \sigma(h_w(x)) = \frac{1}{1 + e^{-w^\top x}},$$

- This is called **discriminative** model, because we only care about discriminating examples of the two classes.

Discriminative model vs. generative model

- Recall: in logistic regression, we model directly $p(y|x)$:

$$p(y = 1|x; w) \triangleq \sigma(h_w(x)) = \frac{1}{1 + e^{-w^\top x}},$$

– This is called **discriminative** model, because we only care about discriminating examples of the two classes.

- Another way to model $p(y)$ and $p(x|y)$ and then use the Bayes Rule:

$$\begin{aligned} p(y = 1|x) &= \frac{p(x, y = 1)}{p(x)} \\ &= \frac{p(x|y = 1)p(y = 1)}{p(x|y = 1)p(y = 1) + p(x|y = 0)p(y = 0)} \end{aligned}$$

– This is called **generative** model, because we can actually use the model to generate data.

Bayes classifier for continuous features

- ▶ Idea: Use the training data to estimate $p(y)$ and $p(x|y)$
- ▶ $p(y)$ can be estimated by counting the number of data points of each class.
- ▶ How to estimate $p(x|y)$?
 - Need additional assumptions (for continuous inputs) – multivariate Gaussian with mean $\mu \in \mathbb{R}^n$, and covariance $\Sigma \in \mathbb{R}^{n \times n}$
 - Each class has mean μ_c and covariance Σ_c , $c \in \{0, 1\}$

Examples of multivariate Gaussian distribution

$$\Sigma = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \quad \Sigma = \begin{bmatrix} 1 & 0.5 \\ 0.5 & 1 \end{bmatrix} \quad \Sigma = \begin{bmatrix} 1 & 0.8 \\ 0.8 & 1 \end{bmatrix}$$




Figure: 2D Gaussian distributions with different Σ

Figure credit: Doina Precup

Gaussian discriminant analysis

- For 2 classes:

$$p(y = 1) = \theta; \quad p(y = 0) = 1 - \theta$$

$$p(x|y = 1) = \frac{1}{(2\pi)^{\frac{n}{2}} |\Sigma_1|^{\frac{1}{2}}} e^{-\frac{1}{2}(x-\mu_1)^\top \Sigma_1^{-1}(x-\mu_1)}$$

$$p(x|y = 0) = \frac{1}{(2\pi)^{\frac{n}{2}} |\Sigma_0|^{\frac{1}{2}}} e^{-\frac{1}{2}(x-\mu_0)^\top \Sigma_0^{-1}(x-\mu_0)}$$

- The parameters to estimate are: $\theta, \mu_1, \Sigma_1, \mu_0, \Sigma_0$

- For C classes:

$$p(y = c) = \theta_c; \quad \text{s.t.} \quad \sum_{c=1}^C \theta_c = 1$$

$$p(x|y = c) = \frac{1}{(2\pi)^{\frac{n}{2}} |\Sigma_c|^{\frac{1}{2}}} e^{-\frac{1}{2}(x-\mu_c)^\top \Sigma_c^{-1}(x-\mu_c)}$$

- The parameters to estimate are: $\{\theta_c, \mu_c, \Sigma_c\}_{c=1}^C$

Estimate the parameters

- ▶ We can write down the likelihood function, like linear regression and logistic regression
- ▶ Compute the gradient with respect to the parameters and set them to 0.
 - The parameter θ_c is given by $\theta_c = \frac{n_c}{n}$, where n_c is the number of instances of class c .
 - The mean μ_c is given by

$$\mu_c = \frac{1}{n_c} \sum_{x_i: y_i=c} x_i$$

- The covariance matrix Σ_c is given by

$$\Sigma_c = \frac{1}{n_c} \sum_{x_i: y_i=c} (x_i - \mu_c)(x_i - \mu_c)^\top$$

Other variants to simplify the model

- ▶ If we assume the same covariance matrix Σ for all the classes, the maximum likelihood estimation of Σ is

$$\Sigma = \frac{n_c}{n} \sum_{c=1}^C \Sigma_c$$

- ▶ Covariance matrix can be restricted to diagonal, or mostly diagonal with few off-diagonal elements, based on prior knowledge.
- ▶ Covariance matrix can even be identity matrix.
- ▶ The shape of the covariance is influenced both by assumptions about the domain and by the amount of data available.
- ▶ If the covariance matrices are different for the class, the model is called **quadratic discriminant analysis (QDA)**; if the covariance matrices are the same, the model is called **linear discriminant analysis (LDA)**; if the covariance matrices are diagonal, the model is called **naive Bayes classifier (NBC)**.

Classification using quadratic discriminant analysis

Recall:

$$p(y = c) = \theta_c; \quad p(x|y = c) = \frac{1}{(2\pi)^{\frac{n}{2}} |\Sigma_c|^{\frac{1}{2}}} e^{-\frac{1}{2}(x-\mu_c)^\top \Sigma_c^{-1}(x-\mu_c)}$$

Using the Bayes rule, we have

$$p(y = c|x) \propto \theta_c |2\pi \Sigma_c|^{-\frac{1}{2}} e^{-\frac{1}{2}(x-\mu_c)^\top \Sigma_c^{-1}(x-\mu_c)}$$

Predict class label as the most probable label:

$$y = \arg \max_c p(y = c|x)$$




Figure credit: Kevin Murphy

Classification using linear discriminant analysis

If we assume the covariance matrices are the same for all the classes:

$$\begin{aligned} p(y = c|x) &\propto \theta_c e^{-\frac{1}{2}(x-\mu_c)^\top \Sigma^{-1}(x-\mu_c)} \\ &= e^{\mu_c^\top \Sigma^{-1}x - \frac{1}{2}\mu_c^\top \Sigma \mu_c + \log \theta_c} \cdot e^{-\frac{1}{2}x^\top \Sigma^{-1}x} \\ &\propto e^{\mu_c^\top \Sigma^{-1}x - \frac{1}{2}\mu_c^\top \Sigma \mu_c + \log \theta_c} \end{aligned}$$

Let $w_c = \Sigma^{-1}\mu_c$, and $b_c = -\frac{1}{2}\mu_c^\top \Sigma \mu_c + \log \theta_c \Rightarrow$ we get a linear model!




Figure credit: Kevin Murphy

Bayes classifier for discrete features

- ▶ Idea: Use the training data to estimate $p(y)$ and $p(x|y)$
- ▶ $p(y)$ can be estimated in the same way as for continuous features
- ▶ How to estimate $p(x|y)$ for discrete values?

- Assume $x = [x_1, \dots, x_n]^T \in \mathbb{R}^n$ has n features. Then using the chain rule, we have

$$p(x|y) = p(x_1|y)p(x_2|y, x_1) \cdots p(x_n|y, x_1, \dots, x_{n-1})$$

– even for binary features, it requires $\mathcal{O}(2^n)$ numbers to describe the model!

- If we assume that the features x_j 's are conditionally independent given y : $p(x_j|y) = p(x_j|y, x_k), \forall i, j$, then we have

$$\begin{aligned} p(x|y) &= p(x_1|y)p(x_2|y, x_1) \cdots p(x_n|y, x_1, \dots, x_{n-1}) \\ &= p(x_1|y)p(x_2|y) \cdots p(x_n|y) \end{aligned}$$

– only requires $\mathcal{O}(n)$ numbers to describe the model!

Conditional independence: an example

- ▶ A box contains two coins: a regular coin (R) and one fake two-headed coin (F). I choose a coin at random and toss it twice. Define the following two events:
 - A = First coin toss results in a head
 - B = Second coin toss results in a head

Are A and B independent?

Conditional independence: an example

- ▶ A box contains two coins: a regular coin (R) and one fake two-headed coin (F). I choose a coin at random and toss it twice. Define the following two events:
 - A = First coin toss results in a head
 - B = Second coin toss results in a head

Are A and B independent?

- ▶ $p(A) = p(B) = p(\text{head}) = p(\text{head}|R) \times p(R) + p(\text{head}|F) \times p(F) = \frac{1}{2} \times \frac{1}{2} + \frac{1}{2} = \frac{3}{4}$
 $p(A, B) = p(\text{head}, \text{head}) = p(\text{head}, \text{head}|R) \times p(R) + p(\text{head}, \text{head}|F) \times p(F) = \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} + \frac{1}{2} = \frac{5}{8}$
 $p(A)p(B) \neq p(A, B) \Rightarrow A \text{ and } B \text{ are dependent!}$

Conditional independence: an example

- ▶ A box contains two coins: a regular coin (R) and one fake two-headed coin (F). I choose a coin at random and toss it twice. Define the following two events:
 - A = First coin toss results in a head
 - B = Second coin toss results in a head

Are A and B independent?

- ▶ $p(A) = p(B) = p(\text{head}) = p(\text{head}|R) \times p(R) + p(\text{head}|F) \times p(F) = \frac{1}{2} \times \frac{1}{2} + \frac{1}{2} = \frac{3}{4}$
 $p(A, B) = p(\text{head}, \text{head}) = p(\text{head}, \text{head}|R) \times p(R) + p(\text{head}, \text{head}|F) \times p(F) = \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} + \frac{1}{2} = \frac{5}{8}$
 $p(A)p(B) \neq p(A, B) \Rightarrow A \text{ and } B \text{ are dependent!}$

- ▶ Consider an additional event:
 - C = Coin R (regular) has been selected.

Then it is easy to show that $p(A|R)p(B|R) = p(A, B|R) \Rightarrow A \text{ and } B \text{ are conditionally independent given } C!$

Naive Bayes classifier for binary features

- ▶ The model parameters are $\{\theta_c = p(y = c)\}_{c=1}^C$ and $\{\beta_{j,c} = p(x_j = 1|y = c)\}_{j,c=1}^{n,C}$
- ▶ Predict class label as the most probable label:

$$y = \arg \max_c \left[p(y = c) \prod_{j=1}^n p(x_j|y = c) \right]$$

- ▶ In practice, using the **log trick** to avoid the numerical issue:

$$\begin{aligned} y &= \arg \max_c \log \left[p(y = c) \prod_{j=1}^n p(x_j|y = c) \right] \\ &= \arg \max_c \log p(y = c) + \sum_{j=1}^n \log p(x_j|y = c) \end{aligned}$$

Maximum likelihood estimation for Naive Bayes

- ▶ The log-likelihood function is

$$\log L \left(\{\theta_c\}_{c=1}^C, \{\beta_{j,c}\}_{j,c=1}^{n,C} \right) = \sum_{i=1}^m \left(\log p(y_i) + \sum_{j=1}^n \log p(x_{i,j}|y_i) \right)$$

- ▶ Computing the gradient with respect to θ_c and setting it to 0 gives us:

$$\theta_c = \frac{n_c}{n}$$

- ▶ Computing the gradient with respect to $\beta_{j,c}$ and setting it to 0 gives us:

$$\begin{aligned} \beta_{j,c} &= p(x_j = 1 | y = c) \\ &= \frac{\text{number of the instances for which } x_{i,j} = 1 \text{ and } y_i = c}{\text{number of the instances for which } y_i = c} \end{aligned}$$

Training Naive Bayes

Estimate $P(X_j | Y)$ and $P(Y)$ directly from the training data by counting!

<u>Sky</u>	<u>Temp</u>	<u>Humid</u>	<u>Wind</u>	<u>Water</u>	<u>Forecast</u>	<u>Play?</u>
sunny	warm	normal	strong	warm	same	yes
sunny	warm	high	strong	warm	same	yes
rainy	cold	high	strong	warm	change	no
sunny	warm	high	strong	cool	change	yes

$$P(\text{play}) = ?$$

$$P(\text{Sky} = \text{sunny} \mid \text{play}) = ?$$

$$P(\text{Humid} = \text{high} \mid \text{play}) = ?$$

...

$$P(\neg \text{play}) = ?$$

$$P(\text{Sky} = \text{sunny} \mid \neg \text{play}) = ?$$

$$P(\text{Humid} = \text{high} \mid \neg \text{play}) = ?$$

...

Training Naive Bayes

Estimate $P(X_j | Y)$ and $P(Y)$ directly from the training data by counting!

<u>Sky</u>	<u>Temp</u>	<u>Humid</u>	<u>Wind</u>	<u>Water</u>	<u>Forecast</u>	<u>Play?</u>
sunny	warm	normal	strong	warm	same	yes
sunny	warm	high	strong	warm	same	yes
rainy	cold	high	strong	warm	change	no
sunny	warm	high	strong	cool	change	yes

$$P(\text{play}) = ?$$

$$P(\text{Sky} = \text{sunny} \mid \text{play}) = ?$$

$$P(\text{Humid} = \text{high} \mid \text{play}) = ?$$

...

$$P(\neg \text{play}) = ?$$

$$P(\text{Sky} = \text{sunny} \mid \neg \text{play}) = ?$$

$$P(\text{Humid} = \text{high} \mid \neg \text{play}) = ?$$

...

Training Naive Bayes

Estimate $P(X_j | Y)$ and $P(Y)$ directly from the training data by counting!

Sky	Temp	Humid	Wind	Water	Forecast	Play?
sunny	warm	normal	strong	warm	same	yes
sunny	warm	high	strong	warm	same	yes
rainy	cold	high	strong	warm	change	no
sunny	warm	high	strong	cool	change	yes

$$P(\text{play}) = 3/4$$

$$P(\text{Sky} = \text{sunny} \mid \text{play}) = ?$$

$$P(\text{Humid} = \text{high} \mid \text{play}) = ?$$

...

$$P(\neg \text{play}) = 1/4$$

$$P(\text{Sky} = \text{sunny} \mid \neg \text{play}) = ?$$

$$P(\text{Humid} = \text{high} \mid \neg \text{play}) = ?$$

...

Training Naive Bayes

Estimate $P(X_j | Y)$ and $P(Y)$ directly from the training data by counting!

<u>Sky</u>	<u>Temp</u>	<u>Humid</u>	<u>Wind</u>	<u>Water</u>	<u>Forecast</u>	<u>Play?</u>
sunny						yes
sunny						yes
rainy	cold	high	strong	warm	change	no
sunny						yes

$$P(\text{play}) = 3/4$$

$$P(\text{Sky} = \text{sunny} \mid \text{play}) = ?$$

$$P(\text{Humid} = \text{high} \mid \text{play}) = ?$$

...

$$P(\neg \text{play}) = 1/4$$

$$P(\text{Sky} = \text{sunny} \mid \neg \text{play}) = ?$$

$$P(\text{Humid} = \text{high} \mid \neg \text{play}) = ?$$

...

Training Naive Bayes

Estimate $P(X_j | Y)$ and $P(Y)$ directly from the training data by counting!

Sky	Temp	Humid	Wind	Water	Forecast	Play?
sunny						yes
sunny						yes
rainy	cold	high	strong	warm	change	no
sunny						yes

$$P(\text{play}) = 3/4$$

$$P(\text{Sky} = \text{sunny} \mid \text{play}) = 1$$

$$P(\text{Humid} = \text{high} \mid \text{play}) = ?$$

...

$$P(\neg \text{play}) = 1/4$$

$$P(\text{Sky} = \text{sunny} \mid \neg \text{play}) = ?$$

$$P(\text{Humid} = \text{high} \mid \neg \text{play}) = ?$$

...

Training Naive Bayes

Estimate $P(X_j | Y)$ and $P(Y)$ directly from the training data by counting!

<u>Sky</u>	<u>Temp</u>	<u>Humid</u>	<u>Wind</u>	<u>Water</u>	<u>Forecast</u>	<u>Play?</u>
sunny	warm	normal	strong	warm	same	yes
sunny	warm	high	strong	warm	same	yes
rainy						no
sunny	warm	high	strong	cool	change	yes

$$P(\text{play}) = 3/4$$

$$P(\text{Sky} = \text{sunny} \mid \text{play}) = 1$$

$$P(\text{Humid} = \text{high} \mid \text{play}) = ?$$

...

$$P(\neg \text{play}) = 1/4$$

$$P(\text{Sky} = \text{sunny} \mid \neg \text{play}) = ?$$

$$P(\text{Humid} = \text{high} \mid \neg \text{play}) = ?$$

...

Training Naive Bayes

Estimate $P(X_j | Y)$ and $P(Y)$ directly from the training data by counting!

<u>Sky</u>	<u>Temp</u>	<u>Humid</u>	<u>Wind</u>	<u>Water</u>	<u>Forecast</u>	<u>Play?</u>
sunny	warm	normal	strong	warm	same	yes
sunny	warm	high	strong	warm	same	yes
rainy						no
sunny	warm	high	strong	cool	change	yes

$$P(\text{play}) = 3/4$$

$$P(\text{Sky} = \text{sunny} \mid \text{play}) = 1$$

$$P(\text{Humid} = \text{high} \mid \text{play}) = ?$$

...

$$P(\neg \text{play}) = 1/4$$

$$P(\text{Sky} = \text{sunny} \mid \neg \text{play}) = 0$$

$$P(\text{Humid} = \text{high} \mid \neg \text{play}) = ?$$

...

Training Naive Bayes

Estimate $P(X_j | Y)$ and $P(Y)$ directly from the training data by counting!

Sky	Temp	Humid	Wind	Water	Forecast	Play?
		normal				yes
		high				yes
rainy	cold	high	strong	warm	change	no
		high				yes

$$P(\text{play}) = 3/4$$

$$P(\text{Sky} = \text{sunny} \mid \text{play}) = 1$$

$$P(\text{Humid} = \text{high} \mid \text{play}) = ?$$

...

$$P(\neg \text{play}) = 1/4$$

$$P(\text{Sky} = \text{sunny} \mid \neg \text{play}) = 0$$

$$P(\text{Humid} = \text{high} \mid \neg \text{play}) = ?$$

...

Training Naive Bayes

Estimate $P(X_j | Y)$ and $P(Y)$ directly from the training data by counting!

Sky	Temp	Humid	Wind	Water	Forecast	Play?
		normal				yes
		high				yes
rainy	cold	high	strong	warm	change	no
		high				yes

$$P(\text{play}) = 3/4$$

$$P(\neg \text{play}) = 1/4$$

$$P(\text{Sky} = \text{sunny} | \text{play}) = 1$$

$$P(\text{Sky} = \text{sunny} | \neg \text{play}) = 0$$

$$P(\text{Humid} = \text{high} | \text{play}) = 2/3 \quad P(\text{Humid} = \text{high} | \neg \text{play}) = ?$$

...

...

Training Naive Bayes

Estimate $P(X_j | Y)$ and $P(Y)$ directly from the training data by counting!

<u>Sky</u>	<u>Temp</u>	<u>Humid</u>	<u>Wind</u>	<u>Water</u>	<u>Forecast</u>	<u>Play?</u>
sunny	warm	normal	strong	warm	same	yes
sunny	warm	high	strong	warm	same	yes
		high				no
sunny	warm	high	strong	cool	change	yes

$$P(\text{play}) = 3/4$$

$$P(\neg \text{play}) = 1/4$$

$$P(\text{Sky} = \text{sunny} \mid \text{play}) = 1$$

$$P(\text{Sky} = \text{sunny} \mid \neg \text{play}) = 0$$

$$P(\text{Humid} = \text{high} \mid \text{play}) = 2/3$$

$$P(\text{Humid} = \text{high} \mid \neg \text{play}) = ?$$

...

...

Training Naive Bayes

Estimate $P(X_j | Y)$ and $P(Y)$ directly from the training data by counting!

<u>Sky</u>	<u>Temp</u>	<u>Humid</u>	<u>Wind</u>	<u>Water</u>	<u>Forecast</u>	<u>Play?</u>
sunny	warm	normal	strong	warm	same	yes
sunny	warm	high	strong	warm	same	yes
		high				no
sunny	warm	high	strong	cool	change	yes

$$P(\text{play}) = 3/4$$

$$P(\neg \text{play}) = 1/4$$

$$P(\text{Sky} = \text{sunny} \mid \text{play}) = 1$$

$$P(\text{Sky} = \text{sunny} \mid \neg \text{play}) = 0$$

$$P(\text{Humid} = \text{high} \mid \text{play}) = 2/3$$

$$P(\text{Humid} = \text{high} \mid \neg \text{play}) = 1$$

...

...

Laplace smoothing

- ▶ Notice that some probabilities estimated by counting might be zero!
- ▶ Instead of the maximum likelihood estimate:

$$\beta_{j,c} = \frac{\text{number of the instances for which } x_{i,j} = 1 \text{ and } y_i = c}{\text{number of the instances for which } y_i = c}$$

use:

$$\beta_{j,c} = \frac{(\text{number of the instances for which } x_{i,j} = 1 \text{ and } y_i = c) + 1}{(\text{number of the instances for which } y_i = c) + C}$$

– add 1 to each count

- ▶ If a feature appears a lot of times, this estimate is only slightly different from maximum likelihood.

Training Naive Bayes with Laplace smoothing

Estimate $P(X_j | Y)$ and $P(Y)$ directly from the training data by counting with Laplace smoothing:

<u>Sky</u>	<u>Temp</u>	<u>Humid</u>	<u>Wind</u>	<u>Water</u>	<u>Forecast</u>	<u>Play?</u>
sunny						yes
sunny						yes
rainy	cold	high	strong	warm	change	no
sunny						yes

$$P(\text{play}) = 3/4$$

$$P(\text{Sky} = \text{sunny} \mid \text{play}) = 4/5$$

$$P(\text{Humid} = \text{high} \mid \text{play}) = ?$$

...

$$P(\neg \text{play}) = 1/4$$

$$P(\text{Sky} = \text{sunny} \mid \neg \text{play}) = ?$$

$$P(\text{Humid} = \text{high} \mid \neg \text{play}) = ?$$

...

Training Naive Bayes with Laplace smoothing

Estimate $P(X_j | Y)$ and $P(Y)$ directly from the training data by counting with Laplace smoothing:

<u>Sky</u>	<u>Temp</u>	<u>Humid</u>	<u>Wind</u>	<u>Water</u>	<u>Forecast</u>	<u>Play?</u>
sunny	warm	normal	strong	warm	same	yes
sunny	warm	high	strong	warm	same	yes
rainy						no
sunny	warm	high	strong	cool	change	yes

$$P(\text{play}) = 3/4$$

$$P(\text{Sky} = \text{sunny} \mid \text{play}) = 4/5$$

$$P(\text{Humid} = \text{high} \mid \text{play}) = ?$$

...

$$P(\neg \text{play}) = 1/4$$

$$P(\text{Sky} = \text{sunny} \mid \neg \text{play}) = 1/3$$

$$P(\text{Humid} = \text{high} \mid \neg \text{play}) = ?$$

...

Training Naive Bayes with Laplace smoothing

Estimate $P(X_j | Y)$ and $P(Y)$ directly from the training data by counting with Laplace smoothing:

<u>Sky</u>	<u>Temp</u>	<u>Humid</u>	<u>Wind</u>	<u>Water</u>	<u>Forecast</u>	<u>Play?</u>
		normal				yes
		high				yes
rainy	cold	high	strong	warm	change	no
		high				yes

$$P(\text{play}) = 3/4$$

$$P(\neg \text{play}) = 1/4$$

$$P(\text{Sky} = \text{sunny} \mid \text{play}) = 4/5$$

$$P(\text{Sky} = \text{sunny} \mid \neg \text{play}) = 1/3$$

$$P(\text{Humid} = \text{high} \mid \text{play}) = 3/5$$

$$P(\text{Humid} = \text{high} \mid \neg \text{play}) = ?$$

...

...

Training Naive Bayes with Laplace smoothing

Estimate $P(X_j | Y)$ and $P(Y)$ directly from the training data by counting with Laplace smoothing:

<u>Sky</u>	<u>Temp</u>	<u>Humid</u>	<u>Wind</u>	<u>Water</u>	<u>Forecast</u>	<u>Play?</u>
sunny	warm	normal	strong	warm	same	yes
sunny	warm	high	strong	warm	same	yes
		high				no
sunny	warm	high	strong	cool	change	yes

$$P(\text{play}) = 3/4$$

$$P(\neg \text{play}) = 1/4$$

$$P(\text{Sky} = \text{sunny} \mid \text{play}) = 4/5$$

$$P(\text{Sky} = \text{sunny} \mid \neg \text{play}) = 1/3$$

$$P(\text{Humid} = \text{high} \mid \text{play}) = 3/5$$

$$P(\text{Humid} = \text{high} \mid \neg \text{play}) = 2/3$$

...

...

Generative model summary

► Advantages:

- Easy to train
- Can handle streaming data well
- Can handle both real and discrete data

► Disadvantages:

- Requires additional assumptions (e.g., Gaussian distribution, conditional independence of features)
- Cannot handle high-dimensional data very well