


5-AMINO-1MQ

NNMT Inhibitor | 5mg & 10mg x 10 Vials

NAD+ Metabolism | Fat Cell Metabolism | SubQ Injection


Purity: 99.8% | Endotoxin-Free | Research Use Only

5-Amino-1MQ (5-amino-1-methylquinolinium) is a small-molecule NNMT (Nicotinamide N-methyltransferase) inhibitor. NNMT is upregulated in fat cells and reduces NAD+ availability for SIRT1 activation. By inhibiting NNMT, 5-Amino-1MQ raises intracellular NAD+, activates SIRT1/AMPK signalling, promotes fat cell browning (beige adipogenesis), and increases metabolic rate.



Parameter	Specification
Class	NNMT inhibitor -- small molecule, not a peptide
Vial Sizes	5mg x 10 10mg x 10 Purity 99.8%
Reconstitution	1 mL BAC Water: 5mg->5mg/mL 10mg->10mg/mL
Standard Dose	100-500 mcg per injection
Frequency	1-2x daily SubQ timing flexible (not fasting-dependent)
Mechanism	NNMT inhibition -> elevated NAD+ -> SIRT1/AMPK activation -> fat browning
Cycle	8-12 weeks on, 4 weeks off
Storage	2-8 deg C Reconstituted: use within 28 days

MECHANISM OF ACTION & RESEARCH BENEFITS



Pathway	Mechanism	Benefit
NNMT Inhibition	Blocks NNMT enzyme that methylates nicotinamide; preserves NAD+ pool; raises intracellular NAD+ in adipocytes	Restored NAD+ availability for metabolic enzymes
SIRT1/AMPK Activation	Elevated NAD+ activates SIRT1 (NAD+-dependent deacetylase) and AMPK; SIRT1 deacetylates PGC-1alpha; drives mitochondrial biogenesis and fat oxidation	Increased metabolic rate; fat oxidation; mitochondrial function
Adipocyte Browning	NNMT inhibition promotes conversion of white adipocytes to beige phenotype; upregulates UCP1 (uncoupling protein); increases thermogenesis	Fat burning via thermogenesis; fat cell metabolism research

Benefit	Context
NAD+ Restoration	Raises intracellular NAD+ levels in adipocytes, restoring SIRT1 and AMPK signalling that declines with age and obesity.
Fat Cell Browning	Promotes beige adipogenesis in white fat depots, increasing metabolic rate via UCP1-mediated thermogenesis.
No Fasting Required	Unlike GH secretagogues, 5-Amino-1MQ mechanism is independent of insulin/GH axis -- inject any time.

RECONSTITUTION & DOSE CALCULATION

All vials use 1 mL BAC Water. Room temperature. Swirl gently -- never shake.



1 mL BAC Water per vial. Roll gently -- NEVER shake.

Step	Action	Detail
1	Gather	Vial, BAC Water, insulin syringe (29-31G), swabs, sharps.
2	Clean stoppers	Wipe both stoppers. Air-dry 10-15 sec.
3	Draw 1mL BAC	Exactly 1 mL. Expel air bubbles.
4	Inject into vial	Aim at GLASS WALL. Slowly. Never onto powder.
5	Swirl & label	Roll 20-30 sec. Clear within 60 sec. Never shake. Label date. 2-8 deg C. Use within 28 days.



1mL Insulin Syringe | 29-31G | SubQ

Formula: $\text{Vol(mL)} = \frac{\text{Dose(mcg)}}{\text{Conc(mcg/mL)}}$ | **Units:** $\text{Vol} \times 100$


5mg x 10 -- 1mL BAC -> 5.0mg/mL (5,000mcg/mL)

Dose(mcg)	100mcg	200mcg	300mcg	400mcg	500mcg
Vol(mL)	0.0200	0.0400	0.0600	0.0800	0.1000
Units	2.0U	4.0U	6.0U	8.0U	10.0U
Doses/Vial	50	25	16	12	10

10mg x 10 -- 1mL BAC -> 10.0mg/mL (10,000mcg/mL)

Dose(mcg)	100mcg	200mcg	300mcg	400mcg	500mcg
Vol(mL)	0.0100	0.0200	0.0300	0.0400	0.0500
Units	1.0U	2.0U	3.0U	4.0U	5.0U
Doses/Vial	100	50	33	25	20

INJECTION TECHNIQUE, PROTOCOLS & GOLDEN RULES



8-Point Rotation Map -- Rotate Every Injection



Step	Action	Detail
1	Wash hands	20 sec with soap. Dry completely.
2	Draw dose	Draw exact volume per dose table. Expel air bubbles.
3	Inject SubQ	Pinch 2-3cm. 45 deg. Release. Inject slowly.
4	Rotate & dispose	Press site. Rotate 8-point map. Sharps container.

Protocol	Dose	Frequency	Duration	Notes
Starting	100-200 mcg	1x daily	8wk on 4wk off	Start low. Assess tolerance.
Standard	300-500 mcg	1-2x daily	8-12wk	Research standard dose.

Side Effect	Likelihood	Management
Injection site reaction	Common	Rotate sites. Fresh needle.
Mild nausea	Occasional	Inject with food if persistent.
Fatigue (early)	Occasional	Metabolic adaptation. Resolves in 1-2 weeks.

Contraindications: Cancer history, pregnancy, under 18.

1. 1 mL BAC Water per vial: 5mg->5mg/mL | 10mg->10mg/mL.

2. No fasting required -- 5-Amino-1MQ is not GH-axis dependent.
3. Start at 100-200 mcg and escalate. Inject 1-2x daily SubQ.
4. Rotate injection sites through 8-point abdomen map.
5. Cycle 8-12 weeks on, 4 weeks off.
6. Store 2-8 deg C. Use within 28 days. Never shake.
7. Contraindicated in cancer history. Not for pregnancy or under 18.

DISCLAIMER:

research use only. Not for human therapeutic use. Consult a qualified professional.

assumes no liability.

|

.com | Research Use Only