

BPC-157

Body Protection Compound | 5mg & 10mg x 10 Vials

GI Repair | Tendon/Ligament Healing | Systemic Cytoprotection | SubQ


Purity: 99.8% | Endotoxin-Free | Research Use Only

BPC-157 (Body Protection Compound-157) is a synthetic pentadecapeptide derived from a protective protein found in gastric juice. It exhibits extraordinary cytoprotective effects across multiple organ systems -- particularly GI tract, tendons, ligaments, bone, and vascular tissue. BPC-157 has a remarkable safety profile with no known LD50 in animal models.



Parameter	Specification
Class	Synthetic pentadecapeptide -- gastric cytoprotective peptide fragment
Vial Sizes	5mg x 10 10mg x 10 Purity 99.8%
Reconstitution	1 mL BAC Water: 5mg->5mg/mL 10mg->10mg/mL
Standard Dose	100-1,000 mcg per injection SubQ 250-500 mcg most common
Frequency	1-2x daily SubQ no fasting required
GI Route	Oral BPC-157 also effective for GI tract -- injectable for systemic
Cycle & Storage	4-12 weeks on, 2-4 weeks off 2-8 deg C 28 days

MECHANISM OF ACTION & RESEARCH BENEFITS



Pathway	Mechanism	Benefit
GI Cytoprotection	Promotes gastric mucosal healing; protects against NSAID and ethanol-induced gastric lesions; heals intestinal anastomoses; anti-ulcer; activates growth factor receptors in GI epithelium	GI repair; gastric ulcer; inflammatory bowel research
Tendon/Ligament Repair	Upregulates tendon fibroblast growth factor receptors; promotes collagen synthesis; accelerates tendon-to-bone healing; increases growth hormone receptor expression in fibroblasts	Tendon/ligament repair; musculoskeletal injury research
Angiogenesis & Nitric Oxide	Upregulates eNOS and NO production; promotes VEGF expression; angiogenic without tumorigenic properties; vascular protection; neuroprotection via NO pathway	Vascular repair; neurological research; systemic protection

Benefit	Context
Unmatched Safety	No known LD50 in animal models -- BPC-157 has one of the safest profiles of any research peptide, even at very high doses.
GI & Systemic	Simultaneously repairs GI tissue (particularly effective for IBD, leaky gut, ulcers) while providing systemic organ protection.
Tendon/Ligament	Accelerates healing of tendons, ligaments, and bone -- frequently combined with TB-500 for synergistic musculoskeletal repair.

RECONSTITUTION & DOSE CALCULATION

All vials use 1 mL BAC Water. Room temperature. Swirl gently -- never shake.



1 mL BAC Water per vial. Roll gently -- NEVER shake.

Step	Action	Detail
1	Gather	Vial, BAC Water, insulin syringe (29-31G), swabs, sharps.
2	Clean stoppers	Wipe both stoppers. Air-dry 10-15 sec.
3	Draw 1mL BAC	Exactly 1 mL. Expel air bubbles.
4	Inject into vial	Aim at GLASS WALL. Slowly. Never onto powder.
5	Swirl & label	Roll 20-30 sec. Clear within 60 sec. Never shake. Label date. 2-8 deg C. Use within 28 days.



1mL Insulin Syringe | 29-31G | SubQ

Formula: Vol(mL)=Dose(mcg)/Conc(mcg/mL) | **Units:** Vol x 100


5mg x 10 -- 1mL BAC -> 5.0mg/mL (5,000mcg/mL)

Dose(mcg)	100mcg	250mcg	500mcg	750mcg	1000mcg
Vol(mL)	0.0200	0.0500	0.1000	0.1500	0.2000
Units	2.0U	5.0U	10.0U	15.0U	20.0U
Doses/Vial	50	20	10	6	5

10mg x 10 -- 1mL BAC -> 10.0mg/mL (10,000mcg/mL)

Dose(mcg)	100mcg	250mcg	500mcg	750mcg	1000mcg
Vol(mL)	0.0100	0.0250	0.0500	0.0750	0.1000
Units	1.0U	2.5U	5.0U	7.5U	10.0U
Doses/Vial	100	40	20	13	10

INJECTION TECHNIQUE, PROTOCOLS & GOLDEN RULES



8-Point Rotation Map -- Rotate Every Injection



Step	Action	Detail
1	Wash & prepare	Wash hands. 29-31G syringe. No fasting required.
2	Draw dose	250-500 mcg standard. Draw exact volume.
3	Inject SubQ	Near injury site (localised) or abdomen (systemic). 45 deg. Rotate.
4	Rotate & dispose	8-point rotation. Sharps container.

Protocol	Dose	Frequency	Duration	Notes
Standard	250-500 mcg	1-2x daily	4-12 wk 2-4 off	Most researched dose range.
High (Research)	750-1,000 mcg	1-2x daily	4-8 wk	Research maximum. Monitor response.
+TB-500 Stack	BPC: 500mcg TB: 2mg	2-3x/week	4-8 wk	Gold standard tissue repair stack.

Side Effect	Likelihood	Management
Injection site reaction	Rare	BPC-157 is exceptionally well tolerated.
Nausea (rare)	Rare	Inject with food if occurs.
Dizziness	Very Rare	Dose-related. Reduce dose.

Contraindications: Cancer history, pregnancy, under 18.

1. 5mg: 1mL BAC->5mg/mL 10mg: 1mL->10mg/mL.
2. No fasting required. Inject any time -- morning or evening.
3. Standard dose: 250-500 mcg per injection, 1-2x daily.
4. Inject near the target injury site for localised tissue repair effect.
5. Stack with TB-500 (2mg, 2-3x/week) for synergistic musculoskeletal repair.
6. BPC-157 has no known LD50 -- one of the safest research peptides available.
7. Oral BPC-157 also effective for GI-specific protocols (skip reconstitution, dissolve in water).
8. Rotate injection sites. Store 2-8 deg C, use within 28 days.
9. Contraindicated in cancer history. Not for pregnancy or under 18.

DISCLAIMER:

research use only. Not for human therapeutic use. Consult a qualified professional.

assumes no liability.

|

.com | Research Use Only