

# GHRP-2

GH Secretagogue | 5mg & 10mg x 10 Vials



Strong GHS -- Fasted 2-3x Daily | Stack with GHRH for Max GH

Purity: 99.8% | Endotoxin-Free | Research Use Only

GHRP-2 (Growth Hormone Releasing Peptide-2) is a synthetic hexapeptide GH secretagogue and ghrelin receptor agonist. It produces strong GH pulses with a mild cortisol and prolactin elevation (unlike Ipamorelin). Highly effective in the GHRH + GHRP synergy stack. Must be injected fasted.


**FASTED REQUIRED:** Inject at least 1-2 hours after last meal. Insulin blunts GH release. Wait 30 min before eating post-injection.

## GHRP-2 Vial Comparison



| Parameter      | Specification                                                            |
|----------------|--------------------------------------------------------------------------|
| Class          | GH Secretagogue (GHS) -- ghrelin receptor agonist (GHSR-1a)              |
| Vial Sizes     | 5mg x 10   10mg x 10   Purity 99.8%                                      |
| Reconstitution | 1 mL BAC Water: 5mg->5mg/mL   10mg->10mg/mL                              |
| Standard Dose  | 100-300 mcg per injection -- always fasted                               |
| Frequency      | 2-3x daily fasted   morning + pre-bed + optional midday                  |
| vs Ipamorelin  | Stronger GH pulse; slight cortisol/prolactin spike; mild hunger increase |
| Cycle          | 12 weeks on, 4 weeks off                                                 |
| Storage        | 2-8 deg C   Reconstituted: use within 28 days                            |

MECHANISM OF ACTION & RESEARCH BENEFITS



| Pathway             | Mechanism                                                                                                                                                          | Benefit                                                |
|---------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------|
| GHSR-1a             | Binds ghrelin receptor on pituitary somatotrophs and hypothalamus; activates Gq/IP3/calcium cascade; strong GH granule exocytosis; larger GH pulse than Ipamorelin | Potent GH release; effective muscle building protocol  |
| Cortisol/ Prolactin | Unlike Ipamorelin, GHRP-2 has mild ACTH/cortisol and prolactin co-stimulation; usually minor at standard doses; relevant consideration for longer cycles           | Monitor for elevated cortisol/prolactin at high doses  |
| GHRH Synergy        | Stacked with CJC-1295 or Sermorelin: GHRH primes pituitary + GHRP-2 triggers powerful release; amplified GH 3-10x vs either alone                                  | Maximum GH when stacked; best body composition results |

| Benefit          | Context                                                                                                                         |
|------------------|---------------------------------------------------------------------------------------------------------------------------------|
| Strong GH Output | GHRP-2 produces larger GH pulses than Ipamorelin, making it preferred for aggressive muscle-building and performance protocols. |
| GHRH Synergy     | Combined with CJC-1295 or Sermorelin, GHRP-2 produces amplified GH pulses via dual-pathway activation.                          |
| Fasted Timing    | Fasted injection is mandatory. Avoid eating within 30 min post-injection for maximum GH effect.                                 |

## RECONSTITUTION & DOSE CALCULATION

All vials use 1 mL BAC Water. Room temperature. Swirl gently -- never shake.



1 mL BAC Water per vial. Roll gently -- NEVER shake.

| Step | Action           | Detail                                                                                       |
|------|------------------|----------------------------------------------------------------------------------------------|
| 1    | Gather           | Vial, BAC Water, insulin syringe (29-31G), swabs, sharps.                                    |
| 2    | Clean stoppers   | Wipe both stoppers. Air-dry 10-15 sec.                                                       |
| 3    | Draw 1mL BAC     | Exactly 1 mL. Expel air bubbles.                                                             |
| 4    | Inject into vial | Aim at GLASS WALL. Slowly. Never onto powder.                                                |
| 5    | Swirl & label    | Roll 20-30 sec. Clear within 60 sec. Never shake. Label date. 2-8 deg C. Use within 28 days. |



1mL Insulin Syringe | 29-31G | SubQ

**Formula:** Vol(mL)=Dose(mcg)/Conc(mcg/mL) | **Units:** Vol x 100


**5mg x 10 -- 1mL BAC -> 5.0mg/mL (5,000mcg/mL)**

| Dose(mcg)  | 100mcg | 150mcg | 200mcg | 250mcg | 300mcg |
|------------|--------|--------|--------|--------|--------|
| Vol(mL)    | 0.0200 | 0.0300 | 0.0400 | 0.0500 | 0.0600 |
| Units      | 2.0U   | 3.0U   | 4.0U   | 5.0U   | 6.0U   |
| Doses/Vial | 50     | 33     | 25     | 20     | 16     |

**10mg x 10 -- 1mL BAC -> 10.0mg/mL (10,000mcg/mL)**

| Dose(mcg)  | 100mcg | 150mcg | 200mcg | 250mcg | 300mcg |
|------------|--------|--------|--------|--------|--------|
| Vol(mL)    | 0.0100 | 0.0150 | 0.0200 | 0.0250 | 0.0300 |
| Units      | 1.0U   | 1.5U   | 2.0U   | 2.5U   | 3.0U   |
| Doses/Vial | 100    | 66     | 50     | 40     | 33     |

## INJECTION TECHNIQUE, PROTOCOLS & GOLDEN RULES



8-Point Rotation Map -- Rotate Every Injection



| Step | Action         | Detail                                                                |
|------|----------------|-----------------------------------------------------------------------|
| 1    | Confirm fasted | 1-2 hr after last meal. Mandatory.                                    |
| 2    | Draw dose      | Draw exact volume. If stacking with GHRH, can combine in one syringe. |
| 3    | Inject SubQ    | Abdomen preferred. Pinch 2-3cm. 45 deg. Rotate sites.                 |
| 4    | Wait 30 min    | Before eating. Record site. Sharps container.                         |

| Protocol    | Dose                    | Frequency         | Duration        | Notes                                       |
|-------------|-------------------------|-------------------|-----------------|---------------------------------------------|
| Standard    | 200 mcg                 | 2-3x daily fasted | 12wk on 4wk off | Both injections must be fasted.             |
| +GHRH Stack | 200mcg GHRP-2 +CJC-1295 | 2x daily fasted   | 12wk on 4wk off | Combine in syringe. 3-10x GH amplification. |

| Side Effect             | Likelihood | Management                                                        |
|-------------------------|------------|-------------------------------------------------------------------|
| Injection site reaction | Common     | Rotate sites. Fresh needle.                                       |
| Mild cortisol/prolactin | Occasional | Monitor at high doses/long cycles. Use Ipamorelin if sensitivity. |
| Hunger increase         | Common     | Ghrelin receptor activation. Manage with diet.                    |
| Water retention         | Occasional | GH-mediated. Reduce dose if significant.                          |

**Contraindications: Cancer history, pregnancy, under 18.**

|                                                                                            |
|--------------------------------------------------------------------------------------------|
| 1. Always inject fasted (1-2hr after last meal). Insulin blunts GH release.                |
| 2. 1 mL BAC Water per vial: 5mg->5mg/mL   10mg->10mg/mL.                                   |
| 3. Stack with CJC-1295 or Sermorelin for 3-10x amplified GH pulse.                         |
| 4. GHRP-2 causes mild cortisol/prolactin elevation -- switch to Ipamorelin if sensitivity. |
| 5. Inject 2-3x daily: morning fasted + pre-bed are optimal windows.                        |
| 6. Wait 30 min post-injection before eating. Rotate 8-point abdomen map.                   |
| 7. Cycle 12 weeks on, 4 weeks off. Store 2-8 deg C, 28 days max.                           |

---

DISCLAIMER: research use only. Not for human therapeutic use. Consult a qualified professional. assumes no liability.  
| .com | **Research Use Only**