

GLOW

BPC-157 10mg + TB-500 10mg + GHK-Cu 50mg | 70mg x 10 Vials

Advanced Tissue Repair, Collagen Synthesis & Regeneration Stack

1mL BAC Water -> 70mg/mL total | SubQ or IM | No Fasting Required

Purity: 99.8% | Endotoxin-Free | Research Use Only

GLOW is a pre-mixed triple-peptide stack combining BPC-157, TB-500, and GHK-Cu into a single 70mg vial. Each component addresses a distinct aspect of tissue repair and regeneration: BPC-157 for GI and vascular protection, TB-500 for cell migration and anti-inflammation, and GHK-Cu for collagen synthesis, gene modulation, and antioxidant support. Together they form a comprehensive tissue repair and rejuvenation protocol in one injection.

GLOW -- Active Components per Vial

BPC-157
10mg

#065f46

TB-500
10mg

#0f766e

GHK-Cu
50mg

#c9a84c

All components pre-mixed in one vial | One injection delivers full stack

Parameter	Specification
BPC-157	10mg per vial -- GI protection, angiogenesis (eNOS/NO), tendon repair, neuroprotection
TB-500	10mg per vial -- actin regulation, cell migration, anti-inflammatory (NF-kB), cardiac/neural repair
GHK-Cu	50mg per vial -- collagen I/III synthesis, wound healing, 4,000+ gene modulation, antioxidant
Total	70mg per vial x 10 vials per kit 700mg total
Reconstitution	1 mL BAC Water -> 70mg/mL total (1mL = full 70mg dose)
Standard Dose	0.5-1.0 mL per injection (35-70mg) SubQ or IM
Frequency	2-3x per week no fasting required inject near target tissue
Cycle & Storage	4-12 weeks on, 2-4 weeks off 2-8 deg C use within 28 days

GLOW | MECHANISM OF ACTION & SYNERGY



Complementary, non-overlapping pathways -- synergistic tissue repair

Component	Mechanism	Research Effect
BPC-157	GI mucosal healing; eNOS/nitric oxide pathway; VEGF angiogenesis; tendon fibroblast activation; no known LD50 -- exceptional safety	GI repair, vascular protection, tendon/ligament healing
TB-500	Thymosin Beta-4 fragment; sequesters G-actin enabling cell migration; NF-kB anti-inflammatory; VEGF angiogenesis; cardiomyocyte survival	Cell migration to injury, anti-inflammatory, cardiac/neural repair
GHK-Cu	Copper tripeptide; activates TGF-beta and collagen I/III synthesis; modulates 4,000+ human genes toward younger phenotype; Cu/Zn-SOD antioxidant; nerve regeneration	Collagen synthesis, wound healing, anti-aging, gene regulation

Benefit	Research Context
Triple-Pathway Synergy	BPC-157 (NO/angiogenesis) + TB-500 (cell migration) + GHK-Cu (collagen synthesis) act through completely non-overlapping pathways -- comprehensive repair.
Skin & Soft Tissue Repair	GHK-Cu drives collagen remodelling while BPC-157 and TB-500 accelerate vascular and cellular aspects of wound healing -- ideal for skin, fascia and soft tissue research.
Single Injection Simplicity	All three peptides pre-mixed in one vial. One injection delivers the complete GLOW stack -- no separate reconstitution or dosing required.


GLOW | RECONSTITUTION & DOSE GUIDE

70mg vial (70mg total) -- Add 1 mL BAC Water -> 70mg/mL total. One injection of 1 mL delivers the full dose of all components.



1 mL BAC Water per vial. Roll gently -- NEVER shake.


Step	Action	Detail
1	Gather supplies	Vial, BAC Water, 29-31G insulin syringe, alcohol swabs, sharps container.
2	Clean stoppers	Wipe both vial stoppers with alcohol swab. Air-dry 10-15 seconds.
3	Draw 1 mL BAC Water	Exactly 1 mL BAC Water. Expel all air bubbles.
4	Inject into vial	Aim stream at GLASS WALL -- never onto powder. Inject slowly.
5	Swirl gently & label	Roll between palms 20-30 sec. Clear within 60 sec. Never shake. Label with date. Store 2-8 deg C. Use within 28 days.




1mL Insulin Syringe | 29-31G | SubQ

At 70mg/mL: 1 mL = full 70mg dose | 0.5 mL = half dose | Each vial contains 10 doses at 0.1 mL (1/10 dose per injection)

Injection Volume	0.1 mL (1/10 dose)	0.25 mL (1/4 dose)	0.5 mL (1/2 dose)	1.0 mL (full dose)
Syringe Units	10U	25U	50U	100U
Total Peptide	7mg	18mg	35mg	70mg
Doses/Vial	10	4	2	1



8-Point Rotation Map -- Rotate Every Injection



Step	Action	Detail
1	Reconstitute	Add 1 mL BAC Water. Swirl gently 20-30 sec. Never shake. Clear solution required.
2	Draw dose	0.5 mL (half dose) or 1.0 mL (full dose). Expel air bubbles.
3	Select site	Inject near target tissue for localised repair, or abdomen for systemic benefit.
4	Inject SubQ/IM	SubQ: pinch 2-3cm, 45 deg, release, inject slowly. IM: 23-25G, 90 deg, aspirate.
5	Rotate & dispose	Record site. 8-point rotation. Press 5 sec. Sharps container.

Protocol	Dose	Frequency	Duration	Notes
Tissue Repair	1.0 mL (full dose)	2-3x/week	4-8 wk	Near injury site. Acute repair phase.
Skin/ Collagen	0.5-1.0 mL	2-3x/week	8-12 wk	Collagen synthesis & rejuvenation protocol.
Maintenance	0.5 mL (half dose)	2x/week	Ongoing	Lower maintenance after acute phase.

Side Effect	Likelihood	Management
Injection site reaction	Common	Mild redness. Rotate sites. Fresh needle each injection.
Temporary skin darkening	Occasional	GHK-Cu copper effect. Not harmful. Fades on cessation.

Mild fatigue	Occasional	Systemic repair response. Usually resolves within 1 week.
--------------	------------	---

Contraindications: Active cancer history, pregnancy, lactation, under 18.

1. GLOW = BPC-157 10mg + TB-500 10mg + GHK-Cu 50mg in one vial.
2. 1 mL BAC Water -> 70mg/mL. Full dose = 1 mL. Half dose = 0.5 mL.
3. No fasting required. Inject any time of day.
4. Inject near target tissue (injury, skin area) for localised benefit.
5. Each vial contains 10 full doses (1 mL each) or 20 half doses.
6. All three peptides work through complementary, non-overlapping pathways.
7. Rotate injection sites through 8-point abdomen map every session.
8. Swirl gently to reconstitute -- never shake. Store 2-8 deg C, 28 days max.
9. Contraindicated in active cancer history. Not for pregnancy or under 18.

DISCLAIMER: This document is produced by for informational and research reference purposes only. GLOW is a research product not approved for human therapeutic use. This guide does not constitute medical advice. assumes no liability for misuse.

| .com | Research Use Only